

Contents

1	Random Variables and Distributions	1
1.1	Probability Primer (quick but complete)	1
1.2	Random Variables, CDF, PDF/PMF, Support	1
1.3	Most-Used Distributions (pdf/cdf/mean/variance)	2
1.4	Joint Distributions, Marginals, Conditionals, Independence	4
1.5	Distributions of Functions of Random Variables	4
1.6	Worked Examples	5
1.7	Practice: Compute and Reason (with solutions)	6
2	Distributions and the Transport Theorem	8
2.1	Random Variables and Distributions	8
2.1.1	CDF and PDF	8
2.1.2	Expectation and Variance	8
2.2	The Transport Theorem (Change of Variables)	8
2.3	Multivariate Transport Theorem	9
2.4	Worked Examples	10
2.5	Exercises	10
3	Expectation and Variance	12
3.1	Definition of Expectation	12
3.2	Basic Properties	12
3.3	Conditional Expectation	13
3.4	Key Inequalities	14
3.5	Worked Examples	14
3.6	Exercises	15
4	Convergence and Limit Theorems	16
4.1	Modes of Convergence	16
4.2	The Law of Large Numbers (LLN)	16
4.3	Central Limit Theorem (CLT)	17
4.4	Delta Method	17
4.5	Slutsky's Theorem and Continuous Mapping	18
4.6	Worked Examples	18
4.7	Exercises	18
5	Summary and Formula Sheet	20
5.1	Probability Basics	20
5.2	Transformation and Transport Theorem	20
5.3	Expectation and Variance Properties	20
5.4	Inequalities	21
5.5	Modes of Convergence	21

5.6	Limit Theorems	21
5.7	Classical Distributions (Quick Reference)	22
5.8	MGF and CF Reference	22
5.9	Convergence Toolbox	22

Chapter 1

Random Variables and Distributions

1.1 Probability Primer (quick but complete)

Definition

A probability space is $(\Omega, \mathcal{F}, \mathbb{P})$ where \mathcal{F} is a σ -algebra on Ω and $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$ is countably additive with $\mathbb{P}(\Omega) = 1$.

Basic rules.

$$\mathbb{P}(A^c) = 1 - \mathbb{P}(A), \quad \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B), \quad \mathbb{P}\left(\bigcup_i A_i\right) \leq \sum_i \mathbb{P}(A_i).$$

Proposition

Total probability and Bayes. If $(B_i)_i$ is a partition with $\mathbb{P}(B_i) > 0$,

$$\mathbb{P}(A) = \sum_i \mathbb{P}(A \mid B_i) \mathbb{P}(B_i), \quad \mathbb{P}(B_k \mid A) = \frac{\mathbb{P}(A \mid B_k) \mathbb{P}(B_k)}{\sum_i \mathbb{P}(A \mid B_i) \mathbb{P}(B_i)}.$$

1.2 Random Variables, CDF, PDF/PMF, Support

Definition

A real random variable is a measurable $X : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$. Its distribution is the pushforward measure $\mathbb{P}_X(B) = \mathbb{P}(X \in B)$. The **CDF** is $F_X(x) = \mathbb{P}(X \leq x)$.

Properties of the CDF: nondecreasing, right-continuous; $F_X(-\infty) = 0$, $F_X(\infty) = 1$. If F_X is (a.e.) differentiable, the **PDF** is $f_X(x) = F'_X(x) \geq 0$ with $\int_{\mathbb{R}} f_X = 1$. For discrete X , the **PMF** is $p_X(x) = \mathbb{P}(X = x)$ with $\sum_x p_X(x) = 1$.

Proposition

Expectation (LOTUS) and variance. For measurable g ,

$$\mathbb{E}[g(X)] = \int_{\mathbb{R}} g(x) dF_X(x) = \begin{cases} \sum_x g(x) p_X(x), & \text{discrete,} \\ \int g(x) f_X(x) dx, & \text{continuous.} \end{cases}$$

Mean $\mu = \mathbb{E}[X]$, variance $\text{Var}(X) = \mathbb{E}[(X - \mu)^2] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$.

Proposition

Core algebra. For $a, b \in \mathbb{R}$:

$$\mathbb{E}[aX + b] = a \mathbb{E}[X] + b, \quad \text{Var}(aX + b) = a^2 \text{Var}(X).$$

If $X \perp\!\!\!\perp Y$, then $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ and $\text{Var}(X + Y) = \text{Var} X + \text{Var} Y$.

1.3 Most-Used Distributions (pdf/cdf/mean/variance)

Discrete

Bernoulli (p)

$$\mathbb{P}(X = 1) = p, \mathbb{P}(X = 0) = 1 - p; \quad \mathbb{E}[X] = p, \text{Var}(X) = p(1 - p).$$

Binomial (n, p)

$$\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, k = 0, \dots, n; \quad \mathbb{E}[X] = np, \text{Var}(X) = np(1 - p).$$

Geometric (p) (trials until first success)

$$\mathbb{P}(X = k) = (1 - p)^{k-1} p, k = 1, 2, \dots; \quad \mathbb{E}[X] = 1/p, \text{Var}(X) = (1 - p)/p^2; \text{ memoryless.}$$

Poisson (λ)

$$\mathbb{P}(X = k) = e^{-\lambda} \lambda^k / k!, k = 0, 1, 2, \dots; \quad \mathbb{E}[X] = \text{Var}(X) = \lambda.$$

Continuous**Uniform (a, b)**

$$f(x) = \frac{1}{b-a} \text{ on } (a, b); \quad F(x) = \frac{x-a}{b-a} \text{ on } (a, b); \quad \mathbb{E}[X] = \frac{a+b}{2}, \quad \text{Var}(X) = \frac{(b-a)^2}{12}.$$

Exponential (λ)

$$f(x) = \lambda e^{-\lambda x} \text{ for } x > 0; \quad F(x) = 1 - e^{-\lambda x}; \quad \mathbb{E}[X] = 1/\lambda, \quad \text{Var}(X) = 1/\lambda^2; \text{ memoryless.}$$

Gamma (k, θ) (shape–scale)

$$f(x) = \frac{x^{k-1} e^{-x/\theta}}{\Gamma(k) \theta^k} \text{ for } x > 0; \quad \mathbb{E}[X] = k\theta, \quad \text{Var}(X) = k\theta^2.$$

Normal (μ, σ^2)

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right); \quad \mathbb{E}[X] = \mu, \quad \text{Var}(X) = \sigma^2; \quad F(x) = \Phi\left(\frac{x-\mu}{\sigma}\right).$$

Beta (α, β) on $(0, 1)$

$$f(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}; \quad \mathbb{E}[X] = \frac{\alpha}{\alpha+\beta}, \quad \text{Var}(X) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}.$$

Chi-square (k)

$$\text{If } Z_i \sim \mathcal{N}(0, 1) \text{ i.i.d., then } \sum_{i=1}^k Z_i^2 \sim \chi^2(k); \quad \mathbb{E}[X] = k, \quad \text{Var}(X) = 2k.$$

Useful relationships.

- Binomial(n, p) with $n \rightarrow \infty, p \rightarrow 0, np \rightarrow \lambda \Rightarrow \text{Poisson}(\lambda)$ (Poisson limit).
- Sums: independent Poissons add; independent Gammas with same scale add; Normal sums stay Normal.
- Affine change: if X has density f_X , then $Y = aX + b$ has $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$ and $\mathbb{E}[Y] = a\mathbb{E}[X] + b, \text{Var}(Y) = a^2 \text{Var}(X)$.

1.4 Joint Distributions, Marginals, Conditionals, Independence

Definition

For (X, Y) continuous, the joint density $f_{X,Y}$ satisfies $f_{X,Y} \geq 0$ and $\iint f_{X,Y} = 1$.

Marginals:

$$f_X(x) = \int f_{X,Y}(x, y) dy, \quad f_Y(y) = \int f_{X,Y}(x, y) dx.$$

For discrete variables, replace integrals by sums.

Definition

Conditional distribution. If $f_X(x) > 0$,

$$f_{Y|X}(y | x) = \frac{f_{X,Y}(x, y)}{f_X(x)}, \quad \mathbb{E}[Y | X = x] = \int y f_{Y|X}(y | x) dy.$$

Definition

Independence. $X \perp\!\!\!\perp Y$ iff $f_{X,Y}(x, y) = f_X(x)f_Y(y)$ (continuous) or $p_{X,Y}(x, y) = p_X(x)p_Y(y)$ (discrete).

Proposition

Total expectation and total variance.

$$\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X | Y]], \quad \text{Var}(X) = \mathbb{E}[\text{Var}(X | Y)] + \text{Var}(\mathbb{E}[X | Y]).$$

Probabilities involving several variables. For independent continuous X, Y ,

$$\mathbb{P}(X < Y) = \int_{-\infty}^{\infty} f_X(x) F_Y(x) dx = \int f_Y(y) [1 - F_X(y)] dy.$$

More generally, $\mathbb{P}((X, Y) \in A) = \iint_A f_{X,Y}(x, y) dx dy$.

1.5 Distributions of Functions of Random Variables

One variable (transport theorem)

Theorem

If $Y = \phi(X)$ with ϕ continuously differentiable and strictly monotone, then

$$f_Y(y) = f_X(\phi^{-1}(y)) \left| \frac{d}{dy} \phi^{-1}(y) \right|.$$

Many-to-one case. If ϕ is m -to-1 on relevant parts of the support,

$$f_Y(y) = \sum_{i=1}^m f_X(x_i(y)) \left| \frac{d}{dy} x_i(y) \right| \quad \text{where } \phi(x_i(y)) = y.$$

Example

$X \sim \text{Unif}(-1, 1)$, $Y = X^2$. For $y \in (0, 1)$, the inverse branches are $x_{\pm}(y) = \pm\sqrt{y}$, so

$$f_Y(y) = f_X(\sqrt{y}) \frac{1}{2\sqrt{y}} + f_X(-\sqrt{y}) \frac{1}{2\sqrt{y}} = \frac{1}{2} \cdot \frac{1}{2\sqrt{y}} + \frac{1}{2} \cdot \frac{1}{2\sqrt{y}} = \frac{1}{2\sqrt{y}}.$$

Two variables (Jacobian)

Theorem

If $(U, V) = T(X, Y)$ is a C^1 bijection with inverse $(x(u, v), y(u, v))$ and Jacobian $J_{T^{-1}}$, then

$$f_{U,V}(u, v) = f_{X,Y}(x(u, v), y(u, v)) |\det J_{T^{-1}}(u, v)|.$$

Example

Polar coordinates. $X = R \cos \Theta$, $Y = R \sin \Theta$ has $|\det J| = r$. If $X, Y \sim \mathcal{N}(0, 1)$ i.i.d., then

$$f_{R,\Theta}(r, \theta) = \frac{1}{2\pi} e^{-r^2/2} r,$$

so R and Θ are independent, R is Rayleigh ($re^{-r^2/2}$ for $r > 0$), and $\Theta \sim \text{Unif}(0, 2\pi)$.

Common constructions.

- *Sum (convolution):* $Z = X + Y$ (indep. continuous) has $f_Z(z) = \int f_X(x) f_Y(z - x) dx$. In transform language, MGFs/CFs multiply.
- *Max/min (i.i.d. continuous):* $F_{\max}(x) = [F(x)]^n$, $F_{\min}(x) = 1 - [1 - F(x)]^n$.
- *Ratio/scaling:* If $X > 0$ and $Y = cX$, then $f_Y(y) = \frac{1}{|c|} f_X(y/c)$. Ratios like $X/(X + Y)$ often lead to Beta laws (see Example 3 below).

1.6 Worked Examples

Worked Solution

Example 1 (sum of uniforms). $X, Y \sim \text{Unif}(0, 1)$ independent; $Z = X + Y$. Then

$$f_Z(z) = \int_0^1 \mathbf{1}_{0 < x < 1} \mathbf{1}_{0 < z - x < 1} dx = \begin{cases} z, & 0 < z < 1, \\ 2 - z, & 1 \leq z < 2, \\ 0, & \text{else.} \end{cases}$$

Worked Solution

Example 2 (who is larger?). $X \sim \text{Exp}(\lambda)$, $Y \sim \text{Exp}(\mu)$ independent. Then

$$\mathbb{P}(X < Y) = \int_0^\infty f_X(x) \mathbb{P}(Y > x) dx = \int_0^\infty \lambda e^{-\lambda x} e^{-\mu x} dx = \frac{\lambda}{\lambda + \mu}.$$

Worked Solution

Example 3 (Gamma–Beta trick). $X \sim \Gamma(\alpha, \theta)$, $Y \sim \Gamma(\beta, \theta)$ independent. Let $U = X + Y$, $V = \frac{X}{X+Y}$. The Jacobian of $(x, y) \mapsto (u, v)$ is $|J| = u$. Then

$$f_{U,V}(u, v) = \frac{u^{\alpha+\beta-1} e^{-u/\theta}}{\Gamma(\alpha)\Gamma(\beta)\theta^{\alpha+\beta}} v^{\alpha-1} (1-v)^{\beta-1},$$

which factorizes: $U \sim \Gamma(\alpha + \beta, \theta)$ and $V \sim \text{Beta}(\alpha, \beta)$, *independent*.

Worked Solution

Example 4 (absolute value). $X \sim \mathcal{N}(0, 1)$, $Y = |X|$. Many-to-one transform with branches $\pm y$ yields

$$f_Y(y) = f_X(y) + f_X(-y) = \frac{2}{\sqrt{2\pi}} e^{-y^2/2}, \quad y > 0.$$

1.7 Practice: Compute and Reason (with solutions)

Exercises

- E1.** (Basics) Show that F_X is right-continuous and $\lim_{x \rightarrow -\infty} F_X(x) = 0$, $\lim_{x \rightarrow \infty} F_X(x) = 1$.
- E2.** (Expectation) If $X \sim \text{Unif}(a, b)$, compute $\mathbb{E}[X^k]$ for $k \in \mathbb{N}$.
- E3.** (Joint \rightarrow marginal/conditional) $f_{X,Y}(x, y) = c(x + y)$ on $0 < x < 1$, $0 < y < 1$. Find c , f_X , f_Y , $\mathbb{E}[X]$, $\mathbb{E}[Y]$, $\text{Cov}(X, Y)$, and $f_{Y|X}$.
- E4.** (Convolution) $X, Y \sim \text{Exp}(\lambda)$ independent. Find the pdf of $Z = X + Y$ and its mean/variance.
- E5.** (Transport) $X \sim \text{Unif}(-1, 1)$ and $Y = X^2$. Find F_Y and f_Y .
- E6.** (Who is larger?) $X \sim \text{Exp}(2)$, $Y \sim \text{Exp}(3)$ independent. Compute $\mathbb{P}(X < Y)$.
- E7.** (Order stats) $U_i \sim \text{Unif}(0, 1)$ i.i.d. Find F_{\max} and $\mathbb{E}[\max_i U_i]$.

Solution sketches

Worked Solution

(1) From measure properties of distribution functions.

$$(2) \mathbb{E}[X^k] = \frac{1}{b-a} \int_a^b x^k dx = \frac{b^{k+1} - a^{k+1}}{(k+1)(b-a)}.$$

(3) $1 = \int_0^1 \int_0^1 c(x+y) dy dx \Rightarrow c = 1$. $f_X(x) = x + \frac{1}{2}$, $f_Y(y) = y + \frac{1}{2}$. $\mathbb{E}[X] = \mathbb{E}[Y] = 7/12$, $\mathbb{E}[XY] = 1/4$, $\text{Cov} = -13/144$. $f_{Y|X}(y|x) = \frac{x+y}{x+1/2}$.

(4) $Z \sim \Gamma(2, 1/\lambda)$ with $f_Z(z) = \lambda^2 z e^{-\lambda z}$ ($z > 0$), $\mathbb{E}[Z] = 2/\lambda$, $\text{Var}(Z) = 2/\lambda^2$.

(5) $F_Y(y) = \mathbb{P}(|X| \leq \sqrt{y}) = \frac{\sqrt{y} - (-\sqrt{y})}{2} = \sqrt{y}$ on $(0, 1)$; $f_Y(y) = \frac{1}{2\sqrt{y}}$.

$$(6) \mathbb{P}(X < Y) = \frac{2}{2+3} = \frac{2}{5}.$$

(7) $F_{\max}(x) = x^n$ on $(0, 1)$, so $f_{\max}(x) = nx^{n-1}$ and $\mathbb{E}[\max] = n/(n+1)$.

What to master from this chapter: how to read and use CDF/PDF/PMF; the shape and parameters of core distributions; marginalization and conditioning; independence; computing probabilities involving several variables; and transforming variables (1D transport and 2D Jacobians).

Chapter 2

Distributions and the Transport Theorem

2.1 Random Variables and Distributions

Definition

A **random variable** is a measurable function $X : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$. Its **distribution** is the measure $\mathbb{P}_X(B) = \mathbb{P}(X \in B)$ for all $B \in \mathcal{B}$.

2.1.1 CDF and PDF

The cumulative distribution function (CDF) is

$$F_X(x) = \mathbb{P}(X \leq x),$$

with properties F_X nondecreasing, right-continuous, $F_X(-\infty) = 0$, $F_X(\infty) = 1$. If F_X is differentiable, its derivative $f_X(x) = F'_X(x)$ is the **probability density function (PDF)**.

For a discrete variable, $p_X(x) = \mathbb{P}(X = x)$ and $F_X(x) = \sum_{t \leq x} p_X(t)$.

2.1.2 Expectation and Variance

For an integrable function g , **LOTUS** (Law of the Unconscious Statistician) states:

$$\mathbb{E}[g(X)] = \int g(x) f_X(x) dx,$$

for continuous X , or $\sum_x g(x) p_X(x)$ for discrete X .

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}X)^2].$$

2.2 The Transport Theorem (Change of Variables)

Theorem

Let X be a random variable with PDF f_X and let $Y = \phi(X)$ be a continuously differentiable, monotone function. Then the PDF of Y is

$$f_Y(y) = f_X(\phi^{-1}(y)) \left| \frac{d}{dy} \phi^{-1}(y) \right|.$$

Sketch. If ϕ is strictly increasing, $\mathbb{P}(Y \leq y) = \mathbb{P}(X \leq \phi^{-1}(y)) = F_X(\phi^{-1}(y))$. Differentiating both sides gives the formula. \square

Example

Linear Transformations: If $Y = aX + b$ with $a \neq 0$, then

$$F_Y(y) = F_X\left(\frac{y-b}{a}\right), \quad f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right).$$

Example

Polynomial Transformation: Let $X \sim \text{Unif}(0, 1)$ and $Y = X^2$. Then $F_Y(y) = P(Y \leq y) = P(X \leq \sqrt{y}) = \sqrt{y}$ for $0 < y < 1$. Hence $f_Y(y) = \frac{1}{2\sqrt{y}}$.

Example

Trigonometric Transformation: If $\Theta \sim \text{Unif}(0, 2\pi)$ and $X = \cos \Theta$, then the PDF of X is

$$f_X(x) = \frac{1}{\pi\sqrt{1-x^2}}, \quad x \in (-1, 1).$$

2.3 Multivariate Transport Theorem

Theorem

If (X, Y) is a continuous random vector with joint PDF $f_{X,Y}$ and $(U, V) = T(X, Y)$ is a one-to-one, differentiable transformation with inverse T^{-1} , then

$$f_{U,V}(u, v) = f_{X,Y}(x(u, v), y(u, v)) \left| \det J_{T^{-1}}(u, v) \right|,$$

where $J_{T^{-1}}$ is the Jacobian matrix of T^{-1} .

Example

Polar Transformation: If (X, Y) has joint PDF $f_{X,Y}(x, y)$, define $R = \sqrt{X^2 + Y^2}$, $\Theta = \arctan(Y/X)$. Then

$$|\det J| = r, \quad f_{R,\Theta}(r, \theta) = f_{X,Y}(r \cos \theta, r \sin \theta) r.$$

If $(X, Y) \sim \mathcal{N}(0, 1)$ i.i.d., then $f_{R,\Theta}(r, \theta) = \frac{1}{2\pi} e^{-r^2/2} r$, so R and Θ are independent.

2.4 Worked Examples**Worked Solution**

Example 1 — Absolute Value: Let $X \sim \mathcal{N}(0, 1)$, $Y = |X|$. Find f_Y .

$$f_Y(y) = f_X(y) + f_X(-y) = \frac{2}{\sqrt{2\pi}} e^{-y^2/2}, \quad y > 0.$$

Worked Solution

Example 2 — Exponential Scaling: Let $X \sim \text{Exp}(1)$ and $Y = 2X$. Then $f_Y(y) = \frac{1}{2} e^{-y/2}$ for $y > 0$.

Worked Solution

Example 3 — Sum of Two Uniforms: If $X, Y \sim \text{Unif}(0, 1)$ independent, then $Z = X + Y$ has PDF

$$f_Z(z) = \begin{cases} z, & 0 < z < 1, \\ 2 - z, & 1 \leq z < 2, \\ 0, & \text{otherwise.} \end{cases}$$

2.5 Exercises

E1. $X \sim \text{Unif}(-1, 1)$. Find the PDF of $Y = X^2$.

E2. If $X \sim \text{Exp}(\lambda)$, find the distribution of $Y = aX + b$.

E3. Let $X \sim \text{Unif}(0, \pi)$, $Y = \sin X$. Compute f_Y .

E4. If (X, Y) has joint PDF $f(x, y) = e^{-(x+y)}$ for $x, y > 0$, find the PDF of $(U, V) = (X + Y, X/(X + Y))$.

E5. (Challenge) Derive the CDF and PDF of the maximum of n i.i.d. $\text{Unif}(0, 1)$.

Worked Solution

Solutions Sketch: (1) $Y \in (0, 1)$, $f_Y(y) = \frac{1}{2\sqrt{y}}$. (2) Linear transform formula. (3) $f_Y(y) = \frac{1}{\pi\sqrt{1-y^2}}$ for $y \in (0, 1)$. (4) Use Jacobian $|J| = u$. (5) $f_{X_{(n)}}(x) = nx^{n-1}$.

Chapter 3

Expectation and Variance

3.1 Definition of Expectation

Definition

For a real random variable X with CDF F_X , the **expectation** of a measurable function $g(X)$ is

$$\mathbb{E}[g(X)] = \begin{cases} \sum_x g(x) p_X(x), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx, & \text{if } X \text{ is continuous.} \end{cases}$$

Proposition

(LOTUS — Law of the Unconscious Statistician) For any measurable g ,

$$\mathbb{E}[g(X)] = \int_{\mathbb{R}} g(x) dF_X(x).$$

Example

If $X \sim \text{Unif}(0, 1)$, then $\mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$.

3.2 Basic Properties

Proposition

Let $a, b \in \mathbb{R}$. Then:

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b, \quad \text{Var}(aX + b) = a^2 \text{Var}(X).$$

If X, Y are independent: $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$, hence $\text{Var}(X + Y) = \text{Var } X + \text{Var } Y$.

Definition

The **variance** of X is $\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2]$. It measures dispersion around the mean.

Proposition

Useful identities:

$$\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2, \quad \text{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y].$$

The **correlation** is $\rho(X, Y) = \text{Cov}(X, Y)/(\sigma_X \sigma_Y) \in [-1, 1]$.

Example

If $X \sim \text{Exp}(\lambda)$, then $\mathbb{E}[X] = 1/\lambda$ and $\text{Var}(X) = 1/\lambda^2$.

3.3 Conditional Expectation

Definition

The **conditional expectation** of Y given X is the random variable $\mathbb{E}[Y|X]$ satisfying

$$\mathbb{E}[g(X)\mathbb{E}[Y|X]] = \mathbb{E}[g(X)Y] \quad \text{for all measurable } g.$$

For discrete (X, Y) , $\mathbb{E}[Y|X = x] = \sum_y y \mathbb{P}(Y = y|X = x)$. For continuous (X, Y) , $\mathbb{E}[Y|X = x] = \int y f_{Y|X}(y|x) dy$.

Proposition

Tower Property: $\mathbb{E}[\mathbb{E}[Y|X]] = \mathbb{E}[Y]$. **Law of Total Variance:**

$$\text{Var}(Y) = \mathbb{E}[\text{Var}(Y|X)] + \text{Var}(\mathbb{E}[Y|X]).$$

Example

If $X \sim \text{Unif}(0, 1)$ and $Y|X = x \sim \text{Exp}(x)$, then

$$\mathbb{E}[Y|X = x] = 1/x, \quad \mathbb{E}[Y] = \int_0^1 \frac{1}{x} dx = \infty$$

(diverges — expectation does not always exist).

3.4 Key Inequalities

Proposition

Markov's Inequality. If $X \geq 0$ and $a > 0$, then

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}[X]}{a}.$$

Proposition

Chebyshev's Inequality. For any X with finite variance and $t > 0$,

$$\mathbb{P}(|X - \mathbb{E}[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2}.$$

Proposition

Jensen's Inequality. For convex ϕ ,

$$\phi(\mathbb{E}[X]) \leq \mathbb{E}[\phi(X)].$$

Equality holds iff X is a.s. constant or ϕ is linear.

3.5 Worked Examples

Worked Solution

Example 1 — Expectation of a nonlinear transform. Let $X \sim \text{Unif}(0, 1)$, find $\mathbb{E}[\sqrt{X}]$ and $\text{Var}(\sqrt{X})$.

$$\mathbb{E}[\sqrt{X}] = \int_0^1 x^{1/2} dx = \frac{2}{3}, \quad \mathbb{E}[X] = \frac{1}{2}, \quad \mathbb{E}[X^{1/2}] = \frac{2}{3}.$$

Then $\mathbb{E}[(\sqrt{X})^2] = \mathbb{E}[X] = 1/2$ and $\text{Var}(\sqrt{X}) = 1/2 - (2/3)^2 = 1/18$.

Worked Solution

Example 2 — Covariance and independence. Let X, Y be independent $\text{Unif}(0, 1)$. Then $\text{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = 1/4 - 1/4 = 0$. Independence \Rightarrow zero covariance.

Worked Solution

Example 3 — Conditional expectation as regression. Suppose (X, Y) have joint density $f_{X,Y}(x, y) = 2$ on $0 < y < x < 1$. Then $f_{Y|X}(y|x) = 1/x$ for $0 < y < x$, and $\mathbb{E}[Y|X = x] = x/2$. The regression line is $\mathbb{E}[Y|X] = X/2$.

Worked Solution

Example 4 — Chebyshev bound. If X has mean μ and variance σ^2 , then $\mathbb{P}(|X - \mu| \geq 2\sigma) \leq 1/4$ (at most 25

3.6 Exercises

- E1.** Prove $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$ directly from the integral definition.
- E2.** For $X \sim \text{Exp}(\lambda)$, compute $\mathbb{E}[X^2]$ and deduce $\text{Var}(X)$.
- E3.** Let $X \sim \text{Unif}(0, 1)$, find $\mathbb{E}[\ln X]$ and $\text{Var}(\ln X)$.
- E4.** Verify the law of total variance for the model $Y|X \sim \text{Bernoulli}(X)$, $X \sim \text{Unif}(0, 1)$.
- E5.** Show that Jensen's inequality implies $\mathbb{E}[e^X] \geq e^{\mathbb{E}[X]}$ for all real X .

Worked Solution

Solution sketches: (1) Expand integrals linearly. (2) $\mathbb{E}[X^2] = 2/\lambda^2$, $\text{Var} = 1/\lambda^2$. (3) $\mathbb{E}[\ln X] = \int_0^1 \ln x \, dx = -1$, $\mathbb{E}[(\ln X)^2] = 2$, $\text{Var} = 1$. (4) $\text{Var}(Y) = \mathbb{E}[X(1 - X)] + \text{Var}(X) = \frac{1}{6}$. (5) Follows from convexity of e^x .

Chapter 4

Convergence and Limit Theorems

4.1 Modes of Convergence

Definition

Let (X_n) be a sequence of random variables and X another random variable.

- **Almost sure (a.s.) convergence:** $X_n \rightarrow X$ a.s. if $\mathbb{P}(\lim_{n \rightarrow \infty} X_n = X) = 1$.
- **Convergence in probability:** $X_n \xrightarrow{p} X$ if for all $\varepsilon > 0$, $\mathbb{P}(|X_n - X| > \varepsilon) \rightarrow 0$.
- **Convergence in distribution:** $X_n \xrightarrow{d} X$ if $F_{X_n}(x) \rightarrow F_X(x)$ for all continuity points x of F_X .

Proposition

Implications.

$$X_n \xrightarrow{a.s.} X \Rightarrow X_n \xrightarrow{p} X \Rightarrow X_n \xrightarrow{d} X.$$

The converses are false in general.

Example

Let X_n be the average of n i.i.d. Bernoulli(1/2) variables. Then $X_n \xrightarrow{p} 1/2$ by the Law of Large Numbers (next section).

4.2 The Law of Large Numbers (LLN)

Theorem

Weak Law of Large Numbers (WLLN). Let X_1, \dots, X_n be i.i.d. with mean μ and finite variance σ^2 . Then

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{p} \mu.$$

Sketch. Since $\mathbb{E}[\bar{X}_n] = \mu$ and $\text{Var}(\bar{X}_n) = \sigma^2/n$, Chebyshev's inequality gives $\mathbb{P}(|\bar{X}_n - \mu| > \varepsilon) \leq \sigma^2/(n\varepsilon^2) \rightarrow 0$. \square

Theorem

Strong Law of Large Numbers (SLLN). If (X_i) are i.i.d. with $\mathbb{E}[|X_i|] < \infty$, then

$$\bar{X}_n \xrightarrow{a.s.} \mathbb{E}[X_1].$$

Interpretation: both LLNs formalize that empirical averages stabilize to their expected value.

4.3 Central Limit Theorem (CLT)

Theorem

Classical CLT. If X_1, \dots, X_n are i.i.d. with mean μ and variance $\sigma^2 > 0$, then

$$Z_n = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow{d} \mathcal{N}(0, 1).$$

Idea. Use moment generating functions. $M_{Z_n}(t) = (M_{(X_i - \mu)/\sigma}(t/\sqrt{n}))^n \rightarrow e^{t^2/2}$, the MGF of $\mathcal{N}(0, 1)$. \square

Example

If $X_i \sim \text{Bernoulli}(p)$, \bar{X}_n has mean p and variance $p(1-p)/n$. Then

$$\mathbb{P}(\bar{X}_n \leq x) \approx \Phi\left(\frac{x - p}{\sqrt{p(1-p)/n}}\right)$$

for large n , where Φ is the standard normal CDF.

4.4 Delta Method

Theorem

If $\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$ and g is differentiable at θ_0 , then

$$\sqrt{n}(g(\hat{\theta}_n) - g(\theta_0)) \xrightarrow{d} \mathcal{N}(0, [g'(\theta_0)]^2 \sigma^2).$$

Example

If \bar{X}_n estimates μ with variance σ^2/n , then $g(\bar{X}_n) = \bar{X}_n^2$ satisfies

$$\sqrt{n}(\bar{X}_n^2 - \mu^2) \xrightarrow{d} \mathcal{N}(0, 4\mu^2 \sigma^2).$$

4.5 Slutsky's Theorem and Continuous Mapping

Proposition

Slutsky's Theorem. If $X_n \xrightarrow{d} X$ and $Y_n \xrightarrow{p} c$ (constant), then:

$$X_n + Y_n \xrightarrow{d} X + c, \quad X_n Y_n \xrightarrow{d} cX, \quad X_n/Y_n \xrightarrow{d} X/c \quad (c \neq 0).$$

Proposition

Continuous Mapping Theorem. If $X_n \xrightarrow{d} X$ and g is continuous, then $g(X_n) \xrightarrow{d} g(X)$.

4.6 Worked Examples

Worked Solution

Example 1 — LLN for Exponentials. If $X_i \sim \text{Exp}(1)$, then $\mathbb{E}[X_i] = 1$ and $\text{Var}(X_i) = 1$. By WLLN, $\bar{X}_n \xrightarrow{p} 1$. By CLT, $\sqrt{n}(\bar{X}_n - 1) \xrightarrow{d} \mathcal{N}(0, 1)$.

Worked Solution

Example 2 — Normal Approximation of Binomial. If $S_n \sim \text{Binomial}(n, p)$, then

$$\frac{S_n - np}{\sqrt{np(1-p)}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Worked Solution

Example 3 — Delta Method. If $X_i \sim \text{Exp}(\lambda)$ and $\hat{\lambda} = 1/\bar{X}_n$, then $\sqrt{n}(\hat{\lambda} - \lambda) \xrightarrow{d} \mathcal{N}(0, \lambda^2)$. By the Delta Method, for $g(x) = 1/x$,

$$\sqrt{n}(g(\bar{X}_n) - g(1/\lambda)) \xrightarrow{d} \mathcal{N}(0, \lambda^2).$$

4.7 Exercises

E1. Let X_1, \dots, X_n i.i.d. with mean μ , variance σ^2 . Prove $\mathbb{E}[\bar{X}_n] = \mu$, $\text{Var}(\bar{X}_n) = \sigma^2/n$.

E2. Verify the WLLN using Chebyshev's inequality for $X_i \sim \text{Exp}(1)$.

E3. Use the CLT to approximate $\mathbb{P}(S_{100} \geq 60)$ when $S_{100} \sim \text{Binomial}(100, 0.5)$.

E4. Apply the Delta Method to $g(x) = \ln x$ for \bar{X}_n estimating $\mu > 0$.

E5. (Challenge) State a version of the multivariate CLT for $\mathbf{X}_i \in \mathbb{R}^d$.

Worked Solution

Solutions Sketch: (1) Direct from linearity. (2) $\mathbb{P}(|\bar{X}_n - 1| > \varepsilon) \leq 1/(n\varepsilon^2)$. (3) Mean = 50, SD = $\sqrt{25} = 5$, so $\mathbb{P}(S_{100} \geq 60) \approx 1 - \Phi(2) = 0.0228$. (4) $g'(\mu) = 1/\mu$, hence asymptotic variance (σ^2/μ^2n) . (5) Multivariate: $\sqrt{n}(\bar{\mathbf{X}}_n - \boldsymbol{\mu}) \xrightarrow{d} \mathcal{N}(0, \Sigma)$.

Worked Solution

Takeaways:

- LLN — averages converge to expectation (consistency).
- CLT — fluctuations around the mean are Gaussian.
- Delta Method — propagate asymptotic normality through smooth transformations.

Chapter 5

Summary and Formula Sheet

5.1 Probability Basics

Worked Solution

CDF:	$F_X(x) = \mathbb{P}(X \leq x)$, nondecreasing, right-continuous.
PDF:	$f_X(x) = F'_X(x)$, $f_X \geq 0$, $\int f_X = 1$.
Discrete:	$p_X(x) = \mathbb{P}(X = x)$, $\sum_x p_X(x) = 1$.
Expectation:	$\mathbb{E}[g(X)] = \int g(x)f_X(x)dx$ or $\sum g(x)p_X(x)$.
Variance:	$\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$.
Covariance:	$\text{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$.
Correlation:	$\rho = \text{Cov}(X, Y)/(\sigma_X\sigma_Y) \in [-1, 1]$.

5.2 Transformation and Transport Theorem

Worked Solution

1D:	$f_Y(y) = f_X(\phi^{-1}(y)) \left \frac{d}{dy} \phi^{-1}(y) \right $.
Many-to-one:	$f_Y(y) = \sum_i f_X(x_i) (\phi^{-1})'_i(y) $.
2D (Jacobian):	$f_{U,V}(u, v) = f_{X,Y}(x(u, v), y(u, v)) \det J_{T^{-1}} $.
Example:	Polar coords: $X = R \cos \Theta$, $Y = R \sin \Theta$, $ J = r$.

5.3 Expectation and Variance Properties

Worked Solution

Linearity	$\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
Scaling	$\text{Var}(aX + b) = a^2 \text{Var}(X)$
Sum of indep.	$\text{Var}(X + Y) = \text{Var } X + \text{Var } Y$
Total Variance	$\text{Var}(X) = \mathbb{E}[\text{Var}(X Y)] + \text{Var}(\mathbb{E}[X Y])$
Tower Property	$\mathbb{E}[X] = \mathbb{E}[\mathbb{E}[X Y]]$

5.4 Inequalities

Worked Solution

Markov:	$\mathbb{P}(X \geq a) \leq \mathbb{E}[X]/a$ for $X \geq 0$.
Chebyshev:	$\mathbb{P}(X - \mu \geq t) \leq \text{Var}(X)/t^2$.
Jensen:	$\phi(\mathbb{E}[X]) \leq \mathbb{E}[\phi(X)]$ for convex ϕ .
Cauchy–Schwarz:	$ \mathbb{E}[XY] \leq \sqrt{\mathbb{E}[X^2]\mathbb{E}[Y^2]}$.

5.5 Modes of Convergence

Worked Solution

Almost sure	$\mathbb{P}(\lim X_n = X) = 1$
In probability	$\mathbb{P}(X_n - X > \varepsilon) \rightarrow 0$
In distribution	$F_{X_n}(x) \rightarrow F_X(x)$ (at continuity points)
Relation	$\text{a.s.} \Rightarrow \text{in } p \Rightarrow \text{in } d$

5.6 Limit Theorems

Worked Solution

WLLN:	$\bar{X}_n \xrightarrow{p} \mu.$
SLLN:	$\bar{X}_n \xrightarrow{\text{a.s.}} \mu.$
CLT:	$\sqrt{n}(\bar{X}_n - \mu)/\sigma \xrightarrow{d} \mathcal{N}(0, 1).$
Delta Method:	$\sqrt{n}(g(\hat{\theta}_n) - g(\theta_0)) \rightarrow \mathcal{N}(0, [g'(\theta_0)]^2 \sigma^2).$
Slutsky:	If $X_n \rightarrow_d X$, $Y_n \rightarrow_p c$, then $X_n + Y_n \rightarrow_d X + c$.
Continuous Mapping:	$g(X_n) \rightarrow_d g(X)$ if g continuous.

5.7 Classical Distributions (Quick Reference)

Worked Solution

Distribution	PDF / PMF	Mean, Variance
Uniform(a, b)	$1/(b - a)$	$(a + b)/2, (b - a)^2/12$
Bernoulli(p)	$p^x(1 - p)^{1-x}$	$p, p(1 - p)$
Binomial(n, p)	$\binom{n}{k}p^k(1 - p)^{n-k}$	$np, np(1 - p)$
Poisson(λ)	$e^{-\lambda}\lambda^k/k!$	λ, λ
Exponential(λ)	$\lambda e^{-\lambda x} \ (x > 0)$	$1/\lambda, 1/\lambda^2$
Normal(μ, σ^2)	$\frac{1}{\sqrt{2\pi\sigma^2}}e^{-(x-\mu)^2/2\sigma^2}$	μ, σ^2
Gamma(k, θ)	$x^{k-1}e^{-x/\theta}/(\Gamma(k)\theta^k)$	$k\theta, k\theta^2$
Beta(α, β)	$\frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}$	$\alpha/(\alpha + \beta), \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$

5.8 MGF and CF Reference

Worked Solution

MGF:	$M_X(t) = \mathbb{E}[e^{tX}]$, when finite.
CF:	$\varphi_X(t) = \mathbb{E}[e^{itX}]$, always exists.
MGF of Normal(μ, σ^2):	$e^{\mu t + \frac{1}{2}\sigma^2 t^2}$.
MGF of Exponential(λ):	$\frac{\lambda}{\lambda - t}, t < \lambda$.
MGF of Bernoulli(p):	$(1 - p) + pe^t$.
MGF of Poisson(λ):	$\exp[\lambda(e^t - 1)]$.

5.9 Convergence Toolbox

Worked Solution

Chebyshev LLN:	$\mathbb{P}(\bar{X}_n - \mu > \varepsilon) \leq \sigma^2/(n\varepsilon^2)$.
MGF proof of CLT:	$M_{Z_n}(t) \rightarrow e^{t^2/2}$.
Slutsky usage:	Replace estimated variance by true one in test stats.
Delta method:	Propagate asymptotic normality via derivative.

*With these tables and the three core chapters mastered, you're fully equipped for the midterm
— both theory and computation.*