



Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM



# Métricas de resumen

Campus Miguelete, 18 de marzo de 2024





- Exploración del dataset
- Manipulación (orden, filtrado, agrupamiento)

# Algunas preguntas





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

“¿Cuánto salía un pasaje en el Titanic?”




“¿Cuánto paga por el seguro médico una persona de 42 años no fumadora?”



# Puntos usando una categórica como x (sí, ¿por qué no?)



**Licenciatura en  
Ciencia de Datos**  
ECYT\_UNSAM




# Puntos usando una categórica como x (sí, ¿por qué no?)





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM


Tarifas en el Titanic





# Puntos usando una categórica como x (sí, ¿por qué no?)




**Licenciatura en  
Ciencia de Datos**  
ECYT\_UNSAM







## Tarifas en el Titanic








## Tarifas en el Titanic





```
P <- ggplot(df, aes(x=class, y=fare, color=class)) +  
  geom_jitter(height=0) + . . .
```

OR


```
p <- ggplot(df, aes(x=class, y=fare, color=class)) +  
  geom_point(position=position_jitter(height=0, seed=1234)) + . . .
```









## Tarifas en el Titanic




# Resumir los datos por clase





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic




# Resumir los datos por clase



Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic



# Cómo se calcula



Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex       | age  | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|-----------|------|---------------------------------|--|--------|------------------------|-------|-------|-------|
|    | row names |      |                                 |  |        |                        |       |       |       |
| 1  | male      | 39.0 | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male      | 20.0 | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male      | 33.0 | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male      | 61.0 | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male      | 34.5 | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male      | 43.0 | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male      | 34.0 | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male      | 18.0 | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male      | 19.0 | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female    | 18.0 | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male      | NA   | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |




# Cómo se calcula



```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex    | age  | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|--------|------|---------------------------------|--|--------|------------------------|-------|-------|-------|
| 1  | male   | 39.0 | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male   | 20.0 | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male   | 33.0 | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male   | 61.0 | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male   | 34.5 | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male   | 43.0 | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male   | 34.0 | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male   | 18.0 | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male   | 19.0 | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female | 18.0 | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male   | NA   | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |



|   | class  | mean_fare |
|---|--------|-----------|
| 1 | First  | 85.35230  |
| 2 | Second | 21.48816  |
| 3 | Third  | 13.85217  |

# Cómo se calcula





```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex    | age  | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|--------|------|---------------------------------|--|--------|------------------------|-------|-------|-------|
| 1  | male   | 39.0 | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male   | 20.0 | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male   | 33.0 | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male   | 61.0 | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male   | 34.5 | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male   | 43.0 | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male   | 34.0 | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male   | 18.0 | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male   | 19.0 | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female | 18.0 | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male   | NA   | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |

```
df_summary <- df %>%  
  group_by(class) %>%  
  summarise(mean_fare = mean(fare, na.rm=T))
```


|   | class  | mean_fare |
|---|--------|-----------|
| 1 | First  | 85.35230  |
| 2 | Second | 21.48816  |
| 3 | Third  | 13.85217  |

# Alguna idea de la extensión





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic






# Alguna idea de la extensión





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM


## Tarifas en el Titanic





# Alguna idea de la extensión



# Licenciatura en Ciencia de Datos ECYT\_UNSAM




# Cuantiles





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic




# Cuantiles





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic



```
p + stat_summary(fun=mean,  
  fun.min = ~quantile(.x, probs=0.25),  
  fun.max = ~quantile(.x, probs=0.75),  
  size=1, color='black')
```


# Cuantiles



# Licenciatura en Ciencia de Datos

ECYT\_UNSAM

# Tarifas en el Titanic



# Cómo se calcula



Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex    | age       | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|--------|-----------|---------------------------------|--|--------|------------------------|-------|-------|-------|
|    |        | row names |                                 |  |        |                        |       |       |       |
| 1  | male   | 39.0      | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male   | 20.0      | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male   | 33.0      | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male   | 61.0      | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male   | 34.5      | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male   | 43.0      | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male   | 34.0      | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male   | 18.0      | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male   | 19.0      | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female | 18.0      | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male   | NA        | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |




# Cómo se calcula



```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex    | age  | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|--------|------|---------------------------------|--|--------|------------------------|-------|-------|-------|
| 1  | male   | 39.0 | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male   | 20.0 | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male   | 33.0 | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male   | 61.0 | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male   | 34.5 | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male   | 43.0 | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male   | 34.0 | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male   | 18.0 | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male   | 19.0 | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female | 18.0 | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male   | NA   | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |



| class  | mean_fare | Q25     | Q75  |
|--------|-----------|---------|------|
| First  | 85.35230  | 32.6156 | 93.5 |
| Second | 21.48816  | 13.0000 | 26.0 |
| Third  | 13.85217  | 7.7500  | 15.5 |



# Cómo se calcula





```
P + stat_summary(fun=mean, color='black', size=1)
```

|    | sex    | age       | siblings<br>of the<br>passenger | parents<br>/<br>children<br>aboard<br>the<br>Titanic | fare   | Port of<br>Embarkation | class | who   | alone |
|----|--------|-----------|---------------------------------|--|--------|------------------------|-------|-------|-------|
|    |        | row names |                                 |  |        |                        |       |       |       |
| 1  | male   | 39.0      | 0                               | 0  | 7.1800 | S                      | First | man   | TRUE  |
| 2  | male   | 20.0      | 0                               | 0  | 4.0125 | C                      | Third | man   | TRUE  |
| 3  | male   | 33.0      | 0                               | 0  | 5.0000 | S                      | First | man   | TRUE  |
| 4  | male   | 61.0      | 0                               | 0  | 6.2375 | S                      | Third | man   | TRUE  |
| 5  | male   | 34.5      | 0                               | 0  | 6.4375 | C                      | Third | man   | TRUE  |
| 6  | male   | 43.0      | 0                               | 0  | 6.4500 | S                      | Third | man   | TRUE  |
| 7  | male   | 34.0      | 0                               | 0  | 6.4958 | S                      | Third | man   | TRUE  |
| 8  | male   | 18.0      | 1                               | 0  | 6.4958 | S                      | Third | man   | FALSE |
| 9  | male   | 19.0      | 0                               | 0  | 6.7500 | Q                      | Third | man   | TRUE  |
| 10 | female | 18.0      | 0                               | 0  | 6.7500 | Q                      | Third | woman | TRUE  |
| 11 | male   | NA        | 0                               | 0  | 6.8583 | Q                      | Third | man   | TRUE  |

```
df_summary <- df %>%  
  group_by(class) %>%  
  summarise(mean_fare = mean(fare, na.rm=T),  
             Q25 = quantile(fare, probs=0.25, na.rm=T),  
             Q75 = quantile(fare, probs=0.75, na.rm=T))
```


| class  | mean_fare | Q25     | Q75  |
|--------|-----------|---------|------|
| First  | 85.35230  | 32.6156 | 93.5 |
| Second | 21.48816  | 13.0000 | 26.0 |
| Third  | 13.85217  | 7.7500  | 15.5 |

# Cuantiles





Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM

## Tarifas en el Titanic






# Cuantiles




Licenciatura en  
Ciencia de Datos  
ECYT\_UNSAM



## Tarifas en el Titanic






## Tarifas en el Titanic








## Tarifas en el Titanic










## Tarifas en el Titanic








## Tarifas en el Titanic









## Tarifas en el Titanic





## Tarifas en el Titanic

