

## ORIGINAL ARTICLE

## Adair-based hemoglobin equilibrium with oxygen, carbon dioxide and hydrogen ion activity

MAREK MATEJÁK, TOMÁŠ KULHÁNEK &amp; STANISLAV MATOUŠEK

*The Institute of Pathological Physiology, First Faculty of Medicine, Charles University in Prague, Czech Republic***Abstract**

As has been known for over a century, oxygen binding onto hemoglobin is influenced by the activity of hydrogen ions ( $H^+$ ), as well as the concentration of carbon dioxide ( $CO_2$ ). As is also known, the binding of both  $CO_2$  and  $H^+$  on terminal valine-1 residues is competitive. One-parametric situations of these hemoglobin equilibria at specific levels of  $H^+$ ,  $O_2$  or  $CO_2$  are also well described. However, we think interpolating or extrapolating this knowledge into an 'empirical' function of three independent variables has not yet been completely satisfactory. We present a model that integrates three orthogonal views of hemoglobin oxygenation, titration, and carbamination at different temperatures. The model is based only on chemical principles, Adair's oxygenation steps and Van't Hoff equation of temperature dependences. Our model fits the measurements of the Haldane coefficient and  $CO_2$  hemoglobin saturation. It also fits the oxygen dissociation curve influenced by simultaneous changes in  $H^+$ ,  $CO_2$  and  $O_2$ , which makes it a strong candidate for integration into more complex models of blood acid-base with gas transport, where any combination of mentioned substances can appear.

**Key Words:** Acid-base equilibrium, blood gas analysis, carboxyhemoglobin, hemoglobin A, oxyhemoglobins

**Abbreviations:** ODC, hemoglobin oxygen dissociation curve; 2,3-DPG, 2,3-diphosphoglycerate;  $[X]$ , molar concentration of X in  $\text{mol}\cdot\text{m}^{-3}$ ;  $aH^+$ , activity of hydrogen ions, where  $\text{pH} = -\log_{10}(aH^+)$ ;  $\alpha_{O_2}$ ,  $O_2$  solubility in  $\text{mol}\cdot\text{m}^{-3}\cdot\text{Pa}^{-1}$ ;  $\alpha_{CO_2}$ ,  $CO_2$  solubility in  $\text{mol}\cdot\text{m}^{-3}\cdot\text{Pa}^{-1}$ ;  $p_{O_2}$ , partial pressure of  $O_2$  in Pa;  $p_{CO_2}$ , partial pressure of  $CO_2$  in Pa;  $Hb_u$ , hemoglobin alpha or beta subunit;  $Hb_{uD}$ , deoxygenated  $Hb_u$ ;  $Hb_{uO}$ , oxygenated  $Hb_u$ ;  $Hb_uNH_3^+$ ,  $Hb_u$  with protonated Nterminus;  $Hb_{uD}NH_3^+$ ,  $Hb_{uD}$  with protonated Nterminus;  $Hb_{uO}NH_3^+$ ,  $Hb_{uO}$  with protonated Nterminus;  $Hb_uNH_2$ ,  $Hb_u$  with  $-NH_2$  form of Nterminus;  $Hb_{uD}NH_2$ ,  $Hb_{uD}$  with  $-NH_2$  form of Nterminus;  $Hb_{uO}NH_2$ ,  $Hb_{uO}$  with  $-NH_2$  form of Nterminus;  $Hb_uCOO^-$ ,  $Hb_u$  with carboxylated Nterminus;  $Hb_{uD}COO^-$ ,  $Hb_{uD}$  with carboxylated Nterminus;  $Hb_{uO}COO^-$ ,  $Hb_{uO}$  with carboxylated Nterminus;  $Hb_uAH$ ,  $Hb_u$  with protonated side-chains;  $Hb_{uD}AH$ ,  $Hb_{uD}$  with protonated side-chains;  $Hb_{uO}AH$ ,  $Hb_{uO}$  with protonated side-chains;  $Hb_uA^-$ ,  $Hb_u$  with deprotonated side-chains;  $Hb_{uD}A^-$ ,  $Hb_{uD}$  with deprotonated side-chains;  $Hb_{uO}A^-$ ,  $Hb_{uO}$  with deprotonated side-chains;  $Hb_uA^-NH_2$ , selected normalized form of  $Hb_u$  with deprotonated side-chains and  $NH_2$  form of Nterminus;  $Hb_{uD}A^-NH_2$ , deoxygenated form of  $Hb_uA^-NH_2$ ;  $Hb_{uO}A^-NH_2$ , oxygenated form of  $Hb_uA^-NH_2$ ;  $f_{nD}$ , fraction of  $Hb_{uD}A^-NH_2$  from  $Hb_{uD}$ ;  $f_{nO}$ , fraction of  $Hb_{uO}A^-NH_2$  from  $Hb_{uO}$ ;  $f_{zCD}$ , fraction of  $Hb_{uD}NH_2$  form  $Hb_{uD}$ ;  $f_{zCO}$ , fraction of  $Hb_{uO}NH_2$  form  $Hb_{uO}$ ;  $f_{hD}$ , fraction of  $Hb_{uD}A^-$  form  $Hb_{uD}$ ;  $f_{hO}$ , fraction of  $Hb_{uO}A^-$  form  $Hb_{uO}$ ;  $\Delta H^+_h$ , change of valence (charge) on side-chains during deoxygenation per one  $Hb_u$ ;  $\Delta H^+_z$ , protonation of  $-NH_2$  form of Nterminus during deoxygenation per one  $Hb_u$ ;  $\Delta H^+_c$ , decarboxylation of carboxylated Nterminus during deoxygenation per one  $Hb_u$ ;  $\Delta H^+$ , Haldane coefficient per hemoglobin subunit;  $Hb_t$ , hemoglobin tetramer without bound  $O_2$  molecules;  $(O_2)_iHb_t$ , hemoglobin tetramer with the number of i bound  $O_2$  molecules;  $(O_2)_iHb_m$ , hemoglobin tetramer composed only of  $Hb_uA^-NH_2$  subunit forms with the number of i bound  $O_2$  molecules;  $s_{O_2}$ ,  $O_2$  saturation of hemoglobin;  $s_{CO_2D}$ ,  $CO_2$  saturation of deoxyhemoglobin;  $s_{CO_2O}$ ,  $CO_2$  saturation of oxyhemoglobin;  $s_{CO_2}$ ,  $CO_2$  saturation of hemoglobin;  $t_{CO_2}$ , total concentration of  $CO_2 = \text{free dissolved } CO_2 + HCO_3^- + CO_3^{2-} + Hb_uCOO^-$ ; dTH, shift of titration curve with the change of both  $p_{O_2}$  and  $p_{CO_2}$  to zero, which equals to how many moles of strong acid must be added to one mole of  $Hb_u$  to reach the pH of the chosen standard titration curve ( $O_2$  and  $CO_2$  free); pH, acidity/basicity of hemoglobin solution (e.g. inside erythrocytes);  $pH_p$ , pH in plasma (e.g. acidity/basicity of blood).

Correspondence: Marek Mateják, U Nemocnice 5, Prague 2, 128 53, Czech Republic. Tel: +420 776 301 395. Fax: +420 224 965 916. E-mail: marek.matejak@lf1.cuni.cz

(Received 20 June 2014; accepted 2 November 2014)

ISSN 0036-5513 print/ISSN 1502-7686 online © 2015 Informa Healthcare  
DOI: 10.3109/00365513.2014.984320

## Introduction

Human hemoglobin A is one of the most extensively studied protein macromolecules. The composition and 3D conformation of both  $\alpha$  and  $\beta$  chains is known [1,2] and the binding of  $O_2$ ,  $H^+$  and 2,3-DPG has been described [2,3]. Since the 1930s it has generally been believed that  $CO_2$  binding to Hb occurs by carbamination of the amino-terminus, i.e. forming a carboxylate compound [4–6]. This hemoglobin carbamination was verified by Morrow et al. [7], who used nuclear magnetic resonance of  $^{13}CO_2$  to find its exact binding sites at valine-1. The shift of titration between oxygenated and deoxygenated forms is also well known. Called the Bohr or Haldane effect [8–10], it is caused by the same valine-1 side and by more than 10 other acid-base residues [2,11].

The most common descriptions of the hemoglobin oxygen dissociation curve (ODC) are the allosteric models [12,13], the model based on Hill equation [14,15], and Adair's four-step model [16]. Some of the ODC models [15,17,18] also include the effects of varying pH and  $CO_2$  concentration. However, these models operate only with interpolation or extrapolation of ODC from normal pH or normal  $pCO_2$  and do not take into account the chemical dependences between titration [8] and carbamination [19]. They fail when both pH and  $CO_2$  are not at normal values, especially in the alkaline pH range.

Based on these findings, we propose a hemoglobin binding equilibrium model that starts with a description of hemoglobin-oxygen dissociation using a slightly modified version of Adair's approach [16]. We continue by describing the relationship between  $CO_2$  and  $H^+$  as competitive inhibition in the amino group of terminal valine-1 residue on each chain (suffixes  $z$  and  $c$ ), which is in accordance with the known facts [4,6,20]. Finally, we lump all other acid-base side chains residues in a hemoglobin subunit into one Bohr proton-binding site of the side chain residues. These are denoted by the suffix  $h$  in the article.

## Methods

The model is built in Mathematica 9.0 (Wolfram, Champaign, IL, USA) and also in Dymola FD01 2014 (Dassault Systemes, Paris, France) as an example of chemical package in open-source Modelica library Physiobrary 2.2.0 (<http://www.physiolibrary.org>) [21,22] according to the model structure defined in the following section. The model contains only physiological parameters such as gas solubilities and dissociation coefficients of defined reactions. The unknown parameters are fitted to the data of Siggaard-Andersen [23], Bauer and Schröder [24], Severinghaus [18], Matthew et al. [1] and Reeves [25]

using Mathematica function FindFit and also using parameter estimation method suggested by Kulhanek et al. [26].



Free dissolved concentrations of  $[O_2]$  and  $[CO_2]$  in red blood cells are calculated from gas partial pressures using Henry's law ( $[O_2] = \alpha_{O_2} \cdot p_{O_2}$  and  $[CO_2] = \alpha_{CO_2} \cdot p_{CO_2}$ ) with solubility coefficients  $\alpha_{O_2} = 1.005 \cdot 10^5 \text{ mol} \cdot \text{m}^{-3} \cdot \text{Pa}^{-1}$  measured at  $38^\circ\text{C}$  by Sendroy et al. [27], and  $\alpha_{CO_2} = 2.3 \cdot 10^4 \text{ mol} \cdot \text{m}^{-3} \cdot \text{Pa}^{-1}$  at  $37^\circ\text{C}$ , as proposed by Maas et al. [28]. These solubilities are slightly different from solubilities in pure water or in plasma because of the effects of salts and proteins inside erythrocyte [27].

## Model structure

The model is built upon several simplifying assumptions. Firstly, each of four hemoglobin tetramer subunits is treated as identical, even though (slight) differences are known to exist between  $\alpha$  and  $\beta$  subunits. Secondly,  $CO_2$  or  $H^+$  binding is supposed not to affect the  $CO_2$  or  $H^+$  affinities in the other three subunits. Thus, the interaction between the subunits is modeled purely through the varying affinities of oxygen in each oxygenation step. The third simplifying assumption has already been mentioned; it is lumping all Bohr proton binding sites of subunit into two (first for the valine-1 amino-terminus and second for the side chains residues), a simplification first suggested by Antonini [29].

## Model structure of hemoglobin subunit

The three reactions that participate in the Bohr and Haldane effect of each subunit are as follows:



where the reactions  $z$  and  $c$  are competitive on the valine-1 amino-terminus, and the reaction  $h$  is independent of  $z$  or  $c$ .

The chemical equilibrium equations of these three reactions are Equations 1–3, where  $K_x$  is the equilibrium dissociation coefficients of the reaction  $x$  (i.e.  $z$ ,  $c$  or  $h$ ).

$$\text{Reaction } z: K_z = \frac{[\text{Hb}_u\text{NH}_2] \cdot a_{\text{H}^+}}{[\text{Hb}_u\text{NH}_3^+]} \quad (1)$$

$$\text{Reaction } c: K_c = \frac{[\text{Hb}_u\text{COO}^-] \cdot a_{\text{H}^+}}{[\text{CO}_2][\text{Hb}_u\text{NH}_2]} \quad (2)$$

$$\text{Reaction } h: K_h = \frac{[\text{Hb}_u\text{A}^-] \cdot a_{\text{H}^+}}{[\text{Hb}_u\text{AH}]} \quad (3)$$

These dissociation coefficients are different between oxy and deoxy subunits, which are distinguished by the subscripts O and D in the following text. They can be also written in their logarithmic

form, where  $\mathbf{pK}_x$  means  $-\log_{10}(K_x)$ . Thus, for instance,  $\mathbf{pK}_{zO}$  denotes the equilibrium coefficient of the reaction  $z$  for the oxy form of the hemoglobin subunit. Similar notation is used for describing the activity of hydrogen ions (acidity), where  $\mathbf{pH} = -\log_{10}(\mathbf{aH}^+)$ .

Using Equations 1–3 it is possible to express fractions of chosen species for deoxy and oxy subunits. We label these fractions as follows:  $\text{Hb}_u\text{NH}_2$  fractions are called  $\mathbf{f}_{zCD}$  ( $\mathbf{f}_{zCO}$ ),  $\text{Hb}_u\text{A}^-$  fractions  $\mathbf{f}_{hD}$  ( $\mathbf{f}_{hO}$ ),  $\text{Hb}_u\text{A}^-\text{NH}_2$  fractions  $\mathbf{f}_{nD}$  ( $\mathbf{f}_{nO}$ ) and  $\text{Hb}_u\text{COO}^-$  fractions  $\mathbf{sCO}_{2D}$  ( $\mathbf{sCO}_{2O}$ ) as in Equation 4–7. The selection of form  $\text{Hb}_u\text{A}^-\text{NH}_2$  from the orthogonal division into  $\text{Hb}_u\text{NH}_2$  and  $\text{Hb}_u\text{A}^-$  is also illustrated in Figure 1A, B.

$$\mathbf{f}_{zCD} = \frac{[\text{Hb}_{uD}\text{NH}_2]}{[\text{Hb}_{uD}]} = \frac{1}{1 + 10^{\mathbf{pK}_{zD} - \mathbf{pH}} + [\text{CO}_2]10^{\mathbf{pH} - \mathbf{pK}_{cD}}} \quad (4)$$

$$\mathbf{f}_{hD} = \frac{[\text{Hb}_{uD}\text{A}^-]}{[\text{Hb}_{uD}]} = \frac{1}{10^{\mathbf{pK}_{hD} - \mathbf{pH}} + 1} \quad (5)$$

$$\mathbf{f}_{nD} = \frac{[\text{Hb}_{uD}\text{A}^-\text{NH}_2]}{[\text{Hb}_{uD}]} = \mathbf{f}_{zCD} \times \mathbf{f}_{hD} \quad (6)$$

$$\mathbf{sCO}_{2D} = \frac{[\text{Hb}_{uD}\text{COO}^-]}{[\text{Hb}_{uD}]} = 10^{\mathbf{pH} - \mathbf{pK}_{cD}} \mathbf{f}_{zCD} [\text{CO}_2] \quad (7)$$

We define a titration shift as the amount of acid that must be added to achieve the same pH after full

deoxygenation of the hemoglobin subunit. This change of subunit charge during deoxygenation (by the Bohr protons) is also called Haldane coefficient  $\Delta\mathbf{H}^+$ . The coefficient can be divided into contributions of the previously mentioned reactions  $\Delta\mathbf{H}^+_{h^+}$ ,  $\Delta\mathbf{H}^+_{z^+}$  and  $\Delta\mathbf{H}^+_{c^+}$ , as is algebraically expressed by Equations 8–11.

$$\Delta\mathbf{H}^+_h = -\frac{[\text{Hb}_{uD}\text{A}^-] - [\text{Hb}_{uO}\text{A}^-]}{[\text{Hb}_u]} = -(\mathbf{f}_{hD} - \mathbf{f}_{hO}) \quad (8)$$

$$\begin{aligned} \Delta\mathbf{H}^+_z &= \frac{[\text{Hb}_{uD}\text{NH}_3^+] - [\text{Hb}_{uO}\text{NH}_3^+]}{[\text{Hb}_u]} \\ &= \left( \frac{10^{\mathbf{pK}_{zD}}}{10^{\mathbf{pH}}} \mathbf{f}_{zCD} - \frac{10^{\mathbf{pK}_{zO}}}{10^{\mathbf{pH}}} \mathbf{f}_{zCO} \right) \end{aligned} \quad (9)$$

$$\begin{aligned} \Delta\mathbf{H}^+_c &= -\frac{[\text{Hb}_{uD}\text{COO}^-] - [\text{Hb}_{uO}\text{COO}^-]}{[\text{Hb}_u]} \\ &= -(\mathbf{sCO}_{2D} - \mathbf{sCO}_{2O}) \end{aligned} \quad (10)$$

$$\Delta\mathbf{H}^+ = \Delta\mathbf{H}^+_z + \Delta\mathbf{H}^+_c + \Delta\mathbf{H}^+_h \quad (11)$$

#### Model structure of hemoglobin tetramer

Chemical speciation of the hemoglobin tetramer molecule can be considered at various levels of detail; the one chosen as appropriate in our approach is indicated in Figure 1C. The possible forms include different combinations of oxygenated and deoxygenated





Figure 1. Schema of hemoglobin calculation. (A) Possible forms of deoxyhemoglobin subunit (the blue circles). The gray circles represent hydrogen ions, and the green circles represent carbon dioxide. The arrows with dissociation coefficients represent the reactions  $z$ ,  $c$  and  $h$ . (B) Schema of chemical speciation of deoxyhemoglobin subunit. (C) Schema of chemical speciation and Adair's oxygenation steps of hemoglobin tetramer. This Figure is reproduced in color in the online version of *The Scandinavian Journal of Clinical & Laboratory Investigation*.

hemes, protonated and deprotonated oxygen-linked acid groups, and free or carboxylated amino endings of each chain. Yet because of the law of detailed balance in equilibrium [30], it is not necessary to calculate all reactions between these forms. As a result, only four oxygenation reactions need be selected for the calculation. This is done in accordance with Figure 1C, by selecting a tetramer form  $(O_2)_i Hb_m$  composed only of four subunits in the form  $Hb_u A^- NH_2$  ( $H^+$  and  $CO_2$  free), and modeling oxygen binding to these forms with Adair-type coefficients of the following reactions:



where dissociation coefficients are defined by Equation 12, which are also represented by vertical arrows at Figure 1C.

$$K_i = \frac{[(O_2)_{i-1} Hb_m] \cdot [O_2]}{[(O_2)_i Hb_m]} \quad (12)$$

For the next calculation, all forms can be expressed as a fraction of the deoxy-tetramer form  $(O_2)_0 Hb_m$ , as shown in Equation 13.

$$[(O_2)_i Hb_m] = \frac{[(O_2)_0 Hb_m] \cdot [O_2]^i}{\prod_{j=1}^i K_j} \quad (13)$$

Let us move the attention from specific forms of  $Hb_m$  to the description of the equilibrium within the whole group of  $Hb_t$ . Looking at Figure 1C, one can see that for each oxygenation step we can calculate the equilibrium in each horizontal line using Equation 14.

$$[(O_2)_i Hb_t] \cdot f_{n_D}^{4-i} \cdot f_{n_O}^i = [(O_2)_i Hb_m] \quad (14)$$

The  $CO_2$ - and pH-dependent oxygen saturation equation in the Adair style (Equation 15) is algebraically derived from Equation 13–14, where  $a_i = \frac{1}{(\prod_{j=1}^i K_j)}$   
 $x = (f_{n_D}(pH, [CO_2]) / f_{n_O}(pH, [CO_2])) \cdot [O_2]$ .

$$s_{O_2} = \frac{a_1 x + 2 a_2 x^2 + 3 a_3 x^3 + 4 a_4 x^4}{4 + 4 a_1 x + 4 a_2 x^2 + 4 a_3 x^3 + 4 a_4 x^4} \quad (15)$$

Hemoglobin saturation with  $CO_2$  ( $s_{CO_2}$ ) is calculated separately in oxygenated and deoxygenated subunits forms ( $s_{CO_{2O}}$  and  $s_{CO_{2D}}$ ) using Equation 16.

$$s_{CO_2} = s_{O_2} \cdot s_{CO_{2O}} + (1 - s_{O_2}) \cdot s_{CO_{2D}} \quad (16)$$

Finally, the shift of titration after deoxygenation and decarbamination of the hemoglobin subunit can be expressed by Equation 17.

$$dTH = s_{O_2} \cdot \Delta H^+ + s_{CO_{2D}} + \frac{s_{CO_{2D}}}{1 + 10^{pH - pK_{zD}}} \quad (17)$$

### Temperature dependences

The temperature dependences are integrated using the Van't Hoff equation (Equation 18), where  $R = 8.314 \text{ J.K}^{-1}.\text{mol}^{-1}$  is gas constant,  $\Delta H^\theta$  is the standard enthalpy change, i.e. an amount of heat consumed by a reaction changing one mole of substrates to products, and  $K_2$  and  $K_1$  are Henry's coefficients of solution or the dissociation coefficient of the chemical reaction at temperature  $T_2$  and  $T_1$  (expressed in Kelvin).

$$\ln\left(\frac{K_2}{K_1}\right) = \frac{-\Delta H^\theta}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right) \quad (18)$$

### Results

The Adair's coefficients are fitted to the collection of ODC measurements by Severinghaus [18] at  $pCO_2 = 0 \text{ Pa}$  and  $pH_p = 7.4$ , see Figure 2A and Table I. The dissociation coefficients of carboxylation are determined in close agreement with Bauer and Schröder using their data [24]; the resulting fit can be seen in Figure 2B, the coefficients are in Table II. The lumped acid dissociation coefficients for the side-chains  $pK_{hD}$  and  $pK_{hO}$  are estimated by optimization of Siggaard-Andersen's data [23] at DPG/Hbt = 0.84,  $pH = 6.5\text{--}8.0$  and  $37^\circ\text{C}$ ; the resulting fit can be seen in Figure 2C, and the coefficients complete Table II.

We conclude that the model also describes the ODC shifts by comparing with Naerea et al.'s [31] oxygen saturation measurements at different plasma pH and  $CO_2$  levels, see Figure 2D. The recalculation of data from plasma  $pH_p$  to intracellular erythrocyte pH uses the equation  $pH = 7.2464 + 0.796(pH_p - 7.4)$ , as presented by Siggaard-Andersen and Salling [32].

The gas solubility as Henry's coefficient at different temperatures can be recalculated using the enthalpy of gas solution ( $-14 \text{ kJ.mol}^{-1}$  for  $O_2$ , and  $-20 \text{ kJ.mol}^{-1}$  for  $CO_2$  [33]). If we assume that the examined hemoglobin of Matthew et al. [1] is at compatible conditions, but at a temperature of  $30^\circ\text{C}$  with  $pK_{zD} = 7.53$ ,  $pK_{zO} = 7.28$ ,  $pK_{cD} = 4.77$  and  $pK_{cO} = 5.20$  as plotted in Figure 2E, then the enthalpies of these reactions are  $-51$ ,  $8$ ,  $59$  and  $-41 \text{ kJ.mol}^{-1}$ .

Atha and Ackers measured the heat of hemoglobin oxygenation under conditions independent of carbon dioxide and Bohr protons and not including the oxygen heat of solution, coming to the value of  $59 \text{ kJ.mol}^{-1}$  [34]. Using this value for each Adair oxygenation step, one can optimize other enthalpies to fit the Reeves data [25] measured at  $19\text{--}43^\circ\text{C}$  (Figure 2F). Resulting enthalpies are  $59 \text{ kJ.mol}^{-1}$  for the deoxy and  $127 \text{ kJ.mol}^{-1}$  for the oxy version of reaction h.






Figure 2. Model curves and measured data points. (A) Points are from Severinghaus's oxygen dissociation curve collection [18] at  $p\text{CO}_2 = 0$  Pa, erythrocyte intracellular  $\text{pH} = 7.25$  and temperature  $37^\circ\text{C}$ . Lines are  $s\text{O}_2$  defined by Equation 15. (B) Carboxylation of hemoglobin measured and estimated by Bauer and Schröder [24]. The lines (Equation 7) are  $s\text{CO}_{2D}$  and  $s\text{CO}_{2O}$  at  $p\text{CO}_2 = 5.33$  kPa. (C) Bohr protons released during oxygenation of one hemoglobin subunit. Dots are data measured by Siggaard-Andersen [23] in erythrolysate at  $37^\circ\text{C}$ ,  $\text{DPG}/\text{Hb}_t = 0.84$ . Lines are calculated from Equation 11 using coefficients from Table II. From top to bottom data are plotted for different  $p\text{CO}_2 = 0$  kPa, 1.47 kPa, 4.53 kPa, 10.67 kPa and 26.66 kPa. (D) Naeraa et al.'s [31] oxygen saturation data comparison. The two groups of lines (Equation 15) are determined mainly by  $p\text{O}_2$  around 6.7 kPa and 11.7 kPa. The small nuances in each group are caused by different  $p\text{CO}_2$  changed from 0–12 kPa. (E) Matthew et al.'s [1] carbamino data measured at  $30^\circ\text{C}$  with constant content of all carbonates  $55 \text{ mmol.L}^{-1}$ . Fitted dissociation constants, which determine the lines, are used to estimate enthalpies of their reactions to be consistent with Bauer and Schröder [24]. (F) Reeves [25] oxygen saturation data at different temperatures. Enthalpies of reaction h and specific Adair oxygenation step enthalpy is estimated (lines) to fit the data (circles). This Figure is reproduced in color in the online version of *The Scandinavian Journal of Clinical & Laboratory Investigation*.

## Discussion

Having a precise quantitative description of hemoglobin behavior is a crucial aspect in describing the behavior of whole blood, which has numerous uses in medicine today. It is important in areas such as blood gas analysis [15,34], the building of complex

models [35], simulation for teaching purposes [36], or rebreathing-based methods for cardiac output estimation, which represent a more specific use [37]. For instance, the precision of the latter methods is crucially dependent on the exact calculation of the total amount of carbon dioxide in blood for various

Table I. Estimated form-specific Adair's coefficients [ $\text{mol.m}^{-3}$ ] at  $37^\circ\text{C}$ .

$K_1$	$K_2$	$K_3$	$K_4$
0.0121	0.0117	0.0871	0.000386

Table II. Estimated acid dissociation coefficients at  $37^\circ\text{C}$ .

Reaction z	Reaction c	Reaction h
$pK_{zD} = 7.73$	$pK_{cD} = 7.54$	$pK_{hD} = 7.52$
$pK_{zO} = 7.25$	$pK_{cO} = 8.35$	$pK_{hO} = 6.89$

(abnormal) patient conditions, as has been recently pointed out [38].

This model offers a precise description of various phenomena that take place with hemoglobin, based on relatively simple starting points, such as competitive binding of  $H^+$  and  $CO_2$  at valine-1 amino terminus or the law of detailed balance [30]. Even with a simple structure, it offers a remarkably good fit to the data of hemoglobin oxygenation, titration, and carbamination at different temperatures, as shown in Figure 2A–E. These can be calculated with any combination of oxygen, carbon dioxide and hydrogen ions defined as open system (lungs), where the partial pressures are equilibrated, and also in a closed system (tissues), where mass conservation laws and the total amount of substances take place, as was also modeled by Rees and Andreassen [39], among others.

The results of our model (Figure 2A–E) show strong nonlinear dependences between variables, which is in agreement with the known data [1,23–25,40]. Looking at Figure A, one can compare sets of ODCs, where each color represents a different pH. Various curves of each color represent ODCs for various levels of  $pCO_2$  for a given pH. As can be seen, the effect of  $pCO_2$  on the ODC is stronger at high (alkaline) pH, which is in agreement with the data [23]. Similarly, one can compare the curves within the sets of solid or dashed lines, where each set represents dissociation curves for various values of pH at a given level of  $pCO_2$ . The variation between the curves of each line type represents the Bohr effect for the given level of  $CO_2$ , this effect can also be appreciated from data of Figure 2C, which shows the average amount of released  $H^+$  upon oxygenation of one hemoglobin subunit.

The model uses enthalpies to calculate temperature dependences of hemoglobin behavior (Figure 2F, E), which allows examination of the heat transfers during single chemical processes. For instance, binding of aqueous oxygen onto hemoglobin produces 30–40 kJ/mol of heat [41–44], and the same amount of heat is consumed by deoxygenation process in metabolically active tissues, thus helping to cool them down. When the hemoglobin model is used in the standard conditions of a large-scale Hum-Mod model [35], the resulting heat transfer due to the exothermy of the hemoglobin oxygen reaction is 4–7% of the total heat produced by muscle, which is in agreement with experimental results [41,43,44]. It is interesting to note that this heat transfer occurs without any increase in blood temperature.

The integration of chemical processes in a macromolecule requires a precise view into their underlying principles. Some physiologists use the elementary chemical equations [39], but do not implement the principle of detailed balance [30]. Other physiologists make empirically-based equations with a raw linear gradient approximation of possible combinations of model values [15]. We feel that it is almost

impossible to see the problem as this type of black-box function with more than two inputs. Instead, it is better to have an integrated model of oxygenation, titration and carbamination.

Today, allosteric hemoglobin oxygenation models do exist that seem more in agreement with the structural knowledge of hemoglobin [12,45–47]. These models take into account two or more structurally different forms: relaxed and tensed. However, these models have so far been limited to hemoglobin oxygenation only. Our Adair-based model can explain not only oxygen and carbon dioxide saturation, but also their cooperation with acid-base buffering properties of hemoglobin. All three of these connected phenomena fit to measured data in physiological ranges.

As any work, the presented model has its limitations. First of all, it does not include the effects of changes in Hb tetramer conformations. Also the model could be extended with dependences on electrolytes such as chloride, 2,3-bisphosphoglycerate or other organic phosphates and their binding reactions, as many research studies show these interactions [48–51]. The next extension of this model could be performed by the integration of an intracellular red cell environment to calculate with phosphate acid-base buffers, and finally the membrane changes with blood plasma, where the chloride shift reaches a Gibbs-Donnan equilibrium and establishes chloride, bicarbonate and hydrogen ion activity ratios. Having an integrated model of blood gases and acid-base is crucial if we want the precise computational algorithms of the current state of a patient. These calculations could be used, for example, inside the next generation of medical devices to estimate not only blood properties, but also the connected properties of circulation [37] or metabolic functions [39].

In this article, we present a hemoglobin model that integrates  $O_2$ ,  $CO_2$  and  $H^+$  binding. The model is not just empirical, but is based on sound theoretical principles, such as the competitive binding of  $CO_2$  and  $H^+$  on the valine-1  $NH_2$  terminus, the Bohr and Haldane effect [9,52] and on the principle of detailed balance. The principle of detailed balance is used for the first time in the Adair type of model. The advantages of this approach include explicit formulation of mass and energy conservation principles: The model accumulates substances and heat inside hemoglobin forms, making it very useful for integration in higher-scale dynamic models.

## Acknowledgements

In addition, the authors want to thank doc. MUDr. Jiří Kofránek, CSc, and Prof. MUDr. Emanuel Nečas, DrSc, for critical review and Austin Schaefer for English language editing.

This work was supported by grants from the Ministry of Industry and Trade of the Czech Republic: MPO TIP TI3/869 – Virtual patient for medical education; and by The Ministry of Education, Youth and Sports via the grant SVV-2014-260033.

**Declaration of interest:** The authors report no conflict of interest. The authors alone are responsible for the content and writing of the paper.

## References

- [1] Matthew JB, Morrow J, Wittebort RJ, Gurd F. Quantitative determination of carbamino adducts of alpha and beta chains in human adult hemoglobin in presence and absence of carbon monoxide and 2, 3-diphosphoglycerate. *J Biol Chem* 1977;252:2234–44.
- [2] Perutz M. Species adaptation in a protein molecule. *Mol Biol Evol* 1983;1:1–28.
- [3] Perutz M, Kilmartin J, Nishikura K, Fogg J, Butler P, Rollemma H. Identification of residues contributing to the Bohr effect of human haemoglobin. *J Mol Biol* 1980;138: 649–68.
- [4] Rossi-Bernardi L, Roughton F. The specific influence of carbon dioxide and carbamate compounds on the buffer power and Bohr effects in human haemoglobin solutions. *J Physiol* 1967;189:1–29.
- [5] Forster R, Constantine H, Craw MR, Rotman H, Klocke R. Reaction of CO<sub>2</sub> with human hemoglobin solution. *J Biol Chem* 1968;243:3317–26.
- [6] Stadie WC, O'Brien H. The carbamate equilibrium I. The equilibrium of amino acids, carbon dioxide, and carbamates in aqueous solution; with a note on the Ferguson-Roughton carbamate method. *J Biol Chem* 1936;112:723–58.
- [7] Morrow J, Matthew J, Wittebort R, Gurd F. Carbon 13 resonances of 13CO<sub>2</sub> carbamino adducts of alpha and beta chains in human adult hemoglobin. *J Biol Chem* 1976;251:477–84.
- [8] Bohr C, Hasselbalch K, Krogh A. Concerning a biologically important relationship—the influence of the carbon dioxide content of blood on its oxygen binding. *Skand Arch Physiol* 1904;16:402.
- [9] Siggaard-Andersen O, Garby L. The Bohr effect and the Haldane effect. *Scand J Clin Lab Invest* 1973;31:1–8.
- [10] Tyuma I. The Bohr effect and the Haldane effect in human hemoglobin. *Jpn J Physiol* 1984;34:205.
- [11] Matthew J, Hanania G, Gurd F. Electrostatic effects in hemoglobin: hydrogen ion equilibria in human deoxy- and oxyhemoglobin A. *Biochemistry* 1979;18:1919.
- [12] Monod J, Wyman J, Changeux J-P. On the nature of allosteric transitions: a plausible model. *J Mol Biol* 1965;12:88–118.
- [13] Eaton WA, Henry ER, Hofrichter J, Bettati S, Viappiani C, Mozzarelli A. Evolution of allosteric models for hemoglobin. *IUBMB Life* 2007;59:586–99.
- [14] Hill AV. The combinations of haemoglobin with oxygen and with carbon monoxide. I. *Biochem J* 1913;7:471.
- [15] Siggaard-Andersen O, Siggaard-Andersen M. The oxygen status algorithm: a computer program for calculating and displaying pH and blood gas data. *Scand J Clin Lab Invest* 1990;50:29–45.
- [16] Adair GS. The hemoglobin system VI. The oxygen dissociation curve of hemoglobin. *J Biol Chem* 1925;63:529–45.
- [17] Dash RK, Bassingthwaite JB. Erratum to: Blood HbO<sub>2</sub> and HbCO<sub>2</sub> dissociation curves at varied O<sub>2</sub>, CO<sub>2</sub>, pH, 2, 3-DPG and temperature levels. *Ann Biomed Eng* 2010; 38:1683–701.
- [18] Severinghaus JW. Simple, accurate equations for human blood O<sub>2</sub> dissociation computations. *J Appl Physiol* 1979; 46:599–602.
- [19] Christiansen J, Douglas C, Haldane J. The absorption and dissociation of carbon dioxide by human blood. *J Physiol* 1914;48:244–71.
- [20] Perrella M, Bresciani D, Rossi-Bernardi L. The binding of CO<sub>2</sub> to human hemoglobin. *J Biol Chem* 1975;250: 5413–8.
- [21] Mateják M, Kulhánek T, Šilar J, Privitzer P, Ježek F, Kofránek J. Physiobase-Modelica library for Physiology. 10th International Modelica Conference; 2014 March 12; Lund, Sweden: Linköping University Electronic Press; 2014. pp. 499–505.
- [22] Mateják M. Physiology in Modelica. *Mefanet J* 2014; 2:10–4.
- [23] Siggaard-Andersen O. Oxygen-linked hydrogen ion binding of human hemoglobin. effects of carbon dioxide and 2, 3-diphosphoglycerate I. Studies on erythrolysate. *Scand J Clin Lab Invest* 1971;27:351–60.
- [24] Bauer C, Schröder E. Carbamino compounds of haemoglobin in human adult and foetal blood. *J Physiol* 1972;227: 457–71.
- [25] Reeves RB. The effect of temperature on the oxygen equilibrium curve of human blood. *Respir Physiol* 1980; 42:317–28.
- [26] Kulhanek T, Matejak M, Silar J, Kofranek J. Parameter estimation of complex mathematical models of human physiology using remote simulation distributed in scientific cloud. Biomedical and Health Informatics (BHI), 2014 IEEE-EMBS International Conference on 1–4 June 2014; pp. 712–5.
- [27] Sendroy J, Dillon RT, Van Slyke DD. Studies of gas and electrolyte equilibria in blood XIX. The solubility and physical state of uncombined oxygen in blood. *J Biol Chem* 1934;105:597–632.
- [28] Maas A, Rispens P, Siggaard-Andersen O, Zijlstra W. On the reliability of the Henderson-Hasselbalch equation in routine clinical acid-base chemistry. *Ann Clin Biochem* 1984;21:26.
- [29] Antonini E, Wyman J, Brunori M, Fronticelli C, Bucci E, Rossi-Fanelli A. Studies on the relations between molecular and functional properties of hemoglobin V. The influence of temperature on the Bohr effect in human and in horse hemoglobin. *J Biol Chem* 1965;240:1096–103.
- [30] Sauro HM. Enzyme kinetics for systems biology. Future Skill Software (Ambrosius Publishing); 2011. pp. 20–1.
- [31] Naeraa N, Petersen ES, Boye E. The influence of simultaneous, independent changes in pH and carbon dioxide tension on the in vitro oxygen tension-saturation relationship of human blood. *Scand J Clin Lab Invest* 1963;15:141–51.
- [32] Siggaard-Andersen O, Salling N. Oxygen-linked hydrogen ion binding of human hemoglobin. Effects of carbon dioxide and 2, 3-diphosphoglycerate. II. Studies on whole blood. *Scand J Clin Lab Invest* 1971;27:361–6.
- [33] Lide DR. CRC handbook of chemistry and physics. Boca Raton, FL: CRC Press; 2004.
- [34] Atha DH, Ackers GK. Calorimetric determination of the heat of oxygenation of human hemoglobin as a function of pH and the extent of reaction. *Biochemistry* 1974;13:2376–2382.
- [35] Hester RL, Brown AJ, Husband L, Iliescu R, Pruett D, Summers R, Coleman TG. HumMod: a modeling environment for the simulation of integrative human physiology. *Front Physiol* 2011;2.
- [36] Kofranek J, Matousek S, Ruz J, Stodulka P, Privitzer P, Matejak M, Tribula M. The Atlas of Physiology and Pathophysiology: Web-based multimedia enabled interactive simulations. *Comput Methods Programs Biomed* 2011;104: 143–53.
- [37] Jarvis SS, Levine BD, Prisk GK, Shykoff BE, Elliott AR, Rosow E, Blomqvist CG, Pawelczyk JA. Simultaneous determination of the accuracy and precision of closed-circuit cardiac output rebreathing techniques. *J Appl Physiol* 2007; 103:867–74.

- [38] Sun X-G, Hansen JE, Ting H, Chuang M-L, Stringer WW, Adame D, Wasserman K. Comparison of exercise cardiac output by the Fick Principle using oxygen and carbon dioxide. *CHEST J* 2000;118:631–40.
- [39] Rees SE, Andreassen S. Mathematical models of oxygen and carbon dioxide storage and transport: the acid-base chemistry of blood. *Crit Rev Biomed Eng* 2005;33.
- [40] Imai K, Yonetani T. PH dependence of the Adair constants of human hemoglobin. Nonuniform contribution of successive oxygen bindings to the alkaline Bohr effect. *J Biol Chem* 1975;250:2227–31.
- [41] Adair G, Cordero N, Shen T. The effect of temperature on the equilibrium of carbon dioxide and blood and the heat of ionization of haemoglobin. *J Physiol* 1929;67:288–98.
- [42] Chipperfield J, Rossi-Bernardi L, Roughton F. Direct calorimetric studies on the heats of ionization of oxygenated and deoxygenated hemoglobin. *J Biol Chem* 1967;242:777–83.
- [43] Weber RE, Campbell KL. Temperature dependence of haemoglobin-oxygen affinity in heterothermic vertebrates: mechanisms and biological significance. *Acta Physiolog* 2011;202:549–62.
- [44] Weber RE, Fago A, Campbell KL. Enthalpic partitioning of the reduced temperature sensitivity of O<sub>2</sub> binding in bovine hemoglobin. *Comparat Biochem Physiol A: Molec Integrat Physiol* 2014;176:20–5.
- [45] Szabo A, Karplus M. A mathematical model for structure-function relations in hemoglobin. *J Mol Biol* 1972;72:163–97.
- [46] Brunori M, Coletta M, Di Cera E. A cooperative model for ligand binding to biological macromolecules as applied to oxygen carriers. *Biophys Chem* 1986;23:215–22.
- [47] Henry ER, Bettati S, Hofrichter J, Eaton WA. A tertiary two-state allosteric model for hemoglobin. *Biophys Chem* 2002;98:149–64.
- [48] de Bruin SH, Rollema HS, Janssen LH, van Os GA. The interaction of chloride ions with human hemoglobin. *Biochem Biophys Res Commun* 1974;58:210–5.
- [49] Rollema HS, de Bruin SH, Janssen L, Van Os G. The effect of potassium chloride on the Bohr effect of human hemoglobin. *J Biol Chem* 1975;250:1333–9.
- [50] Benesch R, Benesch RE, Yu CI. Reciprocal binding of oxygen and diphosphoglycerate by human hemoglobin. *Proc Natl Acad Sci USA* 1968;59:526.
- [51] Bunn HF, Briehl RW. The interaction of 2, 3-diphosphoglycerate with various human hemoglobins. *J Clin Invest* 1970;49:1088.
- [52] Siggaard-Andersen O, Salling N, Nørgaard-Pedersen B, Rørth M. Oxygen-linked hydrogen ion binding of human hemoglobin. effects of carbon dioxide and 2, 3-diphosphoglycerate: III. Comparison of the Bohr effect and the Haldane effect. *Scand J Clin Lab Invest* 1972;29:185–93.