

1 The complex network of trophic interactions in a subAntarctic oceanic Marine Protected Area

2 Tomás I. Marina<sup>1,\*</sup>, Irene R. Schloss<sup>1,2,3</sup>, Gustavo A. Lovrich<sup>1</sup>, Claudia C. Boy<sup>1</sup>, Daniel O.  
3 Bruno<sup>1,3</sup>, Fabiana L. Capitanio<sup>4,5</sup>, Sergio M. Delpiani<sup>6</sup>, Juan Martín Díaz de Astarloa<sup>6</sup>, Cintia  
4 Frayse<sup>1</sup>, Virginia A. García Alonso<sup>4,5</sup>, Andrea Raya Rey<sup>1,3,7</sup>, Laura Schejter<sup>8</sup>, Mariela L.  
5 Spinelli<sup>4,5</sup>, Marcos Tatián<sup>9,10</sup>, Diego Urteaga<sup>11</sup>, Luciana Riccialdelli<sup>1</sup>

6 <sup>1</sup>Centro Austral de Investigaciones Científicas (CADIC-CONICET), 9410, Ushuaia,  
7 Argentina

8 <sup>2</sup>Instituto Antártico Argentino (IAA), 1650, CABA, Argentina

9 <sup>3</sup>Instituto de Ciencias Polares, Ambiente y Recursos Naturales (ICPA), Universidad de Tierra  
10 del Fuego (UNTDF), 9410, Ushuaia, Argentina

11 <sup>4</sup>Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA), C1428EGA,  
12 CABA, Argentina

13 <sup>5</sup>Instituto de Biodiversidad y Biología Experimental y Aplicada (IBBEA), Universidad de  
14 Buenos Aires-CONICET, C1428EGA, CABA, Argentina

15 <sup>6</sup>Instituto de Investigaciones Marinas y Costeras (IIMYC), Universidad Nacional de Mar del  
16 Plata-CONICET, 7600, Mar del Plata, Argentina

17 <sup>7</sup>Wildlife Conservation Society, Argentina

18 <sup>8</sup>Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), Consejo Nacional de  
19 Investigaciones Científicas y Técnicas (CONICET), 7600, Mar del Plata, Argentina

20 <sup>9</sup>Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba (UNC),  
21 5000, Córdoba, Argentina

22 <sup>10</sup>Instituto de Diversidad y Ecología Animal (IDEA-CONICET), 5000, Córdoba, Argentina

23 <sup>11</sup>Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, C1405DJR, CABA,  
24 Argentina

25 Abstract: The total area of the world ocean designated under marine protection has increased  
26 recently. Most Marine Protected Areas (MPAs) target vulnerable, keystone, charismatic and/  
27 or endemic species. In the sub-Antarctic, ocean protection is associated to oceanic islands,  
28 except for MPAs Namuncurá - Burdwood Bank I and II (MPA N-BB, ~53°–55°S and ~56°–  
29 62°W), which are associated to a submarine plateau and its southern adjacent deep slope,  
30 respectively. Here, we present the first analysis of the network of predator-prey interactions  
31 for the MPA N-BB. We applied a network approach to characterise the complexity and  
32 structure of the food web, and identify the species' role in such a framework. The MPA N-BB  
33 food web consisted of 1788 interactions and 379 species, with a connectance of 0.01. Almost  
34 half of the consumers were omnivores (0.48), and the network displayed a small-world  
35 pattern. These suggest that the ecosystem might be vulnerable to perturbations targeting  
36 highly connected species, although other properties might provide resilience and resistance,  
37 resulting in a rearranged structure that preserves its original functions. Several species arose  
38 as important in terms of different aspects of trophic structure and functioning, and response to  
39 perturbations. Generalist species, mainly fishes, play a crucial role in the ecosystem's benthopelagic coupling and should be considered as relevant energy transfers for the ecosystem. We  
40 argue that the diversity of species, including both the benthic and pelagic habitats, is  
41

### 3 TROPHIC INTERACTIONS IN A MARINE PROTECTED AREA

42 responsible for securing the connectivity within the food web against perturbations, therefore  
43 contributing to the structure and stability of the ecosystem.

44 Keywords: Food web; Complexity; Structure; Burdwood Bank; Southwest Atlantic.

## 45 1. INTRODUCTION

46 The evidence of benefits provided by Marine Protected Areas (MPAs) as well as the urgent  
47 need for ocean protection have driven an unprecedented increase in the number of MPAs  
48 worldwide in recent years (Roberts et al. 2017, Sala et al. 2018). Globally, the total area of  
49 the world ocean designated under marine protection adds up to approximately 29,600,000  
50 km<sup>2</sup>, distributed across nearly 18,444 MPAs and covering 8.16% of the ocean's surface  
51 (IUCN 2023), and therefore approaching the 10% goal of the Convention of Biological  
52 Diversity (Secretariat of the Convention on Biological Diversity 2004). Despite this progress,  
53 recent reports have shown that actual protection has been overestimated because it includes  
54 areas that are not yet effectively protected (only declared) as well as areas that allow  
55 significant extractive activities (Sala et al. 2018).

56 In the sub-Antarctic region, the level of ocean protection is mainly associated to oceanic  
57 islands, such as the South Georgias and South Sandwich, Bouvet, Prince Edward, and  
58 Macquarie islands (IUCN 2023). Interestingly, the case of the MPAs Namuncurá - Burdwood  
59 Bank I and II (MPA N-BB, ~53°–55°S ~56°–62°W), which is the focus of this work, is unique  
60 since these MPAs are associated to a submarine plateau and its southern adjacent deep slope  
61 region, respectively (Falabella 2017, Schejter et al. 2020). In addition, such MPAs are part of  
62 a network of protected areas in the sub-Antarctic area (jointly with MPA Yaganes) that aims  
63 to protect this southern region in order to contribute to global ocean health.

64 Many of these MPAs focus on the presence of particularly vulnerable, keystone, or  
65 charismatic species, large numbers (or proportions) of endemic species, and/or high  
66 biodiversity across taxonomic levels (Hogg et al. 2016). Indeed, the MPA N-BB was created  
67 to protect a potentially sensitive and biodiverse benthic habitat that was only barely known  
68 (Schejter et al. 2016, Falabella 2017). The benthic community is featured by high biomass of  
69 vulnerable and fragile species (mainly Porifera, Bryozoa and Cnidaria) that considered with

their environment meet the characteristics of vulnerable marine ecosystems (Schejter & Albano 2021), here defined as sites that present densities of Indicator Taxa of  $> 10 \text{ kg per } 1200 \text{ m}^2$  (CCAMLR 2009). Also, the benthic realm provides habitat to several small-sized species (Lopez-Gappa et al. 2018, Martin Siritto 2019, Schejter & Bremec 2019), and has an important role in the life history of fishes as a food source, refuge and nursery area (García Alonso et al 2018, Vazquez et al 2018, Delpiani et al. 2020, Troccoli et al. 2020, Covatti Ale et al. 2022, Fischer et al. 2022, Matusevich et al. 2023). The maintenance of this singular community is related to local and regional oceanographic processes, including the circulation of the rich Malvinas (Falkland) current in the area (Piola & Gordon 1989, Guerrero et al. 1999) and the upwelling and mixing phenomena (Matano et al. 2019). The input of nutrients from the Malvinas (Falkland) current also supports a diverse plankton community (Guinder et al. 2020).

Overall, 811 benthic and plankton species have been identified for the MPA N-BB ecosystem, where 349 were reported for the first time in the area in recent years (APN 2022). Identifying the main species involved in the maintenance of ecosystem services and health as well as for management and conservation is essential. Recently, the structure of the southwestern South Atlantic Ocean has been proposed to be under a 'wasp-waist' control, meaning that the structure and dynamics of the ecosystem are regulated primarily by mid-trophic level species (e.g., fishes, crustaceans) (Padovani et al. 2012, Saporiti et al. 2015, Riccialdelli et al. 2020). In particular, the ecosystem of the MPA N-BB shows a more pronounced 'wasp-waist' structure, meaning a shorter food chain and a greater trophic overlap and redundancy, than other sub-Antarctic areas, such as the continental shelf off Tierra del Fuego. The Fuegian sprat *Sprattus fuegensis* and longtail southern cod *Patagonotothen ramsayi* are considered the most plausible 'wasp-waist' species (Riccialdelli et al. 2020).

High-latitude marine ecosystems, such as the MPA N-BB, are complex systems in terms of

biodiversity and ecological interactions (Day et al. 2013, Kortsch et al. 2019, Cordone et al. 2020, Trathan et al. 2021). Although there is a robust knowledge about the complexity considering the richness of the benthic and plankton communities in the MPA N-BB ecosystem (Schejter et al. 2016, Guinder et al. 2020, Schejter et al. 2020, APN 2022), a better understanding of species interactions' complexity and structure is needed.

This aspect can be tackled by analysing one of the most frequent relationships between species: the predator-prey interaction (Bascompte 2009). The sum of predator-prey or trophic interactions of a particular region is referred to as a food web, representing the roadmap for matter and energy flow in an ecosystem.

In recent years, network approaches have been successfully applied to study complex high-latitude marine ecosystems, improving our knowledge on structure, functioning, and response to environmental/anthropogenic changes (Kortsch et al. 2015, Cordone et al. 2018, Funes et al. 2022, Marina et al. 2023). Among anthropogenic threats, it is worth mentioning that contaminants like mercury and microplastics have been recently reported as important threats to the MPA N-BB region (Cossi et al. 2021, Di Mauro et al. 2022, Fioramonti et al. 2022); also fishing vessels are allowed to operate in the western section of the MPA N-BB (i.e. Marine National Reserve category), altering the stocks of commercially important fish species (Martinez et al. 2021, APN 2022). Moreover, there is a potential hazard related to the effects of offshore activities (exploration and exploitation) to the west of the MPA N-BB (APN 2022).

In the present work, we present the first detailed analysis of the network of predator-prey interactions, hereafter food web, for the MPA N-BB ecosystem. For this, we applied a network approach to a highly resolved food web. The objective was twofold: characterise the food web in terms of complexity and structure, and identify the species' role in the network.

## 2. MATERIALS & METHODS

## 120 2.1. STUDY AREA

121 The MPAs Namuncurá - Burdwood Bank I and II, created by National Laws 26.875 in 2013  
122 and 27.490 in 2017, comprise a shallow submarine plateau called Burdwood Bank (BB) and a  
123 deep slope that reaches 4000 m in depth, N-BB I and N-BB II, respectively (Tombesi et al.  
124 2020, APN 2022) (Figure 1). They are located 150 km east of Isla de los Estados and 200 km  
125 south of Malvinas/Falkland Islands.

126 The MPA N-BB I comprises nearly 28,900 km<sup>2</sup> circumscribed by the 200 m isobath,  
127 between ~54°–55°S and ~56°–62°W, with a slight slope extended nearly 370 km east–west.  
128 Physical features in the BB are fairly stable, with salinity averaging 34 all year round and  
129 temperature ranging between 4 and 8°C (Guerrero et al. 1999, Acha et al. 2004, Piola &  
130 Falabella 2009). The BB is surrounded by steep flanks of up to 4000 m depth through which  
131 strong currents circulate (Piola & Gordon 1989, Reta 2014, Matano et al. 2019). The N-BB II  
132 includes such a deep slope, protecting about 32,000 km<sup>2</sup> (~55°–56°S, ~58°–62°W). Intense  
133 upwelling and mixing occur in relation with the slope, entraining deep nutrient-rich waters  
134 into the photic layer (Piola & Falabella 2009, Matano et al. 2019) and resulting in a fairly  
135 homogeneous water column both spatially and temporally (Glorioso & Flather 1995,  
136 Guerrero et al. 1999, Matano et al. 2019).

137 Given the evidence collected during several research cruises about the oceanographic and  
138 ecological processes connecting MPAs N-BB I and II (references in APN 2022), a joint  
139 management plan was recently proposed (APN 2022). This is why the study area of the  
140 present work includes both MPAs.

## 141 2.2. NETWORK CONSTRUCTION

142 In order to build the network of predator-prey interactions, we reviewed more than 170  
143 references considering published articles, Ph.D. theses, public databases, and reports  
144 belonging to 16 research cruises conducted in the MPAs N-BB I and II during 2014–2019. It

145 is noteworthy that the sampling effort was greater in the MPA N-BB I. Furthermore, we took  
 146 into account personal communications from experts belonging to the working group of the  
 147 study area (<https://www.pampazul.gob.ar/tag/banco-burdwood/>). The diversity of the authors'  
 148 expertise contributing to the present study was a key factor in enhancing the quality of the  
 149 network, and inherently improved the network representation. A list of the references used to  
 150 build the network is presented in Supplementary Material (Table S1).

151 Due to a lack of trophic data resolution for some species inhabiting the study area, we  
 152 followed the concept of trophic species, here defined as follows: taxa collapsed into a single  
 153 node in the network. In most cases, we followed this concept when specific data on species,  
 154 in the taxonomic sense, were not available. In some cases, we collapsed species when taxa  
 155 shared the same set of predators and prey (trophic similarity, Martinez 1991), one of the  
 156 aggregation methods that better preserve food web functional properties (Gauzens et al.  
 157 2013). In addition, for endemic species (e.g. bryozoan *Burdwoodipora paguricola*) and other  
 158 species with no trophic studies so far, we inferred their feeding interactions applying a  
 159 conservative approach that assumes that the set of prey and predators are at some point  
 160 preserved in time. In those cases we gathered information from upper taxonomic levels (i.e.  
 161 Genus, Family, Order, Class, Phylum) as a good proxy variable (Morales-Castilla et al. 2015,  
 162 Pomeranz et al. 2019). Details about this can be found in Supplementary Material (Table S2).  
 163 Furthermore, we considered non-living food sources, such as detritus and necromass, as prey  
 164 species in the food web context.

165 With the gathered trophic data, we constructed a matrix of pairwise interactions; a value of 1  
 166 or 0 was assigned to each element  $a_{ij}$  of the matrix depending on whether the j-species preyed  
 167 or not on the i-species. Then we transformed such a matrix into an oriented graph with L  
 168 trophic interactions between S nodes or species. The orientation or direction of the graph  
 169 follows the flow of energy and matter in the network, from prey to predator.

## 170 2.3. NETWORK ANALYSIS

171 We analysed the MPA N-BB network of trophic interactions, or food web, at two levels: A)  
 172 network, considering species and interactions of the whole network; and B) species,  
 173 considering interactions and species related to a particular species (Table 1).

174 The network-level analysis aims to characterise the food web in terms of complexity and  
 175 structure. For this, we calculated several network properties commonly used to describe  
 176 empirical food webs (Pascual & Dunne 2005): (1) number of species  $S$ ; (2) number of  
 177 interactions or links  $L$ ; (3) link density  $L/S$ ; (4) connectance  $L/S^2$ ; (5) omnivory  $Omn$ ; and (6)  
 178 small-world pattern. In order to explore the small-world phenomenon, we analysed the  
 179 characteristic path length ( $CPL$ ) and the clustering coefficient ( $CC$ ). The  $CPL$  is the average  
 180 shortest path length between all pairs of nodes (Watts & Strogatz 1998). Here,  $CPL$  was  
 181 calculated as the average number of nodes in the shortest path  $CPL_{Min}(i,j)$  between all pairs of  
 182 nodes  $S(i,j)$  in a network averaged over  $S(S-1)/2$  nodes:

$$CPL = \frac{2}{S(S-1)} \sum_{i=1}^S \sum_{j=1}^S CPL_{Min}(i, j)$$

183 The  $CC$  quantifies the local interconnectedness of the network and it is defined as the fraction  
 184 of the number of existing links between neighbours of node  $i$  among all possible links  
 185 between these neighbours. In this study, the  $CC$  was determined as the average of the  
 186 individual clustering coefficients  $CC_i$  of all the nodes in the network. Individual  $CC_i$  were  
 187 determined as follows:

$$CC_i = \frac{2 E_i}{K_i(K_i - 1)}$$

188 where  $E_i$  is the effective number of interactions between  $K_i$  nearest-neighbour nodes of node  $i$   
 189 and the maximal possible number of such interactions (Newman 2003).

190 To test whether the food web presented the small-world pattern, we compared the empirical



191 values of *CPL* and *CC* with those resulting from 1000 randomly generated networks with the  
 192 same size (*S*) and number of interactions (*L*), following the method proposed by Marina et al.  
 193 2018b.

194 Also, we estimated the (7) degree distributions for the food web, prey and predators, and each  
 195 functional group (e.g., Amphipoda, Ascidiacea, Bivalvia, fish, marine mammals, seabirds,  
 196 among others). The prey and predator distributions indicate the frequency of prey among  
 197 predators, and vice versa; the functional group's degree shows the distribution of interactions  
 198 within groups.

199 The species-level analysis aims to describe the species' role in the food web. For this, we  
 200 considered the following properties: betweenness *B<sub>tw</sub>*, closeness *C<sub>l</sub>*, trophic similarity *T<sub>S</sub>*,  
 201 topological role *T<sub>R</sub>*, and trophic level *T<sub>L</sub>* (Table 1). Topological roles refer to the fact that  
 202 food webs tend to naturally organize in non-random, modular patterns, where modules are  
 203 defined as a group of species that interact more frequently among themselves than with  
 204 species that are not members of the module (Guimerà & Nunes Amaral 2005). Species can  
 205 play different roles in this respect, according to the pattern of interactions within their own  
 206 module and/or across modules. We computed the topological role for each species, classified  
 207 as module hub, species with a relatively high number of interactions, but most within its own  
 208 module; module specialist, species with relatively few interactions and most within its own  
 209 module; module connector, species with relatively few interactions mainly between modules;  
 210 and network connector, species with high connectivity between and within modules (Guimerà  
 211 & Nunes Amaral 2005).

212 We also studied the relationship between species *T<sub>L</sub>* and the other species properties by  
 213 performing linear regression analyses. Thus, we considered the *T<sub>L</sub>* as the dependent variable  
 214 and the given property (i.e. betweenness, closeness, trophic similarity) as the independent  
 215 variable and obtained the coefficients (slope and intercept) for the linear model. Models were

216 fitted using the least squares approach. We also explored the topological role categories with  
 217 the species TL. These species-level properties provide an appropriate description of species'  
 218 role in empirical complex food webs (Cirtwill et al. 2018).

219 All network analyses and graphs were performed in R version 4.2.2 (R Core Team 2022),  
 220 mainly using 'igraph' (Csardi & Nepusz 2006) and 'multiweb' (Saravia 2022) packages. The  
 221 source code and data are available at <https://github.com/TomasMarina/Banco-Burdwood>.

## 222 3. RESULTS

### 223 3.1. NETWORK-LEVEL PROPERTIES

224 In terms of complexity, the MPA Namuncurá - Burdwood Bank food web consisted of 1788  
 225 predator-prey interactions and 379 species, where 93% of them were defined at the species  
 226 taxonomical level (Figure 2, Table S2). The food web presented a link density (e.g., the  
 227 average number of interactions per species) of 4.72, and a connectance of 0.01. Almost half  
 228 of the consumers were omnivores (0.48), feeding on sources at different trophic levels. The  
 229 food web displayed a small-world pattern, meaning that the path length was lower and the  
 230 clustering coefficient higher than the random networks (Table 2).

231 The degree distribution of the food web showed an asymmetric frequency in the number of  
 232 interactions, where most of the species had a relatively low number of interactions and few  
 233 species concentrated most of them (Figure 3A). The distribution of prey among predators  
 234 showed that most consumers fed on a low number of prey whereas few had multiple prey  
 235 (Figure 3B). The top-five predators in number of prey were: yellowfin notothen  
 236 *Patagonotothen guntheri* (Notothenioid fish, 50 prey), rock cod *Patagonotothen ramsayi*  
 237 (Notothenioid fish, 49 prey), broad nose skate *Bathyraja brachyurops* (Chondrichthyan, 33  
 238 prey), Patagonian toothfish *Dissostichus eleginoides* (Notothenioid fish, 30 prey), and  
 239 graytail skate *Bathyraja griseocauda* (Chondrichthyan, 28 prey). Following the same  
 240 distribution pattern, few prey presented multiple predators (Figure 3C). The top-five prey (or

241 food sources) in number of predators were: Detritus (Non-living, 153 predators), the three  
 242 categories of Diatoms considered (benthic, centric and pennate, 72.5 predators on average),  
 243 and species of the genus *Euphausia* (Zooplankton, 46 predators). Finally, taking into account  
 244 the interactions within each functional group, most interactions were concentrated in a few  
 245 species (Figure 3D). The most evident species were: *Doryteuthis gahi* (Cephalopoda),  
 246 *Grimothea (=Munida) gregaria* (Decapoda), *P. ramsayi*, *P. guntheri* and *D. eleginoides*  
 247 (benthic-pelagic fish), *Sprattus fuegensis* and *Micromesistius australis* (pelagic fish), and  
 248 species of *Euphausia* and *Themisto gaudichaudii* (Zooplankton). Overall, there is an evident  
 249 asymmetry in the distribution of interactions among species at different levels in the MPA N-  
 250 BB food web.

251 A list of the distribution of interactions per species is presented in Supplementary Material  
 252 (Table S3).

### 253 3.2. SPECIES-LEVEL PROPERTIES

254 We found different relationships between the species trophic level (TL) and the rest of the  
 255 analysed species-level properties (Figure 4A-D). The most evident significant relationship  
 256 was with trophic similarity, i.e. the higher the species' TL, the lower the trophic similarity or  
 257 the higher the uniqueness in terms of trophic role (Figure 4C). Here it is noteworthy to  
 258 highlight those high-trophic level species ( $TL > 3.1$ ) with low values of trophic similarity:  
 259 *Bathyraja macloviana* and *Squalus acanthias* (Chondrichthyans), *Diplopteraster clarki* and  
 260 *Pteraster* sp. (echinoderms), *Daption capense* and *Eudyptes chrysocome* (seabirds), Ziphiidae  
 261 and *Lagenorhynchus cruciger* (marine mammals) (Table S3).

262 We also found a significant negative relationship between TL and closeness, however less  
 263 evident, meaning that low-TL species are relatively closer to any other species in the food  
 264 web (Figure 4B). Detritus, species of genera *Calanus* and *Euphausia*, and Foraminifera, all  
 265 with  $TL < 3$ , registered the highest closeness values (Table S3). Notably, species of mid-TLs

(3-4.2) showed the highest values of betweenness, meaning that those species participated in the highest number of shortest paths between species (Figure 4A). The following are the species with the highest values (descending order): *P. ramsayi*, *Salilota australis*, *D. eleginoides* (fishes), *D. gahi* (Cephalopoda), and *P. guntheri* (Notothenioid fish) (Table S3).

Considering the topological role, 'module specialist' species were the most frequent and presented a wide TL range (1 - 4.78), as well as 'module hub' species (TL = 1 - 3.92); 'module connector' was constrained to mid-TLs (2 - 3.86); and 'network connector', was represented by only one trophic species: detritus (Figure 4D, see Figure S2 for species' topological roles in a food web graph framework). Here it is important to highlight the two latter topological roles because they are responsible for linking modules and maintaining the connectivity of the food web: 42 species (1 network connector + 41 module connectors) from 19 different functional groups with a TL range = 1 - 3.86. The 41 species with a module connector role represented these functional groups: Amphipoda, Bivalvia, Brachiopoda, Bryozoa, Hydrozoa (as 'Cnidaria\_benthic'), Copepoda, Cumacea, Decapoda, Echinodermata, fish (benthic-pelagic and demersal Osteichthyes, and Chondrichthyes), Foraminifera, Polychaeta, Porifera, Pycnogonida (as 'Benthos\_Misc') and zooplankton (see Supplementary Material Table S3 for the identity of the species).

An exhaustive list of the species-level properties is presented in Supplementary Material (Table S3).

## 285 4. DISCUSSION

### 286 4.1. COMPLEXITY AND STRUCTURE OF THE FOOD WEB

The food web of the MPA N-BB ecosystem analysed in this study is one of the most highly-resolved networks of trophic interactions ever studied, not only for a high-latitude open-ocean ecosystem but also for any marine protected area worldwide to our knowledge. It is of paramount importance to consider the complexity of species interactions in order to gain

291 insights into the structure and functioning of the ecosystem, since the aggregation of species  
292 might mask food web properties and produce type II errors (false positives) (Martinez 1993,  
293 Gauzens et al. 2013).

294 Food web connectance is a feature that resumes the complexity of the network, but more  
295 importantly, it is an emergent property of pairwise species interactions (Poisot & Gravel  
296 2014). It contains information regarding how interactions within an ecological network are  
297 distributed and predicts reasonably well key dynamical properties of ecological networks  
298 (Dunne et al. 2002a). Complex marine food webs (i.e. with more than 25 trophic species)  
299 show connectance values ranging from 0.01 - 0.27 (Marina et al. 2018a). In particular, food  
300 webs from high-latitude regions tend to exhibit a connectance closer to the minimum  
301 (between 0.01 and 0.05) (Santana et al. 2013, Kortsch et al. 2015, Rodríguez et al. 2022).  
302 Whether food webs display a low or a high connectance helps to better comprehend  
303 ecosystem's synthetic properties like robustness. In this sense, empirical analyses support the  
304 notion that highly-connected ecological networks are robust against external perturbations  
305 such as the introduction of new (e.g., invasive) species (Smith-Ramesh et al. 2017) as well as  
306 species removal (e.g., local extinction) (Dunne et al. 2002b, Montoya & Solé 2003). The  
307 connectance of the food web of the MPA Namuncurá - Burdwood Bank (0.01) is one of the  
308 lowest reported so far for these regions; in particular, it appears to be much lower than that of  
309 Beagle Channel (0.05), an adjacent coastal area (Rodríguez et al. 2022).

310 The degree distribution, the distribution of the number of interactions per species, is the core  
311 of the structure of species interactions, which influences the opportunities for multiple  
312 species to persist in the long term and, therefore, their coexistence (Godoy et al. 2018). The  
313 food web for the MPA N-BB presents an asymmetric degree distribution. This pattern was  
314 identified at different levels of analysis: food web, predator, prey, and functional group. Such  
315 asymmetry is a well-known feature in empirical complex food webs in particular (Dunne et

al. 2002a, Montoya & Solé 2003, Stouffer et al. 2005), and has received great attention in complex networks in general (Albert & Barabási 2002, Newman 2003). The degree distribution affects the resilience of complex food webs against random failures and pressure on a particular component of the web: food webs showing right-skewed distributions, like the one described in this study, are more vulnerable to the removal of the most connected species or hubs, with the potential of producing secondary extinctions and a catastrophic fragmentation of the network (Albert et al. 2000, Dunne et al. 2002b, Eklöf & Ebenman 2006).

It is suggested that the small-world pattern, i.e., a network with short path length and high clustering coefficient, is not frequent in complex marine food webs, mainly due to a low clustering coefficient compared to random networks (Dunne et al. 2002c, Marina et al. 2018b). However, the food web of the MPA N-BB does display a small-world pattern. Consequences of this could be of great importance in recognizing species evolutionary paths and the vulnerability to perturbations (Montoya & Solé 2002). On the one hand, a short path length implies a rapid spread of an impact (e.g., contaminant, population fluctuation, local extinction) throughout the network but, at the same time, more potentially adaptive dynamics in the face of external perturbations (Montoya & Solé 2002, Williams et al. 2002). On the other hand, a high clustering coefficient indicates the formation of subnetworks composed only by the neighbours of particular species. This translates into a greater resistance of the network due to the confinement of perturbations mainly within subnetworks and not spreading between them (Kortsch et al. 2019, Heer et al. 2020]. Overall, a small-world topology provides ecological networks with greater resilience and resistance (Bornatowski et al. 2017, Dormann et al. 2017).

Omnivory acts as a buffer to changes as the ecosystem presents alternative energy pathways in the face of perturbations, i.e., reducing the risk of cascading extinctions following the

primary loss of species (Borrvall et al. 2000). Omnivores are species able to adapt faster and to a broader range of environmental conditions by changing their foraging habits to feed on the most abundant prey (Fagan 1997). Furthermore, omnivory can be analysed from the interaction point of view: theoretical studies have identified omnivorous interactions as a possible candidate for a keystone interaction, sensu Kadoya et al. 2018, highlighting the importance of omnivory in stabilizing food web dynamics (McCann & Hastings 1997, Neutel et al. 2002). The high proportion of omnivory in the food web of the MPA N-BB suggests that the network might be robust to variations in prey abundances, which could increase food web's persistence and stability (Stouffer & Bascompte 2010).

In summary, the food web of the MPA N-BB presents a combination of network properties that makes it unique in terms of network resolution, complexity, and structural pattern. All this suggests that the food web might be fragile to external perturbations targeting highly connected species, which in turn coincides to be commercially exploited species as fishes (Laptikhovsky et al. 2013, Martínez et al. 2015, Winter & Arkhipkin 2023). However, structural properties might provide resilience and resistance with the final outcome of a rearranged structure maintaining its functions.

#### 4.2. DOMINANT CONSUMERS AND FOOD SOURCES

The degree distribution allows identifying important species, such as potential keystone species (i.e. highly connected) (Solé & Montoya 2001, Dunne et al. 2002b), generalist/specialist species, and dominant food sources (Kondoh et al. 2010).

We have identified that most of the consumers in the food web of the MPA N-BB either have a narrow diet or are specialists, while few present a broad or generalist diet. The most evident generalist species are *Patagonotothen guntheri* (Covatti Ale et al. 2022), *Patagonotothen ramsayi* (Fischer et al. 2022), juveniles of *Dissostichus eleginoides* (Troccoli et al. 2020), *Bathyraja brachyurops* (Belleggia et al. 2008), and *Bathyraja griseocauda* (Belleggia et al.

2014), with more than 25 potential prey. Since these species present mid-trophic positions in the food web (with the exception of adults of *D. eleginoides* that are top predators), acting as predator and prey, they might be important links between lower and higher trophic levels. This result is in agreement with the sole analysis, using stable isotopes, that exists so far for the trophic structure of the MPA N-BB (Riccialdelli et al. 2020), and resembles other high-latitude marine systems of the Southwest Atlantic and Antarctic regions (Arkhipkin & Laptikhovsky 2013, Marina et al. 2018a). The importance of these particular generalist species also arises since they feed in the benthic and pelagic habitats (Troccoli et al. 2020, Covatti Ale et al. 2022, Fischer et al. 2022), linking these realms and contributing to the vertical carbon flow.

On the other hand, a low number of prey are consumed by many predators in the food web of the MPA N-BB. This suggests that there are dominant food sources on which most consumers depend and from where the ecosystem energy is being transferred to the upper trophic levels. The most demanded source we identified in this study (i.e. detritus) supports the abundant benthic community of filter-feeders (Schejter et al. 2016), components of the animal forest (Schejter et al. 2020), likely feeding on detritus that is constantly resuspended from the bottom (Martin & Flores-Melo 2021). Furthermore, we found that the second and third-most consumed prey were diatoms and species of *Euphausia*, respectively, which are essential sources for the diverse zooplankton community (Spinelli et al. 2020), mid-TL consumers like the Fuegian sprat *Sprattus fuegensis* (Padovani et al. 2021) and *P. ramsayi* (Fischer et al. 2022), and top predators such as the black-browed and grey-headed albatrosses (*Thalassarche melanophris* and *Thalassarche chrysostoma*, respectively) (Catry et al. 2004), and baleen whales (species of the genera *Balaenoptera* and *Eubalaena*) (Valenzuela et al. 2018).

#### 4.3. SPECIES' ROLE RELATED TO THEIR TROPHIC LEVEL



391 Describing species' roles in food webs provides a toolbox to assess the significance of  
 392 species in terms of community's functioning and overall stability (Thebault & Fontaine 2010,  
 393 Cirtwill et al. 2018). We used a range of descriptors to characterise the dynamic and  
 394 multifaceted nature of the species forming the MPA N-BB food web.

395 Closeness and betweenness are defined as "mesoscale" properties because they consider  
 396 direct and indirect interactions, therefore describing the focal species' ability to influence the  
 397 rest of the species of the food web (Lai et al. 2012). Closeness quantifies how many steps  
 398 away species *i* is from all other species in the food web, and is proportional to how rapidly  
 399 the indirect effects of the focal species can spread to other species in the network (Scotti &  
 400 Jordán 2010). In the food web of the MPA N-BB, low-TL consumers arise as important in  
 401 this regard: species of the zooplankton community, *Calanus* and *Euphausia*, *Zygochlamys*  
 402 *patagonica* (Bivalvia), and Brachiopoda. Any perturbation affecting these species, such as the  
 403 recently confirmed contaminants mercury (Fioramonti et al. 2022) and microplastics (Cossi et  
 404 al. 2021, Di Mauro et al. 2022), should be of concern since it might reach many other species  
 405 in the food web. Otherwise, betweenness measures the number of shortest paths between  
 406 species, providing information on the importance of species as "bridges" for energy transfer:  
 407 a species with high betweenness takes part in more food chains and therefore affects more  
 408 energy flows (Scotti & Jordán 2010). We have identified the longtail southern cod *P. ramsayi*  
 409 as the most important species in this sense. Moreover, in light of our analysis, other species  
 410 like the Patagonian toothfish *D. eleginoides* (juveniles), the Patagonian cod *Salilota australis*,  
 411 the yellowfin notothenioid *P. guntheri*, and the Patagonian longfin squid *Doryteuthis gahi*  
 412 should be considered as relevant in the energy transfer in the ecosystem. All these species  
 413 have a mid-trophic position in the food web, supporting the 'wasp-waist' control hypothesis  
 414 for the MPA N-BB (Riccialdelli et al. 2020).

415 Ecosystems with a pronounced 'wasp-waist' structure are suggested to present a high trophic

416 redundancy, since many species would show similar trophic habits (Cury et al. 2000). The  
 417 significant negative relationship between trophic similarity and trophic level enhances the  
 418 hypothesis of functional similarity at low and mid-TL species compared to higher TL species  
 419 for the MPA N-BB food web (Ricciardelli et al. 2020). At the same time, our results highlight  
 420 the uniqueness in terms of the trophic role of high-TL predators. Here, not only the expected  
 421 pelagic animals such as marine mammals and seabirds arise as relevant, but also demersal  
 422 vertebrate (chondrichthyans *Bathyrja macloviana* and *Squalus acanthias*) and benthic  
 423 invertebrate species (echinoderms *Diplopteraster clarki* and *Pteraster sp.*) are noteworthy.  
 424 The role that such species play in the MPA N-BB ecosystem is unique and perturbations on  
 425 them might result in unprecedented changes at the trophic structure and functioning level. In  
 426 this regard, we should mention the potential threat of the fisheries operating in the western  
 427 section of the MPA N-BB, where this activity is allowed and mostly focuses on the  
 428 Patagonian toothfish *D. eleginoides* and the southern blue whiting *Micromesistius australis*  
 429 (Martínez et al. 2015). Although the fishing effort is concentrated outside the limits of the  
 430 MPA N-BB, the impact on the MPA ecosystem should not be neglected (Martínez et al.  
 431 2021).

432 Species' role can also be assessed in a module-based context. Among the varying numbers of  
 433 topological roles in which species can be divided, two are remarkable: 'module connector'  
 434 and 'network connector'. Here, our results point out that there are several species, belonging  
 435 to a wide range of trophic positions (1 to 3.86) and representing 17 different functional  
 436 groups, that should be considered as influential species for the connectivity of the food web.  
 437 Thus, we propose that the diversity of species (benthic and pelagic) maintains the  
 438 connectivity of the food web, therefore contributing to the trophic structure and ecosystem's  
 439 stability.

#### 440 4.4. CAVEATS AND FUTURE PERSPECTIVE

441 The food web studied in the present work might be more representative of the shallow  
442 ecosystem of the submarine plateau called Burdwood Bank, on which most of the research  
443 was focused as the MPA N-BB I was first created. This is related to the sampling effort that  
444 was conducted during the research cruises in the former MPA compared to the MPA N-BB II  
445 (i.e. deep flanks to the south). As a consequence, most of the data we used to build the  
446 network come from studies performed in the MPA N-BB I. Despite this fact, we decided to  
447 build the food web considering both MPAs due to the tight oceanographic and ecological  
448 connection that exists among them (APN 2022 and references therein).

449 It's important to note that we did not consider quantitative data (i.e. abundance, biomass) to  
450 assess the species' role in the food web. Although there exists such data for some species  
451 (Schejter & Albano 2021), it would not be possible to include it in the food web framework  
452 described here due to a taxonomical resolution mismatch. In this regard, we should mention  
453 the case of *Zygochlamys patagonica* (Bivalvia) and Brachiopoda that are highlighted by our  
454 species-level analyses though they have been found in low abundances in the area (Schejter  
455 & Albano 2021).

456 Some species of sessile suspension feeders in high-latitude marine ecosystems, such as  
457 sponges, ascidians and octocorals, avoid predation by producing secondary metabolites that  
458 function as a chemical defense (Núñez-Pons et al. 2010, Moles et al. 2015, Prieto et al. 2022).  
459 Although this was not yet recorded at the MPA N-BB, there are a few studies that reported it  
460 in other locations in species that inhabit the MPA N-BB (Rojo de Almeida et al. 2010).

461 The MPA N-BB I presents complex oceanographic conditions that generate an internal  
462 spatial heterogeneity, mainly along its longitudinal axis (Matano et al. 2019). So far this  
463 heterogeneity has been reflected in phytoplankton and zooplankton communities (Bértola et  
464 al. 2018, García Alonso et al. 2020, Spinelli et al. 2020), and in fish assemblages (Delpiani et  
465 al. 2020). Moreover, seasonal variations also occur in some physical and biological aspects of

the MPA N-BB I (García Alonso et al. 2018, Matano et al. 2019). Considering both MPAs (N-BB I and II), a seasonal variation in the community composition of marine mammals and seabirds was recorded recently (Dellabianca et al. 2023). The spatial and seasonal variations in the plankton community might affect the energy and matter flow to higher levels of the food web. This has been recently studied in the vicinity of the MPA N-BB I, in the Beagle Channel, where a differential energy flow pattern of the plankton community has been recognised in two micro-basins of the Channel separated by a sill, each with different physicochemical properties (Giesecke et al. 2021), nutrient concentration (Latorre et al. 2023) as well as in the dominant component of the plankton community (Bruno et al. 2023, Presta et al. 2023). Although we were aware of the above, we decided to characterise a food web representing the whole MPA N-BB I year round since this is the first study of its type in the area.

Taking into account the mentioned caveats, and with the aim of improving the knowledge regarding the structure, functioning and stability of the MPA N-BB, we suggest that the future perspectives should: 1) incorporate spatial heterogeneity among MPA N-BB I and II (Schejter & Albano 2021), which might lead to distinct food web properties in terms of structure and functioning (Kortsch et al. 2019, Cordone et al. 2020); 2) include species traits, like body size and mass, since they are known to be important drivers in predator-prey interactions (Brose et al. 2019); 3) simulate the anthropogenic impacts already present in the MPA N-BB ecosystem (e.g. microplastics, mercury) (Cossi et al. 2021, Di Mauro et al. 2022, Fioramonti et al. 2022) as perturbations within the framework of the described complex food web; and 4) estimate the interaction strength of each predator-prey relationship in the food web considering species and interaction traits (i.e. body size, body mass, interaction dimensionality), and species density data (Nilsson & McCann 2016, Pawar et al. 2012).

## 5. CONCLUSION

491 We compiled information on the species and trophic diversity of the oceanic Marine  
492 Protected Area Namuncurá - Burdwood Bank, generating an unprecedented, well-resolved  
493 network of trophic interactions for a sub-Antarctic ecosystem, identifying the complexity and  
494 structure of the system, and the main species role in a network framework. Particular  
495 properties at the network level allowed us to identify the ecosystem's vulnerability and  
496 potential response to perturbations in the presence of highly-connected species, with a  
497 rearranged structure maintaining their functions due to its potential resilience and resistance.  
498 We identified several species as important regarding different aspects of trophic structure and  
499 functioning, and response to perturbations (i.e. environmental/anthropogenic changes). On  
500 the one hand, we suggest that generalist species, mainly fishes, play a crucial role in the  
501 ecosystem's benthic-pelagic coupling process. At the same time, we propose that other species  
502 besides the longtail southern cod *Patagonotothen ramsayi* and the Fuegian sprat *Sprattus*  
503 *fuegensis* should be considered relevant energy transfers for the ecosystem. Finally, we argue  
504 that it is the diversity of species, representing the benthic and pelagic habitats, that maintains  
505 the connectivity of the food web against perturbations, therefore contributing to the structure  
506 and stability of the ecosystem.

## 507 ACKNOWLEDGEMENTS

508 We are indebted to all those experts of the working group 'Banco Burdwood'  
509 who humbly provided their knowledge to enhance the quality of the present  
510 research. Although most of them are authors of the present work, it is worth  
511 to mention the following researchers: Brenda L. Doti (IBBEA, CONICET-UBA;  
512 Universidad de Buenos Aires, Argentina), Sofía L. Callá (Museo Argentino de

513 Ciencias Naturales “Bernardino Rivadavia”, Argentina), Sandra Gordillo  
 514 (IDACOR-CONICET; Universidad Nacional de Córdoba, Argentina), Mariano  
 515 I. Martinez (Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”,  
 516 Argentina) and Luciano Padovani (Instituto Nacional de Investigación y  
 517 Desarrollo Pesquero, INIDEP, Argentina). We thank the MPA Namuncurá –  
 518 Burdwood Bank administration. Research cruises were funded by national  
 519 funds by the Law 26.875. This study was funded by Consejo Nacional de  
 520 Investigaciones Científicas y Técnicas (CONICET) and Agencia Nacional de  
 521 Promoción Científica y Tecnológica (PICT 2020-SERIEA-01617), Argentina.  
 522 This work is contribution no. XX of the MPA Namuncurá (Law 26.875).

## 523 LITERATURE CITED

524 Acha EM, Mianzan HW, Guerrero RA, Favero M, Bava J (2004) Marine fronts at the  
 525 continental shelves of austral South America: Physical and ecological processes. *Journal of*  
 526 *Marine Systems* 44:83–105.  
 527 Administración de Parques Nacionales (2022) Plan de gestión AMP Namuncurá Banco  
 528 Burdwood.  
 529 Albert R, Barabási A-L (2002) Statistical mechanics of complex networks. *Reviews of*  
 530 *Modern Physics* 74:47–97.  
 531 Albert R, Jeong H, Barabási A-L (2000) Error and attack tolerance of complex networks.  
 532 *Nature* 406:378–382.  
 533 Arkhipkin A, Laptikhovsky V (2013) From gelatinous to muscle food chain: Rock cod  
 534 *Patagonotothen ramsayi* recycles coelenterate and tunicate resources on the Patagonian  
 535 Shelf. *Journal of Fish Biology* 83:1210–1220.  
 536 Bascompte J (2009) Disentangling the Web of Life. *Science* 325:416–419.  
 537 Belleggia M, Mabragaña E, Figueroa DE, Scenna LB, Barbini SA, Díaz de Astarloa JM,  
 538 (2008) Food habits of the broad nose skate, *Bathyraja brachyurops* (Chondrichthyes,  
 539 Rajidae), in the south-west Atlantic. *Scientia Marina* 72:701–710.  
 540 Belleggia M, Scenna L, Barbini SA, Figueroa DE, Díaz de Astarloa JM (2014) The diets of  
 541 four *Bathyraja* skates (Elasmobranchii, Arhynchobatidae) from the Southwest Atlantic.  
 542 *Cybium* 4:314–318.

- 543 Bértola G, Olguín Salinas H, Alder VA (2018) Distribución espacial de *Rhizosolenia crassa*,  
 544 ¿especie clave del banco burdwood? In: Alder V, Saraceno M, Capitanio F (eds) Libro de  
 545 Resúmenes X Jornadas Nacionales de Ciencias Del Mar, p 73.
- 546 Bornatowski H, Barreto R, Navia AF, de Amorim AF (2017) Topological redundancy and  
 547 “small world” patterns in a food web in a subtropical ecosystem of Brazil. *Marine Ecology*  
 548 38:e12407.
- 549 Borrvall C, Ebenman B, Jonsson TJ (2000) Biodiversity lessens the risk of cascading  
 550 extinction in model food webs. *Ecology Letters* 3:131–136.
- 551 Brose U, Archambault P, Barnes AD, Bersier L-F, Boy T, Canning-Clode J, Conti E, Dias M,  
 552 Digel C, Dissanayake A, Flores AAV, Fussmann K, Gauzens B, Gray C, Häussler J, Hirt  
 553 MR, Jacob U, Jochum M, Kéfi S, McLaughlin O, MacPherson MM, Latz E, Layer-Dobra K,  
 554 Legagneux P, Li Y, Madeira C, Martinez ND, Mendonça V, Mulder C, Navarrete SA,  
 555 O’Gorman EJ, Ott D, Paula J, Perkins D, Piechnik D, Pokrovsky I, Raffaelli D, Rall BC,  
 556 Rosenbaum B, Ryser R, Silva A, Sohlström EH, Sokolova N, Thompson MSA, Thompson  
 557 RM, Vermandele F, Vinagre C, Wang S, Wefer JM, Williams RJ, Wieters E, Woodward G,  
 558 Iles AC (2019) Predator traits determine food-web architecture across ecosystems. *Nature*  
 559 *Ecology & Evolution* 3:919–927.
- 560 Bruno DO, Valencia-Carrasco C, Paci MA, Leonarduzzi E, Castro L, Riccialdelli L, Iachetti  
 561 CM, Cadaillon A, Giesecke R, Schloss IR, Berghoff CF, Martín J, Diez M, Cabreira A,  
 562 Presta ML, Capitanio FL, Boy CC (2023) Spring plankton energy content by size classes in  
 563 two contrasting environments of a high latitude ecosystem: The Beagle Channel. *Journal of*  
 564 *Marine Systems* 240:103876.
- 565 Catry P, Phillips RA, Phalan B, Silk JRD, Croxall JP (2004) Foraging strategies of grey-  
 566 headed albatrosses *Thalassarche chrysostoma*: Integration of movements, activity and feeding  
 567 events. *Marine Ecology Progress Series* 280:261–273.
- 568 Cirtwill AR, Dalla Riva GV, Gaiarsa MP, Bimler MD, Cagua EF, Coux C, Dehling DM  
 569 (2018) A review of species role concepts in food webs. *Food Webs* 16:e00093.
- 570 Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) (2009)  
 571 Vulnerable Marine Ecosystem taxa identification guide.
- 572 Cordone G, Marina TI, Salinas V, Doyle SR, Saravia LA, Momo FR (2018) Effects of  
 573 macroalgae loss in an Antarctic marine food web: Applying extinction thresholds to food web  
 574 studies. *PeerJ* 6:e5531.
- 575 Cordone G, Salinas V, Marina TI, Doyle SR, Pasotti F, Saravia LA, Momo FR (2020) Green  
 576 vs brown food web: Effects of habitat type on multidimensional stability proxies for a highly-  
 577 resolved Antarctic food web. *Food Webs* 25:e00166.
- 578 Cossi PF, Ojeda M, Chiesa IL, Rimondino GN, Fraysse C, Calcagno J, Pérez AF (2021) First  
 579 evidence of microplastics in the Marine Protected Area Namuncurá at Burdwood Bank,  
 580 Argentina: A study on *Henricia obesa* and *Odontaster penicillatus* (Echinodermata:  
 581 Asteroidea). *Polar Biology* 44:2277–2287.
- 582 Covatti Ale M, Fischer L, Deli Antoni M, Díaz de Astarloa JM, Delpiani G (2022) Trophic  
 583 ecology of the yellowfin notothen, *Patagonotothen guntheri* (Norman, 1937) at the Marine  
 584 Protected Area Namuncurá-Burdwood Bank, Argentina. *Polar Biology* 45:549–558.
- 585 Csardi G, Nepusz T (2006) The igraph software package for complex network research.  
 586 *InterJournal, Complex Systems* 1695. <https://igraph.org>

- 587 Cury P, Bakun A, Crawford RJM, Jarre A, Quiñones RA, Shannon LJ, Verheye HM (2000)  
 588 Small pelagics in upwelling systems: Patterns of interaction and structural changes in “wasp-  
 589 waist” ecosystems. *ICES Journal of Marine Science* 57:603–618.
- 590 Day RH, Weingartner TJ, Hopcroft RR, Aerts LAM, Blanchard AL, Gall AE, Gallaway BJ,  
 591 Hannay DE, Holladay BA, Mathis JT, Norcross BL, Questel JM, Wisdom SS (2013) The  
 592 offshore northeastern Chukchi Sea, Alaska: A complex high-latitude ecosystem. *Continental*  
 593 *Shelf Research*, Seasonal and interannual dynamics of the northeastern Chukchi Sea  
 594 Ecosystem 67:147–165.
- 595 Dellabianca NA, Torres MA, Ordoñez C, Fioramonti N, Raya Rey A (2023) Marine  
 596 protected areas in the southern south-west Atlantic: Insights from marine top predator  
 597 communities. *Aquatic Conservation: Marine and Freshwater Ecosystems* 33:472–487.
- 598 Delpiani SM, Bruno DO, Vazquez DM, Llopart F, Delpiani GE, Fernández DA, Rosso JJ,  
 599 Mabrugaña E, Díaz de Astarloa JM (2020) Structure and distribution of fish assemblages at  
 600 Burdwood Bank, the first Sub-Antarctic Marine Protected Area “Namuncurá” in Argentina  
 601 (Southwestern Atlantic Ocean). *Polar Biology* 43:1783–1793.
- 602 Di Mauro R, Castillo S, Pérez A, Iachetti CM, Silva L, Tomba JP, Chiesa IL (2022)  
 603 Anthropogenic microfibers are highly abundant at the Burdwood Bank seamount, a protected  
 604 sub-Antarctic environment in the Southwestern Atlantic Ocean. *Environmental Pollution*  
 605 306:119364.
- 606 Dormann CF, Fründ J, Schaefer HM (2017) Identifying Causes of Patterns in Ecological  
 607 Networks: Opportunities and Limitations. *Annual Review of Ecology, Evolution, and*  
 608 *Systematics* 48:559–584.
- 609 Dunne JA, Williams RJ, Martinez ND (2002a) Food-web structure and network theory: The  
 610 role of connectance and size. *Proceedings of the National Academy of Sciences* 99:12917–  
 611 12922.
- 612 Dunne JA, Williams RJ, Martinez ND (2002b) Network structure and biodiversity loss in  
 613 food webs: Robustness increases with connectance. *Ecology Letters* 5:558–567.
- 614 Dunne JA, Williams RJ, Martinez ND (2002c) Small Networks but not Small Worlds:  
 615 Unique Aspects of Food Web Structure. Santa Fe Institute Working paper.
- 616 Eklöf A, Ebenman B (2006) Species loss and secondary extinctions in simple and complex  
 617 model communities. *Journal of Animal Ecology* 75:239–246.
- 618 Fagan WF (1997) Omnivory as a Stabilizing Feature of Natural Communities. *The American*  
 619 *Naturalist* 150:554–567.
- 620 Falabella V (2017) (Ed) Área Marina Protegida Namuncurá-Banco Burdwood.  
 621 Contribuciones para la línea de base y el plan de manejo. Jefatura de Gabinete de Ministros.  
 622 Buenos Aires, Argentina.
- 623 Fioramonti NE, Ribeiro Guevara S, Becker YA, Riccialdelli L (2022) Mercury transfer in  
 624 coastal and oceanic food webs from the Southwest Atlantic Ocean. *Marine Pollution Bulletin*  
 625 175:113365.
- 626 Fischer L, Covatti Ale M, Deli Antoni M, Díaz de Astarloa JM, Delpiani G (2022) Feeding  
 627 ecology of the longtail southern cod, *Patagonotothen ramsayi* (Regan, 1913)  
 628 (Notothenioidei) in the Marine Protected Area Namuncurá-Burdwood Bank, Argentina. *Polar*  
 629 *Biology* 45:1483–1494.
- 630 Matusevich F, Vazquez DM, Gabbanelli V, Díaz de Astarloa JM, Mabrugaña E (2023)



- Chondrichthyans from the southern tip of South America with emphasis on the marine protected area Namuncurá-Burdwood Bank: Exploring egg nursery grounds. *Polar Biology* 46:357–372.
- Funes M, Saravia LA, Cordone G, Iribarne OO, Galván DE (2022) Network analysis suggests changes in food web stability produced by bottom trawl fishery in Patagonia. *Scientific Reports* 12:10876.
- García Alonso VA, Brown D, Martín J, Pájaro M, Capitanio FL (2018) Seasonal patterns of Patagonian sprat *Sprattus fuegensis* early life stages in an open sea Sub-Antarctic Marine Protected Area. *Polar Biology* 41:2167–2179.
- García Alonso VA, Brown DR, Pájaro M, Capitanio FL (2020) Growing Up Down South: Spatial and Temporal Variability in Early Growth of Fuegian Sprat *Sprattus fuegensis* From the Southwest Atlantic Ocean. *Frontiers in Marine Science* 7:322.
- Gauzens B, Legendre S, Lazzaro X, Lacroix G (2013) Food-web aggregation, methodological and functional issues. *Oikos* 122:1606–1615.
- Giesecke R, Martín J, Piñones A, Höfer J, Garcés-Vargas J, Flores-Melo X, Alarcón E, Durrieu de Madron X, Bourrin F, González HE (2021) General Hydrography of the Beagle Channel, a Subantarctic Interoceanic Passage at the Southern Tip of South America. *Frontiers in Marine Science* 8:621822.
- Glorioso PD, Flather RA (1995) A barotropic model of the currents off SE South America. *Journal of Geophysical Research: Oceans* 100:13427–13440.
- Godoy O, Bartomeus I, Rohr RP, Saavedra S (2018) Towards the Integration of Niche and Network Theories. *Trends in Ecology & Evolution* 33:287–300.
- Guerrero RA, Baldoni AG, Benavides HR (1999) Oceanographic conditions at the southern end of the argentine continental slope. *INIDEP Documento Científico* 5:7-22.
- Guimerà R, Nunes Amaral LA (2005) Functional cartography of complex metabolic networks. *Nature* 433:895–900.
- Guinder VA, Malits A, Ferronato C, Krock B, Garzón-Cardona J, Martínez A (2020) Microbial plankton configuration in the epipelagic realm from the Beagle Channel to the Burdwood Bank, a Marine Protected Area in Sub-Antarctic waters. *PLOS ONE* 15:e0233156.
- Heer H, Streib L, Schäfer RB, Ruzika S (2020) Maximising the clustering coefficient of networks and the effects on habitat network robustness. *PLOS ONE* 15:e0240940.
- Hogg OT, Huvenne VAI, Griffiths HJ, Dorschel B, Linse K (2016) Landscape mapping at sub-Antarctic South Georgia provides a protocol for underpinning large-scale marine protected areas. *Scientific Reports* 6:33163.
- IUCN (2023) The world database on protected areas (WDPA). <https://www.protectedplanet.net/en/thematic-areas/wdpa?tab=WDPA> (accessed 14 July 2023).
- Kadoya T, Gellner G, McCann KS (2018) Potential oscillators and keystone modules in food webs. *Ecology Letters* 21:1330–1340.
- Kondoh M, Kato S, Sakato Y (2010) Food webs are built up with nested subwebs. *Ecology* 91:3123–561 3130.
- Kortsch S, Primicerio R, Aschan M, Lind S, Dolgov AV, Planque B (2019) Food-web structure varies along environmental gradients in a high-latitude marine ecosystem. *Ecography* 42:295–308.

- 675 Kortsch S, Primicerio R, Fossheim M, Dolgov AV, Aschan M (2015) Climate change alters  
 676 the structure of arctic marine food webs due to poleward shifts of boreal generalists.  
 677 Proceedings of the Royal Society B: Biological Sciences 282:20151546.
- 678 Lai S-M, Liu W-C, Jordán F (2012) On the centrality and uniqueness of species from the  
 679 network perspective. Biology Letters 8:570–573.
- 680 Laptikhovsky V, Arkhipkin A, Brickle P (2013) From small bycatch to main commercial  
 681 species: Explosion of stocks of rock cod *Patagonotothen ramsayi* (Regan) in the Southwest  
 682 Atlantic. Fisheries Research 147:399–403.
- 683 Latorre MP, Berghoff CF, Giesecke R, Malits A, Pizarro G, Iachetti CM, Martin J, Flores-  
 684 Melo X, Gil MN, Iriarte JL, Schloss IR (2023) Plankton metabolic balance in the eastern  
 685 Beagle Channel during spring. Journal of Marine Systems 240:103882.
- 686 López-Gappa J, Liuzzi MG, Zelaya DG (2018) A new genus and species of cheilostome  
 687 bryozoan associated with hermit crabs in the subantarctic Southwest Atlantic. Polar Biology  
 688 41:733–741.
- 689 Marina TI, Salinas V, Cordone G, Campana G, Moreira E, Deregibus D, Torre L, Sahade R,  
 690 Tatián M, Barrera Oro E, De Troch M, Doyle S, Quartino ML, Saravia LA, Momo FR  
 691 (2018a) The Food Web of Potter Cove (Antarctica): Complexity, structure and function.  
 692 Estuarine, Coastal and Shelf Science 200:141–151.
- 693 Marina TI, Saravia LA, Cordone G, Salinas V, Doyle SR, Momo FR (2018b) Architecture of  
 694 marine food webs: To be or not be a “small-world”. PLOS ONE 13:e0198217.
- 695 Marina TI, Saravia LA, Kortsch S (2023) New insights into the Weddell Sea ecosystem  
 696 applying a quantitative network approach. In: Workshop on Species interactions in the  
 697 Southern Ocean. Ghent, Belgium.
- 698 Martin J, Flores-Melo X (2021) Área Marina Protegida Namuncurá Banco Burdwood:  
 699 Aspectos físicos y biogeoquímicos. In: Lovrich G (eds) Taller científico sobre el Área Marina  
 700 Protegida Namuncurá - Banco Burdwood. p 1-2.
- 701 Martin Srito S (2019) Fauna asociada a corales (Octocorallia) e hidroides (Hydrozoa) del  
 702 Área Marina Protegida “Namuncurá” (Banco Burdwood) y zonas profundas adyacentes. PhD  
 703 dissertation, Universidad Nacional de Mar del Plata, Mar del Plata, Argentina.
- 704 Martinez ND (1991) Artifacts or Attributes? Effects of Resolution on the Little Rock Lake  
 705 Food Web. Ecological Monographs 61:367–392.
- 706 Martinez ND (1993) Effects of Resolution on Food Web Structure. Oikos 66:403–412.
- 707 Martínez PA, Wöhler OC, Troccoli GH, Di Marco EJ (2021) Análisis del impacto potencial  
 708 provocado por el establecimiento de las áreas marinas protegidas Namuncurá-Banco  
 709 Burdwood I, II y Yaganes en la pesquería argentina de merluza negra (*Dissostichus*  
 710 *eleginoides*). Technical report No. 23/2021. Instituto Nacional de Investigación y Desarrollo  
 711 Pesquero (INIDEP), Argentina.
- 712 Martínez PA, Wöhler OG, Troccoli GH (2015) La evolución de la pesquería de merluza  
 713 negra (*Dissostichus eleginoides*) en el espacio marítimo argentino. Periodo 2003- 2014.  
 714 Technical report No. 11/15. Instituto Nacional de Investigación y Desarrollo Pesquero  
 715 (INIDEP), Argentina.
- 716 Matano RP, Palma ED, Combes V (2019) The Burdwood Bank Circulation. Journal of  
 717 Geophysical Research: Oceans 124:6904–6926.
- 718 McCann K, Hastings A (1997) Re-evaluating the omnivory–stability relationship in food

- 719 webs. Proceedings of the Royal Society of London. Series B: Biological Sciences 264:1249–  
720 1254.
- 721 Moles J, Núñez-Pons L, Taboada S, Figuerola B, Cristobo J, Avila C (2015) Anti-predatory  
722 chemical defenses in Antarctic benthic fauna. Marine Biology 162:1813–1821.
- 723 Montoya JM, Solé RV (2002) Small World Patterns in Food Webs. Journal of Theoretical  
724 Biology 214:405–412.
- 725 Montoya JM, Solé RV (2003) Topological properties of food webs: From real data to  
726 community assembly models. Oikos 102:614–622.
- 727 Morales-Castilla I, Matias MG, Gravel D, Araújo MB (2015) Inferring biotic interactions  
728 from proxies. Trends in Ecology & Evolution 30:347–356.
- 729 Neutel A-M, Heesterbeek JAP, de Ruiter PC (2002) Stability in Real Food Webs: Weak  
730 Links in Long Loops. Science 296:1120–1123.
- 731 Newman MEJ (2003) The Structure and Function of Complex Networks. SIAM Review  
732 45:167–256.
- 733 Nilsson KA, McCann KS (2016) Interaction strength revisited—clarifying the role of energy  
734 flux for food web stability. Theoretical Ecology 9:59–71.
- 735 Núñez-Pons L, Forestieri R, Nieto RM, Varela M, Nappo M, Rodríguez J, Jiménez C,  
736 Castelluccio F, Carbone M, Ramos-Espla A, Gavagnin M, Avila C (2010) Chemical defenses  
737 of tunicates of the genus Aplidium from the Weddell Sea (Antarctica). Polar Biology  
738 33:1319–1329.
- 739 Padovani LN, Álvarez N, Farías A (2021) Alimentación de la sardina fueguina (*Sprattus*  
740 *fuegensis*) en la región patagónica austral durante la época reproductiva. Technical report.  
741 Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), Argentina.
- 742 Padovani LN, Viñas MD, Sánchez F, Mianzan H (2012) Amphipod-supported food web:  
743 *Themisto gaudichaudii*, a key food resource for fishes in the southern Patagonian Shelf.  
744 Journal of Sea Research 67:85–90.
- 745 Pascual M, Dunne JA (2005) Ecological Networks: Linking Structure to Dynamics in Food  
746 Webs. Oxford University Press.
- 747 Pawar S, Dell AI, Savage VM (2012) Dimensionality of consumer search space drives  
748 trophic interaction strengths. Nature 486:485–489.
- 749 Piola AR, Falabella V (2009) El mar Patagónico. In: Falabella V, Campagna C, Croxall J  
750 (eds) Atlas del Mar Patagónico. Especies y Espacios. Wildlife Conservation Society y  
751 Birdlife Internacional, Buenos Aires, Argentina, p 54–75.
- 752 Piola AR, Gordon AL (1989) Intermediate waters in the southwest South Atlantic. Deep Sea  
753 Research Part A. Oceanographic Research Papers 36:1–16.
- 754 Poisot T, Gravel D (2014) When is an ecological network complex? Connectance drives  
755 degree distribution and emerging network properties. PeerJ 2:e251.
- 756 Pomeranz JPF, Thompson RM, Poisot T, Harding JS (2019) Inferring predator–prey  
757 interactions in food webs. Methods in Ecology and Evolution 10:356–367.
- 758 Presta ML, Riccialdelli L, Bruno DO, Castro LR, Fioramonti NE, Florentín OV, Berghoff  
759 CF, Capitano FL, Lovrich GA (2023) Mesozooplankton community structure and trophic  
760 relationships in an austral high-latitude ecosystem (Beagle Channel): The role of bottom-up  
761 and top-down forces during springtime. Journal of Marine Systems 240:103881.
- 762 Prieto IM, Paola A, Pérez M, García M, Blustein G, Schejter L, Palermo JA (2022)

- Antifouling Diterpenoids from the Sponge *Dendrilla Antarctica*. *Chemistry & Biodiversity* 19:e202100618.
- Reta R (2014) Oceanografía del Banco Burdwood: Estado Actual del Conocimiento y Perspectivas. Technical report No. 15. Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), Argentina.
- Riccialdelli L, Becker YA, Fioramonti NE, Torres M, Bruno DO, Raya Rey A, Fernández DA (2020) Trophic structure of southern marine ecosystems: A comparative isotopic analysis from the Beagle Channel to the oceanic Burdwood Bank area under a wasp-waist assumption. *Marine Ecology Progress Series* 655:1–27.
- Roberts CM, O’Leary BC, McCauley DJ, Cury PM, Duarte CM, Lubchenco J, Pauly D, Sáenz-Arroyo A, Sumaila UR, Wilson RW, Worm B, Castilla JC (2017) Marine reserves can mitigate and promote adaptation to climate change. *Proceedings of the National Academy of Sciences* 114:6167–6175.
- Rodríguez ID, Marina TI, Schloss IR, Saravia LA (2022) Marine food webs are more complex but less stable in sub-Antarctic (Beagle Channel, Argentina) than in Antarctic (Potter Cove, Antarctic Peninsula) regions. *Marine Environmental Research* 174:105561.
- Rojo de Almeida MT, Siless GE, Perez CD, Veloso MJ, Schejter L, Puricelli L, Palermo JA (2010) Dolabellane Diterpenoids from the South Atlantic Gorgonian *Convexella Magelhaenica*. *Journal of Natural Products* 73:1714–1717.
- Sala E, Lubchenco J, Grorud-Colvert K, Novelli C, Roberts C, Sumaila UR (2018) Assessing real progress towards effective ocean protection. *Marine Policy* 91:11–13.
- Santana de CN, Rozenfeld AF, Marquet PA, Duarte CM (2013) Topological properties of polar food webs. *Marine Ecology Progress Series* 474:15–26.
- Saporiti F, Bearhop S, Vales DG, Silva L, Zenteno L, Tavares M, Crespo EA, Cardona L (2015) Latitudinal changes in the structure of marine food webs in the Southwestern Atlantic Ocean. *Marine Ecology Progress Series* 538:23–34.
- Saravia LA (2022) Multiweb: R package for Multiple Interactions Ecological Networks. Zenodo, <https://doi.org/10.5281/zenodo.3370396>
- Schejter L, Albano M (2021) Benthic communities at the marine protected area Namuncurá/Burdwood bank, SW Atlantic Ocean: Detection of vulnerable marine ecosystems and contributions to the assessment of the rezoning process. *Polar Biology* 44:2023–2037.
- Schejter L, Bremec CS (2019) Stony corals (Anthozoa: Scleractinia) of Burdwood Bank and neighbouring areas, SW Atlantic Ocean. *Scientia Marina* 83:247–260.
- Schejter L, Genzano G, Gaitán E, Perez CD, Bremec CS (2020) Benthic communities in the Southwest Atlantic Ocean: Conservation value of animal forests at the Burdwood Bank slope. *Aquatic Conservation: Marine and Freshwater Ecosystems* 30:426–439.
- Schejter L, Rimondino C, Chiesa I, Díaz de Astarloa JM, Doti B, Elías R, Escolar M, Genzano G, López-Gappa J, Tatián M, Zelaya DG, Cristobo J, Perez CD, Cordeiro RT, Bremec CS (2016) Namuncurá Marine Protected Area: An oceanic hot spot of benthic biodiversity at Burdwood Bank, Argentina. *Polar Biology* 39:2373–2386.
- Scotti M, Jordán F (2010) Relationships between centrality indices and trophic levels in food webs. *Community Ecology* 11:59–67.
- Secretariat of the Convention on Biological Diversity (2004) Technical advice on the establishment and management of a national system of marine and coastal protected areas.

- 807 Technical Report No. 13. p 40.
- 808 Smith-Ramesh LM, Moore AC, Schmitz OJ (2017) Global synthesis suggests that food web  
809 connectance correlates to invasion resistance. *Global Change Biology* 23:465–473.
- 810 Solé RV, Montoya M (2001) Complexity and fragility in ecological networks. *Proceedings of*  
811 *the Royal Society of London. Series B: Biological Sciences* 268:2039–2045.
- 812 Spinelli ML, Malits A, García Alonso VA, Martín J, Capitanio FL (2020) Spatial gradients of  
813 spring zooplankton assemblages at the open ocean sub-Antarctic Namuncurá Marine  
814 Protected Area/Burdwood Bank, SW Atlantic Ocean. *Journal of Marine Systems*  
815 210:103398.
- 816 Stouffer DB, Bascompte J (2010) Understanding food-web persistence from local to global  
817 scales. *Ecology Letters* 13:154–161.
- 818 Stouffer DB, Camacho J, Guimerà R, Ng CA, Nunes Amaral LA (2005) Quantitative Patterns  
819 in the Structure of Model and Empirical Food Webs. *Ecology* 86:1301–1311.
- 820 R Core Team (2022) R: A Language and Environment for Statistical Computing.
- 821 Thébault E, Fontaine C (2010) Stability of Ecological Communities and the Architecture of  
822 Mutualistic and Trophic Networks. *Science* 329:853–856.
- 823 Tombesi ML, Rabuffetti F, Lovrich GA (2020) Las áreas marinas protegidas en la Argentina.  
824 La Lupa Colección Fuegoína De divulgación científica 16:2–7.
- 825 Trathan PN, Fielding S, Hollyman PR, Murphy EJ, Warwick-Evans V, Collins MA (2021)  
826 Enhancing the ecosystem approach for the fishery for Antarctic krill within the complex,  
827 variable, and changing ecosystem at South Georgia. *ICES Journal of Marine Science*  
828 78:2065–2081.
- 829 Troccoli GH, Aguilar E, Martínez PA, Belleggia M (2020) The diet of the Patagonian  
830 toothfish *Dissostichus eleginoides*, a deep-sea top predator off Southwest Atlantic Ocean.  
831 *Polar Biology* 43:1595–1604.
- 832 Valenzuela LO, Rowntree VJ, Sironi M, Seger J (2018) Stable isotopes ( $\delta^{15}\text{N}$ ,  $\delta^{13}\text{C}$ ,  $\delta^{34}\text{S}$ )  
833 in skin reveal diverse food sources used by southern right whales *Eubalaena australis*.  
834 *Marine Ecology Progress Series* 603:243–255.
- 835 Vazquez DM, Belleggia M, Schejter L, Mabragaña E (2018) Avoiding being dragged away:  
836 Finding egg cases of *Schroederichthys bivi* (Chondrichthyes: Scyliorhinidae) associated  
837 with benthic invertebrates. *Journal of Fish Biology* 92:248–253.
- 838 Watts DJ, Strogatz SH (1998) Collective dynamics of “small-world” networks. *Nature*  
839 393:440–442.
- 840 Williams RJ, Berlow EL, Dunne JA, Barabási A-L, Martinez ND (2002) Two degrees of  
841 separation in complex food webs. *Proceedings of the National Academy of Sciences*  
842 99:12913–12916.
- 843 Winter A, Arkhipkin A (2023) Opportunistic Survey Analyses Reveal a Recent Decline of  
844 Skate (Rajiformes) Biomass in Falkland Islands Waters. *Fishes* 8:24.

## TROPHIC INTERACTIONS IN A MARINE PROTECTED AREA

Table 1: List of network and species-level properties analysed, definitions, and relevant ecological implications related to food web complexity and structure.

Name	Definition	Implications	Reference
Number of species	Number of trophic species in a food web.	It represents the species diversity and has implications for the persistence of the ecosystem.	May 1973, Tilman 1996
Number of interactions	Total number of trophic interactions in a food web.	It represents the number of pathways along which matter and energy can flow.	Dunne et al. 2002a
Link density	Ratio of interactions to species in a food web.	It represents the average number of interactions per species; informs about how connected species are in the food-web.	Dunne et al. 2002a
Connectance	Proportion of potential links among species that are actually realized. Range = 0 - 1.	It measures the probability of interactions and is a fundamental measure of network complexity. Connectance can be negatively or positively associated with food web robustness, depending on the network structure (random vs non-random) or how the strength of the interactions are distributed.	Martinez 1992
Degree distribution	Frequency of trophic species that have k or more interactions.	It suggests on the vulnerability of complex food webs against random failures and intentional attacks (i.e. species extinctions).	Albert & Barabási 2002
Omnivory	Species feeding on prey from more than one trophic level.	It influences food web's stability; intermediate levels of omnivory may stabilize it and may diffuse top-down effects thus reduce the probability of trophic cascades.	McCann & Hastings 1997
Small-world pattern	A network with short path length (distance between nodes) and high clustering coefficient (formation of compartments) compared to random networks.	Consequences of this structural pattern in food webs are of great importance in recognizing evolutionary paths and the vulnerability to perturbations.	Watts & Strogatz 1998, Montoya & Solé 2002
Betweenness	Number of shortest paths going through a species.	Species with high betweenness act as "bridges"; if removed, would have rapidly spreading effects in the food web.	Freeman 1978, Lai et al. 2012
Closeness	Number of steps required to reach every other species from a	The removal of a species with high closeness will affect the most other	Freeman 1978, Lai et al. 2012

## TROPHIC INTERACTIONS IN A MARINE PROTECTED AREA

	given species.	species in the food web.	
Trophic similarity	Trophic overlap based on shared and unique resources (prey) and consumers (predators).	It measures one of the most important aspects of species' niches, the trophic niche, and functional aspects of biodiversity.	Martinez 1992
Topological role	Species role according to interactions within and across modules (subgroups of species).	Four roles are defined: module hub, module specialist, module connector and network connector. Network connector and module connector roles maintain the connectivity of the food web.	Guimerà & Nunes Amaral 2005

Table 2: Network-level properties of the MPA Namuncurá - Burdwood Bank food web. CPL: Characteristic Path Length; CC: Clustering Coefficient; SW: Small-World pattern. See table 1 for definitions and ecological relevance.

Species	Interactions	Density	Connectance	Omnivory	CPL	CC	SW
379	1788	4.72	0.01	0.48	2.99	0.08	TRUE



## TROPHIC INTERACTIONS IN A MARINE PROTECTED AREA




Figure 1: Marine Protected Areas Namuncurá - Burdwood Bank I (MNR and MNP, northern section) and II (MNR and SMNR, southern section). Acronyms indicate categories according to the management plan: MNR - Marine National Reserve, MNP - Marine National Park and RMNR - Restricted Marine National Reserve.




Figure 2: Graph of the food web for the MPA Namuncurá - Burdwood Bank. Circles represent species and arrows trophic interactions. Circle diameter is relative to the number of interactions. Colour gradient indicates the trophic level.



# TROPHIC INTERACTIONS IN A MARINE PROTECTED AREA




Figure 3: Degree distributions for the (A) food web, for (B) prey among predators, (C) predators among prey, and (D) for each functional group. Groups are vertically ordered by increasing trophic level (following coloration of figure 2); groups with less than 3 species

were not plotted (e.g., pelagic fish). All functional groups and the species that comprise them are shown in Supplementary Material (Table S3).




Figure 4: Species-level properties by trophic level: (A) betweenness, (B) closeness, (C) trophic similarity, and (D) topological role. Each point represents a species. Linear regressions for betweenness ( $y = 74.97x - 117.35$ ,  $R^2 = 0.05$ ,  $p$ -value  $< 0.01$ ), closeness ( $y = 9.33e - 06x - 9.31e - 4$ ,  $R^2 = 0.003$ ,  $p$ -value = 0.15) and trophic similarity ( $y = -0.02x + 0.11$ ,  $R^2 = 0.07$ ,  $p$ -value  $< 0.01$ ). Note that for panels A, B and C only species with TLs equal or greater than 2 were considered.