

Measures of Variability: Takeaways

by Dataquest Labs, Inc. - All rights reserved © 2024

Syntax

- Writing a function that returns the range of an array:

```
def find_range(array):  
    return max(array) - min(array)
```

- Writing a function that returns the mean absolute deviation of an array:

```
def mean_absolute_deviation(array):  
    reference_point = sum(array) / len(array)  
    distances = []  
    for value in array:  
        absolute_distance = abs(value - reference_point)  
        distances.append(absolute_distance)  
    return sum(distances) / len(distances)
```

FINDING THE VARIANCE OF AN ARRAY

- If the the array is a series object:

```
sample_variance = Series.var(ddof=1)  
population_variance = Series.var(ddof=0)
```

- If the array is not a series object:

```
from numpy import var  
sample_variance = var(a_sample, ddof=1)  
population_variance = var(a_population, ddof=0)
```

FINDING THE STANDARD DEVIATION OF AN ARRAY

- If the array is a series object:

```
sample_stdev = Series.std(ddof=1)  
population_stdev = Series.std(ddof=0)
```

- If the array is not a series object

```
from numpy import std  
sample_stdev = std(a_sample, ddof=1)  
population_stdev = std(a_population, ddof=0)
```

Concepts

- There are many ways we can measure the **variability** of a distribution. These are some of the measures we can use:
 - **The range.**
 - **The mean absolute deviation.**

- **The variance.**
- **The standard deviation.**

- Variance and standard deviation are the most used metrics to measure variability. To compute the standard deviation σ and the variance σ^2 for a **population**, we can use the formulas:

$$\sigma =$$

$$\sqrt{\frac{\sum_{i=1}^N (x_i - \mu)^2}{N}}$$

$$\sigma^2 = \frac{\sum_{i=1}^N (x_i - \mu)^2}{N}$$

- To compute the standard deviation s and the variance s^2 for a **sample**, we need to add the **Bessel's correction** to the formulas above:

$$s =$$

$$\sqrt{\frac{\sum_{i=1}^n (x_i - \mu)^2}{n-1}}$$

$$s^2 = \frac{\sum_{i=1}^n (x_i - \mu)^2}{n-1}$$

- **Sample variance** s^2 is the only unbiased estimator we learned about, and it's unbiased only when we sample with replacement.

Resources

- [An intuitive introduction to variance and standard deviation.](#)
- Useful documentation:
 - `numpy.var()`
 - `numpy.std()`
 - `Series.var()`
 - `Series.std()`