

Meta-Awareness of Dysregulated Emotional Attention

Ruimi, Liad, M.A.

Hadash, Yuval, M.A.

Zvielli, Ariel, Ph.D..

Amir, Iftach, M.A.

Goldstein, Pavel, Ph.D.

Bernstein, Amit, Ph.D. *

University of Haifa

[In Press, *Clinical Psychological Science*]

Author Note:

Correspondence regarding study should be sent to Amit Bernstein, PhD; Director, The International Research Collaborative on Anxiety. Dr. Bernstein can be contacted at the University of Haifa, Department of Psychology, Mount Carmel, Haifa, 31905, Israel, 972-4-828-8863 (phone), 972-4-824-0966 (facsimile). Electronic mail: abernstein@psy.haifa.ac.il. Dr. Bernstein recognizes the funding support from the Israel Science Foundation and the Caesarea-Rothschild Foundation. Mr. Ruimi recognizes the support from the University of Haifa President's Doctoral Fellowship Program. We want to thank research assistants, Noam Asherovich and Elicar Carsenti for their help in carrying out the study.

Abstract

We explore the human capacity for, and the function(s) of, meta-awareness for biased attentional processing of emotional information (MAB) sub-serving mental (ill)health. We do so by integrating probe-caught sampling methods, signal detection theory, and multi-level modeling of cognitive-experimental laboratory data among daily smokers (N=75) known to exhibit biased attentional processing of reward-related (drug) cues in addiction. We found (a) evidence of the capacity for and individual differences in MAB; (b) momentary MAB was most likely observed in the event of the most extreme micro-expressions of biased attentional processing; and (c) momentary micro-expressions of biased attention *without* MAB were more likely followed by attentional dysregulation, whereas momentary micro-expressions of biased attention *with* MAB were more likely followed by more balanced attentional expression or greater attentional control. We discuss the implications for basic and clinical science of meta-awareness.

Key Words: Attentional Bias; Cognitive Control; Emotional Attention; Meta-Awareness; Smoking

Meta-Awareness of Dysregulated Emotional Attention

By selecting from a sea of potential environmental inputs in a manner that is context-sensitive, flexible, and goal-relevant, attention enables adaptive pursuit of reward and escape from threat (Vuilleumier, 2005). Accordingly, dysregulation in attentional processing of emotional information, commonly referred to as attentional bias (AB), has been implicated in the etiology and maintenance of prevalent forms of maladaptation including, anxiety, depression and addiction; and thus, is a promising intervention target (Heeren, De Raedt, Koster, & Philippot, 2013; Van Bockstaele et al., 2014). Yet the field has faced growing concerns about the weak psychometric properties of cognitive-experimental measures of AB, mixed evidence of the role of AB in psychopathology, and mixed evidence of efforts to experimentally and clinically modify AB (Bar-Haim et al., 2007; Schmukle, 2005; Van Bockstaele et al., 2014).

Accordingly, Zvielli et al (2015) questioned the decades-old paradigm conceptualizing and studying AB as a *static* trait. They proposed that AB may be better conceptualized, studied and ultimately modified as a *dynamic process* that fluctuates toward *and* away from motivationally-relevant stimuli from moment-to-moment in time (Zvielli, Bernstein, & Koster, 2014; Zvielli et al., 2015) (see Figure 1). To begin to do so, they proposed a novel computational procedures designed to quantify and thereby study dynamic fluctuations of biased attention (e.g., Trial Level Bias Scores (TL-BS)) in extant cognitive-experimental tasks of emotional attention (e.g., dot probe, spatial cueing, visual search). They also called for the development of novel measurement methods designed to capture temporal dynamics of emotional attention.

Recent studies have demonstrated convergent, incremental, known-group criterion, and predictive validity of the dynamic features of AB. This evidence of validity has been

reported in multiple tasks (e.g., dot probe task, spatial cueing task) and with respect to a variety of important outcomes, including: spider phobia, addiction behavior (e.g., smoking rate), social anxiety, depression vulnerability (e.g., remitted depression status, depressive episode history), PTSD risk (prospective-longitudinal prediction of PTSD symptom development), as well as PTSD symptom severity and trauma-related behavioral avoidance in highly traumatized refugees (Amir, Zvielli, & Bernstein, 2016; Davis et al., 2016; Iacoviello et al., 2014; Yuval, Zvielli, & Bernstein, 2016; Zvielli, Vrijssen, Koster, & Bernstein, 2016). Furthermore, in contrast to poor reliability of traditional aggregated mean bias scores (i.e., Spearman-Brown prophecy corrected $r = .11$ (Schmukle, 2005; Waechter, Nelson, Wright, Hyatt, & Oakman, 2014), key features of the temporal dynamics of AB (TL-BS bias dynamics parameters) have demonstrated acceptable to excellent split-half reliability (Rodebaugh et al., 2016; Schäfer et al., 2016; Zvielli et al., 2015). Furthermore, the dynamic process approach facilitates empirical examination of the phenomenon from moment-to-moment in time. As a first such step, Amir and colleagues (2016) found that the real-time, dynamic expressions of overt (via eye-movement) and covert (via behavioral reaction time) attentional processing of threatening information were significantly associated from trial-to-trial; and dissociated by voluntary inhibition of overt attention, in anxious adults. These findings are consistent with long-standing basic cognitive- and neuro-science of attention (de Haan, Morgan, & Rorden, 2008; Posner, Snyder, & Davidson, 1980). Thus, the new-found capacity to model the real-time expression of biased attentional processing of emotional information enables the development of theory and empirical study of processes that subserve the temporal dynamics of AB.

Yet, we know strikingly little about this phenomenon despite its theorized functions for human (mal)adaptation. Specifically, we know little about the mechanisms that may contribute to or transact with moment-to-moment expression and fluctuations of biased

attentional processing. These fluctuations may reflect moment-to-moment interactions or conflict between: (1) bottom-up processes driving attention toward motivationally-relevant information; and (2) top-down processes to maintain attention on task-relevant information and down-regulate bottom-up processes (Corbetta & Shulman, 2002). Yet, for such top-down control processes to be functionally deployed, they require moment-to-moment feedback (Mansell & Marken, 2015). This type of feedback mechanisms may monitor real-time allocation of attention, the object of attention, the source of that object of attention, or the relevance or interference of attentional allocation for task performance.

Here we argue that although a number of mechanisms may be important to understanding temporal dynamics of AB, one particularly promising start-point is meta-awareness. Indeed, Meta-Awareness of Bias (MAB) may provide a feedback function facilitating cognitive/top-down control of biased allocation of attention (Bernstein & Zvielli, 2014). Meta-awareness has been conceptualized as awareness of the processes occurring in consciousness (e.g., thinking, feeling, sensing)^a (Bernstein et al., 2015; Dahl, Lutz, & Davidson, 2015). Seminal work on meta-awareness offered a powerful framework for understanding and studying meta-awareness (Schooler, 2002). Likewise, meta-awareness is a core target and mediating process of several psychological interventions (Mindfulness-Based Interventions, Cognitive Therapy (Bernstein et al., 2015a; Teasdale et al., 2002)), and its development is thought to contribute to mental health (Bernstein et al., 2015a).

Although research to-date has yet to focus on AB per se, seminal work on meta-awareness did focus on another important form of attentional dysregulation – mind-wandering. This work has documented individual differences in capacity for meta-awareness of mind-wandering as well as moment-to-moment fluctuations in meta-awareness of mind-wandering (Schooler et al., 2011). Importantly, mind-wandering with and without meta-awareness are differentiated processes with distinct neural (Christoff, Gordon, Smallwood,

Smith, & Schooler, 2009; Hasenkamp, Wilson-Mendenhall, Duncan, & Barsalou, 2012; Smith et al., 2006), and cognitive correlates (Smallwood & Schooler, 2015). Notably, meta-awareness of mind-wandering reduces the occurrence of mind-wandering (Smallwood & Schooler, 2015), and meta-awareness during mind-wandering buffers the detrimental effects of mind-wandering on attentional/executive dysregulation (e.g., response inhibition, capacity to construct mental models) (Smallwood, McSpadden, & Schooler, 2007b). This work is furthermore important in illustrating the utility of real-time experience sampling methods for capturing the moment-to-moment expression and function of meta-awareness of attentional wandering.

Much like meta-awareness of mind-wandering, we posit that MAB may similarly enable self-regulatory control, and thereby reduce the occurrence and harmful functional consequences, of attentional dysregulation. Indeed, scholars studying meta-awareness foresaw the potential for phenomena such as MAB (Baird et al., 2013; Franklin et al., 2014). Accordingly, we hypothesized that: (1) people have a (potential) capacity to monitor and have meta-awareness of their well-rehearsed, and often involuntary dynamic expression of (biased) attentional allocation, although we may expect individual differences in MAB (Farb et al., 2007; Hölzel et al., 2011; Schooler et al., 2011; Shonin, Van Gordon, & Griffiths, 2014; Van Gordon, Shonin, Sumich, Sundin, & Griffiths, 2014); and (2) that MAB will enable self-regulatory attentional control (Ansorge, Fuchs, Khalid, & Kunde, 2011; Koivisto, Kainulainen, & Revonsuo, 2009; Posner & Rothbart, 1998; Ridderinkhof, van den Wildenberg, Wery PM, Segalowitz, & Carter, 2004; Ridderinkhof, Ullsperger, Crone, & Nieuwenhuis, 2004), and thereby remedy AB and its consequences (Bishop, 2008; Derryberry & Reed, 2002; Eysenck, Derakshan, Santos, & Calvo, 2007; Wiers, Gladwin, Hofmann, Salemink, & Ridderinkhof, 2013). Accordingly, findings from a novel intervention methodology provide indirect evidence consistent with the hypothesized capacity for and

proposed functions of MAB. Indeed, real-time feedback on the momentary expression of AB – Attention Feedback Awareness and Control Training (A-FACT) designed to train MAB – led to reduced attentional dysregulation and reduced consequent emotional reactivity in response to threatening information (Bernstein & Zvielli, 2014a; Sanchez Lopez, Everaert, & Koster, 2016; Zvielli, Amir, Goldstein, & Bernstein, 2016; see also deBettencourt, Norman, & Turk-Browne, 2015), for related work on feedback-based intervention methodologies).

In the present study, we therefore aimed to estimate capacity for, and individual differences in, MAB; and to explore the hypothesized function of real-time MAB on real-time micro-expressions dysregulated attentional processing of emotional information. We did so by means of a novel measurement methodology – the Probe-Caught Meta-Awareness of Bias task (PC-MAB). First, probe-sampling methods, used to study meta-awareness of mind wandering (Smallwood et al., 2007b), were applied to estimate real-time “subjective” AB. Then, trial-level bias-scores (Zvielli et al., 2015), were applied to estimate trial-level “objective” AB scores. Finally, Signal Detection Theory (SDT), applied in earlier work on awareness and meta-cognition (Galvin, Podd, Drga, & Whitmore, 2003; Maniscalco & Lau, 2012), was applied to empirically contrast the objective and subjective estimates of AB to quantify MAB as the agreement between these estimates. We focused here on AB to reward stimuli by measuring AB to smoking cues among daily cigarette smokers. This methodology ensures the motivational-relevance of emotional information. This is important to maximize the probability of observing biased attentional processing of target stimuli and thereby the ability to study of MAB (Baker, Piper, McCarthy, Majeskie, & Fiore, 2004; Field & Cox, 2008; Pergamin-Hight, Naim, Bakermans-Kranenburg, van IJzendoorn, & Bar-Haim, 2015; Zvielli et al., 2015).

First, we hypothesized that signal detection analysis would reveal evidence of capacity for and individual differences in MAB. Second, we hypothesized that if this

potential capacity for MAB is observed and substantive, then real-time or momentary MAB would be most likely observed concurrent with the micro-expressions of attention that are most biased. Indeed, greater momentary levels of AB would be more phenomenologically prominent and thereby detected – for example, by interfering with task performance or by cueing some concurrent epi-phenomenon linked to AB (e.g., emotional responding) thereby facilitating MAB. Finally, we hypothesized that momentary MAB would buffer the unfolding of dysregulated emotional attending or attentional reactivity to biased processing of emotional information. Specifically, we hypothesized that the momentary micro-expression of AB in the *absence* of MAB would more likely immediately precede self/emotion-regulatory attentional reactivity or dyscontrol; whereas, momentary micro-expression of AB *with* MAB would more likely immediately precede cognitive/attentional control.

Methods

Participants

Seventy-five adults^b (62.7% female, $M(SD) = 26.8(3.59)$ years-old) were recruited through public advertisements from the University of Haifa. Participants were eligible if: (a) daily cigarette smokers (~10 cigarettes/day) and (b) ≥ 18 years old. Potential participants were ineligible if: (a) diagnosed with ADHD or an autism spectrum disorder; (b) uncorrected vision problems; (c) self-reported past or current participation in a meditation practice (to rule-out past systematic training in meta-awareness); (d) current/history of psychotic disorders; (e) current use of psycho-pharmacology treatment; or (f) not fluent in Hebrew. Participants were daily, long-term, and non-deprived smokers. Specifically, participants smoked at least 5 cigarettes per day (range 5 to 40 cigarettes/day). Overall, participants reported smoking $M(SD) = 11.51(6.31)$ cigarettes/day – 33% of participants smoked 5 to 9 cigarettes/day and 66% smoked 10 more more cigarettes/day. Furthermore, participants

initiated smoking regularly at $M(SD) = 18.12(2.85)$ years of age, smoked for $M(SD) = 8.20(4.00)$ years, made $M(SD) = 1.75(1.77)$ life-time quite attempts, and reported smoking their first cigarette each morning $M(SD) = 76.21(98.2)$ minutes after waking. These smoking characteristics are consistent with a variety of published community samples of regular daily smokers (Goodwin et al., in press). Finally, consistent with the likelihood that participants smoked prior to the lab session, participants reported a relatively low level of craving to smoke at the start of the lab session $M(SD) = 2.93(1.46)$ ($1 = not at all$ to $7 = very much$); and a significantly elevated level of craving to smoke at the end of the lab session $M(SD) = 4.73(1.66)$ ($t(54) = 8.1, p < .001$).

Procedure

Eligible participants completed a brief self-report battery (not relevant to the present report). Participants then completed the biochemical verification of smoking status, followed by the PC-MAB to measure AB and MAB^c.

Materials & Apparatus

Biochemical Smoking Status Assessment. Participants completed a carbon monoxide biochemical verification of smoking status ($PPM \geq 8$; CMD/CO Carbon Monoxide Monitor (Model 3110, Spirometrics, Inc. Auburn, ME).

Attentional Bias Measurement. The visual emotional dot probe task (MacLeod, Mathews, & Tata, 1986), was used to measure AB to appetitive cues. Participants were presented with a fixation cross (500 ms), followed by 100 ms blank screen, followed by two stimuli presented simultaneously for a duration of 500 ms. One stimulus was presented to the left of the fixation cross and the other to the right. One of the stimuli was immediately replaced by a small black probe per trial (50% per side). One of the presented stimuli was always neutral (e.g. pencils), and the other stimuli was either appetitive (e.g., cigarettes; reward trial; 50% of trials) or neutral (neutral trial; 50% of trials). Participants were

instructed to first focus their gaze on the fixation cross and then, as quickly and accurately as possible, press one of two (left or right) response box buttons corresponding to the location of the probe. A random interval of 500-1500 ms preceded the next trial. On congruent trials (CT; 25% of trials) the probe appeared in the location of the appetitive stimulus, and on incongruent trials (IT; 25% of trials), the probe appeared in the opposite location of the appetitive stimulus (location of neutral stimulus). Concurrent with the dynamic process perspective on AB (Zvielli et al., 2015), we computed TL-BS by matching and then subtracting temporally contiguous pairs of reward-neutral trial's RT by subject. Each reward trial (CT or IT) was matched with the first next neutral trial (NT), to reduce any carry-over effects of other rewarding stimuli. The TL-BS computation is thereby used to compute objective trial-level bias scores estimations.

Probe-Caught Meta-Awareness of Bias Task (PC-MAB). Grounded in probe-caught thought sampling methods (Giambra, 1995; Smallwood & Schooler, 2006), participants were repeatedly probed regarding their attention, concurrent with the repeated trial-level expressions of (biased) attentional expression in the dot-probe task. Awareness probes were delivered randomly over the course of the dot-probe task (25% of trials). Immediately following trial response, participants were asked to report whether one of the pictures in the last trial influenced their response. If their response was affirmative, then participants were asked, “How did the picture influence your response” (i.e., "my attention moved away from it" *or* "my attention focused on it"). By contrasting indices of real-time subjective AB estimates (i.e., via probe-caught awareness samples) with real-time objective AB scores (i.e., via TL-BS) we were able to compute an objective index of MAB in real-time (see Figure 1).

A Signal Detection Theory Computation of MAB. On awareness-probed trials, we matched subjective reporting of attention allocation with trial-level bias scores to compute

true- and false- positive/negative rates (TP, FP, TN, FN, correspondingly) (see Table 1). This signal detection-based dichotomization enabled us to quantify agreement between subjective AB estimates (i.e., via answers to awareness-probes) and objective AB scores (i.e., via TL-BS) in one 2 x 2 contingency table for each participant, as depicted in Table 1 (see (Galvin et al., 2003; Maniscalco & Lau, 2012)). The trial-level MAB scores (i.e., true and false positives/negatives) were then used in the computation of a single indicator of MAB – the Diagnostic Odds Ratio (DOR; i.e., the ratio of the odds of disease in test positives relative to the odds of disease in test negatives (Glas, Lijmer, Prins, Bossel, & Bossuyt, 2003), and real-time multi-level analyses (detailed above). DOR provides a valid representation of participant's discriminatory performance, combines estimates of sensitivity and specificity, and has the unique advantage of a single indicator to estimate MAB. Importantly, DOR is also independent of inter-individual differences in the prevalence of AB and in the rate of subjective AB, and thus the most rigorous variable to estimate sample- and inter-individual differences in MAB. The calculation of the DOR can be derived directly from the 2 x 2 contingency tables (see Table 1) and is calculated as follows:

$$\text{DOR} = \frac{\text{TP}}{\text{FP}} / \frac{\text{FN}}{\text{TN}} = \frac{\text{TP} \cdot \text{TN}}{\text{FP} \cdot \text{FN}}$$

To compute the DOR for each participant, and because some 2 x 2 tables contained zeroes (in which DOR would be undefined), we added 0.50 to all values in the table (Littenberg & Moses, 1993). DOR values range from 0 to infinity, with higher values indicating better discriminatory test performance (Glas et al., 2003). DOR = 1 indicates that a participant does not discriminate between AB and non-AB trials (i.e., chance performance, or *no MAB*); similarly, DOR < 1 indicates that a participant does not accurately discriminate between AB and non-AB trials – characterized by higher rates of false negatives and false positive than true positives and true negatives (i.e., *no MAB*). Whereas DOR > 1 indicates that a participant discriminates between AB and non-AB trials at a rate greater than chance

(i.e., *MAB*) (e.g., $DOR = 4$ means that for a particular subject, the odds of correctly detecting AB when AB was objectively expressed (TP) was 4 times greater than the odds for inaccurately detecting AB in the absence of AB (FP)). $DOR=1$ or chance values means that sometimes participants were right/accurate and other times wrong/inaccurate. This pattern of responding is that which you would expect were a person simply guessing (i.e., flipping a coin). In this way, these participants do not demonstrate evidence of MAB. Participants exhibiting $DOR < 1$ were more consistently wrong or inaccurate. In this way, these participants do not demonstrate evidence of MAB. Finally, because the scale of the DOR is exponential (i.e., highly skewed distribution), calculating a log-transformation DOR for each participant (i.e., $\log DOR$) reduced the skew of the distribution – important to reported data analyses of MAB that assume normality (Tabachnick & Fidell, 2007).

Results

Meta-Awareness of Bias: Capacity and Individual Differences

First, we tested evidence for, and individual differences in, MAB via Signal Detection Theory (SDT) analyses. Participants demonstrated some degree of positive *objective* AB (TL-BS) in $M(SD) = 65.65(13.13)\%$ of trials, and positive *subjective* AB (awareness-probes) in $27.66(23.52)\%$ of trials. For each participant, we calculated the diagnostic odds ratio (DOR; i.e., the ratio of the odds of positive objective (TL-BS) AB in positive subjective reports (awareness-probes) relative to the odds of positive objective (TL-BS) AB in negative subjective reports (awareness-probes) – see Methods section below for details) as a single indicator for MAB. We found that 46.7% of participants demonstrated $DOR \leq 1$ (i.e., log transformation of DOR ($\log DOR$) ≤ 0); and 53.3% of participants demonstrated $DOR > 1$ (i.e., $\log DOR > 0$): DOR = ranges from .03 to 17.4; $M(SD) DOR = 2.29(3.31)$, $95\%CI_{\text{mean}} [1.53, 3.05]$; $Md = 1.12$; $\log DOR$ = ranges from -1.54 to 1.24; $M(SD); \log DOR = .04(.53)$,

95%CI_{mean} [-.08,.17]; Md = .04. To more conservatively estimate MAB, so that we do not misclassify values that are negligibly but not meaningfully > chance, we computed the 95%CI of the logDOR_{MEAN} of 0 (95%CI *logDOR*_{MEAN} = -.8 to .17) and operationalized MAB as logDOR values > than .17, the upper limit of the 95% CI of the logDOR_{MEAN} of 0: i.e., logDOR = -.8 to .17 indicate chance performance (i.e., no MAB); logDOR < -.8 indicate no MAB; and logDOR values > .17 reflect levels of MAB > chance. We thus found that 64% of participants demonstrated no MAB (i.e., 24% demonstrated chance performance, 40% demonstrated DOR < chance); and 36% of participants demonstrated some degree of MAB, ranging from low levels just above chance (e.g., DOR = 1.52) to high levels of MAB (e.g., DOR = 17.4).

Second, if this potential capacity for MAB is indeed observed and substantive, then real-time or momentary MAB should be most likely observed concurrent with the micro-expressions of attention that are most biased; moreover, such effects should be uniquely observed among participants for whom SDT-based estimation of MAB via DOR (above) indicates evidence of MAB > chance. Accordingly, to examine the real-time trial-to-trial relations between AB and MAB, we carried out a Mixed Linear Model analysis (MLM) via the PROC MIXED procedure in SAS 9.4 (SAS Institute Inc., 2008). For each participant, each time point (trial) included two scores: real-time objective AB score (momentary TL-BS trial estimate) and real-time subjective AB estimate (momentary response to trial-level awareness probe). Because awareness-probes always immediately follow AB expression, AB amplitude was the predictor in this model. The trial-level AB amplitude predictor was continuous (i.e., absolute TL-BS amplitude) and the trial-level MAB outcome was dichotomous (true positive/negative vs. false positive/negative). LogDOR was included as a moderator. As predicted, we found a significant interaction between momentary AB amplitude and logDOR level with respect to momentary MAB ($F_{(1, 1343)} = 12.74, B = 50.08,$

$SE = 14.03, p = .0004$). To unpack this interaction, we used a contrast within the model. As predicted, we found that momentary AB amplitude predicted momentary MAB ($B = 48.84, SE = 9.70, t = 4.21, p < .0001$) but only for participants who demonstrated evidence of MAB in the SDT analysis (i.e., $\log DOR > .17$). In addition, we found no interaction with the direction of AB direction (i.e., towards or away from reward) ($F_{(2, 1341)} = 1.27, p = .282$), such that the observed association between momentary AB amplitude and momentary MAB did not differ between bias toward and away from reward.

Third, to ensure that our classification of participants with vs. without MAB (i.e., above-chance DORs vs. chance-level and below-chance DORs) is empirically justified, we conducted the above MLM – testing the real-time trial-to-trial relations between AB and MAB – but this time divided the sample into three sub-groups according to their DOR scores: (1) participants expressing DOR levels $>$ chance (i.e., $\log DOR > .17$); (2) participants expressing DOR = chance (i.e., $-.08 \leq \log DOR \leq .17$); and (3) participants expressing DOR $<$ chance (i.e., $\log DOR < -.08$). As expected, we found that participants expressing chance and below-chance DOR scores did not perform differently ($B = -.18, SE = 18.85, t_{1342} = .01, p = 0.9925$); whereas participants demonstrating above-chance DORs (i.e., participants who demonstrated some degree of MAB) perform significantly differently than both sub-groups groups ($B = 40.01, SE = 16.37, t_{1342} = 2.44, p = .0146$). To further examine this effect, we tested these associations with respect to continuous levels of $\log DOR$ – to be certain that these findings are not due to some distributional artifact secondary to the sub-group analysis or due to limited power within the MLM. Again, we found that momentary AB amplitude predicted momentary MAB (DOR scores) only among participants exhibiting above-chance DOR scores ($B = 61.88, SE = 14.39, t_{1343} = 4.3, p < .0001$) and not among participants who demonstrate chance ($B = 11.79, SE = 7.27, t_{1343} = 1.62, p = .11$) or below-chance performance ($B = -13.24, SE = 11.12, t_{1343} = -1.19, p = .23$).

MAB & Attentional (Dys)Control

We next examined the real-time effect of momentary MAB on attentional reactivity or (dys)control. We did so by testing whether momentary micro-expressions of AB without MAB would immediately precede self/emotion-regulatory attentional reactivity or dyscontrol; whereas, momentary micro-expression of AB *with* MAB would immediately precede cognitive/attentional control. We ran a Mixed Linear Model analyses (MLM) via the PROC MIXED procedure in SAS 9.4 (SAS Institute Inc., 2008). For each participant, each time point (trial) included two scores: trial-level true-positive-match of MAB (i.e., true positive between subjective and objective bias in the event of bias toward or away from reward); and degree of change in momentary AB micro-expressions from the trial in which MAB was measured, to the immediately proceeding trial (i.e., $(\text{TL-BS}_{\text{trial } n+1}) - (\text{TL-BS}_{\text{trial } n})$). In this model, momentary MAB (presence or absence of *MAB*) on trial_{*n*} was the predictor and consequent change between the micro-expressions of AB from trial_{*n*} to trial_{*n+1*} the outcome. Accordingly, we were able to test the hypothesized functional role(s) of momentary presence vs. absence of MAB on the immediately proceeding micro-expression of bias – and thereby, model the real-time effect of MAB on cognitive/attentional control. We found that in the event of AB (toward or away from reward), the momentary *absence* of MAB led to a higher degree of attentional reactivity or dyscontrol in which a micro-expression of bias toward reward was immediately proceeded by a fluctuation to bias away from reward, and vice-versa; in contrast, momentary *presence* of MAB reduced this attentional reactivity/dyscontrol, and is more likely followed by a trial characterized by balanced attentional allocation (TL-BS close to zero) ($F_{(1, 29)} = 4.52, p = .0421$; absence of MAB $B = -180.89, SE = 17.05, t = -10.61, p < .0001$ vs. presence of MAB $B = -122.45, SE = 23.94, t = -5.11, p < .0001$) (see Figure 2).

Ruling Out Alternative Explanations and Artifacts

First, we needed to rule out the possibility that positive endorsement on probes in the first half of the task would predict reduced levels of endorsement in the second half of the task. That is, we wanted to rule out a potentially problematic "learning" effect or learned disincentive of positively endorsing question 1 of the probes (i.e., because positively endorsing question 1 of each probe led to additional questions). We thus calculated the ratio of positive vs negative responses to question 1 of the probes during the first (trials 1-80) vs the second (trials 81-160) half of the PC-MAB task trials. We found a significant, strong, and positive correlation between these two ratios, $r = .816$, $p < .001$. Participants showed no evidence of reducing (or changing) their pattern of responding to the probes over the course of repeated trials. Accordingly, we can meaningfully evaluate AB and MAB over the course of the entire block of trials. Second, we wanted to rule out an additional potential carry-over artifact of the probe-sampling methodology wherein a probe on trial n would influence the expression of attentional expression on trial $n+1$. We thus tested whether n probed-trials, relative to n un-probed trials influence levels of AB on trial $n+1$. For each participant, we calculated the aggregated mean of TLBS in $n+1$ trials for trials immediately preceded by probed-trials vs. those immediately preceded by un-probed trials. We found no carryover effect for probes on AB ($t_{74} = 1.04$, $p = .30$). Accordingly, we conclude the awareness probes did not systematically interfere with task performance or attentional processing of emotional information over the course of the task. Finally, analyses were examined among regular daily smokers. We thus wanted to examine and rule out the possibility that heavier or more dependent smokers may demonstrate different levels of MAB or function(s) of MAB on attentional reactivity or (dys)control reported above. Rate of smoking was not associated with levels of MAB ($r = -.10$, $p = .394$), nor did smoking rate moderate the observed effect of MAB on attentional control/reactivity ($F_{(1, 184)} < 1$, $p = .94$). Accordingly, we were able to meaningfully

evaluate levels of MAB and its function on attentional (dys)regulation among the studied sample of regular daily smokers.

Finally, we tested the potential role(s) of craving (i.e., change in subjective reported craving levels from baseline to post-test) on reported effects of MAB. Craving change scores were selected as they best reflect degree of acute subjective craving over the course of the laboratory session during which MAB was measured. In our first MLM analysis (i.e., real-time trial-to-trial relations between AB and MAB), individual differences in subjective craving levels were not related to amplitude of AB ($B = .77$, $SE = 3.68$, $t_{982} = .21$, $p = .8345$). Likewise no interaction was observed between craving levels and our main predictor of MAB (i.e., DOR) ($F_{(1, 982)} = .9$, $B = 4.06$, $SE = 4.28$, $p = .3431$). Thus, the reported effects were not dependent on craving levels over the course of the experiment, nor alternatively explained by craving levels. Finally, when re-testing the real-time trial-to-trial relations between AB and MAB, adding change in craving levels to the model did not change the magnitude or significance of the reported effects. Again, we found a significant interaction between momentary AB amplitude and logDOR level with respect to momentary MAB ($F_{(1, 982)} = 10.84$, $B = 66.31$, $SE = 20.14$, $p = .001$). We then tested the effect of change in craving levels in the second reported MLM model – testing the effect of MAB on cognitive/attentional control (p. 15). Change in craving was, again, nonsignificant ($B = 10.75$, $SE = 11.74$, $t_{127} = .92$, $p = .3616$); craving did not interact with MAB ($B = 3.24$, $SE = 15.09$, $t_{127} = .21$, $p = .8301$); and adding craving levels to the model did not change the reported effect of momentary MAB on attentional/cognitive control ($B = 73.90$, $SE = 35.27$, $t_{21} = 2.1$, $p = .046$). These results did not change when we alternatively tested other craving scores (e.g., baseline alone, post-test alone).

Discussion

We aimed to empirically examine meta-awareness of momentary micro-expressions of biased attentional processing of reward cues among non-deprived daily smokers. First, SDT analysis revealed evidence of the capacity for, and individual differences in MAB. This is the first empirical evidence for MAB. This also represents, to the best of our knowledge, the first attempt to apply SDT (Galvin et al., 2003; Maniscalco & Lau, 2012) to probe-sampling data (Smallwood, McSpadden, & Schooler, 2007a; Teasdale, Segal, & Williams, 1995) to quantify meta-awareness of any process in consciousness more generally. Second, through a novel integration of SDT and multi-level modelling, we found a significant association between the amplitude of momentary (trial-level) AB and momentary (trial-level) MAB. Momentary MAB was most likely observed in the event of the most extreme micro-expressions of AB but only among individuals that our SDT analysis identified as demonstrating some degree of MAB (i.e., DOR > chance). These data provide important initial validation of the methodology to quantify MAB. Finally, as predicted, we found that momentary micro-expression of AB *without* MAB was more likely followed by attentional reactivity/dyscontrol. In contrast, momentary micro-expression of AB *with* MAB was more likely followed by a more balanced (less biased) attentional expression or greater cognitive/attentional control. These data provide promising evidence that MAB may function to regulate attentional processing of emotion or motivationally-relevant information (reward). Finally, we tested and empirically ruled out 4 potential methodological artifacts that could alternatively account for observed effects. There was no evidence of a learned disincentive of positively endorsing question 1 of the probes over the course of the task (i.e., because positively endorsing question 1 of each probe led to additional questions). Awareness probes did not systematically interfere with task performance or attentional processing of emotional information over the course of the task. Rate of smoking was not associated with levels of

MAB or its function on attentional control/reactivity. Degree of craving to smoke during the experimental session did not alternatively account for reported findings.

The present findings have a number of implications for basic and clinical psychological science of attentional dysregulation. First, findings indicate that similar to meta-awareness of other mental processes (Lutz, Jha, Dunne, & Saron, 2015; Schooler et al., 2011), MAB may provide a monitoring-for-regulation function. For example, MAB may reduce dysregulated attentional processing directly by means of top-down executive control that actively inhibits exogenous attentional orienting toward or away from reward cues (Bernstein & Zvielli, 2014a; Schooler et al., 2011). Specifically, MAB may serve a moment-to-moment internal feedback function regarding the allocation of (dysregulated) attention thereby enabling top-down regulation or control of attentional processing (e.g., Mansell & Marken, 2015). Alternatively or additionally, MAB may enable *indirect* regulation or control of attention by directly impacting other internal states that influence or sub-serve attentional processing. For example, MAB may reduce (involuntary) attentional reactivity to internal states elicited by stimulus exposure or it may function to down-regulate the urge or emotional state elicited by stimulus exposure that in turn contributes to attentional orienting (Bernstein et al., 2015b; Herwig, Kaffenberger, Jancke, & Bruhl, 2010; Holzel et al., 2011). Future research designed to test these hypothesized mechanisms may be particularly useful to work on attentional dysregulation as well as the role(s) of meta-awareness for other internal processes more generally.

The present findings may also have interesting translational implications. Indeed, the present findings are broadly in line with the idea that MAB may be a promising intervention target. The findings indicate that people have the (potential) capacity for MAB, and that MAB may have a monitoring-for-regulation function on attentional processing of emotion or motivationally-relevant information important to mental (ill)health. A growing body of work

indicates that meta-awareness broadly, and MAB specifically, is likely malleable and may be targeted via interventions. For example, real-time attentional feedback designed to facilitate MAB may be one such promising approach (Bernstein & Zvielli, 2014b; deBettencourt et al., 2015; Sanchez Lopez et al., 2016; Zvielli, Amir, Goldstein, & Bernstein, 2015). Similarly, mindfulness-based interventions targeting meta-awareness of multiple internal states and processes may also provide powerful means to target MAB specifically or meta-awareness of other internal states important to mental (ill)health more broadly (Bernstein et al., 2015a; Hadash, Plonsker, Vago, & Bernstein, 2016; Lutz et al., 2015). Studies targeting MAB may advance the field's capacity to therapeutically target attentional dysregulation implicated in prevalent forms of mental (ill)health.

The present study has a number of limitations. First, we tested MAB to reward stimuli only among a clinical population for whom addiction-related processes drive biased attentional processing of reward (Field & Cox, 2008; Zvielli et al., 2015). An important next step is to test the capacity for MAB, as well as its functional role(s), with respect to aversive stimuli and respective populations (e.g., patients with anxiety and mood disorders). Second, MAB was not successfully experimentally manipulated in the current study. Our efforts to do so using a brief analogue mindfulness manipulation were unsuccessful. As noted above, future study may test more robust methods to experimentally manipulate MAB so as to evaluate its putative causal function(s) for attentional (dys)regulation. Third, this study examined MAB in the dot-probe task – the most widely used cognitive-experimental task measuring biased attentional processing of emotional information (MacLeod et al., 1986). However, it may also be useful for future study to examine MAB in other experimental tasks measuring covert (e.g., spatial cueing, visual search) and overt (e.g., eye-tracking during free view) attention bias. Fourth, MAB was measured using a probe-caught methodology (Smallwood & Schooler, 2015). Future study may attempt to measure self-initiated MAB in

real-time via a self-caught methodology (Smallwood & Schooler, 2006). Fifth, in the event that our objective index of AB (TLBS) reflected predominantly noise and not “signal” (of biased momentarily attentional allocation) or our subjective index based on the awareness-probes reflected predominantly noise and not “signal” (of MAB), then the observed results would be statistically unlikely. Nevertheless, the greater the signal-to-noise ratio in measurement of meta-awareness on the one hand and momentary bias on the other, the more likely that research on MAB will yield important information and translational implications for intervention. It is important that ongoing work focus on methodological and statistical advances to achieve such aims (Kruijt, Field, & Fox, 2016), such as via application of computational modeling, machine learning or pattern classification to such data as a means of optimizing signal to noise. Finally, we did not test the role of MAB on outcomes related to attentional processing of reward (smoking cues) among the smokers beyond the lab (e.g., cessation attempt outcomes, intervention responding, smoking rate, time since last cigarette, etc.) nor the possible therapeutic effects of targeting MAB for such addiction outcomes. Such work may be an exciting translational direction of this initial laboratory investigation.

In summary, we developed a methodology to objectively estimate MAB. We found initial evidence of the capacity for, and individual differences in MAB, as well as evidence that MAB may function to regulate attentional processing of reward cues. The current study may serve as an empirical proof of concept and initial context to develop and test the utility of the proposed methodological and data analytic framework for the study of meta-awareness. We believe this work may be important in so far as we speculate that meta-awareness may have broad and significant implications for clinical psychological science. The reported findings may contribute to an emerging science of the nature and functions of meta-awareness of various internal processes including (biased) attentional processing of emotional or motivationally-relevant information.

Footnotes

^a This "meta-" level of awareness is distinguished from awareness of the contents of thoughts (i.e., mental representations) without concurrent meta-awareness of the thinking process; or similarly, awareness of the objects producing sense impressions with no meta-awareness of the sense impressions themselves (Bernstein et al., 2015a; Brown & Ryan, 2003; Dahl et al., 2015).

^b Five additional participants were not included in reported analyses because they reported not understanding the task instructions or stopped the experiment repeatedly to ask for assistance from the experimenter; accordingly, their data were identified during data collection and omitted prior to data analysis.

^c In addition to the study aims, we also attempted to elicit meta-awareness by means of a brief recorded focused-attention to breath practice. We randomized participants to one of two groups: (1) meta-awareness enhancement condition (e.g., *"at some point your attention will begin to wander to things unrelated to your breathing. Once you've noticed this wandering, focus your attention back to the sensations that arise during the rise and drop of the abdomen"*); or (2) control condition (i.e., audio description of *"bird migration"*). The manipulation however did not work – there was no between-group differences on perceived meta-awareness [$t_{72}=.3$, N.S.]. All analyses were tested with respect to group. Group had no effect at all on any of the reported analyses. All reported analyses were thus carried out in the total $N=75$ sample, collapsing across conditions to maximize power and the reliability of reported findings.

Table 1

2 x 2 objective and subjective bias contingency table used to compute Meta-Awareness of Bias (MAB)

		<i>Objective (TL-BS)</i>	
		<i>Bias</i>	<i>No Bias</i>
<i>Subjective (PC-MAB)</i>	<i>Bias</i>	True Positives (TP)	False Positives (FP)
		False	True
	<i>No Bias</i>	Negatives (FN)	Negatives (TN)

Note. (1) The abbreviations TP, FP, FN, and TN denote the number of respectively, true positives, false positives, false negatives, and true negatives; (2) TL-BS = trial-level bias scores and PC-MAB = probe-caught meta-awareness of bias task.




Figure 1. Dynamic Process Perspective on Attentional Bias Expression (left) and the Probe-Caught Meta-Awareness of Bias task (right)




Figure 2. Attentional Bias With and Without Meta-Awareness: Temporal Evidence of a Monitoring-For-Control Function.

References

- Amir, I., Zvielli, A., & Bernstein, A. (2016). (De)coupling of our eyes and our mind's eye: A dynamic process perspective on attentional bias. *Emotion (Washington, D.C.)*, 16(7), 978-986. doi:10.1037/emo0000172 [doi]
- Ansorge, U., Fuchs, I., Khalid, S., & Kunde, W. (2011). No conflict control in the absence of awareness. *Psychological Research*, 75(5), 351-365.
- Baird, B., Smallwood, J., Fishman, D. J., Mrazek, M. D., & Schooler, J. W. (2013). Unnoticed intrusions: Dissociations of meta-consciousness in thought suppression. *Consciousness and Cognition*, 22(3), 1003-1012. doi: 10.1016/j.concog.2013.06.009.
- Baker, T. B., Piper, M. E., McCarthy, D. E., Majeskie, M. R., & Fiore, M. C. (2004). Addiction motivation reformulated: An affective processing model of negative reinforcement. *Psychological Review*, 111(1), 33-51. doi:10.1037/0033-295X.111.1.33
- Bar-Haim, Y., Lamy, D., Pergamin, L., Bakermans-Kranenburg, M., & van IJzendoorn, M. H. (2007). Threat-related attentional bias in anxious and nonanxious individuals: A meta-analytic study. *Psychological Bulletin*, 133(1), 1-24. doi:10.1037/0033-2909.133.1.1
- Bernstein, A., & Zvielli, A. (2014a). Attention feedback awareness and control training (A-FACT): Experimental test of a novel intervention paradigm targeting attentional bias. *Behaviour Research and Therapy*, 55, 18-26.
- Bernstein, A., Hadash, Y., Lichtash, Y., Tanay, G., Shepherd, K., & Fresco, D. M. (2015a). Decentering and related constructs: A critical review and metacognitive processes model. *Perspectives on Psychological Science*, 10(5), 599-617. doi:10.1177/1745691615594577
- Bernstein, A., Hadash, Y., Lichtash, Y., Tanay, G., Shepherd, K., & Fresco, D. M. (2015b). Decentering and related constructs: A critical review and meta-cognitive processes

model. *Perspectives on Psychological Science*, 10(5), 599-617.

doi:10.1177/1745691615594577

Bernstein, A., & Zvielli, A. (2014b). Attention feedback awareness and control training (A-FACT): Experimental test of a novel intervention paradigm targeting attentional bias. *Behaviour Research and Therapy*, 55, 18-26.

doi:<http://dx.doi.org/10.1016/j.brat.2014.01.003>

Bishop, S. J. (2008). Trait anxiety and impoverished prefrontal control of attention. *Nature Neuroscience*, 12(1), 92-98.

Brown, K. W., & Ryan, R. M. (2003). The benefits of being present: Mindfulness and its role in psychological well-being. *Journal of Personality and Social Psychology*, 84(4), 822-848. doi:10.1037/0022-3514.84.4.822

Christoff, K., Gordon, A. M., Smallwood, J., Smith, R., & Schooler, J. W. (2009). Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proceedings of the National Academy of Sciences*, 106(21), 8719.

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, 3(3), 201-215.

Dahl, C. J., Lutz, A., & Davidson, R. J. (2015). Reconstructing and deconstructing the self: Cognitive mechanisms in meditation practice. *Trends in Cognitive Sciences*, 19(9), 515-523. doi:<http://dx.doi.org/10.1016/j.tics.2015.07.001>

Davidson, R. J., & Kaszniak, A. W. (2015). Conceptual and methodological issues in research on mindfulness and meditation. *American Psychologist*, 70(7), 581.

Davis, M., Rosenfield, D., Zvielli, A., Bernstein, A., Reinecke, A., Beevers, C., . . . Smits, J. (2016). Attention bias dynamics and symptom severity during and following CBT for social anxiety disorder. *Journal of Consulting and Clinical Psychology*,

- de Haan, B., Morgan, P. S., & Rorden, C. (2008). Covert orienting of attention and overt eye movements activate identical brain regions. *Brain Research, 1204*(0), 102-111.
doi:<http://dx.doi.org/10.1016/j.brainres.2008.01.105>
- deBettencourt, M., Norman, K., & Turk-Browne, N. (2015). Relating sustained attention to visual long-term memory. *Journal of Vision, 15*(12), 444-444.
- Derryberry, D., & Reed, M. A. (2002). Anxiety-related attentional biases and their regulation by attentional control. *Journal of Abnormal Psychology, 111*(2), 225.
- Eysenck, M. W., Derakshan, N., Santos, R., & Calvo, M. G. (2007). Anxiety and cognitive performance: Attentional control theory. *Emotion, 7*(2), 336.
- Farb, N. A. S., Segal, Z. V., Mayberg, H., Bean, J., McKeon, D., Fatima, Z., & Anderson, A. K. (2007). Attending to the present: Mindfulness meditation reveals distinct neural modes of self-reference. *Social Cognitive and Affective Neuroscience, 2*(4), 313.
- Field, M., Munafò, M. R., & Franken, I. H. (2009). A meta-analytic investigation of the relationship between attentional bias and subjective craving in substance abuse. *Psychological bulletin, 135*(4), 589.
- Field, M., & Cox, W. M. (2008). Attentional bias in addictive behaviors: A review of its development, causes, and consequences. *Drug and Alcohol Dependence, 97*(1-2), 1-20.
doi:10.1016/j.drugalcdep.2008.03.030
- Franklin, M. S., Mrazek, M. D., Anderson, C. L., Johnston, C., Smallwood, J., Kingstone, A., & Schooler, J. W. (2014). Tracking distraction: The relationship between mind wandering, meta-awareness, and ADHD symptomatology. *Journal of Attention Disorders, 56*(3):442. doi: 10.1177/1087054714543494.
- Galvin, S. J., Podd, J. V., Drga, V., & Whitmore, J. (2003). Type 2 tasks in the theory of signal detectability: Discrimination between correct and incorrect decisions. *Psychonomic Bulletin & Review, 10*(4), 843-876.

- Giambra, L. M. (1995). A laboratory method for investigating influences on switching attention to task-unrelated imagery and thought. *Consciousness and Cognition*, 4(1), 1-21.
- Glas, A. S., Lijmer, J. G., Prins, M. H., Bonsel, G. J., & Bossuyt, P. M. (2003). The diagnostic odds ratio: A single indicator of test performance. *Journal of Clinical Epidemiology*, 56(11), 1129-1135.
- Goodwin, R., Wall, M., Hu, M. C., Gbedemah, M., Weinberger, A., Galea, S., Zvolensky, M. J., Hangley, E., & Hasin, D. (in press). Nicotine dependence among smokers from 2003-2013: A new test of the hardening hypothesis. *Tobacco Control*.
- Hadash, Y., Plonsker, R., Vago, D. R., & Bernstein, A. (2016). Experiential self-referential and selfless processing in mindfulness and mental health: Conceptual model and implicit measurement methodology. *Psychological Assessment*, 28(7), 856-869.
doi:10.1037/pas0000300
- Hasenkamp, W., Wilson-Mendenhall, C., Duncan, E., & Barsalou, L. W. (2012). Mind wandering and attention during focused meditation: A fine-grained temporal analysis of fluctuating cognitive states. *NeuroImage*, 59(1), 750-760.
doi:10.1016/j.neuroimage.2011.07.008
- Heeren, A., De Raedt, R., Koster, E. H., & Philippot, P. (2013). The (neuro) cognitive mechanisms behind attention bias modification in anxiety: Proposals based on theoretical accounts of attentional bias. *Frontiers in Human Neuroscience*, 7, 119.
doi:10.3389/fnhum.2013.00119
- Herwig, U., Kaffenberger, T., Jancke, L., & Bruhl, A. B. (2010). Self-related awareness and emotion regulation. *50*(2), 734-741.
- Hölzel, B. K., Lazar, S. W., Gard, T., Schuman-Olivier, Z., Vago, D. R., & Ott, U. (2011). How does mindfulness meditation work? proposing mechanisms of action from a

- conceptual and neural perspective. *Perspectives on Psychological Science*, 6(6), 537-559. doi:10.1177/1745691611419671
- Holzel, B. K., Lazar, S. W., Gard, T., Schuman-Olivier, Z., Vago, D. R., & Ott, U. (2011). How does mindfulness meditation work? proposing mechanisms of action from a conceptual and neural perspective. *Perspectives on Psychological Science : A Journal of the Association for Psychological Science*, 6(6), 537-559. doi:10.1177/1745691611419671 [doi]
- Iacoviello, B. M., Wu, G., Abend, R., Murrough, J. W., Feder, A., Fruchter, E., . . . Pine, D. S. (2014). Attention bias variability and symptoms of posttraumatic stress disorder. *Journal of Traumatic Stress*, 27(2), 232-239.
- Koivisto, M., Kainulainen, P., & Revonsuo, A. (2009). The relationship between awareness and attention: Evidence from ERP responses. *Neuropsychologia*, 47(13), 2891-2899.
- Kruijt, Anne-Wil, Andy P. Field, and Elaine Fox. "Capturing dynamics of biased attention: are new attention variability measures the way forward?." *PloS one* 11.11 (2016): e0166600.
- Littenberg, B., & Moses, L. E. (1993). Estimating diagnostic accuracy from multiple conflicting reports: A new meta-analytic method. *Medical Decision Making : An International Journal of the Society for Medical Decision Making*, 13(4), 313-321.
- Lutz, A., Jha, A. P., Dunne, J. D., & Saron, C. D. (2015). Investigating the phenomenological matrix of mindfulness-related practices from a neurocognitive perspective. *American Psychologist*, 70(7), 632.
- MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional bias in emotional disorders. *Journal of Abnormal Psychology*, 95(1), 15-20.

- Maniscalco, B., & Lau, H. (2012). A signal detection theoretic approach for estimating metacognitive sensitivity from confidence ratings. *Consciousness and Cognition*, 21(1), 422-430. doi:<http://dx.doi.org/10.1016/j.concog.2011.09.021>
- Mansell, W., & Marken, R. S. (2015). The origins and future of control theory in psychology. *Review of General Psychology*, 19(4), 425-430. doi:10.1037/gpr0000057
- Pergamin-Hight, L., Naim, R., Bakermans-Kranenburg, M. J., van IJzendoorn, M. H., & Bar-Haim, Y. (2015). Content specificity of attention bias to threat in anxiety disorders: A meta-analysis. *Clinical Psychology Review*, 35, 10-18.
- Posner, M. I., Snyder, C. R., & Davidson, B. J. (1980). Attention and the detection of signals. *Journal of Experimental Psychology: General*, 109(2), 160.
- Posner, M. I., & Rothbart, M. K. (1998). Attention, self-regulation and consciousness. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 353(1377), 1915-1927. doi:10.1098/rstb.1998.0344
- Ridderinkhof, K. R., van den Wildenberg, Wery PM, Segalowitz, S. J., & Carter, C. S. (2004). Neurocognitive mechanisms of cognitive control: The role of prefrontal cortex in action selection, response inhibition, performance monitoring, and reward-based learning. *Brain and Cognition*, 56(2), 129-140.
- Ridderinkhof, K. R., Ullsperger, M., Crone, E. A., & Nieuwenhuis, S. (2004). The role of the medial frontal cortex in cognitive control. *Science (New York, N.Y.)*, 306(5695), 443-447. doi:10.1126/science.1100301
- Rodebaugh, T. L., Scullin, R. B., Langer, J. K., Dixon, D. J., Huppert, J. D., Bernstein, A., . . . Lenze, E. J. (2016). Unreliability as a threat to understanding psychopathology: The cautionary tale of attentional bias. *Journal of Abnormal Psychology*, 125(6), 840.
- Sanchez Lopez, A., Everaert, J., & Koster, E. (2016). Attention training through gaze-contingent feedback: Effects on reappraisal and negative emotions. *Emotion*,

Sayette, M.A., Schooler, J.W., & Reichle, E.D. (2010). Out for a smoke: The impact of cigarette craving on zoning out while reading. *Psychological Science*, 21, 1, 26-30.

doi:10.1177/0956797609354059.

SAS Institute Inc. (2008).

. *SAS / STAT® 9.2 user's guide cary* ()

Schäfer, J., Bernstein, A., Zvielli, A., Höfler, M., Wittchen, H., & Schönfeld, S. (2016).

Attentional bias temporal dynamics predict posttraumatic stress symptoms: A prospective–longitudinal study among soldiers. *Depression and Anxiety*,

Schmukle, S. C. (2005). Unreliability of the dot probe task. *European Journal of Personality*, 19(7), 595-605.

Schooler, J. W. (2002). Re-representing consciousness: Dissociations between experience and meta-consciousness. *Trends in cognitive sciences*, 6(8), 339-344.

Schooler, J. W., Smallwood, J., Christoff, K., Handy, T. C., Reichle, E. D., & Sayette, M. A. (2011). Meta-awareness, perceptual decoupling and the wandering mind. *Trends in Cognitive Sciences*, 15(7), 319-326.

Shonin, E., Van Gordon, W., & Griffiths, M. D. (2014). Meditation awareness training (MAT) for improved psychological well-being: A qualitative examination of participant experiences. *Journal of Religion and Health*, 53(3), 849-863.

Smallwood, J., McSpadden, M., & Schooler, J. W. (2007a). The lights are on but no one's home: Meta-awareness and the decoupling of attention when the mind wanders. *Psychonomic Bulletin & Review*, 14(3), 527-533.

Smallwood, J., & Schooler, J. W. (2015). The science of mind wandering: Empirically navigating the stream of consciousness. *Annual Review of Psychology*, 66, 487-518.

Smallwood, J., McSpadden, M., & Schooler, J. W. (2007b). The lights are on but no one's home: Meta-awareness and the decoupling of attention when the mind wanders.

Psychonomic Bulletin & Review, 14(3), 527-533. doi:10.3758/BF03194102

Smallwood, J., & Schooler, J. W. (2006). The restless mind. *Psychological Bulletin*, 132(6), 946-958. doi:10.1037/0033-2909.132.6.946

Smith, R., Keramatian, K., Smallwood, J., Schooler, J., Luus, B., & Christoff, K. (2006).

Mind-wandering with and without awareness: An fMRI study of spontaneous thought processes. *Proceedings of the 28th Annual Conference of the Cognitive Science Society*, 804-809.

Tabachnick, B. G., & Fidell, L. S. (2007). *Using multivariate statistics (5th ed.)* Boston, MA: Allyn & Bacon/Pearson Education.

Teasdale, J. D., Moore, R. G., Hayhurst, H., Pope, M., Williams, S., & Segal, Z. V. (2002).

Metacognitive awareness and prevention of relapse in depression: Empirical evidence. *Journal of Consulting and Clinical Psychology*, 70(2), 275.

Teasdale, J. D., Segal, Z., & Williams, J. M. G. (1995). How does cognitive therapy prevent depressive relapse and why should attentional control (mindfulness) training help?

Behaviour Research and Therapy, 33(1), 25-39. doi:[http://dx.doi.org/10.1016/0005-7967\(94\)E0011-7](http://dx.doi.org/10.1016/0005-7967(94)E0011-7)

Van Bockstaele, B., Verschuere, B., Crombez, G., De Houwer, J., Tibboel, H., & Koster, E.

H. W. (2014). A review of current evidence for the causal impact of attentional bias on fear and anxiety. *Psychological Bulletin*, 140(3), 682-721.

Van Gordon, W., Shonin, E., Sumich, A., Sundin, E. C., & Griffiths, M. D. (2014).

Meditation awareness training (MAT) for psychological well-being in a sub-clinical sample of university students: A controlled pilot study. *Mindfulness*, 5(4), 381-391.

- Varela, F. J. (1996). Neurophenomenology: A methodological remedy for the hard problem. *Journal of consciousness studies*, 3(4), 330-349.
- Vuilleumier, P. (2005). How brains beware: Neural mechanisms of emotional attention. *Trends in Cognitive Sciences*, 9(12), 585.
- Waechter, S., Nelson, A. L., Wright, C., Hyatt, A., & Oakman, J. (2014). Measuring attentional bias to threat: Reliability of dot probe and eye movement indices. *Cognitive Therapy and Research*, 38(3), 313-333.
- Wiers, R. W., Gladwin, T. E., Hofmann, W., Salemink, E., & Ridderinkhof, K. R. (2013). Cognitive bias modification and cognitive control training in addiction and related psychopathology mechanisms, clinical perspectives, and ways forward. *Clinical Psychological Science*, 1(2), 192-212.
- Yuval, K., Zvielli, A., & Bernstein, A. (2016). Attentional bias dynamics and posttraumatic stress in survivors of violent conflict and atrocities new directions in clinical psychological science of refugee mental health. *Clinical Psychological Science*, , 2167702616649349.
- Zvielli, A., Amir, I., Goldstein, P., & Bernstein, A. (2015). Targeting biased emotional attention to threat as a dynamic process in time attention feedback awareness and control training (A-FACT). *Clinical Psychological Science*, , 2167702615588048.
- Zvielli, A., Bernstein, A., & Koster, E. H. (2014). Dynamics of attentional bias to threat in anxious adults: Bias towards and/or away? *PloS One*, 9(8), e104025.
doi:10.1371/journal.pone.0104025
- Zvielli, A., Vrijnsen, J. N., Koster, E. H. W., & Bernstein, A. (2016). **Attentional bias dynamics in remitted depression.** *Journal of Abnormal Psychology*, 125(6), 768-776.
doi:<http://dx.doi.org/10.1037/abn0000190>

Zvielli, A., Amir, I., Goldstein, P., & Bernstein, A. (2016). Targeting biased emotional attention to threat as a dynamic process in time: Attention feedback awareness and control training (A-FACT). *Clinical Psychological Science*, 4(2), 287-298.

doi:10.1177/2167702615588048

Zvielli, A., Bernstein, A., & Koster, E. H. W. (2015). Temporal dynamics of attentional bias. *Clinical Psychological Science*, 3(5), 772-788. doi:10.1177/2167702614551572