

## PAPER

# Compactness of Finite Union of Regular Patterns and Regular Patterns without Adjacent Variables

Naoto TAKETA<sup>†</sup>, *Nonmember*, Tomoyuki UCHIDA<sup>†</sup>, Takayoshi SHOUDAI<sup>††</sup>, Satoshi MATSUMOTO<sup>†††</sup>,  
Yusuke SUZUKI<sup>†</sup>, and Tetsuhiro MIYAHARA<sup>†</sup>, *Members*

**SUMMARY** A regular pattern is a string consisting of constant symbols and distinct variable symbols. The language  $L(p)$  of a regular pattern  $p$  is the set of all constant strings obtained by replacing all variable symbols in the regular pattern  $p$  with constant strings.  $\mathcal{RP}^k$  denotes the class of all sets consisting at most  $k$  ( $k \geq 2$ ) regular patterns. For sets of regular patterns  $P$  and  $Q$  which are in the class  $\mathcal{RP}^k$ , we write  $P \sqsubseteq Q$  if for any regular pattern  $p \in P$  there exists a regular pattern  $q \in Q$  that is a generalization of  $p$ . In 1998 Sato et al.[1] showed that the finite set  $S_2(P)$  of symbol strings is a characteristic set of  $L(P) = \bigcup_{p \in P} L(p)$ , where  $S_2(P)$  is obtained from  $P \in \mathcal{RP}^k$  by substituting variables with symbol strings of at most length 2. Sato et al.[1] also showed that  $\mathcal{RP}^k$  has compactness with respect to containment, if the number of constant symbols is greater than or equal to  $2k - 1$ . In this paper, we check the results of Sato et al.[1] and correct the error of the proof of their theorem. Further, we consider the set  $\mathcal{RP}_{NAV}^k$  of all non-adjacent regular patterns, which are regular patterns without adjacent variables, and show that the set  $S_2(P)$  obtained from a set  $P$  in the class  $\mathcal{RP}_{NAV}^k$  of at most  $k$  ( $k \geq 1$ ) non-adjacent regular patterns is a characteristic set of  $L(P)$ . Further we show that  $\mathcal{RP}_{NAV}^k$  has compactness with respect to containment if the number of constant symbols is greater than or equal to  $k + 2$ . Thus we show that we can design an efficient learning algorithm of a finite union of pattern languages of non-adjacent regular patterns with the number of constant symbols which is smaller than the case of regular patterns.

**key words:** Regular Pattern Language, Compactness

## 1. Introduction

A pattern is a string consisting of constant symbols and variable symbols. For example, we consider constant symbols  $a, b, c$  and variable symbols  $x, y$ , then  $axbxcy$  is a pattern.  $\mathcal{P}$  denotes the set of all patterns. For a pattern  $p \in \mathcal{P}$ , the pattern language generated by  $p$ , denoted by  $L(p)$ , or simply called a pattern language, is the set of all strings obtained by replacing all variable symbols with constant symbol strings, where the same variable symbol is replaced by the same constant string. For example the pattern language  $L(axbxcy)$  generated by the above pattern  $axbxcy$  denotes  $\{aubucw \mid u \text{ and } w \text{ are constant strings that are not } \varepsilon\}$ . A pattern where each variable symbol appears at most once is called a *regular pattern*. For example, a pattern  $axbxcy$  is not a regular pattern, but a pattern  $axbzcxy$  with variable symbols  $x, y, z$  is a regular pattern.  $\mathcal{RP}$  denotes the set of

all regular patterns. If a pattern  $p \in \mathcal{P}$  is obtained from a pattern  $q \in \mathcal{P}$  by replacing variable symbols in  $q$  with patterns, we say that  $q$  is a *generalization* of  $p$  and denote this by  $p \preceq q$ . For example, a pattern  $q = axz$  is a generalization of a pattern  $p = axbxcy$ , because  $p$  is obtained from  $q$  by replacing the variable  $z$  in  $q$  with a pattern  $bxcy$ . So we write  $p \preceq q$ . For patterns  $p, q \in \mathcal{P}$ , it is obvious that  $p \preceq q$  implies  $L(p) \subseteq L(q)$ . But, the converse, that is, the statement that  $L(p) \subseteq L(q)$  implies  $p \preceq q$  does not always hold. With respect to this statement, Mukouchi[2] showed that if the number of constant symbols is greater than or equal to 3, for any regular pattern  $p, q \in \mathcal{RP}$ ,  $L(p) \subseteq L(q)$  implies  $p \preceq q$ .

We denote by  $\mathcal{RP}^+$  the class of all non-empty finite sets of regular patterns and by  $\mathcal{RP}^k$  the class of at most  $k$  ( $k \geq 2$ ) regular patterns. For a set of regular patterns  $P \in \mathcal{RP}^k$  we define  $L(P) = \bigcup_{p \in P} L(p)$  and consider the class  $\mathcal{RPL}^k$  of regular pattern languages of  $\mathcal{RP}^k$ , where  $\mathcal{RPL}^k = \{L(P) \mid P \in \mathcal{RP}^k\}$ . Let  $P, Q \in \mathcal{RP}^k$  and  $Q = \{q_1, \dots, q_k\}$ . We denote by  $P \sqsubseteq Q$  that for any regular pattern  $p \in P$  there exists a regular pattern  $q_i$  such that  $p \preceq q_i$  holds. From definition, it is obvious that  $P \sqsubseteq Q$  implies  $L(P) \subseteq L(Q)$ . Then Sato et al.[1] shows that if  $k \geq 3$  and the number of constant symbols is  $2k - 1$  then the finite set  $S_2(P)$  of constant symbols obtained from  $P \in \mathcal{RP}^k$  by substituting variable symbols with constant strings of at most 2 length is a characteristic set of  $L(P)$ , that is, for any regular pattern language  $L' \in \mathcal{RPL}^k$ ,  $S_2(P) \subseteq L'$  implies  $L(P) \subseteq L'$ . Thus they show that the following three statements: (i)  $S_2(P) \subseteq L(Q)$ , (ii)  $P \sqsubseteq Q$  and (iii)  $L(P) \subseteq L(Q)$  are equivalent. But the Lemma14 [1], which is used in this results, contains an error. In this paper we correct this lemma and give a correct proof showing the equivalence of the three statements shown in [1]. Sato et al.[1] shows that  $\mathcal{RP}^k$  has compactness with respect to containment if the number of constant symbols is greater than or equal to  $2k - 1$ . On the contrary to this result, we show that the set  $S_2(P)$  obtained from a set  $P$  in the class  $\mathcal{RP}_{NAV}^k$  of at most  $k$  ( $k \geq 1$ ) regular patterns having non-adjacent variables is a characteristic set of  $L(P)$ . Further, we show that if the number of constant symbols is greater than or equal to  $k + 2$  then  $\mathcal{RP}_{NAV}^k$  has compactness with respect to containment. In Table 1 we summarize the all results in this paper.

The results of this paper suggest efficient learning algorithms for the sets of regular patterns representing finite

Manuscript received January 1, 2015.

Manuscript revised January 1, 2015.

<sup>†</sup>Graduate School of Information Sciences, Hiroshima City University

<sup>††</sup>Department of Computer Science and Engineering, Fukuoka Institute of Technology

<sup>†††</sup>Faculty of Science, Tokai University

DOI: 10.1587/trans.E0.??.

**Table 1** The conditions of the number of constant symbols with respect to the compactness of inclusion

$k$	2	$\geq 3$
$\mathcal{RP}^k$	$\geq 4$	$\geq 2k - 1$
$\mathcal{RP}_{NAV}^k$	$\geq k + 2$	

unions of languages and the sets of regular patterns having non-adjacent variables.

This paper is organized as follows. In Sect.2 as preparations, we give definitions of pattern languages, regular pattern languages and compactness, and then introduce the results of Sato et al.[1]. In Sect.3, we show that  $S_2(P)$  is a characteristic set of  $L(P)$  in  $\mathcal{RPL}^k$  and  $\mathcal{RP}^k$  has compactness with respect to containment. In Sect.4, we propose regular patterns having non-adjacent variables, show that  $S_2(P)$  obtained from a set  $P$  in  $\mathcal{RP}_{NAV}^k$  is a characteristic set of  $L(P)$ , and  $\mathcal{RP}_{NAV}^k$  has compactness with respect to containment.

## 2. Preliminaries

Let  $\Sigma$  be a non-empty finite set of constant symbols. Let  $X$  be an infinite set of variable symbols such that  $\Sigma \cap X = \emptyset$  holds. Then, a *string* on  $\Sigma \cup X$  is a sequence of symbols in  $\Sigma \cup X$ . Particularly, the string having no symbol is called the *empty string* and is denoted by  $\varepsilon$ . We denote by  $(\Sigma \cup X)^*$  the set of all strings on  $\Sigma \cup X$  and by  $(\Sigma \cup X)^+$  the set of all strings on  $\Sigma \cup X$  except  $\varepsilon$ , i.e.,  $(\Sigma \cup X)^+ = (\Sigma \cup X)^* \setminus \{\varepsilon\}$ . A *pattern* on  $\Sigma \cup X$  is a string in  $(\Sigma \cup X)^*$ . Note that the empty string  $\varepsilon$  is a pattern on  $\Sigma \cup X$ . A pattern  $p$  is said to be *regular* if each variable symbol appears at most once in  $p$ . The length of  $p$ , denote by  $|p|$ , is the number of symbols in  $p$ . Note that  $|\varepsilon| = 0$  holds. The set of all patterns and regular patterns are denoted by  $\mathcal{P}$  and  $\mathcal{RP}$ , respectively. For a set  $S$ , we denote by  $\#S$  the number of elements in  $S$ . Let  $p, q$  be strings. If  $p$  and  $q$  are equal as strings, we denote it by  $p = q$ . We denote by  $p \cdot q$  the string obtained from  $p$  and  $q$  by concatenating  $q$  after  $p$ . Note that for a string  $p$  and the empty string  $\varepsilon$ ,  $p \cdot \varepsilon = \varepsilon \cdot p = p$ . A substitution  $\theta$  is a mapping from  $(\Sigma \cup X)^*$  to  $(\Sigma \cup X)^*$  such that (1)  $\theta$  is a homomorphism with respect to string concatenation, i.e.,  $\theta(p \cdot q) = \theta(p) \cdot \theta(q)$  holds for patterns  $p$  and  $q$ , (2)  $\theta(\varepsilon) = \varepsilon$  holds, (3) for each constant symbol  $a \in \Sigma$ ,  $\theta(a) = a$  holds, and (4) for each variable symbol  $x \in X$ ,  $|\theta(x)| \geq 1$  holds. Let  $x_1, \dots, x_n$  are variable symbols and  $p_1, \dots, p_n$  non-empty patterns. The notation  $\{x_1 := p_1, \dots, x_n := p_n\}$  denotes a substitution that replaces each variable symbol  $x_i$  with a non-empty pattern  $p_i$  for  $i \in \{1, \dots, n\}$ . For a pattern  $p$  and a substitution  $\theta = \{x_1 := p_1, \dots, x_n := p_n\}$ , we denote by  $p\theta$  a new pattern obtained from  $p$  by replacing variable symbols  $x_1, \dots, x_n$  in  $p$  with patterns  $p_1, \dots, p_n$  according to  $\theta$ , respectively. For a pattern  $p$  and  $q$ , the pattern  $q$  is a *generalization* of  $p$ , or  $p$  is an *instance* of  $q$ , denoted by  $p \preceq q$ , if there exists a substitution  $\theta$  such that  $p = q\theta$  holds. If  $p \preceq q$  and  $q \preceq p$  hold, we denote it by  $p \equiv q$ . The notation  $p \equiv q$  means that  $p$  and  $q$  are equal as strings except for variable symbols. For a pattern  $p$ , the *pattern language*

of  $p$ , denoted by  $L(p)$ , is the set  $\{w \in \Sigma^* \mid w \preceq p\}$ . For patterns  $p$  and  $q$ , it is clear that  $L(p) = L(q)$  if  $p \equiv q$ , and  $L(p) \subseteq L(q)$  if  $p \preceq q$ . Note that  $L(\varepsilon) = \{\varepsilon\}$ . In particular, if  $p$  is a regular pattern, we say that  $L(p)$  is a *regular pattern language*. The set of all pattern languages and regular patterns languages are denoted by  $\mathcal{PL}$  and  $\mathcal{RPL}$ , respectively.

**Lemma 1** (Mukouchi[2]): Let  $p$  and  $q$  be regular patterns. Then  $p \preceq q$  if and only if  $L(p) \subseteq L(q)$ .

Next, we consider unions of pattern languages. The class of all non-empty finite subsets of  $\mathcal{P}$  is denoted by  $\mathcal{P}^+$ , i.e.,  $\mathcal{P}^+ = \{P \subseteq \mathcal{P} \mid 0 < \#P < \infty\}$ . For a positive integer  $k$  ( $k > 0$ ), the class of non-empty sets consisting of at most  $k$  patterns, i.e.,  $\mathcal{P}^k = \{P \subseteq \mathcal{P} \mid 0 < \#P \leq k\}$ . We denote by  $\mathcal{PL}^k$  the class of unions of at most  $k$  pattern languages, i.e.,  $\mathcal{PL}^k = \{L(P) \mid P \in \mathcal{P}^k\}$ , where  $L(P) = \bigcup_{p \in P} L(p)$ . In a similar way, we also define  $\mathcal{RP}^+$ ,  $\mathcal{RP}^k$  and  $\mathcal{RPL}^k$ . For  $P, Q$  in  $\mathcal{P}^+$ , the notation  $P \sqsubseteq Q$  means that for any  $p \in P$  there is a pattern  $q \in Q$  such that  $p \preceq q$  holds. It is clear that  $P \sqsubseteq Q$  implies  $L(P) \subseteq L(Q)$ . However, the converse is not valid in general.

**Definition 1:** Let  $C$  be a class of languages,  $L$  a language in  $C$  and  $S$  a non-empty finite subset of  $L$ . We say that  $S$  is a *characteristic set* of  $L$  within  $C$  if for any  $L' \in C$ ,  $S \subseteq L'$  implies  $L \subseteq L'$ .

Let  $n$  be a positive integer and  $p$  a regular pattern. We denote by  $S_n(p)$  the set of all strings in  $\Sigma^*$  obtained by replacing all variable symbols in  $p$  with strings in  $\Sigma^+$  of length at most  $n$ . Moreover, for a positive integer  $n$  and a set  $P \in \mathcal{RP}^+$ , let  $S_n(P) = \bigcup_{p \in P} S_n(p)$ . It is clear that  $S_n(P) \subseteq S_{n+1}(P) \subseteq L(P)$  for any positive integer  $n$ .

**Theorem 1** (Sato et al.[1]): Let  $k$  be a positive integer and  $P \in \mathcal{RP}^k$ . Then, there exists a positive integer  $n$  such that  $S_n(P)$  is a characteristic set of  $L(P)$  within  $\mathcal{RPL}^k$ .

Sato et al.[1] showed that 2 is sufficient for the number  $n$  in the theorem above, under the assumption that the number of constants is not less than  $2k - 1$ . Hence, in this paper, we consider a characteristic set  $S_2(P)$  of  $L(P)$  within  $\mathcal{RPL}^k$ .

**Theorem 2** (Sato et al.[1]): Let  $p, q, p_1, p_2, q_1, q_2, q_3$  be regular patterns and  $x$  a variable symbol with  $p = p_1xp_2$  and  $q = q_1q_2q_3$ . Then  $p \preceq q$  if the following three conditions are holds:

- (i)  $p_1 \preceq q_1q_2$ , (ii)  $p_2 \preceq q_2q_3$ ,
- (iii)  $q_2$  contains at least one variable symbol.

**Lemma 2** (Sato et al.[1]): Suppose  $\#\Sigma \geq 3$ . Let  $p, p_1, p_2, q$  be regular patterns and  $x$  a variable symbol with  $p = p_1xp_2$ . Let  $a, b$  and  $c$  be mutually distinct constant symbols. If  $p_1ap_2 \preceq q$ ,  $p_1bp_2 \preceq q$  and  $p_1cp_2 \preceq q$ , then  $p \preceq q$  holds.

**Lemma 3** (Sato et al.[1]): Suppose  $\#\Sigma \geq 3$ . Let  $p_1, p_2, q_1, q_2$  be regular patterns and  $x$  a variable symbol. Let  $a, b$  be constant symbols with  $a \neq b$  and  $w$  a string in  $\Sigma^*$ . Let  $p = p_1AwBp_2$  and  $q = q_1AwBq_2$  be regular patterns that

satisfy the following three conditions:

- (i)  $p_1 \preceq q_1$ ,
- (ii)  $p_2 \preceq q_2$ ,
- (iii)  $(A, B) \in \{(a, b), (b, a)\}$ .

If  $p\{x := a\} \preceq q$  and  $p\{x := b\} \preceq q$ , then we have  $p \not\preceq q$ .

From Lemma ??, the following lemma holds.

**Theorem 3** (Sato et al.[1]): Let  $\#\Sigma \geq 2k + 1$ ,  $P \in \mathcal{RP}^+$  and  $Q \in \mathcal{RP}^k$ . Then, the following (i), (ii) and (iii) are equivalent:

- (i)  $S_1(P) \subseteq L(Q)$ , (ii)  $P \sqsubseteq Q$ , (iii)  $L(P) \subseteq L(Q)$ .

Example 1 in [1] is given as a counter-example of Theorem 3.

From Theorem 3, we have the following corollary.

**Corollary 1** (Sato et al.[1]): Let  $\#\Sigma \geq 3$  and  $p, q$  regular patterns. Then, the following (i), (ii) and (iii) are equivalent:

- (i)  $S_1(p) \subseteq L(q)$ , (ii)  $p \preceq q$ , (iii)  $L(p) \subseteq L(q)$ .

### 3. Compactness for Sets of Regular Patterns

In this section, we define the compactness of sets of regular patterns, formally. Then, if  $\#\Sigma \geq 2k - 1$  holds, we show that  $\mathcal{RP}^k$  has compactness with respect to the containment.

**Definition 2:** Let  $C$  be a subset of  $\mathcal{RP}^+$ . For any regular pattern  $p \in \mathcal{RP}$  and any set  $Q \in C$ , the set  $C$  said to have *compactness with respect to containment* if there exists a regular pattern  $q \in Q$  such that  $L(p) \subseteq L(q)$  holds if  $L(p) \subseteq L(Q)$  holds.

**Lemma 4** (Sato et al.[1]): Let  $\Sigma$  be an alphabet with  $\#\Sigma \geq 3$  and  $p, q$  regular patterns on  $\Sigma$ . Let  $D$  be the set of either (i) or (ii) of regular patterns on  $\Sigma$  below: Assume that  $a \neq b$  and that a variable symbol  $y$  does not appear in  $p$ .

- (i)  $\{ay, by\}$  (ii)  $\{ya, yb\}$ .

Then, if  $p\{x := r\} \preceq q$  for all  $r \in D$ , then  $p\{x := xy\} \preceq q$ .

**Proof.** It is obvious if no variable symbol appears in  $p$ . Therefore, let  $p = p_1xp_2$ , where  $p_1, p_2$  are regular patterns and  $x$  is a variable symbol. We assume that  $p\{x := xy\} \not\preceq q$  in order to derive the contradictions.

(i) Case of  $D = \{ay, by\}$  ( $a \neq b$ ):

Since  $p\{x := xy\} \not\preceq q$ ,  $p_1ayp_2 \preceq q$  and  $p_1byp_2 \preceq q$ , there exist regular patterns  $q_1, q_2$  on  $\Sigma$  such that  $q = q_1ay_1wby_2q_2$  or  $q = q_1by_1way_2q_2$  for some variable symbols  $y_1, y_2$  ( $y_1 \neq y_2$ ) and a constant string  $w$  ( $|w| \geq 0$ ) from Theorem 2. When  $q = q_1ay_1wby_2q_2$  holds, the following four conditions (1), (2), (1'), (2') holds:

- (1)  $p_1 \preceq q_1$
- (1')  $p_2 \preceq wby_2q_2$  or  $p_2 \preceq y'wby_2q_2$  ( $y' \in X$ )
- (2)  $p_1 \preceq q_1ay_1w$

$$(2') p_2 \preceq q_2 \text{ or } p_2 \preceq y''q_2 \ (y'' \in X)$$

From the above condition (2), there exist regular patterns  $p'_1, p''_1$  such that  $p_1 = p'_1p''_1$ ,  $p'_1 \preceq q_1a$  and  $p''_1 \preceq y_1w$  hold. Therefore, since  $p = p_1xp_2 = p'_1p''_1xp_2$ , if  $p_2 \preceq wby_2q_2$  holds,  $p \preceq q_1ap''_1xwby_2q_2 \equiv q\{y_1 := p''_1x\}$  holds. Otherwise  $p_2 \preceq y'wby_2q_2$ ,  $p \preceq q_1ap''_1xy'wby_2q_2 = q\{y_1 := p''_1xy'\}$  holds. Hence,  $p \preceq q$  holds. This contradicts the assumption.

(ii) Case of  $D = \{ya, yb\}$  ( $a \neq b$ ): By reversing the strings of  $p$  and  $q$ , we can prove that  $p\{x := xy\} \preceq q$  holds, in a similar way as (i).

In Lemma 14 (ii) of [1], they stated that, when  $\#\Sigma \geq 3$ , for regular patterns  $p, q$ , if  $p\{x := r\} \preceq q$  for any  $r \in D$ , then  $p\{x := xy\} \preceq q$  holds, where  $D = \{a_1b_1, a_2b_2, a_3b_3\}$  ( $a_i \neq a_j$  and  $b_i \neq b_j$  for each  $i, j$  ( $i \neq j, 1 \leq i, j \leq 3$ )). Unfortunately, there exists the following counterexample of Lemma 14 (ii) of [1].

**Example 1:** Assume that  $a_1 = b_3$  and  $a_2 = b_1$  hold. Let  $p = cb_3a_1b_1b_3x'a_1b_1b_3a_2c$  and  $q = xb_3a_1b_1b_3a_2y$ . We have  $p\{x' := a_1b_1\} \preceq q$ ,  $p\{x' := a_2b_2\} \preceq q$ , and  $p\{x' := a_3b_3\} \preceq q$ , because  $p\{x' := a_1b_1\} = q\{x := cb_3a_1b_1, y := b_3a_2c\}$ ,  $p\{x' := a_2b_2\} = q\{x := c, y := b_2a_1b_1b_3a_2c\}$ , and  $p\{x' := a_3b_3\} = q\{x := cb_3a_1b_1b_3a_3, y := c\}$  hold. However, it is clear that  $p\{x := xy\} \not\preceq q$  holds.

The following Lemma 5 corrects completely mistakes of Lemma 14 (ii) of [1].

**Lemma 5:** Let  $\Sigma$  be an alphabet with  $\#\Sigma \geq 3$ ,  $p, q$  regular patterns on  $\Sigma$ . Let  $D$  be the following set of constant strings on  $\Sigma$  whose lengths are just 2:

**Lemma 6:** Let  $\Sigma$  be an alphabet with  $\#\Sigma \geq 3$  and  $p, q$  regular patterns on  $\Sigma \cup X$ . Let  $D$  be the following set of regular patterns on  $\Sigma \cup X$ . Then, if  $p\{x := r\} \preceq q$  for all  $r \in D$ , then  $p\{x := xy\} \preceq q$ :

$$D = \{ya, bc, dy\} \ (b \notin \{a, d\} \text{ and } c \notin \{a, d\}).$$

**Proof.** It is obvious if no variable symbol appears in  $p$ . Thus, for a variable symbol  $x \in X$ , let  $p = p_1xp_2$ , where  $p_1, p_2$  are regular patterns on  $\Sigma \cup X$ . We assume that  $p\{x := xy\} \not\preceq q$  in order to derive the contradiction.

Since  $p\{x := xy\} \not\preceq q$ , the regular pattern  $q$  can be expressed in one of the following forms:

- (i)  $q = q_1AwBw'Cq_2$ , where  $\{A, B, C\} = \{y_1a, bc, dy_2\}$  for distinct variable symbols  $y_1, y_2 \in X$ , and each of  $w, w'$  is either an empty string or a regular pattern on  $\Sigma \cup X$ .
- (ii)  $q = q_1AwBq_2$ , where  $\{A, B\} = \{y_1ay_2, bc\}$  ( $a = d$ ) for distinct variable symbols  $y_1, y_2 \in X$ , and  $w$  is either an empty string or a regular pattern on  $\Sigma \cup X$ .
- (iii)  $q = q_1AwBq_2$ , where  $\{A, B\} = \{dy_1a, bc\}$  for a variable symbol  $y_1 \in X$  and  $w$  is either an empty string or a regular pattern on  $\Sigma \cup X$ .

Firstly, we will prove that for the case (i),  $p\{x := xy\} \preceq q$  holds.

*Claim 1.*  $B \notin \{y_1a, dy_2\}$ .

*Proof of Claim 1.* Suppose that  $(A, B, C) = (dy_2, y_1a, bc)$ . For some  $y'_1, y'_2 \in X$ , the following conditions hold:

- $$\begin{aligned} (1) \ p_1 \preceq q_1 & \quad (1') \ p_2 \preceq wy_1aw'bcq_2, \text{ or} \\ & \quad p_2 \preceq y'_2wy_1aw'bcq_2 \\ (2) \ p_1 \preceq q_1dy_2w, \text{ or} & \quad (2') \ p_2 \preceq w'bcq_2 \\ & \quad p_1 \preceq q_1dy_2wy'_1 \\ (3) \ p_1 \preceq q_1dy_2wy_1aw' & \quad (3') \ p_2 \preceq q_2 \end{aligned}$$

When  $p_2 \preceq wy_1aw'bcq_2$  of (1') holds, let  $q'_1 = q_1dy_2$ ,  $q'_2 = wy_1aw'$ ,  $q'_3 = bcq_2$ . Since  $p_1 \preceq q_1dy_2wy_1aw'$  holds from (3),  $p_1 \preceq q'_1q'_2$  and  $p_2 \preceq q'_2q'_3$  hold and  $q'_2$  contains a variable symbol. When  $p_2 \preceq y'_2wy_1aw'bcq_2$  of (1') holds, let  $q'_1 = q_1d$ ,  $q'_2 = y_2wy_1aw'$ ,  $q'_3 = bcq_2$ . Since  $p_1 \preceq q_1dy_2wy_1aw'$  holds from (3),  $p_1 \preceq q'_1q'_2$  and  $p_2 \preceq q'_2q'_3$  hold and  $q'_2$  contains a variable symbol. In both cases, from Theorem 2,  $p \preceq q$  holds. It contradicts the assumption.

Similarly, we can show that any case of  $(A, B, C) = (y_1a, dy_2, bc)$ ,  $(bc, y_1a, dy_2)$ ,  $(bc, dy_2, y_1a)$  contradicts the assumption. Therefore, we have  $B \notin \{y_1a, dy_2\}$ . (*End of Proof of Claim 1*)

*Claim 2.*  $(A, B, C) = (y_1a, bc, dy_2)$ .

*Proof of Claim 2.* From *Claim 1*, we have  $B = bc$ . Suppose that  $(A, B, C) = (dy_2, bc, y_1a)$  holds. Then, the following conditions hold: for  $y'_1, y'_2 \in X$ ,

- $$\begin{aligned} (1) \ p_1 \preceq q_1 & \quad (1') \ p_2 \preceq wbcw'y_1aq_2, \text{ or} \\ & \quad p_2 \preceq y'_2wbcw'y_1aq_2 \\ (2) \ p_1 \preceq q_1dy_2w & \quad (2') \ p_2 \preceq w'y_1aq_2 \\ (3) \ p_1 \preceq q_1dy_2wbcw', \text{ or} & \quad (3') \ p_2 \preceq q_2 \\ & \quad p_1 \preceq q_1dy_2wbcw'y'_1 \end{aligned}$$

From  $p_1 \preceq q_1dy_2w$  of (2), for some  $p'_1$  and  $p''_1$ ,  $p_1$  is expressed as  $p'_1p''_1$ , where  $p'_1 \preceq q_1d$  and  $p''_1 \preceq y_2w$ . When  $p_2 \preceq wbcw'y_1aq_2$  of (1'), we have  $p = p_1xp_2 = p'_1p''_1xp_2 \preceq q_1dp'_1xwbcw'y_1aq_2 = q\{y_2 := p''_1x\}$ . When  $p_2 \preceq y'_2wbcw'y_1aq_2$  of (1'), we have  $p = p_1xp_2 = p'_1p''_1xp_2 \preceq q_1dp'_1xy'_2wbcw'y_1aq_2 = q\{y_2 := p''_1xy'_2\}$ . Thus,  $p\{x := xy\} \preceq q\{y_2 := p''_1xyy'_2\}$  holds. It contradicts the assumption. Therefore, we have  $(A, B, C) = (y_1a, bc, dy_2)$ . (*End of Proof of Claim 2*)

From *Claim 2*,  $q$  is expressed as  $q_1y_1awbcw'dy_2q_2$ , where  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ . If  $p\{x := xy\} \not\preceq q$  holds, we have the following conditions: for  $y'_1, y'_2 \in X$ ,

- $$\begin{aligned} (1) \ p_1 \preceq q_1 \text{ or } p_1 \preceq q_1y'_1 & \quad (1') \ p_2 \preceq wbcw'dy_2q_2 \\ (2) \ p_1 \preceq q_1y_1aw & \quad (2') \ p_2 \preceq w'dy_2q_2 \\ (3) \ p_1 \preceq q_1y_1awbcw' & \quad (3') \ p_2 \preceq q_2 \text{ or } p_2 \preceq y'_2q_2 \end{aligned}$$

*Claim 3.*  $w$  and  $w'$  contains no variable symbol.

*Proof of Claim 3.* Let  $q'_1 = q_1y_1a$ ,  $q'_2 = wbcw'$ , and  $q'_3 = dy_2q_2$ . From (1') and (3),  $p_1 \preceq q'_1q'_2$  and  $p_2 \preceq q'_2q'_3$ . If  $q'_2$  contains a variable symbol, from Theorem 2,  $p \preceq q$  holds.

It contradicts the assumption. Therefore,  $w$  and  $w'$  contains no variable symbol. (*End of Proof of Claim 3*)

From *Claim 3*,  $w$  and  $w'$  are strings consisting of symbols in  $\Sigma$ . Note that from (1') and (2'),  $wbcw'd$  and  $w'd$  are prefixes of  $p_2$ , and from (2) and (3),  $awbcw'$  and  $aw$  are suffixes of  $p_1$ .

If  $|w| = |w'|$ , then  $c = a$  holds. It contradicts the condition  $c \neq a$ .

If  $|w| = |w'| + 1$ , then  $b = a$  holds. It contradicts the condition  $b \neq a$ .

If  $|w| = |w'| + 2$ , since  $awbcw'$  and  $aw$  are suffixes of  $p_1$ , and since  $|w| \geq 2$ ,  $a$  is a suffix of  $w$ . From (1') and (2'), since  $wbcw'd$  and  $w'd$  are prefixes of  $p_2$ , we have  $w = w'da$ . Since  $awbcw'$  and  $aw$  are suffixes of  $p_1$ , we have  $w = bcw'$ . Therefore,  $w'da = bcw'$  holds. We show the next claim:

*Claim 4.* Let  $w'$  be a string of constant symbols in  $\Sigma$  and  $a, b, c, d$  constant symbols in  $\Sigma$ . Then, if  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ , then  $w'da \neq bcw'$  holds.

*Proof of Claim 4.* When  $|w'| = 0, 1, 2, 3$ , it is easy to see that  $w'da = bcw'$  does not satisfy the conditions  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ . Therefore,  $w'da \neq bcw'$  holds. Let  $n = |w'|$ . When  $n \geq 4$ , we assume that for any string  $w''$  with  $|w''| < n$ , if  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ ,  $w''da \neq bcw''$  holds. Since the string  $w'$  has a prefix  $bc$  and a suffix  $da$ , there exists a string  $w''$  with  $|w''| \geq 0$  such that  $w' = bcw''da$  holds. Since  $w'da = bcw''dada$  and  $bcw' = bcbcw''da$ , if  $w'da = bcw'$  holds, we have  $bcw''dada = bcbcw''da$ . Then we conclude that  $w''da = bcw''$ . It contradicts the induction hypothesis. Thus,  $w'da \neq bcw'$  holds. From the above, for any string  $w'$  with  $|w'| \geq 0$ , if  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ ,  $w'da \neq bcw'$  holds. (*End of Proof of Claim 4*)

Thus, the case of  $|w| = |w'| + 2$  contradicts *Claim 4*.

If  $|w| \geq |w'| + 3$ , from (2) and (3), there exists a string  $w''$  of length  $|w| - |w'| - 2$  such that  $w = w''bcw'$  holds. Moreover, from (2) and (3), since  $|aw| < |wbcw'|$  and  $aw = aw''bcw'$ ,  $aw''$  is a suffix of  $w$ . On the other hand, from (1') and (2'),  $w'd$  is a prefix of  $w$ . Since  $|w'd| + |aw''| = |w'| + |w''| + 2 = |w|$ , we have  $w = w'daw''$ . Therefore,  $w'daw'' = w''bcw'$  holds.

*Claim 5.* Let  $w', w''$  be strings of constant symbols in  $\Sigma$  and  $a, b, c, d$  constant symbols in  $\Sigma$ . Then, if  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ , then  $w'daw'' \neq w''bcw'$  holds.

*Proof of Claim 5.* We assume that the following equation holds:




$$w'daw'' = w''bcw' \quad (1)$$

We prove this claim by an induction on  $|w'| + |w''|$ . W.l.o.g., we suppose that  $|w'| \geq |w''|$  holds.

- (i)  $|w'| \geq 0$  and  $|w''| = 0$ : We have  $w'da = bcw'$  ( $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ ). It contradicts *Claim 4*.

We assume that for constant strings  $u$  and  $v$  with  $|u| + |v| < |w'| + |w''|$ ,  $vbcu \neq udav$  holds. We partition the relations between  $|w'|$  and  $|w''|$  into the following four parts:

- (ii)  $0 < |w''| \leq |w'| \leq |w''| + 1$ : When either  $|w'| = |w''|$  or

**Fig. 1** Subcase  $|w'| = |w''|$  of (ii) of Claim 5 (Lemma 6)**Fig. 2** Subcase  $|w'| = |w''| + 1$  of (ii) of Claim 5 (Lemma 6)**Fig. 3** Case  $|w''| + 2 \leq |w'| \leq 2|w''| - 1$  of (iii) of Claim 5 (Lemma 6)

$|w'| = |w''| + 1$ , Eq. 1 is depicted as shown in Figs. 1, 2. Trivially, these cases contradict the conditions  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ .

- (iii)  $|w''| + 2 \leq |w'| \leq 2|w''| - 1$ : On Eq. 1, since  $|w'daw''| = |w''bcw'| = |w'| + |w''| + 2$ , a suffix of  $w'$  overlaps with a prefix of  $w''$  as shown in Fig. 3. That is, there exists a constant string  $u$  of length  $2|w'| - (|w'| + |w''| + 2) = |w'| - |w''| - 2$  such that  $u$  is a prefix and a suffix of  $w'$ . Since  $uda$  is of length  $|w'| - |w''|$ ,  $uda$  is also a prefix of  $w'$ . Similarly,  $bcu$  is also a suffix of  $w'$ . Since  $|w'| - (|uda| + |bcu|) = 2|w''| - |w'| \geq 1$ , there exist a constant string  $v$  of length  $2|w''| - |w'|$  such that  $w' = udavbcu$  holds. Since  $w''$  is a suffix of  $w'$  and  $|vbcu| = (2|w''| - |w'|) + 2 + (|w'| - |w''| - 2) = |w''|$ , we have  $w'' = vbcu$ . Similarly, we have  $w'' = udav$ . Thus, we have a new equation  $vbcu = udav$ . Since  $|u| = |w'| - |w''| - 2 \leq |w''| - 3 < |w''|$  and  $|v| = 2|w''| - |w'| < |w'|$ , i.e.,  $|u| + |v| < |w'| + |w''|$  holds, it contradicts the induction hypothesis on  $|u| + |v|$ . Therefore, the claim holds.






- (iv)  $2|w''| \leq |w'| \leq 2|w''| + 3$ : When  $|w'| = 2|w''|$ , we easily see that  $w' = w''w''$ . Therefore,  $w''da = bcw''$  holds as shown in Fig. 4. It contradicts Claim 3. When  $|w'| = 2|w''| + i$  ( $i = 1, 2, 3$ ), Eq. 1 is depicted as shown in Figs. 5, 6, and 7. Trivially, these cases contradict the conditions  $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ .

- (v)  $2|w''| + 4 \leq |w'|$ : Since the strings  $w''bc$  and  $adw''$  are a prefix and a suffix of  $w'$ , respectively, and  $|w''bc| + |adw''| = 2|w''| + 4$ , there exists a string  $u$  with  $|u| \geq 0$  such that  $w' = w''bcudaw''$  holds. From Eq. 1,  $w''bcudaw''daw'' = w''bcw''bcudaw''$ , i.e.,  $udaw'' = w''bcu$  holds as shown in Fig. 8. Let  $v = w''$ . Since  $|u| + |v| = |w'| - (2|w''| + 4) + |w''| < |w'| + |w''|$ , it contradicts the induction hypothesis on  $|u| + |v|$ . Therefore, the claim holds.

From the above, we conclude that  $w'daw'' \neq w''bcw'$  holds. (End of Proof of Claim)

Thus, the case of  $|w| \geq |w'| + 3$  contradicts Claim 5.

Next, we suppose that  $|w| < |w'|$  holds. Note again that

**Fig. 4** Subcase  $|w'| = 2|w''|$  of (iv) of Claim 5 (Lemma 6)**Fig. 5** Subcase  $|w'| = 2|w''| + 1$  of (iv) of Claim 5 (Lemma 6)**Fig. 6** Subcase  $|w'| = 2|w''| + 2$  of (iv) of Claim 5 (Lemma 6)**Fig. 7** Subcase  $|w'| = 2|w''| + 3$  of (iv) of Claim 5 (Lemma 6)**Fig. 8** Case  $2|w''| + 4 \leq |w'|$  of (v) of Claim 5 (Lemma 6)

from (1') and (2'),  $wbcw'd$  and  $w'd$  are prefixes of  $p_2$ , and from (2) and (3),  $awbcw'$  and  $aw$  are suffixes of  $p_1$ .

If  $|w'| = |w| + 1$ , since  $|wbc| = |w'd|$ , we have  $c = d$ . This contradicts the condition  $c \neq d$ .

If  $|w'| = |w| + 2$ , since  $|wbc| = |w'|$ ,  $bc$  is a suffix of  $w'$ . Moreover, since  $w'd$  is a prefix of  $wbcw'$ ,  $d$  is the first symbol of  $w'$ . Since  $aw$  is a suffix of  $w'$  and  $|w'| = |aw| + 1$ ,  $a$  is the second symbol of  $w'$ . Therefore, we have  $w' = wbc = daw$ . This contradicts Claim 4.

If  $|w'| \geq |w| + 3$ , there exists a string  $w''$  with  $|w''| \geq 1$  such that  $w' = wbcw''$  holds. Then, since  $wbcw'd$  and  $w'd = wbcw''d$  are prefixes of  $p_2$ ,  $w''d$  is a prefix of  $w'$ . Since  $w'$  and  $aw$  are suffixes of  $p_1$  and  $|w'| = |wbcw''| = |w| + |w''| + 2 > |aw|$ ,  $aw$  is a suffix of  $w'$ . Since  $|w''d| + |aw| = |w'|$ , we have  $w' = w''daw$ . Therefore, we have  $w' = wbcw'' = w''daw$ . This contradicts Claim 5.

Thus, the case of  $A = y_1a$ ,  $B = bc$ , and  $C = dy_2$  implies the contradictions.

From the above, we conclude that if  $p\{x := r\} \preceq q$  for all  $r = \{ya, bc, dy\}$  ( $b \notin \{a, d\}$  and  $c \notin \{a, d\}$ ), then  $p\{x := xy\} \preceq q$  holds.

**Lemma 7:** Let  $\Sigma$  be an alphabet with  $|\Sigma| \geq 3$  and  $p, q$  regular patterns on  $\Sigma$ . Let  $D$  be one of the following sets (i), (ii) of regular patterns on  $\Sigma$ . Then, if  $p\{x := r\} \preceq q$  for all  $r \in D$ , then  $p\{x := xy\} \preceq q$ :

- (i)  $D = \{ya, bc, dy\}$  ( $b = a$ ,  $b \neq d$ , and  $c \notin \{a, d\}$ ),
- (ii)  $D = \{ya, bc, dy\}$  ( $b \notin \{a, d\}$ ,  $c = d$ , and  $c \neq a$ ).

**Proof.** It is obvious if no variable symbol appears in  $p$ . Therefore, let  $p = p_1xp_2$ , where  $p_1, p_2$  are regular patterns and  $x$  is a variable symbol. We assume that  $p\{x := xy\} \not\preceq q$  in order to derive the contradiction.

(i) Let  $D = \{ya, bc, dy\}$  ( $b = a$ ,  $b \neq d$ , and  $c \notin \{a, d\}$ ). Since  $p\{x := r\} \preceq q$  for all  $r \in D$ , there are three strings of length 2 corresponding to  $ya, bc, dy$  in  $q$ . Note that the three strings may appear partly overlapping. The symbols appearing in  $D$  corresponds to a variable or a constant in  $q$ . Let  $y_1, y_2, y_3$  be variable symbols appearing in  $q$ . The strings  $ya$  and  $dy$  must correspond to the strings  $y_1a$  and  $dy_3$  in  $q$ , respectively. There are the following three possibilities of strings in  $q$  which corresponds to  $bc$  in  $p\{x := bc\}$ .

- (a)  $bc$ , (b)  $y_2c$ , (c)  $by_2$ .

(a) Let  $A, B, C$  be strings where  $\{A, B, C\} = \{y_1a, bc, dy_3\}$  and let  $q = q_1AwBw'Cq_2$ . Since  $p\{x := r\} \preceq q$  for all  $r \in D$  and  $p\{x := xy\} \not\preceq q$  hold, the following conditions hold:

- (1)  $p_1 \preceq q_1$  (1')  $p_2 \preceq wBw'Cq_2$
- (2)  $p_1 \preceq q_1Aw$  (2')  $p_2 \preceq w'Cq_2$
- (3)  $p_1 \preceq q_1AwBw'$  (3')  $p_2 \preceq q_2$

Let  $q'_1 = q_1A$ ,  $q'_2 = wBw'$ ,  $q'_3 = Cq_2$ . From (3) and (1'), we have  $p_1 \preceq q'_1q'_2$ ,  $p_2 \preceq q'_2q'_3$ . From Lemma 2, if  $q'_2$  contains a variable,  $p \preceq q$  holds. Therefore,  $B$  must be  $bc$ . If  $A = dy_3$ , from (2),  $p_1 \preceq q_1dy_3w$  holds. Let  $p_1 = p'_1p''_1$ ,  $p'_1 \preceq q_1d$ , and  $p''_1 \preceq y_3w$ . From (1'), we have  $p = p_1xp_2 = p'_1p''_1xp_2 \preceq q_1dp''_1xwbcw'y_1aq_2 = q\{x := p''_1x\}$ . This shows that there is a substitution  $\theta$  such that  $p = q\theta$  holds, and this contradicts the assumption. Therefore, we only need to consider the case where  $A = y_1a$ ,  $B = bc$ , and  $C = dy_3$ .

From the above, we consider two cases: one in which the symbols overlap and the other in which they do not.

- (a-1)  $q = q_1y_1awbcw'dy_3q_2$ ,
- (a-2)  $q = q_1y_1acwdy_3q_2$  ( $a = b$ ).

(a-1) From the proof of Lemma 6,  $p\{x := xy\} \preceq q$  holds. Therefore, it contradicts the assumption.

(a-2) Let  $q = q_1y_1acwdy_3q_2$  ( $a = b$ ). For this  $q$ , the following conditions hold:

- (1)  $p_1 \preceq q_1$  (1')  $p_2 \preceq cwdy_3q_2$
- (2)  $p_1 \preceq q_1y_1$  (2')  $p_2 \preceq wdy_3q_2$
- (3)  $p_1 \preceq q_1y_1acwdy_3$  (3')  $p_2 \preceq q_2$

If  $|w| = 0$ , from (1') and (2'), the prefix of  $p_2$  is  $cd$  and  $d$ . Therefore,  $c = d$ . This contradicts the fact that  $c \neq d$ .

If  $|w| = 1$ , from (1') and (2'), the prefix of  $p_2$  is  $cwd$

and  $wd$ . Therefore,  $w = c = d$ . This contradicts the fact that  $c \neq d$ .

If  $|w| \geq 2$ , then from (1') and (2'), prefixes of  $p_2$  is  $cwd$  and  $wd$ . Let  $w$  be  $w_1w_2w_3 \cdots w_{n-1}w_n$  ( $n \geq 2$ ,  $w_i \in \Sigma$  for  $i = 1, \dots, n$ ). From  $cw = wd$ , a prefix of  $w$  is  $c$  and a suffix of  $w$  is  $d$ . Therefore, we have  $w = cw_2w_3 \cdots w_{n-1}d$ . Since  $cw = cw_2w_3 \cdots w_{n-1}d$ ,  $wd = cw_2w_3 \cdots w_{n-1}dd$ , from  $cw = wd$ ,  $w_i = w_{i+1}$  holds for  $i = 2, \dots, n-2$ . Therefore,  $c = d$ . This contradicts the fact that  $c \neq d$ .

(b) Let  $q = q_1AwBw'Cq_2$  where  $\{A, B, C\} = \{y_1a, y_2c, dy_3\}$ , and let  $q = q_1AwBw'Cq_2$ . Since  $c \neq a$  holds,  $q$  have a substring that is corresponding to (i-2) of Lemma 7. Therefore,  $p\{x := xy\} \preceq q$  holds. This contradicts the assumption.

(c) As in (b), this contradicts the assumption.

(ii) In this case, by reversing the strings  $p$  and  $q$ , we can prove that the assumption  $p\{x := xy\} \preceq q$  is contradicted, as in the case of (i).

When the conditions of both Lemmas 6 and 7 are not satisfied, counterexamples exist as follows:

**Proposition 1:** Let  $\Sigma$  be an alphabet with  $|\Sigma| \geq 3$ . For a variable symbol  $y$ , let  $D = \{ya, bc, dy\}$  ( $b = a$  and  $c = d$ ). There exist regular patterns  $p$  and  $q$  on  $\Sigma$  such that  $p\{x := r\} \preceq q$  for any  $r \in D$ , but  $p\{x := xy\} \not\preceq q$ .

**Proof.** We give an example which shows this proposition. Let  $a, b, c, d, e$  be constant symbols in  $\Sigma$  and  $x, y, y_1, y_2$  variable symbols in  $X$ . Let

$$p = eabcbcadabcbcadaxbcadadabcbcadade,$$


$$q = y_1abcbcadabcbcadady_2 \quad (b = a \text{ and } c = d).$$

Obviously  $p\{x := xy\} \not\preceq q$  holds. For these  $p$  and  $q$ , the condition for Proposition 1 holds as follows (see also Fig. 9):


$$\begin{aligned} p\{x := ya\} &= (eabcbcadabcbcaday)abcbcadadabcbcadade \\ &= q\{y_1 := eabcbcadabcbcaday, y_2 := e\} \\ &\preceq q, \\ p\{x := bc\} &= (eabcbcad)abcbcadabcbcadad(abcbcadade) \\ &= q\{y_1 := eabcbcad, y_2 := abcbcadade\} \\ &\preceq q, \\ p\{x := dy\} &= eabcbcadabcbcadad(ybcadadabcbcadade) \\ &= q\{y_1 := e, y_2 := ybcadadabcbcadade\} \\ &\preceq q. \end{aligned}$$

**Lemma 8:** Let  $k$  be an integer with  $k \geq 1$ . Let  $\Sigma$  be an alphabet with  $|\Sigma| = k + 2$ . Let  $p \in \mathcal{RP}$  in which a variable symbol  $x$  appears, and let  $Q \in \mathcal{RP}^k$ . If for any string  $w \in \Sigma^*$  with  $|w| = 2$ , there exists a regular pattern  $q_w \in Q$  such that  $p\{x := w\} \preceq q_w$  holds, then there exists a regular pattern  $q \in Q$  such that  $p\{x := xy\} \preceq q$  holds, where  $y$  is a variable

$p\{x := ay\} =$	<table><tr><td>e</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>y</td><td>a</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>e</td></tr><tr><td colspan="16"><math>y_1</math></td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td><math>y_2</math></td></tr></table>	e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	y	a	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e	$y_1$																a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$
e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	y	a	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e																																			
$y_1$																a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$																																				
$p\{x := bc\} =$	<table><tr><td>e</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>e</td></tr><tr><td colspan="8"><math>y_1</math></td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td colspan="6"><math>y_2</math></td></tr></table>	e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e	$y_1$								a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$								
e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e																																			
$y_1$								a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$																																												
$p\{x := dy\} =$	<table><tr><td>e</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>y</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td>e</td></tr><tr><td><math>y_1</math></td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>b</td><td>c</td><td>b</td><td>c</td><td>a</td><td>d</td><td>a</td><td>d</td><td colspan="13"><math>y_2</math></td></tr></table>	e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	y	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e	$y_1$	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$															
e	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	y	b	c	a	d	a	d	a	b	c	b	c	a	d	a	d	e																																			
$y_1$	a	b	c	b	c	a	d	a	b	c	b	c	a	d	a	d	$y_2$																																																			

Fig. 9 Substitutions for  $p$  and each correspondence to  $q$ .

**Fig. 10** Let  $\Sigma = \{a, b, c, d, e, f, g\}$ ,  $Q = \{q_1, q_2, q_3, q_4, q_5\}$ . We set  $A(q_1) = \{a\}$  and  $B(q_1) = \{a\}$ , and then  $\sigma_A(q_1) = a$  and  $\sigma_B(q_1) = a$ , and so on. For each regular pattern  $q_i$  ( $i = 1, \dots, 5$ ), we represent a string  $w \in \Sigma \cdot \Sigma$  satisfying that  $p\{x := w\} \preceq q_i$  by the line between the left (first) and right (second) symbols of  $w$ . For example, the leftmost figure shows that  $p\{x := ay\} \preceq q_1$  and  $p\{x := ya\} \preceq q_1$  for a variable symbol  $y$ . We note that these figures may contain more lines than those depicted. From these figures, we get  $\ell_A = 1$ ,  $\ell_B = 0$ , and  $Q^{(\emptyset, \emptyset)} = \{q_5\}$ ,  $Q^{(\emptyset, \cdot)} = \{q_4\}$ ,  $Q^{(\cdot, \emptyset)} = \{q_3\}$ ,  $Q^{(\cdot, \cdot)} = \{q_1, q_2\}$ .



**Fig. 11** In the left figure, we aggregate all of the lines appearing in Fig. 10. For all  $w = a'b' \in A' \cdot B'$ , there must be a regular pattern  $q_i$  ( $1 \leq i \leq 5$ ) that satisfies that  $p\{x := w\} \preceq q_i$ .

symbol that does not appear in  $q$ .

**Proof.** W.l.o.g., we suppose that  $\#Q = k$  holds. Otherwise, for some regular pattern  $q$  already in  $Q$ , we can add a new regular pattern  $q'$  equivalent to  $q$ , i.e.,  $q' \equiv q$ , to  $Q$  repeatedly until  $\#Q = k$  is satisfied. For any  $q \in Q$ , we define the sets  $A(q), B(q) \subseteq \Sigma$  as follows:

$$A(q) = \{a \in \Sigma \mid p\{x := ay\} \preceq q, y \in X\},$$

$$B(q) = \{b \in \Sigma \mid p\{x := yb\} \preceq q, y \in X\}.$$

If there exists  $q \in Q$  such that  $|A(q)| \geq 2$  or  $|B(q)| \geq 2$ ,

from Lemma ??,  $p\{x := xy\} \preceq q$  holds. Below, we suppose that  $|A(q)| \leq 1$  and  $|B(q)| \leq 1$ . Let  $\emptyset$  be a constant symbol that is not a member in  $\Sigma$ . We define the functions  $\sigma_A : Q \rightarrow \Sigma \cup \{\emptyset\}$  and  $\sigma_B : Q \rightarrow \Sigma \cup \{\emptyset\}$  as follows:

$$\sigma_A(q) = \begin{cases} a & \text{if } A(q) = \{a\}, \\ \emptyset & \text{if } A(q) = \emptyset. \end{cases}$$

$$\sigma_B(q) = \begin{cases} b & \text{if } B(q) = \{b\}, \\ \emptyset & \text{if } B(q) = \emptyset. \end{cases}$$

The inverse functions of  $\sigma_A$  and  $\sigma_B$  are denoted by  $\sigma_A^{-1}$  and  $\sigma_B^{-1}$ , respectively. That is, for  $a, b \in \Sigma \cup \{\emptyset\}$ , let  $\sigma_A^{-1}(a) = \{q \in Q \mid \sigma_A(q) = a\}$  and  $\sigma_B^{-1}(b) = \{q \in Q \mid \sigma_B(q) = b\}$ . We give an example in Fig. 10.

$A$  and  $B$  denotes the following subsets of  $\Sigma$ :

$$A = \bigcup_{q \in Q \setminus \sigma_A^{-1}(\emptyset)} A(q), \quad B = \bigcup_{q \in Q \setminus \sigma_B^{-1}(\emptyset)} B(q).$$

Then, let  $A' = \Sigma \setminus A$  and  $B' = \Sigma \setminus B$ . For any  $a, b \in \Sigma$ , we use the following notations:

$$\ell_A = \sum_{a \in A} (\# \sigma_A^{-1}(a) - 1), \quad \ell_B = \sum_{b \in B} (\# \sigma_B^{-1}(b) - 1).$$

These  $\ell_A$  and  $\ell_B$  represent the numbers of excess duplicate

symbols in  $A$  and  $B$ . We easily see the following claim:

*Claim 1.*

- (i)  $\#A + \#A' = \#B + \#B' = k + 2$ ,
- (ii)  $\#A + \ell_A + \#\sigma_A^{-1}(\emptyset) = \#B + \ell_B + \#\sigma_B^{-1}(\emptyset) = k$ .

Since  $\#\Sigma = k + 2$  and  $\#Q = k$ ,  $\#A' \geq 2$  and  $\#B' \geq 2$  hold. We partition  $Q$  into the following subsets:

$$\begin{aligned} Q^{(\emptyset, \emptyset)} &= \sigma_A^{-1}(\emptyset) \cap \sigma_B^{-1}(\emptyset), \\ Q^{(\emptyset, \cdot)} &= \sigma_A^{-1}(\emptyset) \cap (Q \setminus \sigma_B^{-1}(\emptyset)), \\ Q^{(\cdot, \emptyset)} &= (Q \setminus \sigma_A^{-1}(\emptyset)) \cap \sigma_B^{-1}(\emptyset), \\ Q^{(\cdot, \cdot)} &= (Q \setminus \sigma_A^{-1}(\emptyset)) \cap (Q \setminus \sigma_B^{-1}(\emptyset)). \end{aligned}$$

From the condition of this lemma, for any string  $w \in \Sigma^*$  with  $|w| = 2$ , there exists a regular pattern  $q_w \in Q$  such that  $p\{x := w\} \preceq q_w$  holds. In particular, for  $w = a'b' \in A' \cdot B'$ , we must have  $q_w \in Q$  that satisfies that  $p\{x := w\} \preceq q_w$  (Fig. 11). It is easy to see that if  $w \in (A \cdot B) \cup (A' \cdot B) \cup (A \cdot B')$ , there exists a regular pattern  $q_w \in Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)} \cup Q^{(\cdot, \cdot)}$  such that  $p\{x := w\} \preceq q_w$  holds. The following two claims are proven from Lemmas ?? and 5:

*Claim 2.* If there exist  $q \in Q^{(\emptyset, \emptyset)}$  and distinct 5 strings  $w_i \in A' \cdot B'$  ( $1 \leq i \leq 5$ ) such that  $p\{x := w_i\} \preceq q$  holds ( $1 \leq i \leq 5$ ), then  $p\{x := xy\} \preceq q$  holds.

*Claim 3.* If there exist  $q \in Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)}$  and distinct 3 strings  $w_i \in A' \cdot B'$  ( $1 \leq i \leq 3$ ) such that  $p\{x := w_i\} \preceq q$  holds ( $1 \leq i \leq 3$ ), then  $p\{x := xy\} \preceq q$  holds.

If there exist a regular pattern  $q \in Q^{(\emptyset, \emptyset)} \cup Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)}$  and enough strings  $w \in A' \cdot B'$  such that either of the conditions of *Claims 2* and *3* is satisfied, this lemma holds. Then, we assume that it is not the case.

*Assumption 1.* There is no regular pattern  $q \in Q^{(\emptyset, \emptyset)}$  and 5 strings  $w \in A' \cdot B'$  such that the condition of *Claim 2* is satisfied and there is no regular pattern  $q \in Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)}$  and 3 strings  $w \in A' \cdot B'$  such that the condition of *Claim 3* is satisfied.

Let  $\mathcal{L}_1 = \#\{w \in A' \cdot B' \mid \exists q \in Q^{(\emptyset, \emptyset)} \cup Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)} \text{ s.t. } p\{x := w\} \preceq q\}$ . Under *Assumption 1*, each  $q \in Q^{(\emptyset, \emptyset)}$  has at most 4 strings  $w \in A' \cdot B'$  such that the condition of *Claim 2* is satisfied, and each  $q \in Q^{(\emptyset, \cdot)} \cup Q^{(\cdot, \emptyset)}$  has at most 2 strings  $w \in A' \cdot B'$  such that the condition of *Claim 3* is satisfied. Then, by *Claim 1*,

$$\begin{aligned} \mathcal{L}_1 &\leq 4\#Q^{(\emptyset, \emptyset)} + 2\#Q^{(\emptyset, \cdot)} + 2\#Q^{(\cdot, \emptyset)} \\ &= 2(\#Q^{(\emptyset, \emptyset)} + \#Q^{(\emptyset, \cdot)}) + 2(\#Q^{(\emptyset, \emptyset)} + \#Q^{(\cdot, \emptyset)}) \\ &= 2\#\sigma_A^{-1}(\emptyset) + 2\#\sigma_B^{-1}(\emptyset) \\ &= 2(k - \#A - \ell_A) + 2(k - \#B - \ell_B) \\ &= 2(\#A' - \ell_A - 2) + 2(\#B' - \ell_B - 2) \\ &= 2(\#A' + \#B') - 2(\ell_A + \ell_B) - 8. \end{aligned}$$

Next, we partition  $Q^{(\cdot, \cdot)}$  into the following two subsets:

$$Q_1^{(\cdot, \cdot)} = \{q \in Q^{(\cdot, \cdot)} \mid \sigma_A(q) \in B \text{ or } \sigma_B(q) \in A\},$$

$$Q_2^{(\cdot, \cdot)} = \{q \in Q^{(\cdot, \cdot)} \mid \sigma_A(q) \in B' \text{ and } \sigma_B(q) \in A'\}.$$

We show the next two claims on  $Q_1^{(\cdot, \cdot)}$  and  $Q_2^{(\cdot, \cdot)}$ :

*Claim 4.* If there exist  $q \in Q_1^{(\cdot, \cdot)}$  and a string  $a'b' \in A' \cdot B'$  such that  $p\{x := a'b'\} \preceq q$  holds, then  $p\{x := xy\} \preceq q$  holds.

*Proof of Claim 4.* Suppose that both  $\sigma_A(q) \in B$  and  $\sigma_B(q) \in A$  hold. Then, since  $a' \notin \{\sigma_A(q), \sigma_B(q)\} \subseteq A \cap B$  and  $b' \notin \{\sigma_A(q), \sigma_B(q)\} \subseteq A \cap B$ , from Lemma 6,  $p\{x := xy\} \preceq q$  holds. Suppose that  $\sigma_A(q) \in B$  and  $\sigma_B(q) \in A'$ . If  $a' = \sigma_B(q)$ , since  $a' \in B$ ,  $a' \neq b'$  holds. Since  $\sigma_A(q) \in B$ ,  $b' \neq \sigma_A(q)$  holds. That is,  $a' = \sigma_B(q)$ ,  $a' \neq \sigma_A(q)$ , and  $b' \notin \{\sigma_A(q), \sigma_B(q)\}$  hold. Therefore, from Lemma 7,  $p\{x := xy\} \preceq q$  holds. If  $a' \neq \sigma_B(q)$ , since  $b' \neq \sigma_A(q)$ , from Lemma 6,  $p\{x := xy\} \preceq q$  holds. Similarly, the case that  $\sigma_A(q) \in B'$  and  $\sigma_B(q) \in A$  is proven. (*End of Proof of Claim*)

*Claim 5.* If there exist  $q \in Q_2^{(\cdot, \cdot)}$  and a string  $a'b' \in A' \cdot B'$  such that  $(a' \neq \sigma_B(q) \text{ or } b' \neq \sigma_A(q))$  and  $p\{x := a'b'\} \preceq q$  hold, then  $p\{x := xy\} \preceq q$  holds.

*Proof of Claim 5.* When  $a' = b'$ , since  $\sigma_A(q) \neq \sigma_B(q)$ , from Lemma 6, this claim holds. Similarly, when  $a' \neq b'$ , from Lemma 6 or Lemma 7, this holds. (*End of Proof of Claim*)

If there exist a regular pattern  $q \in Q_2^{(\cdot, \cdot)}$  and a string  $w \in A' \cdot B'$  such that the condition of *Claim 5* is satisfied, this lemma holds. Then, we also assume that it is not the case.

*Assumption 2.* There is no  $q \in Q_2^{(\cdot, \cdot)}$  and a string  $a'b' \in A' \cdot B'$  such that the condition of *Claim 5* is satisfied.

Let  $\mathcal{L}_2 = \#\{a'b' \in A' \cdot B' \mid \exists q \in Q_2^{(\cdot, \cdot)} \text{ s.t. } p\{x := a'b'\} \preceq q\}$ . For any  $a'b' \in A' \cdot B'$  and  $q \in Q_2^{(\cdot, \cdot)}$ , if  $a' = \sigma_B(q)$  and  $b' = \sigma_A(q)$  hold (it is the condition of Proposition 1), by considering the duplicate numbers  $\ell_A$  and  $\ell_B$ , we have the following inequality:

$$\mathcal{L}_2 \leq \min\{\#A' + \ell_B, \#B' + \ell_A\}.$$

We show the last claim:

*Claim 6.*  $\#A' \times \#B' - \mathcal{L}_1 - \mathcal{L}_2 \geq 2$ .

*Proof of Claim 6.* First we prove the inequality when  $\#A \leq k - 1$  and  $\#B \leq k - 1$ , i.e.,  $\#A' \geq 3$  and  $\#B' \geq 3$  hold. Since  $\mathcal{L}_2 \leq \frac{1}{2}(\#A' + \#B' + \ell_A + \ell_B)$ ,

$$\begin{aligned} &\#A' \times \#B' - \mathcal{L}_1 - \mathcal{L}_2 \\ &\geq \#A' \times \#B' - (2(\#A' + \#B') - 2(\ell_A + \ell_B) - 8) \\ &\quad - \frac{1}{2}(\#A' + \#B' + \ell_A + \ell_B) \\ &= \#A' \times \#B' - \frac{5}{2}(\#A' + \#B') + \frac{3}{2}(\ell_A + \ell_B) + 8 \\ &= (\#A' - \frac{5}{2})(\#B' - \frac{5}{2}) + \frac{3}{2}(\ell_A + \ell_B) + \frac{7}{4} \geq 2. \end{aligned}$$

When  $\#A = k$  and  $\#B \leq k$ , i.e.,  $\#A' = 2$  and  $\#B' \geq 2$  hold, since  $\ell_A = 0$ ,  $\mathcal{L}_1 \leq 2\#B' - 2\ell_B - 4$  holds. Moreover,  $\mathcal{L}_2 \leq \min\{\#B', \ell_B + 2\}$  holds. From *Claim 1*,  $\ell_B + 2 =$



$k - \# \sigma_B^{-1}(\emptyset) - \#B = \#B' - \# \sigma_B^{-1}(\emptyset)$  holds. Therefore,  $\mathcal{L}_2 \leq \ell_B + 2$  holds. Thus,

$$\begin{aligned} & \#A' \times \#B' - \mathcal{L}_1 - \mathcal{L}_2 \\ & \geq 2\#B' - (2\#B' - 2\ell_B - 4) - (\ell_B + 2) \\ & = \ell_B + 2 \geq 2. \end{aligned}$$

Similarly, the case when  $\#A \leq k$  and  $\#B = k$  is proven. (*End of Proof of Claim*)

Under *Assumptions* 1 and 2, from *Claim* 6, there exist at least two  $w \in A' \cdot B'$  and a regular pattern  $q \in Q_1^{(\cdot, \cdot)}$  such that the condition of *Claim* 4 is satisfied. Therefore, for such a regular pattern  $q$ ,  $p\{x := xy\} \preceq q$  holds.

**Lemma 9** (Sato et al.[1]): Let  $\Sigma$  be a finite alphabet with  $\#\Sigma \geq 3$  and  $p, q$  regular patterns. If there exists a constant symbol  $a \in \Sigma$  such that  $p\{x := a\} \preceq q$  and  $p\{x := xy\} \preceq q$ , then  $p \preceq q$  holds, where  $y$  is a variable symbol that does not appear in  $q$ .

$$\begin{aligned} D &= \{a_1b_1, a_2b_2, a_3b_3\} \\ (a_i &\neq a_j \text{ and } b_i \neq b_j \text{ for each } i, j \ (i \neq j, 1 \leq i, j \leq 3), \\ b_1 &\neq a_2 \text{ and } b_2 \neq a_3) \end{aligned}$$

We assume that a variable symbol  $y$  does not appear in  $p$ . Then, if  $p\{x := r\} \preceq q$  for all  $r \in D$ , then  $p\{x := xy\} \preceq q$ .

**Proof.** It is obvious if the variable symbol  $x$  does not appear in  $p$ . Therefore, let  $p = p_1xp_2$ , where  $p_1, p_2$  are regular patterns. We assume that  $p\{x := xy\} \not\preceq q$  in order to derive the contradictions. Since  $p\{x := r\} \preceq q$  holds for any  $r \in D$ , the regular pattern  $q$  contains  $a_1b_1, a_2b_2$  and  $a_3b_3$ . We remark that  $a_i$  and  $b_j$  may be same for  $i, j \ (1 \leq i, j \leq 3)$ . Since  $p\{x := r\} \preceq q$  for all  $r \in D$  holds, there exist the following 10 cases (i)–(xv) for three regular patterns on  $\Sigma$  contained in  $q$  that correspond to three constant strings in  $D$ : Here,  $y_1, y_2, y_3$  are variable symbols.

- |                                |                                 |
|--------------------------------|---------------------------------|
| (i) $a_1b_1, a_2b_2, a_3b_3$   | (vi) $a_1b_1, y_1b_2, y_2b_3$   |
| (ii) $a_1b_1, a_2b_2, a_3y_1$  | (vii) $y_1b_1, y_2b_2, y_3b_3$  |
| (iii) $a_1b_1, a_2b_2, y_1b_3$ | (viii) $y_1b_1, y_2b_2, a_3y_3$ |
| (iv) $a_1b_1, y_1b_2, a_3y_2$  | (ix) $y_1b_1, a_2y_2, a_3y_3$   |
| (v) $a_1b_1, a_2y_1, a_3y_2$   | (x) $a_1y_1, a_2y_2, a_3y_3$    |

For the cases (iv), we can prove that  $p\{x := xy\} \preceq q$  holds in a similar way as Lemma 4. For the cases (v)–(x), we can prove that  $p\{x := xy\} \preceq q$  holds in a similar way as Lemma 4. Hence, for the cases (i)–(iii), we will prove that  $p\{x := xy\} \preceq q$  holds.

(I) Cases of (i), (ii) and (iii), that are the cases that  $q$  contains  $a_1b_1, a_2b_2$  and  $a_3b_3$ :

We consider the following four cases (I-1)–(I-4) of  $q$  for some regular patterns  $q_1, q_2$  and some constant strings  $w, w'$  ( $|w| \geq 0$  and  $|w'| \geq 0$ ):

- (I-1)  $q = q_1a_1b_1wa_2b_2w'a_3b_3q_2$ ,
- (I-2)  $q = q_1a_1b_1a_3b_3q_2$  ( $b_1 = a_2$  and  $a_3 = b_2$ ),
- (I-3)  $q = q_1a_1b_1b_2wa_3b_3q_2$  ( $b_1 = a_2$ ),
- (I-4)  $q = q_1a_1b_1wa_2b_2b_3q_2$  ( $b_2 = a_3$ ).

(I-1) Case of  $q = q_1a_1b_1wa_2b_2w'a_3b_3q_2$ : Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

- |                                      |                                       |
|--------------------------------------|---------------------------------------|
| (1) $p_1 \preceq q_1$                | (1') $p_2 \preceq wa_2b_2w'a_3b_3q_2$ |
| (2) $p_1 \preceq q_1a_1b_1w$         | (2') $p_2 \preceq w'a_3b_3q_2$        |
| (3) $p_1 \preceq q_1a_1b_1wa_2b_2w'$ | (3') $p_2 \preceq q_2$                |

If  $|w| = |w'|$  holds,  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$  from the above conditions (2) and (3). Then,  $a_1b_1w = a_2b_2w'$ . Hence,  $a_1b_1 = a_2b_2$ . This contradicts the assumption of  $a_1 \neq a_2$  and  $b_1 \neq b_2$ .

If  $|w| + 1 = |w'|$  holds,  $wa_2b_2w'a_3b_3$  and  $w'a_3b_3$  are the prefix of  $p_2$ . If there exists a constant symbol  $w_1$  such that  $w'a_3b_3 = ww_1a_3b_3$ , then  $b_2$  and  $a_3$  are the same symbol from  $wa_2b_2 = ww_1a_3$ . From the above conditions (2) and (3),  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$ . Then, there exists a constant symbol  $w_2$  such that  $w' = w_2w$ , then  $b_2$  and  $a_1$  are the same symbol from  $b_2w_2w = a_1b_1w$ . Hence, from  $b_2 = a_3$ ,  $a_3$  and  $a_1$  are same symbol. This contradicts the assumption of  $a_3 \neq a_1$ .

If  $|w| + 1 < |w'|$ , from the above (2) and (3),  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$ . If there exists a constant string  $w_1$  ( $|w_1| \geq 2$ ) such that  $w' = w_1w$ , then  $a_1b_1$  is the suffix of  $w_1$ . From the above conditions (1') and (2'),  $wa_2b_2w'a_3b_3$  and  $w'a_3b_3$  are the prefix of  $p_2$ . If there exist constant strings  $w_1$  and  $w_2$  such that  $w' = w_1w = ww_2$  holds, then  $a_2b_2$  and  $a_3b_3$  are the suffix of  $w_1$  from  $|w_1| = |w_2|$  and  $|ww_2a_3b_3| = |wa_2b_2w_1|$ . Hence,  $a_1b_1 = a_3b_3$ . This contradicts the assumption of  $a_1 \neq a_3$  and  $b_1 \neq b_3$ .

If  $|w| > |w'|$ , we can prove the contradiction in a similar way as  $|w| \leq |w'|$ .

(I-2) Case of  $q = q_1a_1b_1a_3b_3q_2$  ( $b_1 = a_2$  and  $a_3 = b_2$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

- |                             |                              |
|-----------------------------|------------------------------|
| (1) $p_1 \preceq q_1$       | (1') $p_2 \preceq a_3b_3q_2$ |
| (2) $p_1 \preceq q_1a_1$    | (2') $p_2 \preceq b_3q_2$    |
| (3) $p_1 \preceq q_1a_1b_1$ | (3') $p_2 \preceq q_2$       |

From the above conditions (2) and (3), since  $a_1b_1$  and  $a_1$  are the suffix of  $p_1$ ,  $b_1 = a_1$  holds. From the assumption of  $b_1 = a_2$ ,  $a_1 = a_2$ . This contradicts the assumption of  $a_1 \neq a_2$ .

(I-3) Case of  $q = q_1a_1b_1b_2wa_3b_3q_2$  ( $b_1 = a_2$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

- |                                 |                                  |
|---------------------------------|----------------------------------|
| (1) $p_1 \preceq q_1$           | (1') $p_2 \preceq b_2wa_3b_3q_2$ |
| (2) $p_1 \preceq q_1a_1$        | (2') $p_2 \preceq wa_3b_3q_2$    |
| (3) $p_1 \preceq q_1a_1b_1b_2w$ | (3') $p_2 \preceq q_2$           |

If  $|w| = 0$ , i.e.,  $w$  is the empty string, then  $a_1$  and  $a_1b_1b_2$  are the suffix of  $p_1$  from the above conditions (2) and (3) and  $b_2a_3b_3$  and  $a_3b_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Since  $b_2 = a_1$  and  $b_2a_3 = a_3b_3$ ,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

If  $|w| \geq 1$ ,  $a_1$  and  $a_1b_1b_2w$  are the suffix of  $p_1$  from the

above conditions (2) and (3). Hence, the last symbol of  $w$  is  $a_1$ . Moreover,  $b_2wa_3b_3$  and  $wa_3b_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Hence, the last symbol of  $w$  is  $a_3$ . Therefore,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

(I-4) Case of  $q = q_1a_1b_1wa_2b_2b_3q_2$  ( $b_2 = a_3$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq wa_2b_2b_3q_2 \\ (2) p_1 \preceq q_1a_1b_1w & (2') p_2 \preceq b_3q_2 \\ (3) p_1 \preceq q_1a_1b_1wa_2 & (3') p_2 \preceq q_2 \end{array}$$

If  $|w| = 0$ , i.e.,  $w$  is the empty string, then  $a_1b_1$  and  $a_1b_1a_2$  are the suffix of  $p_1$  from the above conditions (2) and (3) and  $a_2b_2b_3$  and  $b_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Since  $b_1 = a_2$  and  $a_2 = b_3$ , then  $b_1 = b_3$  holds. This contradicts the assumption of  $b_1 \neq b_3$ .

If  $|w| \geq 1$ , since  $a_1b_1w$  and  $a_1b_1wa_2$  are the suffix of  $p_1$  from the above conditions (2) and (3), the first symbol of  $w$  is  $b_1$ . Moreover, since  $wa_2b_2b_3$  and  $b_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'), the first symbol of  $w$  is  $b_3$ . Therefore,  $b_1 = b_3$  holds. This contradicts the assumption of  $b_1 \neq b_3$ .

(II) Case of (iv) that  $q$  contains  $a_1b_1, a_2b_2$  and  $a_3y$ : Let  $A, B, C$  be distinct regular patterns in  $\{a_1b_1, a_2b_2, a_3y\}$  such that  $q = q_1AwBw'Cq_2$ . Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq wBw'Cq_2 \\ (2) p_1 \preceq q_1Aw & (2') p_2 \preceq w'Cq_2 \\ (3) p_1 \preceq q_1AwBw' & (3') p_2 \preceq q_2 \end{array}$$

If  $|w| = |w'|$ , then  $Aw$  and  $AwBw'$  are the suffix of  $p_1$  from the above conditions (2) and (3). Hence,  $Aw = Bw'$  holds. This contradicts the assumption of  $A \neq B$ .

If  $|w| \neq |w'|$ , then we consider the two cases  $A = a_3y$  and  $B = a_3y$ : In the case of  $A = a_3y$ , without losing generality, we assume that  $B = a_1b_1$  and  $C = a_2b_2$ . Then, there exist regular patterns  $p'_1, p''_1$  such that  $p_1 = p'_1p''_1$ ,  $p'_1 \preceq q_1a_3$  and  $p''_1 \preceq yw$  from the above condition (2). Moreover, from the above condition (1'),  $p = p_1xp_2 = p'_1p''_1xp_2 \preceq q_1a_3p'_1xwa_1b_1w'a_2b_2q_2 = q_1a_3ywa_1b_1w'a_2b_2q_2\{y := p''_1x\} = q\{y := p''_1x\}$  holds. Hence,  $p \preceq q$  holds. This contradicts the assumption. In the case of  $B = a_3y$ , without losing generality, we assume that  $A = a_1b_1$  and  $C = a_2b_2$ . Let  $q'_1 = q_1a_1b_1$ ,  $q'_2 = wa_3yw'$ , and  $q'_3 = a_2b_2q_2$  such that  $q'_2$  contains at most one variable symbol. Then, the above conditions (3) and (1') are represented by  $p_1 \preceq q'_1q'_2$  and  $p_2 \preceq q'_2q'_3$ , respectively. From Theorem 2,  $p \preceq q$  holds. This contradicts the assumption.

Next, in the case of  $C = a_3y$ , we consider the following five cases (II-1)–(II-5):

$$\begin{array}{l} \text{(II-1)} q = q_1a_1b_1wa_2b_2w'a_3yq_2, \\ \text{(II-2)} q = q_1a_1b_1b_2yq_2 \text{ } (a_2 = b_1 \text{ and } a_3 = b_2), \\ \text{(II-3)} q = q_1a_1b_1b_2wa_3yq_2 \text{ } (b_1 = a_2), \\ \text{(II-4)} q = q_1a_3ywa_1b_1b_2q_2 \text{ } (b_1 = a_2), \\ \text{(II-5)} q = q_1a_1b_1ywa_2b_2q_2 \text{ } (b_1 = a_3). \end{array}$$

(II-1) Case of  $q = q_1a_1b_1wa_2b_2w'a_3yq_2$ : Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq wa_2b_2w'a_3yq_2 \\ (2) p_1 \preceq q_1a_1b_1w & (2') p_2 \preceq w'a_3yq_2 \\ (3) p_1 \preceq q_1a_1b_1wa_2b_2w' & (3') p_2 \preceq q_2 \end{array}$$

If  $|w| + 1 = |w'|$ , then  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$  from the above conditions (2) and (3). Since there exists a constant symbol  $w_1$  such that  $w' = w_1w$  and  $b_2w_1w = a_1b_1w$  hold, then  $b_2 = a_1$ . Moreover,  $wa_2b_2w'a_3$  and  $w'a_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Since there exists a constant symbol  $w_2$  such that  $w' = ww_2$  and  $wa_2b_2 = ww_2a_3$  hold, then  $b_2 = a_3$ . Thus,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

If  $|w| + 1 < |w'|$ , then  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$  from the above conditions (2) and (3). Hence,  $a_1b_1$  is the suffix of  $w$ . Moreover,  $wa_2b_2w'a_3$  and  $w'a_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Hence, there exist constant symbols  $w_1$  and  $w_2$  such that  $w' = w_1w$ ,  $w' = ww_2$  and  $|a_2b_2w_1| = |w_2a_3| + 1$  hold. Thus, since the second-to-last symbol of  $w_1$  is  $a_3$ ,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

If  $|w| = |w'| + 1$ , then  $wa_2b_2w'a_3$  and  $w'a_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'). Since there exists a constant symbol  $w_1$  such that  $w = w'_w_1$  and  $w'_w_1 = w'a_3$  hold, then  $w_1 = a_3$  holds. Moreover, since  $a_1b_1wa_2b_2w'$  and  $a_1b_1w$  are the suffix of  $p_1$  from the above conditions (2) and (3), there exists a constant symbol  $w_2$  such that  $w = w_2w'$  and  $|w_1a_2b_2w'| = |a_1b_1w_2w'|$  hold. Hence,  $w_1 = a_1$  holds. Thus,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

If  $|w| > |w'| + 1$ , since  $wa_2b_2w'a_3$  and  $w'a_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'), there exists a constant string  $w_1$  such that  $w = w'_w_1$  and the first symbol of  $w_1$  is  $a_3$ . Moreover, since there exists a constant string  $w_2$  such that  $w = w_2w'$  and  $|w_1a_2b_2| = |a_1b_1w_2|$  hold,  $a_1b_1$  is the prefix of  $w_1$ . Thus,  $a_3 = a_1$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

(II-2) Case of  $q = q_1a_1b_1b_2yq_2$  ( $a_2 = b_1$  and  $a_3 = b_2$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq b_2yq_2 \\ (2) p_1 \preceq q_1a_1 & (2') p_2 \preceq yq_2 \\ (3) p_1 \preceq q_1a_1b_1 & (3') p_2 \preceq q_2 \end{array}$$

From the above conditions (2) and (3),  $a_1b_1$  and  $a_1$  are the suffix of  $p_1$ . Hence,  $b_1 = a_1$  holds. Thus, from the assumption of  $b_1 = a_2$ ,  $a_1 = a_2$  holds. This contradicts the assumption of  $a_1 \neq a_2$ .

(II-3) Case of  $q = q_1a_1b_1b_2wa_3yq_2$  ( $b_1 = a_2$ ): Assume

that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq b_2 w a_3 y q_2 \\ (2) p_1 \preceq q_1 a_1 & (2') p_2 \preceq w a_3 y q_2 \\ (3) p_1 \preceq q_1 a_1 b_1 b_2 w & (3') p_2 \preceq q_2 \end{array}$$

If  $|w| = 0$ , i.e.,  $w$  is the empty string, then  $a_1$  and  $a_1 b_1 b_2$  are the suffix of  $p_1$  from the above conditions (2) and (3). Hence,  $a_1 = b_2$  holds. Moreover, since  $b_2 a_3$  and  $a_3$  is the prefix of  $p_2$ ,  $b_2 = a_3$  holds. Thus,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

If  $|w| \geq 1$ , since  $a_1$  and  $a_1 b_1 b_2 w$  are the suffix of  $p_1$  from the above conditions (2) and (3), the last symbol of  $w$  is  $a_1$ . Moreover, since  $b_2 w a_3$  and  $w a_3$  are the prefix of  $p_2$  from the above conditions (1') and (2'), the last symbol of  $w$  is  $a_3$ . Thus,  $a_1 = a_3$  holds. This contradicts the assumption of  $a_1 \neq a_3$ .

(II-4) Case of  $q = q_1 a_3 y w a_1 b_1 b_2 q_2$  ( $b_1 = a_2$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq w a_1 b_1 b_2 q_2 \\ (2) p_1 \preceq q_1 a_3 y w & (2') p_2 \preceq b_2 q_2 \\ (3) p_1 \preceq q_1 a_3 y w a_1 & (3') p_2 \preceq q_2 \end{array}$$

From the above condition (3), there exist regular patterns  $p'_1 \succeq p''_1$  such that  $p_1 = p'_1 p''_1$ ,  $p'_1 \preceq q_1 a_3$  and  $p''_1 \preceq y w a_1$  hold. Hence, since  $p = p_1 x p_2 = p'_1 p''_1 x p_2 \preceq q_1 a_3 p''_1 x w a_1 b_1 b_2 q_2 = q_1 a_3 y x w a_1 b_1 b_2 q_2 \{y := p''_1 x\} = q \{y := p''_1 x\}$ , then  $p \preceq q$  holds. Thus, this contradicts the assumption.

(II-5) Case of  $q = q_1 a_1 b_1 y w a_2 b_2 q_2$  ( $b_1 = a_3$ ): Assume that the following six conditions (1),(2),(3),(1'),(2'),(3') are hold.

$$\begin{array}{ll} (1) p_1 \preceq q_1 & (1') p_2 \preceq y w a_2 b_2 q_2 \\ (2) p_1 \preceq q_1 a_1 & (2') p_2 \preceq w a_2 b_2 q_2 \\ (3) p_1 \preceq q_1 a_1 b_1 y w & (3') p_2 \preceq q_2 \end{array}$$

There exist regular patterns  $q'_1, q'_2, q'_3$  such that  $q'_1 = q_1 a_1 b_1$ ,  $q'_2 = y w$ ,  $q'_3 = a_2 b_2 q_2$ , from the above condition (3)  $p_1 \preceq q'_1 q'_2$  and from the above condition (1')  $p_2 \preceq q'_2 q'_3$  hold. Moreover, since  $q'_2$  contains the variable symbol  $y$ ,  $p \preceq q$  holds from Theorem 2. This contradicts the assumption.

From the Lemma 8 and Lemma 9, we have the following theorem.

**Theorem 4:** Let  $k \geq 3$ ,  $\# \Sigma \geq 2k - 1$ ,  $P \in \mathcal{RP}^+$  and  $Q \in \mathcal{RP}^k$ . Then, the following (i),(ii) and (iii) are equivalent:

$$(i) S_2(P) \subseteq L(Q), (ii) P \sqsubseteq Q, (iii) L(P) \subseteq L(Q).$$

**Proof.** it is clear that (ii) implies (iii) and (iii) implies (i). From Theorem 3, if  $\# \Sigma \geq 2k + 1$ , then (i) implies (ii). Let  $\# Q = k$ ,  $p \in P$ ,  $\# \Sigma = 2k - 1$  or  $2k$ . Then, we show that (i) implies (ii). It suffices to show that  $S_2(p) \subseteq L(Q)$  implies

$P \sqsubseteq Q$  for any regular pattern  $p \in \mathcal{RP}$ . The proof is done by mathematical induction on  $n$ , where  $n$  is the number of variable symbols appears in  $p$ .

In case  $n = 0$ ,  $S_2(p) = \{p\}$ . By (i), we have  $\{p\} = L(Q)$ . Thus,  $p \preceq q$  for some  $q \in Q$ .

For  $n \geq 0$ , we assume that it is valid for any regular pattern  $p$  with  $n$  variable symbols. Let  $p$  be a regular pattern such that  $n + 1$  variable symbols appear in  $p$  and  $S_2(p) \subseteq L(Q)$ .

We assume that  $p \not\sqsubseteq Q$ , that is,  $p \not\preceq q_i$  for any  $i \in \{1, \dots, k\}$ . Let  $Q = \{q_1, \dots, q_k\}$  and  $p_1, p_2$  regular patterns,  $x$  a variable symbol with  $p = p_1 x p_2$ . For  $a, b \in \Sigma$ , let  $p_a = p\{x := a\}$  and  $p_{ab} = p\{x := ab\}$ . Both  $p_a$  and  $p_{ab}$  have  $n$  variable symbols, respectively. Thus,  $S_2(p_a) \subseteq L(Q)$  and  $S_2(p_{ab}) \subseteq L(P)$  hold. By the induction hypothesis, there exist  $i, i' \in \{1, \dots, k\}$  such that  $p_a \preceq q_i$  and  $p_{ab} \preceq q_{i'}$ . Let  $D_i = \{a \in \Sigma \mid p\{x := a\} \preceq q_i\}$  ( $i = 1, \dots, k$ ). We assume that  $\#D_i \geq 3$  for some  $i \in \{1, \dots, k\}$ . By Lemma ??, we have  $p \preceq q_i$ . This contradicts the assumption. Thus, we have  $\#D_i \leq 2$  for any  $i \in \{1, \dots, k\}$ . If  $\# \Sigma = 2k - 1$ , then  $\#D_i = 2$  or  $\#D_i = 1$  for any  $i \in \{1, \dots, k\}$ . Moreover, If  $\# \Sigma = 2k$ , then  $\#D_i = 2$  for any  $i \in \{1, \dots, k\}$ . Since  $k \geq 3$ ,  $2k + 1 \geq k + 2$  holds. By Lemma 8, there exists  $i \in \{1, \dots, k\}$  such that  $p\{x := xy\} \preceq q_i$ . Therefore, by Lemma 9, we have  $p \preceq q_i$ . This contradicts the assumption. Thus, (i) implies (ii).

From Theorem 4, the following corollary holds.

**Corollary 2:** Let  $k \geq 3$ ,  $\# \Sigma \geq 2k - 1$  and  $P \in \mathcal{RP}^+$ . Then,  $S_2(P)$  is a characteristic set for  $L(P)$  within  $\mathcal{RPL}^k$ .

**Lemma 10** (Sato et al.[1]): Let  $k \geq 3$  and  $\# \Sigma \leq 2k - 2$ . Then,  $\mathcal{RP}^k$  does not have compactness with respect to containment.

**Proof.** Let  $\Sigma = \{a_1, \dots, a_{k-1}, b_1, \dots, b_{k-1}\}$  and  $p, q_i$  regular patterns,  $w_i \in \Sigma^*$  ( $i = 1, \dots, k - 1$ ) defined in a similar way to Example ??. Let  $q_k = x_1 a_1 w_1 x y w_1 b_1 x_2$ . Since  $p\{x := a_i\} = x_1 a_1 w_1 a_i w_1 b_1 x_2 \preceq q_i$  and  $p\{x := b_i\} = x_1 a_1 w_1 b_i w_1 b_1 x_2 \preceq q_i$  for any  $i \in \{1, \dots, k - 1\}$ , we have  $S_1(p) \subseteq \bigcup_{i=1}^{k-1} L(q_i)$ . For any  $w \in \{s \in \Sigma^+ \mid |s| \geq 2\}$ ,  $p\{x := w\} = x_1 a_1 w_1 w w_1 b_1 x_2 \preceq q_k$ . Thus, we have  $L(p) \subseteq L(Q)$ . By Theorem 1, since  $p \not\preceq q_i$ ,  $L(p) \not\subseteq L(q_i)$  for any  $i \in \{1, \dots, k\}$ . Therefore,  $\mathcal{RP}^k$  does not have compactness with respect to containment.

From Theorem 4 and Lemma 10, we have the following theorem.

**Theorem 5:** Let  $k \geq 3$  and  $\# \Sigma \geq 2k - 1$ . Then,  $\mathcal{RP}^k$  has compactness with respect to containment.

In case  $k = 2$ , we have the following theorem.

**Theorem 6:** Let  $\# \Sigma \geq 4$ ,  $P \in \mathcal{RP}^+$  and  $Q \in \mathcal{RP}^2$ . The following (i), (ii) and (iii) are equivalent:

$$(i) S_2(P) \subseteq L(Q), (ii) P \sqsubseteq Q, (iii) L(P) \subseteq L(Q).$$

**Proof.** It is clear that (ii) implies (iii), and (iii) implies (i). Thus, we show that (i) implies (ii). It suffices to show that

$S_2(p) \subseteq L(Q)$  implies  $P \sqsubseteq Q$  for any regular pattern  $p \in \mathcal{RP}$ . Let  $Q = \{q_1, q_2\}$ . The proof is done by mathematical induction on  $n$ , where  $n$  is the number of variable symbols appearing in  $p$ . In case  $n = 0$ ,  $p \in \Sigma^+$ . Since  $S_2(p) = \{p\} \subseteq L(Q)$ , we have  $p \preceq q$  for some  $q \in Q$ . For  $n \geq 0$ , we assume that it is valid for any regular pattern  $p$  with  $n$  variable symbols. Let  $p$  be a regular pattern such that  $n + 1$  variable symbols appear in  $p$ , and  $S_2(p) \subseteq L(Q)$ . We assume that  $p \not\preceq q_i$  ( $i = 1, 2$ ). Let  $p_1, p_2$  be regular patterns and  $x$  a variable symbol with  $p = p_1 x p_2$ . For  $a, b \in \Sigma$ , let  $p_a = p\{x := a\}$  and  $p_{ab} = p\{x := ab\}$ . Note that  $p_a$  and  $p_{ab}$  have  $n$  variable symbols. Thus, by the assumption,  $S_2(p_a) \subseteq L(Q)$  and  $S_2(p_{ab}) \subseteq L(Q)$  implies  $p_a \preceq q_i$  and  $p_{ab} \preceq q_{i'}$  for some  $i, i' \in \{1, 2\}$ . Let  $D_i = \{a \in \Sigma \mid p\{x := a\} \preceq q_i\}$  ( $i = 1, 2$ ). By Lemma ??, if  $\#D_i \geq 3$  for some  $i \in \{1, 2\}$ , then  $p \preceq q_i$ . This contradicts that  $p \not\preceq q_i$  ( $i = 1, 2$ ). Thus, we have  $\#D_i \leq 2$  for any  $i \in \{1, 2\}$ . Since  $\#\Sigma \geq 4$ , we consider that  $\#D_1 = 2$  and  $\#D_2 = 2$ . From Lemma 8,  $p\{x := xy\} \preceq q_i$  for some  $i \in \{1, 2\}$ . From Lemma 9, we have  $p \preceq q_i$  for some  $i \in \{1, 2\}$ . This contradicts that  $p \not\preceq q_i$  ( $i = 1, 2$ ). Therefore, (i) implies (ii).

The next example is a counter-example of Theorem 6.

**Example 2:** Let  $\Sigma = \{a, b, c\}$ ,  $p, q_1, q_2$  regular patterns and  $x, x', x''$  variable symbols such that  $p = x' a x b x''$ ,  $q_1 = x' a b x''$  and  $q_2 = x' c x''$ . Let  $w \in \Sigma^+$ . If  $w$  contains  $c$ , then  $p\{x := w\} \preceq q_2$ . On the other hand, if  $w$  does not contain  $c$ , then  $p\{x := w\} \preceq q_1$ . Thus,  $L(p) \subseteq L(q_1) \cup L(q_2)$ . However,  $p \not\preceq q_1$  and  $p \not\preceq q_2$ .

From Theorem 6, we have that following two corollaries.

**Corollary 3:** Let  $\#\Sigma \geq 4$  and  $P \in \mathcal{RP}^+$ . Then,  $S_2(P)$  is a characteristic set for  $L(P)$  within  $\mathcal{RPL}^2$ .

**Corollary 4:** Let  $\#\Sigma \geq 4$ . Then,  $\mathcal{RP}^2$  has compactness with respect to containment.

#### 4. Regular Pattern without Adjacent Variable Symbols

A regular pattern  $p$  is said to be a *non-adjacent variable regular pattern* (NAV regular pattern) if  $p$  does not contain consecutive variable symbols. For example, the regular pattern  $p = axybc$  is not a NAV regular pattern because  $xy$  is appeared in  $p$ . Let  $\mathcal{R}_{NAV}$  be the set of all NAV regular patterns. Let  $\mathcal{RP}_{NAV}^+$  be the set of all finite subsets  $S$  of  $\mathcal{R}_{NAV}$  such that  $S$  is not the empty set, i.e.,  $\mathcal{RP}_{NAV}^+ = \{S \subseteq \mathcal{R}_{NAV} \mid \#S \leq 1\}$ , and  $\mathcal{RP}_{NAV}^k$  the set of all subsets  $P$  of  $\mathcal{RP}_{NAV}^+$  such that  $P$  consists of at most  $k$  ( $k \geq 1$ ) NAV regular patterns, i.e.,  $\mathcal{RP}_{NAV}^k = \{P \in \mathcal{RP}_{NAV}^+ \mid \#P \leq k\}$ . We can define the compactness with respect to containment for  $\mathcal{RP}_{NAV}^k$  in a similar way as Def.2. For any NAV regular pattern  $p \in \mathcal{R}_{NAV}$  and any set  $Q \in \mathcal{RP}_{NAV}^k$  with  $k$  ( $k \geq 1$ ), the set  $\mathcal{RP}_{NAV}^k$  said to have *compactness with respect to containment* if there exists a NAV regular pattern  $q \in Q$  such that  $L(p) \subseteq L(q)$  holds if  $L(p) \subseteq L(Q)$  holds. Then, we have the following Theorem 7.

$p = x' c a d a a d a c b a d a d a a d a x a d a a d a c b a d a d a a d a b x''$ ,  
 $q_1 = x' c a d a a d a c b a d a d a a d a c x''$ ,  
 $q_2 = x' b a d a a d a c x''$ ,  
 $q_3 = x' a a d a d x''$ .

Fig. 12 NAV regular patterns  $p, q_1, q_2$ , and  $q_3$

**Theorem 7:** For an integer  $k$  ( $k \geq 2$ ), let  $\#\Sigma \geq k + 2$ ,  $P \in \mathcal{RP}_{NAV}^+$ ,  $Q \in \mathcal{RP}_{NAV}^k$ . Then, the following (i), (ii) and (iii) are equivalent:

- (i)  $S_2(P) \subseteq L(Q)$ , (ii)  $P \sqsubseteq Q$ , (iii)  $L(P) \subseteq L(Q)$ .

**Proof.** From the definitions of  $\mathcal{RP}_{NAV}^+$  and  $\mathcal{RP}_{NAV}^k$ , it is clear that (ii) implies (iii) and (iii) implies (i). Hence, we will show that (i) implies (ii) by mathematical induction on the number  $n$  of variable symbols that appear in a NAV regular pattern  $p \in P$  as follows: If  $n = 0$ , then we have  $S_2(\{p\}) = \{p\}$ . Hence,  $p \in L(Q)$ . Therefore, there exists  $q \in Q$  such that  $p \preceq q$  holds.

If  $n \geq 0$ , we assume that the proposition holds for any regular NAV regular pattern containing  $n \geq 0$  variable symbols. Let  $p$  be a NAV regular pattern containing  $n + 1$  variable symbols such that  $S_2(\{p\}) \subseteq L(Q)$  holds and  $p$  contains a variable symbol  $x$ . There exist two NAV regular patterns  $p_1, p_2$  such that  $p = p_1 x p_2$  holds. By the induction hypothesis, for any constant string  $w \in \Sigma^*$  with  $|w| = 2$ ,  $\{p\{x := w\}\} \preceq Q$  holds because  $p\{x := w\}$  contains  $n$  variable symbols. Hence, there exists a NAV regular pattern  $q_w \in Q$  such that  $p\{x := w\} \preceq q_w$  holds. From Lemma 8, there exists a regular pattern  $q \in Q$  such that  $p\{x := xy\} \preceq q$  holds, where  $y$  is a variable symbol that does not appear in  $q$ . This contradicts the condition  $Q \in \mathcal{RP}_{NAV}^k$ . Thus, we have that (i) implies (ii).

**Corollary 5:** Let  $k \geq 2$ ,  $\#\Sigma \geq k + 2$  and  $P \in \mathcal{RP}_{NAV}^+$ . Then,  $S_2(P)$  is a characteristic set of  $\mathcal{RPL}_{NAV}^k$ .

**Lemma 11:** Let  $k \geq 2$  and  $\#\Sigma \leq k + 1$ . Then,  $\mathcal{RP}_{NAV}^k$  does not have compactness with respect to containment.

**Proof.** Let  $\Sigma$  be the set of  $k + 1$  constant symbols  $a_1, \dots, a_{k+1}$ , i.e.,  $\Sigma = \{a_1, \dots, a_{k+1}\}$ . We assume that for  $i = 1, 2, \dots, k$ ,  $p\{x := a_i y\} \preceq q_i$  and  $p\{x := y a_{i+1}\} \preceq q_i$  ( $i = 1, 2, \dots, k$ ) hold. If  $p\{x := a_{k+1} a_1\} \preceq q_1$  holds,  $S_2(p) \setminus S_1(p) \subseteq \bigcup_{i=1}^k L(q_i)$  holds. This show that  $L(p) \subseteq L(Q)$  holds. However, for  $i = 1, 2, \dots, k$ , since  $p \not\preceq q_i$  holds, we have that  $L(p) \not\subseteq L(q_i)$  holds. Hence,  $\mathcal{RP}_{NAV}^k$  does not have compactness with respect to containment.

Next, we give an example for Lemma 11 in Example 3.

**Example 3:** Let  $\Sigma$  be the set of four constant symbols  $a, b, c, d$ , i.e.,  $\Sigma = \{a, b, c, d\}$  and  $x, x', x''$  three distinct variable symbols. Let  $p, q_1, q_2, q_3$  be the NAV regular patterns given in Fig. 12. Then, we have  $L(p) \subseteq L(q_1) \cup L(q_2) \cup L(q_3)$ . This show that for  $P = \{p\}$ ,  $Q = \{q_1, q_2, q_3\}$ , (iii) of Theorem 7 holds. However, since  $p \not\preceq q_1$ ,  $p \not\preceq q_2$  and  $p \not\preceq q_3$  hold, we have  $P \not\sqsubseteq Q$ , that

**Table 2** The conditions on the number  $\#\Sigma$  of constant symbols in  $\Sigma$  required for compactness with respect to containment.

Class	$k = 2$	$k \geq 3$
$\mathcal{RP}^k$	$\#\Sigma \geq 4$	$\#\Sigma \geq 2k - 1$
$\mathcal{RP}_{NAV}^k$	$\#\Sigma \geq k + 2$	

is, (ii) of Theorem 7 does not hold.

From Theorem 7 and Lemma 11, we have the following theorem.

**Theorem 8:** Let  $k \geq 2$  and  $\#\Sigma \geq k + 2$ . Then, the set  $\mathcal{RP}_{NAV}^k$  has compactness with respect to containment.

## 5. Conclusion

In this paper, for an integer  $k$  ( $k \geq 2$ ), we have shown the conditions on the number of constant symbols in  $\Sigma$ , summarized in Table 2, required for the classes  $\mathcal{RP}^k$  of all the set of  $k$  regular pattern languages and  $\mathcal{RP}_{NAV}^k$  of all the set of  $k$  NAV regular patterns to have compactness with respect to containment.

## Acknowledgements

This work was partially supported by Grant-in-Aid for Scientific Research (C) (Grant Numbers 19K12102, 19K12103, 20K04973, 21K12021, 22K12172) from Japan Society for the Promotion of Science (JSPS).

## References

- [1] M.Sato, Y.Mukouchi, D.Zheng, Characteristic Sets for Unions of Regular Pattern Languages and Compactness, in Proc. ALT '98, Springer LNAI 1501, pp.220-233, 1998.
- [2] Y. Mukouchi, Characterization of Pattern Languages, in Proc. ALT '91, Ohmsha, pp.93-104, 1991.
- [3] K.Wright, Identification of Unions of Languages Drawn from an Identifiable Class, in Proc. COLT '89, Morgan Kaufmann, pp.328-333, 1989.
- [4] H. Arimura, T. Shinohara, S. Otsuki, Finding Minimal Generalizations for Unions of Pattern Languages and Its Application to Inductive Inference from Positive Data, in Proc. STACS '94, Springer LNCS 775, pp.649-660, 1994.
- [5] Y. Suzuki, T. Shoudai, T. Uchida and T. Miyahara, Ordered Term Tree Languages Which are Polynomial Time Inductively Inferable from Positive Data, Theoretical Computer Science, 350(1):63-90, 2006.
- [6] T. Uchida, T. Shoudai, S. Miyano, Parallel Algorithms for Refutation Tree Problem on Formal Graph Systems, IEICE Trans. Inf. & Syst., E78-D(2):99-112, 1995.