

Lecture – 10

CS 372 (Computer Graphics)



Course Instructure : Dr. S. K. Maji
Asst. Prof.(CSE)

TRANSFORMATIONS IN 3D

Homogeneous representation of a 3D point (x, y, z) is (x_h, y_h, z_h, h) . where h is the homogeneous term.

TRANSLATION

$$\begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

SCALING


$$\begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

can be changed for uniform scaling.

ROTATION

In 2D we were rotating around origin. In 3D rotation will be around an axis.

Rotation around Z-axis
in counter-clock direction.
X & Y changes, Z remains
fixed.



$$\begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rotation around X-axis

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta & 0 \\ 0 & \sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Rotation around Y-axis

$$\begin{bmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

REFLECTION

Reflection around
XY plane, Z value -ve.

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Reflection around
XZ plane, Y -ve

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Reflection around
YZ plane, X -ve

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Around X-axis

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Around Y-axis

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


Around Z-axis

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Around Origin? All diagonal elements, except 'R', -1.

ROTATION OF AN ARBITRARY POINT AROUND AN INCLINED AXIS

1. Translate origin to a, b, c : T_1
2. Rotate s.t. vector (l, m, n) coincides with z -axis, say Z .
 - So, rotate s.t. vector coincides with xZ plane: T_2
 - Then rotate so that it coincides with Z -axis: T_3
3. Then rotate the point by θ : T_4
4. Reverse of 2
5. Reverse of 1.



∴ $T = T_1^{-1} T_2^{-1} T_3^{-1} T_4 T_3 T_2 T_1$

