

Introduction to topology in electronic structure of crystalline solids

Tomáš Rauch

IFTO, FSU Jena

Schedule

- ▶ 24.03. Introduction into topological insulators
- ▶ 14.04. Calculation of topological invariants of realistic materials
- ▶ TBA Further concepts related to topological insulators
- ▶ TBA Topological metals and higher-order topological insulators
- ▶ TBA Applications
- ▶ TBA Unanswered topics?

Literature

- ▶ D. Vanderbilt, Berry phases in electronic structure theory
- ▶ B. A. Bernevig, Topological insulators and topological superconductors
- ▶ J. K. Asboth, L. Oroszlany, A. Pályi, A short course on topological insulators

The Nobel Prize in Physics 2016



© Trinity Hall, Cambridge University. Photo: Kiloran Howard

David J. Thouless

Prize share: 1/2



Photo: Princeton University, Comms. Office, D. Applewhite

F. Duncan M. Haldane

Prize share: 1/4



Ill: N. Elmehed. © Nobel Media 2016

J. Michael Kosterlitz

Prize share: 1/4

for theoretical discoveries of topological phase transitions and
topological phases of matter

Fig. from nobelprize.org

Topology in mathematics

¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia


Topology in mathematics

- ▶ topological invariants

¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in mathematics

- ▶ topological invariants
- ▶ Euler characteristic¹: $\chi = \text{vertices} - \text{edges} + \text{faces} = 2$




$$\chi = 8 - 12 + 6 = 2$$


¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in mathematics

- ▶ topological invariants
- ▶ Euler characteristic¹: $\chi = \text{vertices} - \text{edges} + \text{faces} = 2$



$$\chi = 8 - 12 + 6 = 2$$




$$\chi = (8+1) - (12+4) + (6+3) = 2$$


¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in mathematics

- ▶ topological invariants
- ▶ Euler characteristic¹: $\chi = \text{vertices} - \text{edges} + \text{faces} = 2$



$$\chi = 8 - 12 + 6 = 2$$




$$\chi = (8+1) - (12+4) + (6+3) = 2$$


¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in mathematics

- ▶ topological invariants
- ▶ Euler characteristic¹: $\chi = \text{vertices} - \text{edges} + \text{faces} = 2$



$$\chi = 8 - 12 + 6 = 2$$




$$\chi = (8+1) - (12+4) + (6+3) = 2$$


¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in mathematics


- ▶ topological invariants
- ▶ Euler characteristic¹: $\chi = \text{vertices} - \text{edges} + \text{faces} = 2$



$$\chi = 8 - 12 + 6 = 2$$



$$\chi = (8+1) - (12+4) + (6+3) = 2$$




$$\chi = 24 - 48 + 24 = 0$$

¹H. Eschrig: Topology and Geometry for Physics, Springer (2011)
Figs. from Wikipedia

Topology in solid-state physics (example)²


Quantum Hall effect



²Fig. from von Klitzing *et al.*, Rev. Mod. Phys. **58**, 519 (1986)

Topology in solid-state physics (example)²

Quantum Hall effect



²Fig. from von Klitzing *et al.*, Rev. Mod. Phys. **58**, 519 (1986)

Berryology³

- ▶ time-varying Hamiltonian $H(\mathbf{R})$ with parameters $R_i = R_i(t)$

³M. Berry, Proc. R. Soc. Lond. A **392**, 45 (1984)

Berryology³

- ▶ time-varying Hamiltonian $H(\mathbf{R})$ with parameters $R_i = R_i(t)$
- ▶ adiabatic evolution along path C in the parameter space

³M. Berry, Proc. R. Soc. Lond. A **392**, 45 (1984)

Berryology³

- ▶ time-varying Hamiltonian $H(\mathbf{R})$ with parameters $R_i = R_i(t)$
- ▶ adiabatic evolution along path C in the parameter space
- ▶ instantaneous eigenstates $|n(\mathbf{R})\rangle$ of $H(\mathbf{R})$:

$$H(\mathbf{R}) |n(\mathbf{R})\rangle = E_n |n(\mathbf{R})\rangle$$

- ▶ gauge freedom of arbitrary phase

³M. Berry, Proc. R. Soc. Lond. A **392**, 45 (1984)

Berryology³

- ▶ time-varying Hamiltonian $H(\mathbf{R})$ with parameters $R_i = R_i(t)$
- ▶ adiabatic evolution along path C in the parameter space
- ▶ instantaneous eigenstates $|n(\mathbf{R})\rangle$ of $H(\mathbf{R})$:

$$H(\mathbf{R})|n(\mathbf{R})\rangle = E_n|n(\mathbf{R})\rangle$$

- ▶ gauge freedom of arbitrary phase
- ▶ How will the phase $\theta(t)$ of $|\psi(t)\rangle = e^{-i\theta(t)}|n(\mathbf{R}(t))\rangle$ evolve?
- ▶ Given by

$$H(\mathbf{R}(t))|\psi(t)\rangle = i\hbar \frac{d}{dt} |\psi(t)\rangle$$

³M. Berry, Proc. R. Soc. Lond. A **392**, 45 (1984)

- ▶ Berry phase for a closed loop $C = \partial S$

$$\begin{aligned}\gamma_n &= \int_C \mathbf{A}_n(\mathbf{R}) \cdot d\mathbf{R} \\ &= \iint \mathbf{\Omega}_n(\mathbf{R}) \cdot d\mathbf{S}\end{aligned}$$

- ▶ Berry connection

$$\mathbf{A}_n(\mathbf{R}) = i \langle n(\mathbf{R}) | \nabla_{\mathbf{R}} | n(\mathbf{R}) \rangle$$

- ▶ Berry curvature

$$\begin{aligned}\mathbf{\Omega}_n(\mathbf{R}) &= \nabla_{\mathbf{R}} \times \mathbf{A}_n(\mathbf{R}) \\ &= i \langle \nabla_{\mathbf{R}} n(\mathbf{R}) | \times | \nabla_{\mathbf{R}} n(\mathbf{R}) \rangle \\ &= i \sum_{m \neq n} \frac{\langle n(\mathbf{R}) | \nabla_{\mathbf{R}} H(\mathbf{R}) | m(\mathbf{R}) \rangle \times \langle m(\mathbf{R}) | \nabla_{\mathbf{R}} H(\mathbf{R}) | n(\mathbf{R}) \rangle}{(E_m(\mathbf{R}) - E_n(\mathbf{R})^2)}\end{aligned}$$

⁴M. Berry, Proc. R. Soc. Lond. A **392**, 45 (1984)

Berryology in periodic systems

Berryology in periodic systems

- ▶ periodicity:

Berryology in periodic systems

- ▶ periodicity:

- ▶ wave function $\psi_n(\mathbf{k}, \mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_n(\mathbf{k}, \mathbf{r})$

Berryology in periodic systems

- ▶ periodicity:
 - ▶ wave function $\psi_n(\mathbf{k}, \mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_n(\mathbf{k}, \mathbf{r})$
 - ▶ periodic \mathbf{k} -space \rightarrow Brillouin zone

Berryology in periodic systems

- ▶ periodicity:
 - ▶ wave function $\psi_n(\mathbf{k}, \mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_n(\mathbf{k}, \mathbf{r})$
 - ▶ periodic \mathbf{k} -space \rightarrow Brillouin zone
 - ▶ eff. Schrödinger equation: $H(\mathbf{k}) u_n(\mathbf{k}, \mathbf{r}) = E_n(\mathbf{k}) u_n(\mathbf{k}, \mathbf{r})$
 $\mathbf{k} \dots$ parameter of H

Berryology in periodic systems

- ▶ periodicity:
 - ▶ wave function $\psi_n(\mathbf{k}, \mathbf{r}) = e^{i\mathbf{k} \cdot \mathbf{r}} u_n(\mathbf{k}, \mathbf{r})$
 - ▶ periodic \mathbf{k} -space \rightarrow Brillouin zone
 - ▶ eff. Schrödinger equation: $H(\mathbf{k}) u_n(\mathbf{k}, \mathbf{r}) = E_n(\mathbf{k}) u_n(\mathbf{k}, \mathbf{r})$
 $\mathbf{k} \dots$ parameter of H
 \Rightarrow Berry phase of electrons in solids:

$$\varphi_n(C) = i \int_C \langle u_n(\mathbf{k}) | \nabla_{\mathbf{k}} u_n(\mathbf{k}) \rangle \cdot d\mathbf{k}$$

Example: polarization in 1D⁵

⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Example: polarization in 1D⁵


- ▶ Definition of polarization for lattice periodic systems

⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Example: polarization in 1D⁵

- ▶ Definition of polarization for lattice periodic systems

- ▶ $P = \frac{1}{a} \int_{\text{cell}} x \rho(x) dx$?




⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Example: polarization in 1D⁵

- ▶ Definition of polarization for lattice periodic systems

- ▶ $P = \frac{1}{a} \int_{\text{cell}} x \rho(x) dx$? ... doesn't work




⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Example: polarization in 1D⁵

- ▶ Definition of polarization for lattice periodic systems

- ▶ $P = \frac{1}{a} \int_{\text{cell}} x \rho(x) dx$? ... doesn't work



- ▶ Modern theory:


$$P = -\frac{e}{2\pi} \sum_n^{\text{occ}} \gamma_n = -\frac{e}{2\pi} \sum_n^{\text{occ}} \int_0^{2\pi/a} A_n(k) dk$$

⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Example: polarization in 1D⁵

- ▶ Definition of polarization for lattice periodic systems

- ▶ $P = \frac{1}{a} \int_{\text{cell}} x \rho(x) dx$? ... doesn't work



- ▶ Modern theory:

$$P = -\frac{e}{2\pi} \sum_n^{\text{occ}} \gamma_n = -\frac{e}{2\pi} \sum_n^{\text{occ}} \int_0^{2\pi/a} A_n(k) dk$$

- ▶ defined only modulo e ... Berry connection not gauge invariant

⁵Armitage and Wu, SciPost Phys. **6**, 046 (2019)

Polarization in 1D centrosymmetric crystals

Polarization in 1D centrosymmetric crystals

$$P = -P \bmod e$$

Polarization in 1D centrosymmetric crystals

$$P = -P \bmod e$$

- ▶ two possible solutions:
 - ▶ $P = 0 \bmod e$ $\gamma = 0 \bmod 2\pi \dots$ trivial
 - ▶ $P = \frac{e}{2} \bmod e$ $\gamma = \pi \bmod 2\pi \dots$ non-trivial
- ▶ P is a topological invariant!

Polarization in 1D centrosymmetric crystals

$$P = -P \bmod e$$

- ▶ two possible solutions:
 - ▶ $P = 0 \bmod e$ $\gamma = 0 \bmod 2\pi \dots$ trivial
 - ▶ $P = \frac{e}{2} \bmod e$ $\gamma = \pi \bmod 2\pi \dots$ non-trivial
- ▶ P is a topological invariant!
- ▶ macroscopic charge density $\rho(x) = -\frac{d}{dx}P(x)$

Polarization in 1D centrosymmetric crystals

$$P = -P \bmod e$$

- ▶ two possible solutions:
 - ▶ $P = 0 \bmod e$ $\gamma = 0 \bmod 2\pi \dots$ trivial
 - ▶ $P = \frac{e}{2} \bmod e$ $\gamma = \pi \bmod 2\pi \dots$ non-trivial
- ▶ P is a topological invariant!
- ▶ macroscopic charge density $\rho(x) = -\frac{d}{dx}P(x)$
 - ▶ surface charge:

$$\begin{aligned} Q_{\text{surf}} &= \int_{\text{bulk}}^{\text{vac}} dx \rho(x) \\ &= - \int_{\text{bulk}}^{\text{vac}} dx \frac{d}{dx} P(x) \\ &= -(P_{\text{vac}} - P_{\text{mat}}) \bmod e = P_{\text{mat}} + ne \\ &= ne + \begin{cases} 0 & \text{trivial} \\ e/2 & \text{non-trivial} \end{cases} \end{aligned}$$

Polarization in 1D centrosymmetric crystals

- ▶ includes basic ingredients for a topological classification

Polarization in 1D centrosymmetric crystals

- ▶ includes basic ingredients for a topological classification
 - ▶ symmetry to define (quantized) invariants

Polarization in 1D centrosymmetric crystals

- ▶ includes basic ingredients for a topological classification
 - ▶ symmetry to define (quantized) invariants
 - ▶ different values of invariant $\hat{=}$ different topological phases


Polarization in 1D centrosymmetric crystals

- ▶ includes basic ingredients for a topological classification
 - ▶ symmetry to define (quantized) invariants
 - ▶ different values of invariant $\hat{=}$ different topological phases
 - ▶ phase transition only via breaking of the adiabatic assumption
 \leftrightarrow closing of the band gap

Polarization in 1D centrosymmetric crystals

- ▶ includes basic ingredients for a topological classification
 - ▶ symmetry to define (quantized) invariants
 - ▶ different values of invariant $\hat{=}$ different topological phases
 - ▶ phase transition only via breaking of the adiabatic assumption
 \leftrightarrow closing of the band gap
 - ▶ (mostly) presence of boundary states in the band gap

1D example: SSH model⁶



tight-binding Hamiltonian:

$$H(k) = \begin{pmatrix} 0 & v + we^{-ik} \\ v + we^{ik} & 0 \end{pmatrix} = d_x(k)\sigma_x + d_y(k)\sigma_y$$

with:

$$d_x(k) = v + w \cos(k)$$


$$d_y(k) = w \sin(k)$$

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

⁶Armitage and Wu, SciPost Phys. **6**, 046 (2019)

1D example: SSH model⁷

$$E_{\pm}(k) = \pm \sqrt{v^2 + w^2 + 2vw \cos(k)}$$



⁷Short course on topological insulators, arXiv:1509.02295

1D example: SSH model

Berry phase of the SSH model:

$$\gamma = \int_{-\pi}^{\pi} dk A(k) = \frac{\pi}{2} \left[1 + \operatorname{sgn} \left(\frac{v-w}{v+w} \right) \right]$$

- ▶ $v > w$: $P = 0$ and $Q_{\text{surf}} = 0 \dots$ trivial
- ▶ $v < w$: $P = e/2$ and $Q_{\text{surf}} = e/2 \dots$ non-trivial