

Map, Filter, Reduce

In Lecture

- We have talked about how to scale a **list**

[5, 1, 4, 9, 11, 22, 12, 55]



[10, 2, 8, 18, 22, 44, 24, 110]

- Or square a **list**

[5, 1, 4, 9, 11, 22, 12, 55]



[25, 1, 16, 81, 121, 484, 144, 3025]

How to convert the function for tuple?

- List

```
def seqScaleI(seq,n):  
    output = []  
    for i in seq:  
        output.append(i*n)  
    return output
```

```
def seqScaleR(seq,n):  
    if not seq:  
        return seq  
    return [seq[0]*n]+seqScaleR(seq[1:],n)
```

- Tuple?

- Try it for 5 min
- No need to code from scratch, copy the list version and modify

How to convert the functions for tuple?

- List

- What needed to be changed?

```
def seqScaleI(seq,n):  
    output = []  
    for i in seq:  
        output.append(i*n)  
    return output
```

```
def seqScaleR(seq,n):  
    if not seq:  
        return seq  
    return [seq[0]*n]+seqScaleR(seq[1:],n)
```

- Tuple

```
def seqScaleIT(seq,n):  
    output = ()  
    for i in seq:  
        output += (n*i,)  
    return output
```

```
def seqScaleRT(seq,n):  
    if not seq:  
        return seq  
    return (seq[0]*n,)+seqScaleRT(seq[1:],n)
```

Map()

How to Sum Digits (Not square yet)?

```
>>> sds(123456789)
285
```

- Can we use map()?
 - But map only applies on list?
- Hint:
 - Given an integer N, what is

`list(str(N))`

```
>>> list(str(123456))
['1', '2', '3', '4', '5', '6']
```

- Can use map to apply “square” to them?

```
>>> list(str(123456))
['1', '2', '3', '4', '5', '6']
>>> map(lambda x:x*x, list(str(123456)))
Traceback (most recent call last):
  File "<pyshell#25>", line 1, in <module>
    map(lambda x:x*x, list(str(123456)))
  File "G:/My Drive/Courses/CS1010E/CS1010E TA Folders/Tutor
k 08 map filter fold reduce/Wk08 more about sequence Tutoria
ap
    output.append(f(i))
  File "<pyshell#25>", line 1, in <lambda>
    map(lambda x:x*x, list(str(123456)))
TypeError: can't multiply sequence by non-int of type 'str'
```

- Almost?
 - How?

```
>>> def sds(n):  
    return sum(map(lambda x:int(x),list(str(n))))  
  
>>> sds(123456789)  
45  
>>> sds(1111111111)  
10  
>>> sds(1000000000000)  
1
```

- How to sum digit squares?

```
>>> sdss(22222)  
20
```

- Your try, 5 min.

Challenge!

Write an integrated function for **BOTH** list and tuples in **ONE** single function

Taylor Series with map()


- Can we use it for Taylor series?

Use higher order functions for this?

$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1}$	$= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$	for all x
$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n}$	$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$	for all x
$\tan x = \sum_{n=1}^{\infty} \frac{B_{2n}(-4)^n(1-4^n)}{(2n)!} x^{2n-1}$	$= x + \frac{x^3}{3} + \frac{2x^5}{15} + \dots$	for $ x < \frac{\pi}{2}$
$\sec x = \sum_{n=0}^{\infty} \frac{(-1)^n E_{2n}}{(2n)!} x^{2n}$	$= 1 + \frac{x^2}{2} + \frac{5x^4}{24} + \dots$	for $ x < \frac{\pi}{2}$
$\arcsin x = \sum_{n=0}^{\infty} \frac{(2n)!}{4^n(n!)^2(2n+1)} x^{2n+1}$	$= x + \frac{x^3}{6} + \frac{3x^5}{40} + \dots$	for $ x \leq 1$
$\arccos x = \frac{\pi}{2} - \arcsin x$		
$= \frac{\pi}{2} - \sum_{n=0}^{\infty} \frac{(2n)!}{4^n(n!)^2(2n+1)} x^{2n+1}$	$= \frac{\pi}{2} - x - \frac{x^3}{6} - \frac{3x^5}{40} - \dots$	for $ x \leq 1$
$\arctan x = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} x^{2n+1}$	$= x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$	for $ x \leq 1, x \neq \pm i$

Try sin(x)? Your turn

- Note the sequence

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n}$$

$$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

for all x

- The function is (x is the angle)
 - `cf = lambda n: (x**(2*n) * ((-1)**n)) / factorial(2*n)`
- Just map the function cf to
 $[0, 1, 2, 3, 4, 5\dots]$
- Our target should be
 - `cf(0) + cf(1) + cf(2) + cf(3) + cf(4) + ...`

Try sin(x)? Your turn

- The function is (x is the angle)

- `cf = lambda n: (x**(2*n) * ((-1)**n)) / factorial(2*n)`

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \quad \text{for all } x$$

- Our target should be

- `cf(0) + cf(1) + cf(2) + cf(3) + cf(4) + ...`

```
def myCos(x):  
    def cf(n):  
        return (x**(2*n) * ((-1)**n) / factorial(2*n))  
    return sum(map(cf, range(0, 10)))
```

```
>>> myCos(3.141592654/3)
```

```
0.49999999998815835
```

```
>>> cos(3.141592654/3)
```

```
0.49999999998815835
```

Try cos(x)?

- The function is (x is the angle)

- $\text{sf} = ???$

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \quad \text{for all } x$$

- Our target should be

- $\text{sf}(0) + \text{sf}(1) + \text{sf}(2) + \text{sf}(3) + \text{sf}(4) + \dots$

```
def mySin(x):  
    def sf(n):  
        ??????????????  
    return sum(map(sf, range(0, 10)))  
  
>>> mySin(3.141592654/3)  
0.8660254038528065  
>>> sin(3.141592654/3)  
0.8660254038528065
```

And try other TS! And also log, e^x , and so on!

- Can we use it for Taylor series?

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \quad \text{for all } x$$

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \quad \text{for all } x$$

$$\tan x = \sum_{n=1}^{\infty} \frac{B_{2n}(-4)^n(1-4^n)}{(2n)!} x^{2n-1} = x + \frac{x^3}{3} + \frac{2x^5}{15} + \dots \quad \text{for } |x| < \frac{\pi}{2}$$

$$\sec x = \sum_{n=0}^{\infty} \frac{(-1)^n E_{2n}}{(2n)!} x^{2n} = 1 + \frac{x^2}{2} + \frac{5x^4}{24} + \dots \quad \text{for } |x| < \frac{\pi}{2}$$

$$\arcsin x = \sum_{n=0}^{\infty} \frac{(2n)!}{4^n (n!)^2 (2n+1)} x^{2n+1} = x + \frac{x^3}{6} + \frac{3x^5}{40} + \dots \quad \text{for } |x| \leq 1$$

$$\begin{aligned} \arccos x &= \frac{\pi}{2} - \arcsin x \\ &= \frac{\pi}{2} - \sum_{n=0}^{\infty} \frac{(2n)!}{4^n (n!)^2 (2n+1)} x^{2n+1} = \frac{\pi}{2} - x - \frac{x^3}{6} - \frac{3x^5}{40} - \dots \end{aligned} \quad \text{for } |x| \leq 1$$

$$\arctan x = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} x^{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \quad \text{for } |x| \leq 1, x \neq \pm i$$

Obverse the Function “reduce()”

- What does this following do?

```
>>> reduce(lambda x,y:x+y, [1,2,3])  
6
```

- first = seq[0] (first == 1)
- for i in [2,3]:
 - i = 2:
 - first = 1 + 2
 - i = 3
 - first = 3 + 3
- return first (first == 6)

```
def reduce(f, seq):  
    if not seq:  
        return seq  
    first = seq[0]  
    for i in seq[1:]:  
        first = f(first, i)  
    return first
```

Obverse the Function “reduce()”

- In general, let f be a function that takes two arguments

`reduce(f, [1, 2, 3, 4, 5, 6,])`
↓
`f(f(f(f(f(1, 2), 3), 4), 5), 6)`

```
def reduce(f, seq):  
    if not seq:  
        return seq  
    first = seq[0]  
    for i in seq[1:]:  
        first = f(first, i)  
    return first
```

- If f is the addition function

`reduce(f, [1, 2, 3, 4, 5, 6,])`
↓
`(((((1+2)+3)+4)+5)+6)`

Obverse the Function “reduce()”

```
>>> reduce(lambda x,y:x+y, [1,2,3])  
6
```

- This the summation function:

$$\sum_{i=0}^{i<n} seq[i] = seq[0] + seq[1] + seq[2] \dots$$

- How to write the following?

$$\prod_{i=0}^{i<n} seq[i] = seq[0] \times seq[1] \times seq[2] \dots$$

- Try it yourself?

• `[1,2,3,4,5,6,7]` → `5040`

```
def reduce(f, seq):  
    if not seq:  
        return seq  
    first = seq[0]  
    for i in seq[1:]:  
        first = f(first, i)  
    return first
```

History of “reduce()”

- In 1994, reduce() was a built-in function for Python
- Around 2016, reduce() was moved to a package called functools
 - [The fate of reduce\(\) in Python 3000](#)

```
>>> from functools import reduce
>>> reduce(lambda x,y:x+y, [1,2,3])
6
>>> reduce(lambda x,y:x+y, (1,2,3))
6
>>> reduce(lambda x,y:x+y, ('a','b','c'))
'abc'
```

- In the same document, you can see that two convenient functions was added
 - any(), all()

any() and all()

- If L = [1,2,3,4], is there any number that is greater than 3 in L?

```
>>> L = [1,2,3,4]
>>> any(x > 3 for x in L)
True
```

- If L = [1,2,3,4], is there any number that is greater than 9 in L?

```
>>> any(x > 9 for x in L)
False
```

- Is there any prime number in the lists?

```
>>> any(isPrime(x) for x in [4,6,8,9,99])
False
>>> any(isPrime(x) for x in [4,6,8,9,97,99])
True
```

any() and all()

- If L = [1,2,3,4], are all numbers in L greater than 3 ? (and 0?)

```
>>> all(x > 3 for x in L)
False
>>> all(x > 0 for x in L)
True
```

- Are all the numbers in the lists are prime numbers?

```
>>> all(isPrime(x) for x in [4,6,8,9,99])
False
>>> all(isPrime(x) for x in [3,5,7,11,97])
True
```