

**Q function:**  $Q(x) = \int_x^\infty \frac{1}{\sqrt{2\pi}} e^{t^2/2} dt$

Transform a Gaussian Distribution to Normal Distribution:

$$F_X \sim \mathcal{N}(m, \sigma^2)$$

$$F_X(x) = \Phi\left(\frac{x-m}{\sigma}\right), \text{ where } \Phi \sim \mathcal{N}(0, 1)$$

$$Tail(x) = Q\left(\frac{x-m}{\sigma}\right)$$

Some bounds:  $Q(x) \leq e^{-x^2/2}$

$$Q(x) < \frac{1}{\sqrt{2\pi}x^2} e^{-x^2/2}$$

$$Q(x) > \left(1 - \frac{1}{x^2}\right) \frac{1}{\sqrt{2\pi}x^2} e^{-x^2/2}$$

$$Q(x) \leq \frac{1}{2} e^{-x^2/2}$$

$$\text{Chebyshv: } Q(x) \leq \frac{1}{2x^2}$$

**Markov Inequality:**  $P(x \leq a) \leq \frac{E(x)}{a}$

**Chebyshv Inequality:**  $P(|x - m| \geq a) \leq \frac{\sigma^2}{a^2}$

**Chernoff Bound:**  $P(X \leq a) \leq e^{-\tilde{v}a} E(e^{\tilde{v}X})$

Chernoff Bound holds for  $\tilde{v} > 0$ , the *tightest* bound achieves when

$$\frac{d}{d\tilde{v}} (E(e^{\tilde{v}X})) = E(X e^{\tilde{v}X}) = a E(e^{\tilde{v}X})$$

**Poisson Process:**  $Pr(N(t) = k) = \frac{(\lambda t)^k}{k!} e^{-\lambda t}$

$N(t)$  stands for number of occurrence, and this is a *memoryless* process.

## Real Gaussian Vector

$$\vec{w} \sim \mathcal{N}(0, I)$$

$$\text{Definition: } P(\vec{w}) = \frac{1}{(\sqrt{2\pi})^n} e^{-\left(\frac{\|\vec{w}\|^2}{2}\right)}$$

Each definition of  $\vec{w}$  follows  $\mathcal{N}(0, 1)$

$$\text{For } \vec{x} = A\vec{w} + \vec{b}$$

$$p(\vec{x}) = \frac{1}{(\sqrt{2\pi})^n \sqrt{\det(A^T A)}} e^{\frac{1}{2}(\vec{x}-\vec{b})^T (A^T A)^{-1} (\vec{x}-\vec{b})}$$

Moreover  $\vec{x} \sim \mathcal{N}(\vec{b}, AA^T)$

And  $\forall \vec{c} \in \mathcal{R}^n$ ,  $\vec{c}^T \vec{x} \sim \mathcal{N}(\vec{c}^T \vec{b}, \vec{c}^T AA^T \vec{c})$

For 1D case, if  $X \sim \mathcal{N}(0, \sigma^2)$ , then  $aX + b \sim \mathcal{N}(b, a^2 \sigma^2)$

## Complex Random Variable

$$x \sim \mathcal{CN}(0, 1) \text{ (or } x \sim \mathcal{CN}(0, \sigma^2))$$

*Real* and *Imaginary* components yields

$$Re, Im \sim \mathcal{N}(0, 1/2) \text{ (or } Re, Im \sim \mathcal{N}(0, \sigma^2/2))$$

The *angle* is uniformly distributed on  $[0, 2\pi)$

$|x|$  follows *Rayleigh* Distribution

$$P(r) = 2re^{-r^2} \text{ for } \mathcal{CN}(0, 1)$$

$$\text{(or } P(r) = \frac{r}{\sigma_0^2} e^{-\frac{r^2}{2\sigma_0^2}} \text{ for } \mathcal{CN}(0, \sigma^2),$$

$$\sigma_0^2 = \sigma^2/2 \text{ is for } Re \text{ and } Im \text{ component})$$

## Relations between different distributions

For  $X_1 \sim \mathcal{N}(m_1, \sigma^2)$ ,  $X_2 \sim \mathcal{N}(m_2, \sigma^2)$

If  $m_1 = m_2 = 0$ ,  $X = \sqrt{X_1^2 + X_2^2}$  follows *Rayleigh* Distribution,  $p(x) = \frac{x}{\sigma^2} e^{-\frac{x^2}{2\sigma^2}}$ ,  $x > 0$

If  $m_1, m_2 > 0$ ,  $X$  follows *Ricean* Distribution,

$$p(x) = \frac{x}{\sigma^2} I_0\left(\frac{sx}{\sigma^2}\right) e^{-\frac{x^2+s^2}{2\sigma^2}}, \text{ where } s = \sqrt{m_1^2 + m_2^2},$$

$$I_0(x) = \frac{1}{2\pi} \int_0^{2\pi} e^{xcos\theta} d\theta$$

For  $Z_i \sim \mathcal{N}(0, 1)$ , then  $\sum_{i=1}^k Z_i^2$  follows *Chi-Square* distribution with *degree of freedom* k.

$$p(x, k) = \frac{x^{(k/2-1)} e^{-x/2}}{2^{k/2} \Gamma(k/2)}, x > 0.$$

$$(p(x, 1) = \frac{e^{-x/2}}{\sqrt{2x\pi}},$$

$$\Gamma(t) = \int_0^\infty x^{t-1} e^{-x} dx, \Gamma(t) = t! \text{ if } t \text{ is integer})$$

*Nakagami* Random Variable is typical for Fading Channel.

$$p(x) = \frac{2}{\Gamma(m)} \left(\frac{m}{\Omega}\right)^m x^{2m-1} e^{-mx^2/\Omega}, x > 0, \text{ where}$$

$\Omega = E(x^2)$ , and  $m = \frac{\Omega^2}{E[(x^2 - \Omega)^2]}$ ,  $m \leq 1/2$ , called fading figure.

### Lloyd-Max Algorithm

Notice  $b_0 = 0, b_n = \infty$ , update algorithm:

$$\text{representation point } a_i = \frac{\int_{b_{i-1}}^{b_i} x f_X(x) dx}{\int_{b_{i-1}}^{b_i} f_X(x) dx},$$

which is basically the distribution ‘center’

and the *endpoint*  $b_i = \frac{a_{i-1} + a_i}{2}$

### Nyquist Criterion

$g(t) = p(t) * h(t) * q(t)$  satisfies no *inter-symbol interference* property

$$g(jT - kT) = \begin{cases} 1, & \text{if } j = k \\ 0, & \text{otherwise} \end{cases}$$

in frequency domain,

$$\sum_{m \in \mathbb{Z}} \hat{g}(f - m/T) \text{rect}(fT) = T \text{rect}(fT)$$

which can be simplified to

$$\frac{1}{T} \sum_{m \in \mathbb{Z}} \hat{g}(f - m/T) = 1$$

### Gaussian Process

Definition:  $\{Z(t)\}$ , for  $i$  in a finite set of  $\{n\}$ ,

$\{Z(t_i)\}$  are jointly Gaussian set of r.v.

Basically *i.i.d. Gaussian r.v.* on each time instance.

Let  $\{Z(t)\}$  be White Gaussian Noise Process.

$V = \int g(t)Z(t)dt$  will be Gaussian r.v. with zero mean.

Notice from *Linear Functional of WSS process*

$$E[V^2] = \int_{-\infty}^{\infty} |\hat{g}(f)|^2 S_Z(f) df,$$

where  $S_Z(f)$  is spectral density.

Now note that  $\sigma^2 = E[V^2] - E[V]^2$

Typically,  $E[V] = E[\int g(t)Z(t)dt] = 0$ ,

because  $Z$  is zero-mean

$$E[V^2] = E[\int g(t)Z(t)dt \int g(\tau)Z(\tau)d\tau]$$

$$= \int \int E[Z(t)Z(\tau)]g(t)g(\tau)dt d\tau$$

$$= \int \int \frac{N_0}{2} \delta(t - \tau) g(t)g(\tau)dt d\tau$$

(This is because the gaussian r.v. on different time are independent, and  $E(Z^2) - E(Z)^2 = \sigma^2 = N_0/2$ , assuming spectral density  $\frac{N_0}{2}$ )

### Error Probability

*bit error probability*:  $Pr(e) = Q(\frac{d(a_0, a_1)/2}{\sigma}) = Q(\frac{d(a_0, a_1)}{\sqrt{2N_0}})$ , assuming WGN spectral density  $N_0/2$

### Symbol distance & Symbol Energy

In general, the energy for a point in constellation graph is its *norm*<sup>2</sup>, take mean of them for the average symbol energy

Further, bit energy  $E_b = E_s / \log_2 M$

*FSK*:  $d = \sqrt{2E_s}$  (holds for every two symbols)

$$M\text{-PSK}: d = 2\sqrt{E_s} \sin(\pi/M)$$

$$M\text{-PAM}/M\text{-ASK}: d = \sqrt{\frac{12E_{savg}}{M^2 - 1}}$$

*M-QAM*: just derive from stretch, but for  $M$

$$= 4, 16, 64, 256 \quad d = \sqrt{\frac{6E_{savg}}{M - 1}}$$

$$P_c(M - QAM) = P_c^2(\sqrt{M} - PAM) = (1 - P_e(\sqrt{M} - PAM))^2$$

$$P_e(M - QAM) = 1 - (1 - P_e(\sqrt{M} - PAM))^2$$