Non-Specialist Companion Guide: Emergent Matter, Fermionic Spin, and Induced Gravity from Topological Wave Solitons

Eden Asipov and Eliana Asipov
Independent Researchers

Companion Overview to the Research Manuscript: "Emergent Matter, Fermionic Spin, and Induced Rank-2 Gravity from Topological Wave Solitons in a Polarizable Vacuum Medium"

---

1. Executive Summary: What Is the Big Picture?

Modern physics describes the universe using two different rulebooks that do not fit smoothly together:

1. **Quantum Field Theory (QFT)** treats elementary matter particles (like quarks and electrons) as *infinitely small points* of zero size moving through space. Because their size is assumed to be zero, dividing by distance $r \to 0$ creates mathematical infinities right at the center of every particle.
2. **Einstein's General Relativity (GR)** treats gravity as the smooth bending of empty spacetime geometry. When combined with point-like masses, it predicts infinite-density singularities at the centers of black holes, particles, and the Big Bang.

How This Framework Offers a Unified Perspective

Instead of separating the world into "empty geometry" and "zero-size point particles," this research models the vacuum as a continuous, elastic, relativistic **wave medium**. In this medium, matter particles are not zero-size points—they are **stable, three-dimensional topological wave knots** (called *solitons*).

Thinking of particles as self-trapped wave knots addresses several longstanding puzzles in a single mathematical framework:

- **Why particles come in whole units:** You cannot put "half a twist" into a closed loop of rope. Topology forces wave knots to carry whole-number twists ($B = 1, 2, 3, \dots$), explaining why proton and neutron numbers (baryon number) are strictly conserved.
- **Where mass ($E = mc^2$) and realistic nuclear binding come from:** Over 92% of a proton's mass comes directly from the trapped wave tension inside its finite core ($\sim 0.76 \times 10^{-15}$ m across), without needing point particles. Moreover, when protons and neutrons bind into atomic nuclei like the deuteron ($B = 2$) or helium-4 ($B = 4$), short-range vector-meson repulsion ($\mathcal{L}_6$) balances two-pion attraction so nuclei bind softly (1.98 MeV and 28.6 MeV), matching experimental nuclear physics.
- **Why fractional pieces ("quarks / merons") cannot escape:** A proton knot is built from three fractional 1/3-twist wave centers locked in an equilateral triangle. Pulling one piece away stretches a wave-tension tube whose energy grows with distance—like stretching an unbreakable rubber band—while an isolated 1/3 piece oscillates above the vacuum cutoff frequency and rapidly radiates away into ripples.
- **Why gravity bends light and radiates gravitational waves:** The concentrated wave energy of a soliton polarizes and stretches the surrounding vacuum medium. Near a massive body, this changes the vacuum's effective refractive index and elastic frame, slowing down wave clocks and contracting wave rulers—reproducing Einstein's light bending, planetary orbit precession, spinning frame-dragging, and gravitational waves, while keeping gravity completely finite at $r = 0$.
- **How the universe expands without a Big Bang singularity, a vacuum-energy catastrophe, or mystery dark-matter particles:** On cosmic scales, the uniform dielectric state of the vacuum medium evolves over time ($a(t)$). Because proton and atom wave knots are held at a fixed size by their internal wave stiffness, they do not stretch as the universe expands—whereas free light waves traveling between galaxies stretch and redshift ($1 + z = a_0/a(t)$). Running the clock backward to the early universe, the strong short-range repulsion of overlapping soliton cores ($\mathcal{L}_6$) and spin-torsion prevents an infinite-density Big Bang singularity and triggers a **smooth cosmic bounce and early inflation**.

Meanwhile, because an unconstrained wave medium in static equilibrium has zero net pressure, the naive 10^{120} vacuum-energy catastrophe cancels out, leaving only a tiny residual horizon energy set by the pion wave scale ($\sim$ 2.5 meV) that matches observed dark energy, while topologically neutral ($B = 1 - 1 = 0$) knot–antiknot breather states formed during the early chiral phase transition act as dark matter ($\Omega_{\rm DM}/\Omega_B \approx 5.44$).

---

2. Section-by-Section Guide for Non-Specialists

Guide to Section 2: Why Does a Wave Knot Need Stiffness, and Where Does It Come From?

If you make a wave packet in ordinary water or air, it quickly spreads out and disappears. On the other hand, if you have a simple nonlinear wave that pulls inward without resistance, it collapses down to an infinitely sharp point. To form a stable, finite-sized particle (a soliton), the wave medium must have both **surface tension** (which pulls the knot inward) and **short-distance bending stiffness** (which pushes outward when the knot is squeezed too tightly).

In early soliton models, physicists added higher-derivative "stiffness formulas" ($\mathcal{L}_4$ and $\mathcal{L}_6$) by hand just to prevent collapse. In Section 2 of the paper, we show how those exact stiffness terms arise naturally from standard short-range vector waves (ρ and ω mesons) already known to exist in nuclear physics. Because the strengths of those vector waves are measured independently in meson experiments, the stiffness of the proton knot is tied directly to known physics rather than introduced as an arbitrary free parameter.

Guide to Section 3: How Continuous Waves Lock into Discrete Particles—and Why "Quarks" Are Confined

Imagine taking a flat rubber sheet, pulling all of its outer edges together to form a closed sphere, and wrapping that sheet around a globe. You can wrap it around the globe zero times, once, twice, or three times—but you can never wrap a closed sheet around a globe 1.37 times. That whole number of wrappings is a **topological invariant** ($B \in \mathbb{Z}$). In our 3D wave medium, a proton or neutron is a wave configuration that wraps internal field space once ($B = 1$) across 3D space (illustrated in **Figure C1(a)**). Because the wrapping number cannot change smoothly, the particle cannot simply dissolve away.

Inside a single $B = 1$ baryon knot, projecting the 3-sphere onto its internal symmetry axes (the mathematical **flag manifold** $F_2 = SU(3)/U(1)^2$) splits the integer twist equally among **three fractional wave centers ("merons")**, each carrying $B = 1/3$ of the twist (**Figure C1(b)**). Our 3D wave simulations demonstrate two physical reasons why these 1/3 pieces are never seen alone in nature:

1. **Frequency Trapping:** An isolated 1/3 wave entity oscillates *faster* than the minimum cutoff frequency of free vacuum waves ($1.14 \times m_0$), so it rapidly sheds its energy as radiation and disperses. When three 1/3 entities lock together at 120° phase angles, their mutual attraction slows their collective frequency *below* the vacuum cutoff ($0.795 \times m_0$). Because the surrounding vacuum cannot propagate waves below its cutoff frequency, the 3-entity proton is permanently trapped and stable.
2. **Elastic Flux-Tube Confinement:** If you try to pull one of the 1/3 centers away from the other two, a tube of twisted wave energy stretches between them whose energy grows linearly with distance—acting like an unbreakable elastic string (**Figure C1(b)**).




Figure C1: Anatomy of 3D topological wave knots. **(a)** Unlike a point particle whose field diverges to infinity at $r \to 0$ (red dashed curve), a $B = 1$ wave knot twists smoothly from 180° at its center to 0° at infinity (blue curve), where the classical static soliton core accounts for 92.2% of the proton's rest mass (with the remaining 7.8% from quantum spin rotation) within an RMS radius of $R_0 \approx 0.76$ fm (green shaded region). **(b)** Projecting the $B = 1$ knot onto the flag manifold $F_2 = SU(3)/U(1)^2$ reveals three fractional $B = 1/3$ meron ("quark") cores locked at 120° phase angles and bound by an elastic Y-junction wave-tension flux tube. **(c)** When a wave knot twists only the 2-sphere electromagnetic subfield ($\pi_3(S^2) = \mathbb{Z}$) with zero baryon twist ($B = 0$), its field fibers form a linked toroidal Hopfion loop ($Q_H = 1$) representing the electron.

Guide to Section 4: How a 3D Wave Knot Behaves Like a Spin-1/2 Fermion, Obeys $E = mc^2$, and Binds into Atomic Nuclei

Normal everyday objects (like a basketball) return to their exact starting state when rotated by 360° (2π). Electrons, protons, and neutrons are different: if you rotate a proton by 360°, its quantum wavefunction flips sign ($\Psi \to -\Psi$), and you must rotate it by a full 720° (4π) to bring it back to $+\Psi$ (**Figure C2(a)**). That 720° property is the definition of a **spin-1/2 fermion** and the reason matter obeys the Pauli Exclusion Principle (so atoms have volume and do not collapse).

A 3D topological knot in a continuous wave medium has this exact property built into its geometry (analogous to the "Dirac belt trick"): rotating the core of a $B = 1$ knot by 360° while holding the distant vacuum fixed twists the surrounding field in a way that cannot be untangled without a second 360° rotation (**Figure C2(a)**). Furthermore, when the wave knot moves at velocity v, the finite wave speed c of the medium automatically flattens the knot along its direction of motion (Lorentz contraction) and creates a beat-frequency wave (the de Broglie quantum wavelength, **Figure C2(b)**), yielding Einstein's exact relativistic momentum $p = \gamma mv$ and rest mass $m = E_{\text{rest}}/c^2$. Finally, because short-range vector-meson core repulsion ($\mathcal{L}_6$) balances two-pion attraction, multiple nucleon knots bind together softly (1.98 MeV for the deuteron and 28.6 MeV for helium-4, **Figure C2(c)**) instead of collapsing into an overbound lump.




Figure C2: Emergent spin-1/2 behavior, relativistic motion, and soft nuclear binding. **(a)** The 720° topological "belt trick": rotating a 3D wave knot by 360° twists its surrounding field tethers and flips its quantum sign ($\Psi \to -\Psi$); a full 720° rotation is required to untangle the field and restore $+\Psi$. **(b)** A wave knot moving at $v = 0.75c$ automatically undergoes Lorentz contraction (R_0/γ, green ellipse) and develops de Broglie phase ripples of wavelength $\lambda_{\text{dB}} = h/(\gamma mv)$. **(c)** Short-range ω-meson core stiffness ($\mathcal{L}_6$ repulsion, purple dashed curve) balances two-pion attraction to create a shallow 1.98 MeV deuteron binding pocket (green dot) rather than collapsing.

Guide to Section 5: How a Polarizable Wave Medium Reproduces Gravity, Gravitational Waves, and Cosmic Expansion

In glass or water, light slows down and bends because the material has an optical **refractive index** $n > 1$. In our framework, the vacuum medium's refractive index increases slightly near a heavy soliton knot: $n(r) = e^{-2\Phi(r)/c^2} \approx 1 + \frac{2GM}{rc^2}$ (**Figure C3(a)**).

1. **Time Dilation & Ruler Contraction:** Because every clock is made of oscillating solitons and every ruler is held together by wave forces, a higher refractive index $n(r)$ slows down soliton oscillation frequencies (gravitational time dilation) and shrinks soliton equilibrium sizes (spatial contraction) in the exact 50/50 ratio of Einstein's General Relativity—bending starlight by twice the Newtonian prediction ($\gamma_{\rm PPN} = 1$, **Figure C3(a)**) and precessing Mercury's orbit ($\beta_{\rm PPN} = 1$).
2. **No Infinite Singularity at $r = 0$:** Because a soliton knot is not a zero-size point, its mass is spread smoothly over its core radius R_0. Inside the core, gravity smoothly levels off to zero force at the center rather than diverging to infinity.
3. **Spinning Frame-Dragging & Gravitational Waves:** A single scalar number $n(r)$ is enough for static gravity, but when solitons *spin* or *orbit* each other, they shear the vacuum medium's directional frame. Using Andrei Sakharov's quantum vacuum elasticity mechanism, the medium develops full rank-2 elastic stress-strain dynamics—producing spinning frame-dragging (Gravity Probe B) and the two transverse ripples (h_+ and $h_\times$) of gravitational waves observed by LIGO.
4. **Cosmic Expansion, Non-Singular Bounce, Dark Energy & Neutral $B = 0$ Dark Matter:**
   - **Why Bound Rulers Don't Expand (Cosmological Redshift):** A proton or atom is a self-bound soliton whose size $R_0 \approx 0.76$ fm is locked by local stiff nuclear wave forces ($\mathcal{L}_4 + \mathcal{L}_6$), whereas light traveling between galaxies is an unbound linear wave ripple that stretches with the universal background dielectric factor $a(t)$ (**Figure C3(b)**). Thus, galactic distances and light wavelengths stretch ($1 + z = a_0/a(t)$) while nuclear and atomic rulers remain invariant to 1 part in 10^{83}.
   - **Replacing the Big Bang Singularity with a Smooth Bounce:** Running cosmic expansion backward in time, squeezing topological soliton knots together activates the ultra-stiff repulsive sextic core pressure ($P_{\mathcal{L}_6} \propto a^{-6}$) and spin-torsion repulsion, causing the early universe to smoothly **bounce** at a finite minimum scale factor $a_{\rm min} > 0$ (**Figure C3(c)**) and transition into **Starobinsky R^2 inflation**.
   - **Why Dark Energy Is Tiny Instead of $10^{120}\times$ Too Large:** As shown in condensed-matter physics by Grigory Volovik, any self-sustained medium in static equilibrium automatically cancels its own bulk vacuum pressure ($P^{\rm eq}_{\rm vac} = -\rho^{\rm eq}_{\rm vac} c^2 = 0$). Only because the universe is *expanding* with a finite causal horizon $R_H = c/H_0$ is the pion chiral condensate slightly shifted out of equilibrium—leaving a tiny residual **Chiral-Horizon Dark Energy** $\rho_\Lambda = \frac{\hbar H_0}{(2\pi)^2} f_\pi^2 m_\pi \approx (2.56\ \text{meV})^4$ that matches the observed dark energy density $(2.24\ \text{meV})^4$ to within 14%.
   - **Neutral $B = 0$ Wave-Knot Breathers as Dark Matter ($\Omega_{\rm DM}/\Omega_B \approx 5.44$):** When the early universe cooled through the chiral phase transition ($T_c \approx 160$ MeV), wave-medium domain formation produced not only net-twist baryons ($B = \pm 1$), but also metastable, topologically neutral **skyrmion–antiskyrmion breather knots** ($B = 1 - 1 = 0$) with zero electric charge. Because they carry no net baryon twist or charge, they do not interact with light or nuclear forces, gravitating with an exploratory freeze-out mass ratio $\Omega_{\rm DM}/\Omega_B \approx \pi\sqrt{3} \approx 5.44$ that closely matches the Planck satellite measurement (5.40) without introducing new fundamental particles.




Figure C3: Refractive vacuum gravity and non-singular cosmology. **(a)** A massive soliton polarizes the surrounding vacuum refractive index $n(r)$, bending passing light rays by the exact Einstein angle ($\gamma_{\rm PPN} = 1$) while remaining regular inside the finite core R_0. **(b)** As the cosmic vacuum background expands from $a = 0.5$ to $a_0 = 1.0$, unbound intergalactic light waves stretch and redshift ($1 + z = a_0/a$) while self-bound soliton knots keep a fixed ruler size R_0. **(c)** At extreme early-universe densities, repulsive $\mathcal{L}_6$ core stiffness prevents a Big Bang singularity (red dotted curve) and produces a smooth cosmic bounce at finite scale $a_{\rm min} > 0$ (green solid curve).

Guide to Section 6: How a Wave Model Explains Quantum Entanglement Without Faster-Than-Light Signals

John Stewart Bell proved that if two particles are just two separate waves moving independently through ordinary 3D space, their spin correlations can never exceed a strict limit ($S \leq 2.0$). Yet laboratory experiments consistently measure $S \approx 2.828$ ($2\sqrt{2}$).

In our framework, while a *single* classical soliton lives in 3D space, the **quantum state of a pair of solitons** (Level 0 in our hierarchy) is a joint wave on the **combined 6-dimensional orientation space** of both knots ($SU(2) \times SU(2)$). When two spin-1/2 knots are created together with zero total spin (a "singlet"), they share a single non-separable orientation wavefunction in configuration space—producing the exact quantum correlation $S = 2\sqrt{2} \approx 2.8284$, while any physical energy or signal traveling through 3D space still obeys the local speed of light $c/n(\mathbf{x})$.

Guide to Section 7: Why High-Energy Colliders See Three Hard "Partons," and Where Electrons and Neutrinos Fit In

1. **High-Energy Scattering (Why Protons Look Like 3 Point-Like Quarks at High Energy):** If you bounce low-energy electrons off a proton, it looks like a smooth sphere about 0.8 fm across. However, because the proton is built from **three** $B = 1/3$ **meron cores** coupled to vector mesons (**Figure C1(b)**), relativistic Lorentz contraction at high collider energies flattens the three meron cores along the boost direction, causing high-energy electrons to scatter with a $1/Q^4$ power law—matching accelerator measurements.
2. **Left-Handed vs. Right-Handed Particles (Chiral Fermions):** Nature's weak nuclear force is "left-handed." While the Nielsen–Ninomiya theorem shows that ordinary flat-space waves produce equal numbers of left- and right-handed states, a topological soliton knot twists the background field by 180° from its center ($F(0) = \pi$) to infinity ($F(\infty) = 0$). Its core boundary acts as a **topological domain wall** that traps an unpaired left-handed zero-energy fermion state while gapping the right-handed partner.
3. **Leptons (Electrons, Muons, Taus) as Electromagnetic Hopfion Knots:** Protons and neutrons twist the full 3-sphere of the strong nuclear field ($\pi_3(S^3)$). When a wave knot only twists the 2-sphere of the electromagnetic subfield ($\pi_3(S^2)$), it forms a **linked toroidal knot called a Hopfion** ($Q_H = \pm 1$, **Figure C1(c)**). Because Hopfions have zero baryon number ($B = 0$), they do not feel the strong nuclear force, and 1-loop QED vacuum polarization naturally sets their mass scale $\sim 1836\times$ lighter than the proton ($m_e^{\text{rad}} = 0.509$ MeV vs. 0.511 MeV measured).

Guide to Section 8: Observational Tests, Practical Laboratory Engineering, and Honest Limits

- **Three Sharp Observational & Experimental Tests:**
  1. *Inside Neutron Stars (LIGO & NICER):* Squeezing $B = 1$ wave knots above $3.5\times$ normal nuclear density causes them to transition into a stiffer crystal of half-knots ($B = 1/2$) backed by repulsive $\mathcal{L}_6$ cores—predicting a peak in the nuclear sound speed ($c_s^2/c^2 \approx 0.58$), maximum neutron star masses of $2.22 \pm 0.08\, M_\odot$, and a tidal deformability $\Lambda_{1.4} = 410 \pm 45$.
  2. *High-Energy Electron-Proton Scattering (Jefferson Lab 12 GeV):* Relativistic contraction of the three $B = 1/3$ meron cores predicts that the proton's electric-to-magnetic form-factor ratio $\mu_p G_E^p / G_M^p$ crosses zero at $Q_0^2 \approx 9.6 \pm 1.2$ GeV^2.
  3. *Gravitational-Wave Merger Phase Shift:* The collective skyrmion crystal stiffness predicts a -2.4 rad 5PN tidal phase shift at 1000 Hz in binary neutron-star mergers.
- **Practical Laboratory Engineering:** Even though nuclear solitons are 10^{-15} m across, the *same mathematical equations* govern **magnetic skyrmions and hopfions** in thin-film metals, **3D-printed graded-index (GRIN) metamaterial lenses**, **sub-cutoff multipole-canceling waveguide traps**, and **disorder-protected topological resonators**.
- **Honest Scientific Scope:** The framework is a **non-perturbative effective theory**: it excels at describing bound particles, nuclear forces, spin, and post-Newtonian/induced gravity below its ultraviolet breakdown scale ($Q^2 \lesssim$ 15–25 GeV^2), while at extreme collider energies ($Q^2 \gg 25$ GeV^2) standard perturbative quark-gluon scattering (QCD) remains the appropriate mathematical description.