



---

# Decentralizing Agriculture

**A shift in the industrial paradigm of food production**

Marco Zolla Salazar

AuthorId

Faculty

Major

bei

Employer

**Kontakt:**

marco.zolla@proton.me

ContactTitle

# Contents

|          |                                                                                       |           |
|----------|---------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                                   | <b>3</b>  |
| 1.1      | Motivation . . . . .                                                                  | 3         |
| 1.2      | Project vision . . . . .                                                              | 3         |
| <b>2</b> | <b>The problems of industrial agriculture</b>                                         | <b>4</b>  |
| 2.1      | Land Use . . . . .                                                                    | 4         |
| 2.2      | Biodiversity loss . . . . .                                                           | 5         |
| 2.3      | Soil degradation and erosion . . . . .                                                | 6         |
| 2.4      | Fertilizer use . . . . .                                                              | 7         |
| 2.5      | Pesticide use . . . . .                                                               | 8         |
| 2.6      | Monoculture . . . . .                                                                 | 9         |
| 2.6.1    | The monoculture effect: genetic vulnerability to disease and pest outbreaks . . . . . | 9         |
| 2.6.2    | Historical and contemporary case studies . . . . .                                    | 10        |
| 2.6.3    | Soil degradation . . . . .                                                            | 11        |
| 2.6.4    | Yield penalties and the diversification dividend . . . . .                            | 12        |
| 2.6.5    | Diversification as a systemic alternative . . . . .                                   | 12        |
| 2.6.6    | Structural drivers: market concentration and monoculture . . . . .                    | 13        |
| 2.7      | Freshwater depletion . . . . .                                                        | 14        |
| 2.8      | Greenhouse gas emissions . . . . .                                                    | 16        |
| 2.9      | Produce transportation, storage and disposal . . . . .                                | 17        |
| 2.10     | The existential dependency on a few producers . . . . .                               | 18        |
| 2.11     | Suppression of dietary diversity and nutritional quality . . . . .                    | 19        |
| 2.12     | A systemic crisis . . . . .                                                           | 20        |
|          | <b>References</b>                                                                     | <b>22</b> |

# Abstract

Industrial agriculture is responsible for the majority of global biodiversity loss, roughly one third of all greenhouse gas emissions, and the creation of fragile, highly concentrated food supply chains on which billions of people existentially depend. This paper proposes an open-source controlled environment agriculture (CEA) platform designed to address these problems through decentralisation, automation, and a shared data commons.

The system comprises a sensor array and camera mounted in a small-scale greenhouse, a microcontroller running a PID control loop over environmental actuators, a single-board computer performing edge ML inference, and an optional connection to a cloud-based data flywheel where growth models are continuously retrained on anonymised telemetry from the global network of deployed nodes. A computer vision pipeline provides growth metrics, disease diagnosis, and species recognition. A climate-aware species recommendation engine minimises energy consumption by matching recommended crops to the local ambient environment. All hardware designs, firmware, software, trained models, and collected data are released under open licences, enabling a virtuous cycle: the more nodes deployed, the richer the shared dataset, and the more accurate the growth optimisation.

The project does not aim to replace field agriculture wholesale. Its niche is high-value, perishable, water-intensive crops grown in proximity to consumers — the segment where the environmental and supply-chain benefits of CEA are largest and most defensible. Within that niche, we argue the platform can meaningfully reduce land use pressure, water consumption, fertiliser runoff, pesticide pollution, transport emissions, food waste, and supply-chain concentration, while enabling a diversity of crops that the industrial monoculture model structurally cannot support.

# 1 Introduction

## 1.1 Motivation

The global food system is in a state of structural crisis. Industrial agricultural production — the dominant paradigm for the past century — has delivered extraordinary increases in caloric output, but at costs that are becoming impossible to ignore: ecosystem collapse, soil exhaustion, water depletion, climate destabilisation, and the creation of food supply chains so concentrated that a single geopolitical shock can threaten the nutrition of hundreds of millions of people.

At the same time, the tools to build something better have never been more accessible. Capable micro-controllers cost a few dollars. Machine learning frameworks are open source. Camera modules capable of running computer vision inference are commodity hardware. The question is no longer whether small-scale, intelligent, automated food production is technically feasible — it is — but whether it can be made cheap enough, robust enough, and open enough to displace the incumbent system at meaningful scale.

This paper describes a platform designed to make that possible.

## 1.2 Project vision

The project’s core proposition is that a network of many small, locally operated, data-connected greenhouses can collectively outperform a system of few large, centralised farms — on environmental metrics, on resilience, and eventually on cost — provided that the intelligence required to operate them efficiently is shared as a common good rather than locked up in proprietary systems.

The platform is designed around four interlocking principles:

1. **Edge-first control.** All critical control logic runs locally and offline. The system operates without internet connectivity. The cloud connection is optional and asynchronous, used only for learning and coordination, never for real-time operation.
2. **Open everything.** Hardware reference designs, firmware, software, trained models, and the collected dataset are all released under maximally permissive open licences. Anyone can deploy, modify, and contribute back.
3. **Data flywheel.** Every deployed node, with the user’s consent, contributes anonymised telemetry to a shared dataset. Models trained on this dataset are continuously improved and pushed back to all nodes via over-the-air updates. The network becomes smarter as it grows.
4. **Climate-adapted intelligence.** The system does not prescribe a fixed set of crops. A climate-aware recommendation engine suggests species whose optimal growing conditions most closely match the local ambient environment, minimising the energy required to close the gap between what the environment provides and what each crop needs.

## 2 The problems of industrial agriculture

Industrial agriculture has been organised for roughly a century around a single objective: to maximise the yield of a narrow set of commodity crops per unit of land and labour. By that measure it has succeeded, sustaining a global population larger than at any point in history. The methods that produced this abundance, however, carry consequences that yield figures do not record. The clearing of land for cultivation is among the foremost drivers of global biodiversity loss. The fertilisers and pesticides that sustain high yields contaminate water, soil, and air well beyond the fields to which they are applied. The concentration of production into a few crops, a few firms, and a few exporting countries has built a food supply that is efficient under stable conditions and brittle under shocks. The sector is also among the largest contributors to the climate change that increasingly threatens its own productive base.

These are not separable side effects to be corrected one at a time. They share a common origin in the logic of industrial production: the pursuit of yield through scale, intensification, and uniformity. That logic takes many forms. It clears habitat for ever-larger cultivated areas, substitutes synthetic inputs for functions that healthy soils and diverse landscapes once performed, and favours a few genetically narrow, high-yielding crops over the diversity that once buffered the system against shocks. The same drive toward scale concentrates economic power upstream, binding farmers to proprietary input chains and leaving the system resistant to reform from within. The harms documented below are not independent of one another; they are recurring expressions of how industrial agriculture is organised.

The following sections document these problems in turn, drawing on recent peer-reviewed literature and institutional reports, beginning with the land and ecosystems consumed by agricultural expansion and ending with the system’s effect on the diversity and nutritional quality of human diets.

### 2.1 Land Use

The prevailing paradigm of industrial agricultural production requires the use of vast swathes of land, along with a considerable amount of water, energy and other resources, which inevitably ends up homogenizing the local flora and destroying the habitat of the local fauna, leading to an overall loss of biodiversity. The agricultural expansion that the current system entails is indeed one of the main drivers of global biodiversity loss [1, p. 412], [2, p. 131], and is the most widespread form of land-use change, with cropping and animal husbandry covering over one third of the terrestrial land surface [1, p. XVI]. More than 90% of global biodiversity impacts from land-use change are attributable to agriculture, with crop cultivation accounting for 72% and pastures for 21% — dwarfing the combined impact of mining, urban development, and all other industries [3].

This expansion of cropland has been a constant for decades, with global cropland area growing by 78 million ha (780,000 km<sup>2</sup>) between 2001 and 2023, largely driven by an increase in permanent crops such as cocoa, oil palm and coffee [4, p. 3]. All of these are used intensively for monoculture, as are feedstock crops—as of the early 2000s, only 55% of global crop calories were allocated to direct human consumption, with 36% diverted to animal feed and the remainder to biofuels and industrial uses [5, Table 2]; this particular form of agriculture, which consists in the use of a single crop for large productions, is another potent driver

of biodiversity and habitat loss which will be explored in a subsequent section.

The extension of anthropogenic drivers of change has led to as little as 13% of the ocean and 23% of the land falling under the classification of ‘wilderness’. These few remaining areas, classified as sufficiently free of obvious human impacts to qualify as wilderness, tend to be remote or unproductive, like tundras or oceanic gyres, the most hospitable biomes having been almost or completely modified by humans in most regions [1, p. 206]. With the transformation for human convenience of the landscapes of the world that are the most accommodating to life, displaced wild populations are left without a habitat.

The impact of this intrusion into the wilderness has far-reaching consequences as well. According to the FAO’s State of Food and Agriculture report [6, p. 4], agriculture drives nearly 90 percent of global deforestation, with cropland expansion accounting for almost half of the total deforested area. This leads to changes in both global and local climates through two main mechanisms: the release of stored carbon produces a robust warming signal worldwide, while the removal of forest cover changes surface albedo, evapotranspiration and canopy roughness, which produces a net global warming effect for deforestation in the tropics and mid-latitudes, but a net cooling effect beyond roughly 50°N where increased albedo dominates. At the local scale, deforestation in the tropics raises surface temperatures by 0.2–2.4°C, with smaller increases in temperate zones (0.02–1.0°C), though boreal deforestation can produce modest local cooling [7, pp. 1, 6].

A pan-tropical synthesis estimated total tropical deforestation at 6.5–9.5 million hectares per year between 2011 and 2015, of which agriculture drove the overwhelming majority (90–99%), accounting for 6.4–8.8 million hectares per year [8, pp. 1, 3]. Crucially, only 45–65% of all deforested land was subsequently converted into actively managed cropland, pasture, or tree crops, meaning that a substantial fraction of tropical forest destruction does not result in lasting agricultural production.

## 2.2 Biodiversity loss

Beyond direct habitat destruction, the intensification of agriculture has driven rapid declines among species of agricultural habitats, contributing to sharply rising extinction rates across all taxonomic groups for which robust assessment data exist [1, Section 2.2.5.1]. Such losses have measurable functional consequences: biodiversity loss reduces the efficiency by which ecological communities produce biomass and recycle nutrients, with effects on primary productivity comparable in magnitude to climate warming and drought [9, pp. 60–61]. The spread and intensification of agriculture over the past half century — through greatly expanded scales, monoculturing, increasing pesticide and fertiliser application, and the elimination of hedgerows and other wildlife habitat — is directly related to major documented declines in insect biomass and diversity, with climate change—to which agriculture is itself a leading contributor (Section 2.8)—acting as an additional stressor [10, pp. 1–3]. At the global scale, agriculture has been identified as a threat to 24,000 of the 28,000 (86%) species so far documented as at risk of extinction [11, p. 7].<sup>1</sup>

The impact reaches beyond field boundaries. A 27-year study using standardised traps across 63 German nature protection areas found a seasonal decline of 76% and a mid-summer peak decline of 82% in total flying insect biomass, with losses apparent regardless of habitat type, suggesting large-scale drivers consistent

---

<sup>1</sup>The Chatham House figure draws on IUCN Red List data as compiled by Ritchie and Roser (2019).

with, though not yet definitively attributable to, agricultural intensification [12, pp. 14–15]. This decline is multi-causal and compounding: habitat loss, climate change, pesticide exposure, and other stressors co-occur and interact, with reported rates of annual abundance decline frequently around 1–2% across taxa, though with much variation across time, space, and taxonomic lineage [13, p. 2]. A global analysis of 923 terrestrial insect assemblages monitored across 106 long-term studies found overall abundance declining at approximately 1.5% per year, with losses driven disproportionately by formerly abundant species, though data remain heavily concentrated in Europe and North America [14, Fig. 2]. Spatially, the interaction of intensive agriculture and climate warming is associated with reductions of almost 50% in abundance and 27% in species richness relative to undisturbed habitats [15, Fig. 2]. In the tropics, where long-term monitoring data are scarcer, a 22-year study in protected Costa Rican lowland forest documented declines in 41% of common caterpillar genera, with losses linked to climate variability and, potentially, pesticide drift from adjacent agricultural land [16, pp. 3, 6].

The decline extends well beyond insects. A 37-year monitoring programme spanning 170 common bird species across more than 20,000 sites in 28 European countries found an overall population decline of 25%, with farmland birds suffering the steepest losses at nearly 57%—far exceeding declines in woodland, urban, or other habitat-associated species. Agricultural intensification, measured as the cover of farms with high pesticide and fertiliser inputs, was identified as the dominant pressure in a quasicausal analysis [17, Fig. 2]. Pollinators have been similarly affected: managed honey bee colonies declined by 59% in North America between 1947 and 2005 and by 25% in central Europe between 1985 and 2005, though global managed stocks rose by approximately 45% between 1961 and 2008 due to expansion in countries such as China and Argentina; among wild bees, four bumblebee species have gone extinct across Europe [18, p. 1255957-1]. The mechanistic drivers of pollinator loss—including neonicotinoid exposure, habitat destruction, and pathogen interactions—are examined in Section 2.5.

## 2.3 Soil degradation and erosion

Soils underpin the production of ecosystem goods and services that sustain both natural habitats and human societies. They are a key reservoir of biodiversity, from micro-organisms to flora and fauna; and this biotic community drives the soil functions (nutrient cycling, water regulation, carbon sequestration, etc.) that generate those services. Those functions are controlled primarily by the soil’s chemical composition, biological activity, and physical structure. When any of these attributes deteriorate, the soil’s ability to deliver essential ecosystem services, such as food production, water purification, and climate regulation, is compromised or lost [19, pp. 4–5].

Industrial tillage-based agriculture destroys soil at a rate that outpaces natural soil formation by orders of magnitude. Erosion rates from conventionally ploughed agricultural fields average one to two orders of magnitude greater than rates of soil production, erosion under native vegetation, and long-term geological erosion — making conventional plow-based agriculture fundamentally unsustainable over civilizational timescales [20]. Industrial farming practices such as mono-cropping, minimal use of organic fertilisers<sup>2</sup>, and

---

<sup>2</sup>Organic fertilisers — manure, compost, and crop residues returned to the field — add carbon-rich matter that sustains soil biological activity and structure. They are distinct from the synthetic nitrogen and phosphorus fertilisers whose environmental

removal of crop residues are a primary cause of the decline in soil organic matter, which is essential to long-term soil productivity [21, Table 7].

The IPCC’s Special Report on Climate Change and Land estimates that cropland soils have lost 20–60% of their organic carbon content relative to pre-cultivation states (medium confidence), and that soils under conventional agriculture continue to be a source of greenhouse gas emissions (medium confidence) [22, TS.4].

A second-order meta-analysis synthesizing 230 first-order meta-analyses encompassing more than 25,000 primary studies confirms that land conversion for crop production is a consistently net-negative driver for soil organic carbon (SOC), with mean SOC losses of approximately 25% following conversion from forest or wetland, and 16% from grassland (though the latter carries a wider confidence interval that includes zero), though forest-conversion losses in particular may be underestimated because most underlying studies measure only over years to decades, whereas full equilibration after land-use change may take considerably longer (estimated at around 80 years for grassland-to-cropland conversion). This matters because SOC is foundational to soil health, food production, and climate regulation: global SOC stocks to 2 m depth are estimated at approximately 2,400 Gt C — three times the carbon currently held in the atmosphere — meaning even small proportional losses translate into significant atmospheric CO<sub>2</sub> increases. SOC restoration is possible, but it is understood to be generally slower than depletion [23].

The on-site consequences of this depletion are direct: SOC loss diminishes nutrient availability, water-holding capacity, and soil structural stability, reducing the productive capacity of cropland and driving increased dependence on synthetic inputs to maintain yields [21, pp. 4–5]. The agronomic mechanisms by which monoculture accelerates this degradation, and the yield recovery achievable through diversification, are examined in Section 2.6.3.

## 2.4 Fertilizer use

Excess nitrogen and phosphorus from agricultural fertilizers leach into waterways, triggering eutrophication — a process in which accelerated algal growth reduces water clarity and, when these blooms die, microbial decomposition severely depletes dissolved oxygen, creating hypoxic or anoxic dead zones in which most aerobic organisms cannot survive [24, Consequences]. Cultural eutrophication of this kind has been identified as a leading cause of freshwater ecosystem impairment [25, p. 1] and has driven the exponential spread of coastal dead zones, from over 400 documented systems in 2008 [26, p. 926] to more than 500 by 2018 [27, p. 1].

Between 50 and 70% of applied nitrogen is typically lost to the environment through volatilisation, leaching, surface runoff, and denitrification [28, p. 1]. These pathways drive the eutrophication described above — through nutrient-laden runoff and leachate reaching waterways — and the atmospheric accumulation of nitrous oxide examined below, as excess soil nitrogen undergoes microbial denitrification.

Excess nitrogen in agricultural soils undergoes microbial conversion to nitrous oxide (N<sub>2</sub>O) — a greenhouse gas with a global warming potential 273 times that of CO<sub>2</sub> over 100 years [29, Table 7.15]. Global anthropogenic N<sub>2</sub>O emissions increased by 40% between 1980 and 2020, driven primarily by agricultural nitrogen use [30, Section 4.2.1], and atmospheric concentrations have now exceeded projected levels under

---

impacts are discussed in Section 2.4.



all CMIP5 and CMIP6 scenarios [30, Section 4.1.1].

Critically, the relationship between nitrogen application and  $\text{N}_2\text{O}$  emissions is not linear but accelerating: as fertilizer inputs increase beyond crop demand, the emission factor itself grows, meaning that each additional unit of nitrogen applied in excess generates a disproportionately larger  $\text{N}_2\text{O}$  penalty. As a consequence, small reductions in over-application can yield emissions reductions substantially larger than linear models — such as the IPCC’s default 1% emission factor — would predict [31, pp. 9199–9202]. Notably, this nonlinear amplification is most pronounced for conventional synthetic fertilizers; controlled-release formulations show a significantly attenuated response [31, p. 9201] — consistent with slower nitrogen delivery reducing the excess that drives nonlinear emissions.<sup>3</sup> In soilless systems such as closed-loop hydroponics, the microbial nitrification–denitrification pathway that generates these emissions is largely absent, removing the mechanism entirely rather than merely attenuating it.

## 2.5 Pesticide use

Pesticides contaminate soil, water, and air well beyond the target application zone, harming non-target organisms throughout the food web. An estimated 64% of the world’s agricultural land — covering approximately 24.5 million  $\text{km}^2$  — is at risk of pollution by more than one pesticide active ingredient, with 31% of that area at high risk; five watersheds identified as presenting high combined concern for pesticide pollution risk, water scarcity, and biodiversity include the Orange (South Africa), Huang He (China), Indus (India), Murray (Australia), and Paraná (Argentina) [32, pp. 206–208].

A 2023 meta-analysis synthesizing 54 studies found that pesticides consistently reduced both the abundance and diversity of soil fauna communities. The most damaging scenarios involved combinations of multiple substances, broad-spectrum compounds, and insecticides, which significantly decreased soil invertebrate diversity even at label-recommended application rates. Critically, under the repeated-application regimes typical of conventional agriculture, negative effects on soil fauna did not attenuate: long-term field studies exhibited effect sizes comparable to short-term experiments [33, Sections 3.2–3.3, 4.2].

One of the most publicly visible consequences of the use of pesticides in industrial agriculture is the elevated loss rate of pollinator colonies. Neonicotinoid insecticides — the class most strongly implicated in bee population declines — are systemic compounds widely applied as prophylactic seed treatments, chronically exposing foraging pollinators through contaminated pollen and nectar [18, pp. 1255957-4–1255957-5]. Multiple interacting stressors — encompassing parasites, pathogens, nutritional deficiency, and pesticide exposure — represent the most probable drivers of elevated colony loss rates [18, p. 1255957-5]. Furthermore, the share of global agricultural area devoted to pollinator-dependent crops rose from 19.4% to 32.8% between 1961 and 2016, with expansion rates highest for crops in the moderate- and high-dependence categories [34, p. 3520], raising concern that continued pollinator decline could jeopardize the stability of agricultural food production.

The harm does not stop at the field boundary. A systematic review of epidemiological and toxicological

---

<sup>3</sup>The evidence base for controlled-release attenuation of nonlinear  $\text{N}_2\text{O}$  response remains small ( $n = 8$  site-years in the meta-analysis). Instrumented soil-based units within the network proposed in this paper could contribute longitudinal field data to strengthen or qualify this finding.

evidence links chronic pesticide exposure to a wide spectrum of non-communicable diseases, including cancer, neurological disorders, and endocrine disruption [35, Sections 5–6]. A 2023 review of 63 epidemiological studies found that the strongest human evidence for associations between pesticide exposure and cancer concerned acute myeloid leukemia and colorectal cancer, where high-quality prospective studies observed dose-response relationships, though evidence for other cancer types remained inconsistent; the authors nonetheless concluded that existing evidence is sufficient to justify regulatory action [36, pp. 904–907]. Organochlorine and organophosphate compounds present in pesticides can mimic or antagonize endogenous hormones, with documented associations including reproductive disorders and developmental effects in children exposed *in utero* [35, Sections 5.1–5.2, 6.3]; these compounds have also been linked to breast and prostate cancers, though the epidemiological evidence remains mixed [36, pp. 900–901, 904].

The severity of these harms is not an unavoidable cost of pest management: it is in significant part a structural consequence of industrial open-field agriculture, where large-scale monoculture systems drive heightened pest pressure and perpetuate reliance on synthetic pesticides [37, p. 1].

## 2.6 Monoculture

Monoculture — the cultivation of a single crop species, or a narrow set of genetically near-identical cultivars, over large and continuous areas — is not merely one practice among many in industrial agriculture; it is its organizing principle. The economic logic of scale, mechanization, and commodity-market integration pushes production towards ever-greater uniformity, with cascading consequences for soil health, pest and disease dynamics, ecosystem resilience, and food-system stability. The following subsections document each of these dimensions in turn.

### 2.6.1 The monoculture effect: genetic vulnerability to disease and pest outbreaks

A fundamental result of epidemiological theory is that disease establishment requires host population density above a critical threshold [38, Section 1, pp. 535–536], and that genetically uniform host populations are disproportionately vulnerable to disease [39, pp. 381–382], [40, p. 718]. Monoculture maximizes both risk factors simultaneously: by planting a single crop species across large continuous areas, it concentrates the effective host population for any specialist pathogen or pest, while genetic uniformity ensures that a parasite capable of exploiting one individual can readily transmit to every neighbour, encountering no variation in resistance mechanisms.

This phenomenon — termed the “monoculture effect” — has been empirically quantified in a meta-analysis synthesizing 102 studies comprising over 2,000 effect sizes. In experimental crop systems, cultivar mixtures experienced approximately 50% less parasitism than monocultures on average — though the authors note this reflects a best-case scenario in which mixtures were assembled from cultivars with known resistance phenotypes [41, pp. 21, 23, 26].

The practical power of this effect has been demonstrated at operational scale. In a landmark field experiment in Yunnan Province, China, genetically diversified rice plantings across first five, then ten townships showed that disease-susceptible varieties planted in mixtures with resistant varieties suffered 94% less rice blast (the major disease of rice) severity than when grown in monoculture, and yielded 89% more per hill.

The results were so dramatic that fungicidal sprays were no longer applied by the second year of the programme, though the authors note this may not be achievable in all seasons [40, pp. 718–721, Table 1]. This experiment remains one of the most compelling demonstrations that intraspecific crop diversification can function as a practical, large-scale disease management strategy.

The same vulnerability extends to insect pests. A review of 287 herbivore species found that approximately half (52%) reached higher population densities in monocultures, with only 15% showing the opposite pattern [42, Table 1, pp. 566–567], consistent with Root’s resource concentration hypothesis — that herbivores arriving in concentrated host stands are more likely to remain and reproduce, driving specialist accumulation [43, p. 116]. A more recent global synthesis of 609 studies confirms that increasing plant species diversity consistently suppresses pest performance and promotes crop production across agro-ecosystems, grasslands, and forests<sup>4</sup> [44]. Natural enemy populations also tend to reach higher densities in diversified systems than in monocultures [42, Table 1, p. 567], reducing the biological control that would otherwise help regulate pest populations [45, p. 975]. The combination of elevated pest pressure and diminished biological control drives dependence on pesticides, whose application further disturbs the natural enemy communities that would limit outbreaks [45, pp. 975, 982], creating a self-reinforcing cycle.

## 2.6.2 Historical and contemporary case studies

The vulnerability inherent in genetic uniformity is not theoretical. It has repeatedly materialized in catastrophic losses throughout the history of modern agriculture.

The 1970 Southern Corn Leaf Blight—caused by a single cytoplasmic male-sterility trait (T-cytoplasm) shared by 75–90% of US hybrid maize cultivars [46, pp. 108–109]—destroyed an estimated 15% of the American crop in a single season and caused approximately \$1 billion in losses [47, p. 1113] ( $\geq$ \$6.0 billion in 2015 dollars [48, p. 1218]), a direct consequence of genetic homogeneity [47, p. 1113]. The disease had previously caused less than 1% average annual loss [47, p. 1115]; genetic uniformity converted a background pathogen into a national crisis in a single growing season. The lesson was not absorbed: by 2004, much of US corn germplasm still originated from just seven progenitor lines, and the tendency of breeders to recycle elite inbreds may, over the long term, erode the crop’s genetic base [49, pp. 1193, 1203].

Global banana production depends overwhelmingly on a single cultivar — the Cavendish subgroup — which represents approximately 40% of global production and makes up almost all of the export trade, with an additional 20% of production consisting of closely related varieties sharing a very similar genetic background. The Cavendish itself rose to dominance only because its predecessor, the Gros Michel, was decimated by *Fusarium* wilt Race 1 from the 1920s onward — a disaster enabled by the genetic uniformity of the Gros Michel monoculture [50, pp. 2147, 2149–2150]. History is now repeating: a new strain, *Fusarium oxysporum* f.sp. *cubense* Tropical Race 4 (TR4), is spreading through Cavendish plantations worldwide: since symptoms were first observed in Cavendish in Taiwan in 1967 [50, p. 2150], TR4 has expanded to 23 countries as of 2024, including Colombia, Peru, and Venezuela [51, p. 1]. The pathogen can survive in soil for up to 30 years in the form of chlamydospores [52, p. 653], spreads through contaminated soil, water, and planting material, and there are currently no market-acceptable resistant cultivar replacements [53, p. 761]. More

---

<sup>4</sup>Cited from abstract only; full-text verification pending.

than 400 million people depend on bananas as a staple food or source of income, and the crop’s combined economic value exceeds US\$52 billion per year [50, pp. 2146, 2151]. The global banana industry thus faces an existential threat that is a direct structural consequence of monoculture production: an entire sector built on a single clone, with no viable fallback.

These are not isolated events. Coffee leaf rust — caused by *Hemileia vastatrix* — devastated Sri Lanka’s coffee industry in the nineteenth century and remains a major threat to production worldwide [54, p. 1039]. The vulnerability is again structural: coffee is the main source of income for more than 100 million people [54, p. 1039], and *Coffea arabica*, which accounts for approximately 60% of world production, carries an exceptionally narrow genetic base — severe bottlenecks during domestication mean that a single coffee plant from the Botanical Garden of Amsterdam is believed to be one of the progenitors of most current cultivars [54, p. 1039]. A recent severe epidemic throughout Central America, Colombia, Peru, and Ecuador caused yield losses of up to 35%, directly affecting the livelihoods of hundreds of thousands of farmers [54, p. 1040]. Wheat rust presents the same recurring pattern. Stem rust historically destroyed 19–28% of the US wheat crop in epidemic years from the 1910s to the 1950s [55, pp. 304–305]. A new threat emerged in 1998 with the Ug99 race of the stem rust fungus, which has since spread to 13 countries across Africa and the Middle East in 13 variants; the majority of wheat cultivars grown worldwide lack adequate resistance [55, pp. 305, 309]. Each of these cases — maize, banana, coffee, wheat — confirms the same structural lesson: production systems built on a narrow genetic base remain vulnerable to a single pathogen.

### 2.6.3 Soil degradation

Continuous monoculture degrades the very substrate upon which it depends. Repeated cultivation of the same crop typically depletes specific nutrients, disrupts soil microbial communities, and diminishes soil organic matter (SOM) and active carbon (the readily-decomposable fraction of SOM) [56, Abstract], though conservation practices such as residue retention and reduced tillage can attenuate these effects [56, p. 15]. The productivity consequences of SOM loss are documented in Section 2.3.

The impact on soil biota is also substantial. Monoculture systems exhibit reduced soil bacterial diversity and altered microbial community composition compared to crop-rotation systems [57, Abstract, Fig. 2]. Under continuous monoculture, root exudate chemistry narrows [58, Fig. 3C], and disease-suppressive rhizosphere bacteria such as *Paenibacillus* are depleted while soil-borne pathogens such as *Fusarium oxysporum* accumulate, as documented in continuously-cropped peanut systems [59, Figs. 2f, 4e]. Conversely, crop rotation increases microbial biomass carbon and nitrogen by 13–16% [60, Section 3.2]; an earlier meta-analysis reports MBC +20.7–20.9% and MBN +26.1% [61, Fig. 2]. Rotation also raises bacterial diversity relative to continuous monoculture [60, Section 3.3][57, Abstract, Fig. 2]. These biological improvements translate into measurable structural gains: a meta-analysis of 2,199 paired observations found that rotation improved macroaggregate stability<sup>5</sup> by 7–9% relative to continuous monoculture, with climate and initial soil organic carbon jointly identified as the predominant drivers [63, Table 1, Section 3.5]. This is consistent with the foundational mechanism that macroaggregate stability depends on actively-growing roots and fungal hyphae, sustained under pasture or rotation and depleted by continuous cultivation [62, pp. 143, 158], and

---

<sup>5</sup>Macroaggregates are soil clumps >0.25 mm, stabilised primarily by roots and fungal hyphae [62, pp. 141, 152, 155].

with monoculture’s suppression of soil carbon and microbial communities documented above.

The structural losses above compound the broader pattern of erosion under industrial agriculture documented in Section 2.3.

#### **2.6.4 Yield penalties and the diversification dividend**

Far from maximizing output, monoculture imposes a persistent yield penalty relative to diversified systems. A global meta-analysis synthesizing 3,663 paired field-trial yield observations (1980–2024) found that crop rotation increased subsequent crop yield by 16–23% on average (depending on whether the pre-crop was a non-legume or legume), and that considering the entire rotation sequence, total yields, dietary energy, protein, iron, magnesium, zinc, and revenue all increased by 14–27% relative to continuous monoculture. Notably, win–win relationships among yield, nutrition, and revenue were consistently more frequent (33–54%) than trade-offs [64, Abstract, Fig. 1c, Fig. 3]. The dividend magnitude varies substantially by pre-crop type: gains rise to 60% (CI 44–79%) for cereals following crop mixtures from different functional groups, and to 83% (CI 48–128%) for maize following fodder legumes [64, Fig. 2].

The benefits extend beyond yield quantity to yield stability, a dimension of critical importance for food security. An analysis of five decades of FAO data across 91 nations found that greater effective crop diversity at the national level was associated with significantly increased temporal stability of total national harvest, comparable in magnitude to the destabilizing effect of precipitation variability. This boost in stability is particularly consequential, as nations with low yield stability suffered a sharp harvest loss (>25% decline in total yield) approximately every 8 years; high-stability nations suffered such losses only once every century or more [65, Abstract, Fig. 2].

Long-term field experiments across North America further confirm that more diverse rotations raised maize yields in 9 of 11 sites by 28.1% on average across all growing conditions, with drought-year yield gains of 14.0–88.9% relative to simplified rotations [66, Figs. 1A, 2B]. This evidence is consistent with the broader ecological principle that diversity increases the stability and productivity of biological communities [9, pp. 60–61].

#### **2.6.5 Diversification as a systemic alternative**

The empirical case against monoculture is not limited to individual metrics. A second-order meta-analysis of 69 prior meta-analyses (5,160 original studies, 41,946 comparisons between diversified and simplified practices) found that agricultural diversification enhances biodiversity, pollination, pest control, nutrient cycling, soil fertility, and water regulation without compromising crop yields [67, Abstract, Fig. 2]. In the subset of meta-analyses that measured both yield and other ecosystem services together, 63% of paired comparisons yielded win–win outcomes for both [67, Fig. 3]. These results suggest that the simplification inherent in monoculture is not merely an environmental cost borne for the sake of production; it is a production strategy that more diverse alternatives consistently outperform across multiple ecosystem-service dimensions, without compromising yield.

At the system level, an assessment of two agroindustrial cropping systems (wheat in Kenya and soybean in Bolivia) corroborates the meta-analytic case: the Bolivian soybean system was 89.2% rainfed monoculture

with 5.6% of arable land under vegetation, while the Kenyan wheat system spanned 12 land-cover classes with 41% of land under natural vegetation. These functional-diversity differences drove correspondingly low capacity to withstand climate and economic shocks (extremely low for soybean, low for wheat) [68, Abstract, p. 7].

At the experimental scale, an 8-year on-farm trial across eight farms in south-western France demonstrated substantial environmental gains from crop-diversification relative to the farms' initial conventional systems: mean reductions of 43% in irrigation, 36% in greenhouse-gas emissions, and 30% in on-farm energy use [69, Abstract, Section 3.2].

### **2.6.6 Structural drivers: market concentration and monoculture**

The persistence of monoculture despite its well-documented costs is not primarily a matter of agronomic ignorance; it is the structural outcome of concentrated market power and the economic incentives it creates. Four firms (BASF, Bayer, Corteva, and Syngenta) control 56% of the global commercial seed market and 61% of the global pesticides market: both sectors exceed the four-firm 40%-of-market threshold ETC Group invokes for oligopoly [70, pp. 2, 4]. Even where many farms exist, their effective genetic choices are constrained by a handful of input suppliers, making homogenisation not merely an agronomic preference but the structural outcome of concentrated market power. Consolidation drives a shift in research and development toward a small number of high-yielding proprietary varieties suited to monoculture systems and tied to proprietary agrochemical inputs. Farmers adopt these packages not merely out of preference but under structural compulsion: the agricultural treadmill<sup>6</sup> forces continuous yield maximisation simply to maintain income [71, Chapter 5, pp. 95–96], while institutional lock-in (whereby research funding, career incentives, and scientific culture are organised around the dominant input-intensive paradigm, making it structurally difficult to develop, validate, or disseminate alternatives outside it) renders those alternatives inaccessible even where they demonstrably work [72, Section 5, pp. 980–981]. The resulting genetic uniformity then creates the very pest and disease pressures established in Section 2.6.1, driving a self-reinforcing cycle of dependence on agrochemical inputs.

Concentrated market power enforces itself at the point of seed reproduction through two complementary strategies that prevent farmers from reproducing their own seed. The first is biological: hybrid varieties, whose offspring do not breed true, structurally eliminate the agronomic rationale for seed saving. The second is legal: the commercialization of transgenic seeds in the 1990s introduced full patent protections that prohibit seed saving outright, with violations subject to civil and criminal penalties [73, Section 2.1, p. 1268]. The practical effect has been a two-phase collapse of farm-level seed autonomy in the US: soybean seed saving declined gradually from a 1960 peak of 63% to 33% by 1991, then fell to 10% by 2001 following the 1996 introduction of Monsanto's Roundup Ready soybeans with no-replant licensing clauses [74, pp. 128–

---

<sup>6</sup>The agricultural treadmill, a concept introduced by economist Willard Cochrane in 1958, describes the mechanism by which farmers are compelled to continuously adopt productivity-increasing technologies: early adopters gain a temporary price advantage, but as adoption spreads, oversupply drives prices down, forcing all farmers to adopt the technology simply to maintain the same revenue. Those who cannot keep up exit farming; those who remain must adopt the next round of inputs. The result is an irreversible dependency on purchased inputs and a structural incentive to maximise yield per acre above all other considerations.

129, Figure 1]. These strategies converge with bundling: patented seed traits are paired with proprietary agrochemical inputs in packages designed to foreclose substitution at each step. A 2008 census of US patents covering herbicide-resistant transgenic plants found that the top three firms (Monsanto, DuPont-Pioneer, and Syngenta) held 85.0% of patents on maize and 69.6% of patents on non-maize plants [75, Table 1, pp. 121–122]. This concentration subsequently deepened: between 2015 and 2018, Bayer acquired Monsanto, Dow and DuPont merged to form Corteva Agrisciences, and ChemChina acquired Syngenta, reducing the sector’s major players from six to four [76, Abstract, p. 129].

The concentration dynamics described above have been intensified in recent decades by the financialization of agrifood systems (the growing subordination of corporate strategy to the short-term return imperatives of financial markets), which has encouraged food and agricultural corporations to pursue mergers and acquisitions to generate shareholder value and capture market share. An analysis of 4,449 food system M&A deals between 2001 and 2020 found that the overwhelming majority of activity was horizontal (firms acquiring competitors within the same sub-sector) rather than vertical, producing high levels of internal concentration within sub-sectors [77, Abstract, Section 4]. This process has been compounded by a parallel shift in ownership structure: large asset management firms have emerged as simultaneous major shareholders in dominant agrifood corporations across the sector, a pattern known as common ownership [78, p. 613]. At their height in 2010–2014, equity-related investment funds accounted, by one estimate, for about a third of global financial investment in the agrifood sector [78, p. 605, citing Valoral Advisors]. Common-ownership theory holds that firms with shared major shareholders have little incentive to compete aggressively against one another. Applied to the agrifood sector, this adds a further anti-competitive layer to already concentrated markets [78, pp. 614–615]. The result is a system of mutually reinforcing structural forces (IP enforcement, merger-driven consolidation, and common ownership) that operates without any internal mechanism for self-correction.

The structural character of these overlapping forces has a direct implication for the design of alternatives: technical solutions that operate within the same concentrated input markets will reproduce the same dependencies. Effective alternatives must be architecturally independent of proprietary input chains — which is to say, they must be open by design.

## 2.7 Freshwater depletion

Agriculture is the largest consumer of global freshwater, though the precise magnitude of its share is more uncertain than commonly acknowledged. The most widely cited figure, that irrigation accounts for 70% of global freshwater withdrawals, has been repeated across more than 3,000 scientific and policy documents, but a 2025 analysis drawing on eleven global hydrological models and thirteen independent datasets found that this figure lacks firm empirical support: the available data are consistent with agriculture’s share of withdrawals lying anywhere between 45% and 90%, a range the authors describe as a lower bound on the true uncertainty [79, Abstract, Empirical Support section].

The full scale of agricultural water consumption is partially obscured because it includes the depletion of non-renewable aquifers. Modelled global groundwater depletion increased 22% in just one decade, from 240 km<sup>3</sup>/year in 2000 to 292 km<sup>3</sup>/year in 2010, with five crops (wheat at 22% of global GWD, rice at 17%,

sugar crops at 7%, cotton at 7%, and maize at 5%) dominating depletion in production, and with the crop commodities present in international trade data (the remainder, mostly fodder crops, being absent from such data) accounting for 83% of the global total [80, p. 701, Fig. 1]. Unlike surface water shortages, depletion of fossil aquifers is functionally irreversible on human timescales: some of the groundwater currently consumed to irrigate these crops fell as precipitation thousands of years ago and will not recharge within any planning horizon relevant to food security policy [81, pp. 6–7].

Irreversibility, however, is not the universal condition of every depleted aquifer. A global compilation of in situ measurements from about 170,000 monitoring wells in 1,693 aquifer systems finds that rapid declines (faster than 0.5 m/year) characterise 12% of these systems, and that in roughly 30% of the aquifers for which both periods could be compared, depletion accelerated between the late twentieth and early twenty-first centuries. Among aquifers that declined in both periods, accelerating cases outnumbered decelerating ones by roughly 5 to 2 after excluding cases of negligible change [82, Abstract, pp. 715–717]. The authors also note that more than 80% of the accelerating-decline aquifers experienced declining precipitation, identifying climate variability as a contributing driver of acceleration [82, pp. 717–718]. The same dataset does identify aquifers where declines have slowed, reversed, or never occurred, with reversals attributed to regulatory measures, inter-basin surface-water transfers, and managed aquifer recharge [82, pp. 718–719]. But the asymmetry is severe, and it is evident in more than just counts: water tables fall, on average, four times faster than they rise in the aquifers where they do rise [82, p. 719]. Depletion, not recovery, is the dominant signal.

Recovery is nonetheless possible where the right interventions are sustained. The clearest demonstration is the North China Plain, one of the world’s most depleted aquifer regions. An analysis of about 190,000 measurements from more than 2,000 wells documents an unprecedented reversal beginning in 2020, with groundwater in unconfined aquifers rising at roughly 0.7 m/year [83, Abstract, p. 3]. The preconditions were demanding. Recovery required the central route of the South-to-North Water Diversion to replace groundwater pumping with water from the humid south, the closure of tens of thousands of wells, a sustained reallocation of several cubic kilometres per year to environmental flows and managed aquifer recharge, and the amplifying effect of an unusually wet 2021 whose contribution the authors themselves describe as short-lived [83, Abstract, pp. 5–8]. It also required scaling back the region’s dominant winter wheat-summer maize double-cropping rotation, an intensive groundwater-fed irrigation pattern: the agricultural share of regional water use fell from 70% in 2005 to 50% by 2023 even as regional food production rose [83, pp. 5–6]. Even with all of this, GRACE-derived storage recovery has so far compensated for only about half of the roughly 49 km<sup>3</sup> of total water storage depletion accumulated in the region between 2005 and 2019 [83, p. 7]. Recovery is therefore possible where aquifers are shallow enough to recharge on policy-relevant timescales, where a humid donor basin and the state capacity to operate an inter-basin water transfer exist, and where pumping can be capped against the demand pressures that drove depletion. In cultivated drylands underlain by fossil water, those preconditions are largely absent.



## 2.8 Greenhouse gas emissions

The greenhouse gas contribution of the global agrifood system can be quantified through complementary accounting frameworks that differ primarily in system boundary.<sup>7</sup> The FAOSTAT boundary (covering farm-gate<sup>8</sup> production, land-use change, and pre- and post-production supply-chain activities directly attributable to the agrifood sector) places global emissions at 16.2 Gt CO<sub>2</sub>eq for 2022, equivalent to approximately 30% of total anthropogenic GHG emissions [84, p. 2, Highlights]. The EDGAR-FOOD database, which applies a broader lifecycle boundary that includes upstream fertiliser and pesticide manufacture and downstream household preparation and landfill, estimated 18 Gt CO<sub>2</sub>eq and 34% for 2015 [85, p. 198, Fig. 1]. The IPCC’s Special Report on Climate Change and Land, drawing on multiple estimation methods, places the full food-system contribution at 21–37% of total net anthropogenic emissions for 2007–2016 [22, SPM A.3]. Within the FAOSTAT framework, emissions divide across three domains: farm-gate activities (48%; 7.8 Gt), land-use change (19%; 3.1 Gt), and the supply chain (33%; 5.3 Gt) [84, p. 2, Highlights]. The composition has shifted over time: between 2000 and 2022 supply-chain emissions grew 52%, farm-gate emissions 15%, and land-use-change emissions fell 30%, lifting the supply-chain share of the agrifood total from 24% to 33%, with each of its underlying components (input manufacturing, food processing, packaging and retail with transport, and household consumption) growing more than 80% [84, p. 3–4], consistent with the longer-window trend of doubling pre- and post-production emissions documented in the underlying FAOSTAT methodology [86, Abstract, p. 1799].

The dominant source of food-system emissions is the production of animal-based food (including livestock feed), which accounts for 57% of total food-related GHG emissions [87, p. 724] while providing only 37% of global protein and 18% of calories [88, p. 990]: a disproportionality whose implications for dietary strategy are examined in Section 2.11.

The agrifood sector accounts for a structurally disproportionate share of the most potent non-CO<sub>2</sub> greenhouse gases. The IPCC estimates that agriculture, forestry, and other land use (AFOLU) is responsible for approximately 44% of anthropogenic methane, 81% of anthropogenic nitrous oxide, and 13% of anthropogenic CO<sub>2</sub> globally [22, SPM A.3]. The N<sub>2</sub>O trajectory (its 40% growth in anthropogenic emissions since 1980 and observed concentrations exceeding all CMIP6 projections) is examined in detail in the preceding discussion of fertilizer use [30, Abstract]. Methane is of particular strategic importance because its atmospheric perturbation lifetime is only  $11.8 \pm 1.8$  years [29, Table 7.15], compared with over a century for nitrous oxide and millennia for CO<sub>2</sub>. A sustained reduction in methane emissions therefore translates into measurable cooling within two decades [29, Sections 7.6.1.4–7.6.1.5, Fig. 7.22]. Since ruminants<sup>9</sup> and rice cultivation are the dominant agricultural sources of methane [22, Section 2.3.2.2], shifting food production away from ruminant-dependent systems represents one of the most actionable near-term levers for slowing global warming.

---

<sup>7</sup>A *system boundary* defines which activities along the food chain are counted as emission sources, e.g. whether only on-farm processes are included or also upstream manufacturing and downstream retail.

<sup>8</sup>*Farm-gate* emissions comprise all GHG releases from crop and livestock production activities within the agricultural holding, up to the point the product leaves the farm.

<sup>9</sup>Ruminants are herbivorous mammals (principally cattle, sheep, and goats) that digest plant matter through enteric fermentation in a specialised multi-chambered stomach, a process that releases methane as a metabolic by-product.

The structural implication of these trajectories is severe. Even if all fossil fuel combustion were halted immediately, food-system emissions continuing on a business-as-usual path would alone exhaust the remaining carbon budget for 1.5°C between 2051 and 2063, and would consume over 96% of the budget consistent with a 67% chance of limiting warming to 2°C by end of century<sup>10</sup> [89, p. 706, Fig. 1]. Meeting Paris Agreement targets therefore requires the simultaneous implementation of multiple food-system interventions: dietary shifts, caloric adjustment, yield improvements, waste reduction, and production efficiency gains. No single strategy is sufficient. The agrifood system is therefore not a peripheral contributor to the climate crisis but a structural driver of it, whose reform is a precondition of any plausible stabilisation pathway [89, pp. 706–707, Fig. 2].

## 2.9 Produce transportation, storage and disposal

During 2010–2016, global food loss and waste accounted for an estimated 8–10% of total anthropogenic greenhouse gas emissions. An estimated 25–30% of all food produced is lost or wasted<sup>11</sup>, with losses in developing countries driven by inadequate infrastructure (such as lack of refrigeration) while waste in developed countries is driven by consumer behaviour [22, Section 5.5.2.5].

Beyond the farm gate, food-system emissions from transport, retail, and household consumption have grown substantially. Crippa et al. estimate that food transport contributes 4.8% and retail 4.0% of total food-system emissions, with transport’s share of food-system emissions rising by 67% and retail’s share by 300% between 1990 and 2015 [85, pp. 200–201]. More recent data show that pre- and post-production emissions reached 5.3 Gt CO<sub>2</sub>eq in 2022, or 33% of all agrifood emissions, up from 24% in 2000. Each subcategory (manufacturing of inputs, transport, packaging and retail, and household consumption) grew by more than 80% over the same period [84, pp. 3–4]. Cold chain infrastructure alone (refrigeration across processing, transport, retail, and household stages) accounts for an estimated 1.32 Gt CO<sub>2</sub>eq<sup>12</sup>, or nearly 10% of total agrifood emissions, having more than doubled since 2000 [90, Section 3]. In 61 countries and territories, supply-chain emissions now account for 60% or more of national agrifood emissions [84, p. 7].

Food waste that reaches landfills is a disproportionate source of methane. In the United States, food waste constitutes an estimated 24% of municipal solid waste disposed of in landfills, yet owing to its rapid decay rate, an estimated 58% of fugitive methane emissions from municipal solid waste landfills are attributable to landfilled food waste [91, Table A1, Table 3]<sup>13</sup>. Food waste decomposes faster than other organic materials, and an estimated 61% of the methane it generates escapes to the atmosphere uncaptured [91, p. iii].

<sup>10</sup>Clark et al. report emissions in CO<sub>2</sub> warming-equivalents (CO<sub>2</sub>-we), a metric based on GWP\* that captures the different temporal dynamics of short-lived (CH<sub>4</sub>) and long-lived (CO<sub>2</sub>, N<sub>2</sub>O) climate forcers. Cumulative budgets expressed in the more common GWP<sub>100</sub>-based CO<sub>2</sub>eq would yield different numerical thresholds.

<sup>11</sup>The IPCC assigns this estimate *medium confidence* (SRCCL, Section 5.5.2.5).

<sup>12</sup>Flammini et al. report 30–50% uncertainty on this estimate due to limitations in energy-use and refrigerant-leakage data.

<sup>13</sup>These figures are specific to the United States. No comparable global analysis of food waste’s share of landfill methane emissions exists in the peer-reviewed literature as of early 2026.

## 2.10 The existential dependency on a few producers

The global food system is vulnerable to disruption at multiple levels of concentration. Three global food crises in the past fifty years (1973–1974, 2007–2012, and the 2019–2022 pandemic and conflict disruptions) have each exposed interlocking structural weaknesses: the system’s reliance on a small number of staple grains produced using highly industrialised methods, the geographic concentration of the countries that export them, and the dominance of a small number of firms over global commodity markets [92, pp. 1–2]. In 2021–2022, seven countries plus the European Union accounted for 84% of world wheat exports, and four countries accounted for 83% of world maize exports; by 2024–2025, wheat concentration had risen to 88% [93, pp. 16, 25]. The majority of nations do not produce enough staple grains to meet domestic demand and therefore depend on global markets dominated by these few exporters [92, p. 6].

The concentration risk extends upstream. The oligopolistic control of the commercial seed and pesticide markets documented in Section 2.6.6 is mirrored in agricultural machinery, where the top four firms control 43% of the global market [70, Table 4]. This degree of corporate concentration has been linked to reduced competition (though empirical evidence on pricing effects remains limited by restricted access to proprietary industry data) and to narrower technological choices, as dominant firms focus on high-cost proprietary input packages over less capital-intensive alternatives [94, pp. 404–406].

Fertilizer supply chains compound these vulnerabilities along similar lines. Roughly one-third of seaborne fertilizer trade (about 16 million tonnes per year, dominated by phosphate and urea) passes through the Strait of Hormuz [95, p. 7], and the Middle East accounts for approximately one-quarter of global urea exports and over 15% of global ammonia exports [96, p. 34]. As of mid-2026, the closure of the strait has pushed the World Bank’s fertilizer price index up 12% in a single quarter, with benchmark urea reaching \$725/tonne in March 2026 (a 55% monthly jump and its highest level since 2022) following halted exports from Iran and Qatar [96, p. 34]. The FAO has warned that these disruptions will translate into lower yields and tightening food supplies in the latter half of 2026 and into 2027, with import-dependent countries in Africa, Asia, and parts of the Middle East most exposed [97]. Whether the immediate outcome reaches the worst projections is uncertain, but the structural vulnerability the crisis exposes is real.

At the most local level, food production itself is consolidating into fewer and larger operations. Globally, 0.1% of farms (those exceeding 1 000 hectares) operate over half of all agricultural land, while the 85% of farms smaller than 2 hectares operate just 9% [98, Chapter 3]. Farms exceeding 50 hectares produce more than 55% of crop-derived dietary energy worldwide<sup>14</sup> [98, Chapter 3]. If historical patterns of structural transformation continue, the total number of farms worldwide is projected to decrease by half before the end of the century, although farm numbers in sub-Saharan Africa are expected to rise [98, Chapter 3]. Smallholders (farms under 2 hectares, which still produce 16% of global crop-derived dietary energy) are vital contributors to food supply, yet the prevailing trajectory of consolidation is reducing their number [98, Chapter 3]. The result is a food system in which the vast majority of people (including entire nations) depend for their sustenance on supply chains they neither control nor can easily replace.

---

<sup>14</sup>This estimate is based on crop production data from 77 countries covering approximately 54% of global cropland area (FAO SOFA 2025, Chapter 3).

## 2.11 Suppression of dietary diversity and nutritional quality

Industrialised agriculture prioritises monoculture production of a narrow range of commodity crops, the cheap extracted ingredients of which (oils, starches, sugars) form the basis of ultra-processed foods<sup>15</sup> that now dominate the food supplies of high-income countries and are rising rapidly in middle-income countries [100, pp. 21, 27] [99, Figure 1]. An umbrella review of 14 meta-analyses encompassing nearly 10 million participants found convincing evidence that greater ultra-processed food exposure is associated with higher risks of cardiovascular disease mortality and type 2 diabetes, and highly suggestive evidence of association with obesity<sup>16</sup> [101, Figures 2–3].

The nutritional cost of industrial agriculture extends beyond dietary composition to the crops themselves. Analyses of archived wheat grain from the Broadbalk long-term experiment (1845–2005) show that zinc, copper, and magnesium concentrations declined significantly after the mid-1960s (by 33–49% for zinc, 25–39% for copper, and 20–27% for magnesium), coinciding with the introduction of semi-dwarf high-yield cultivars. Iron concentrations were also significantly lower in short-straw cultivars than in their long-straw predecessors, though the within-period temporal trend was not statistically significant [102, Tables 2, 4]. Soil mineral levels remained stable or increased over the same period, indicating that the decline is a plant-level phenomenon rather than a consequence of soil depletion [102, p. 321]. A broader study of 43 garden crops using USDA composition data from 1950 to 1999 found statistically reliable declines in protein (–6%), calcium (–16%), iron (–15%), and riboflavin (–38%), among others, attributing the pattern to a “dilution effect” in which selection for higher yields reduces the concentration of minerals and other nutrients per unit of harvested mass [103, Table 3]. Similar trends have been documented in UK fruits and vegetables over an 80-year period, with iron declining by approximately 50% and copper by 49%<sup>17</sup> [105, Table 1]. These declines are consequential because iron, zinc, and vitamin A deficiencies constitute the most prevalent forms of micronutrient malnutrition globally, with the heaviest burden falling on populations that depend on staple cereal and legume crops for the bulk of their dietary intake [106, pp. 353–354]; the documented decline in mineral density of precisely these crops compounds an already severe public health problem.

---

<sup>15</sup>Ultra-processed foods are defined under the NOVA classification as branded, commercial formulations made from cheap ingredients extracted or derived from whole foods and combined with additives, most containing little to no whole food [99, Panel 1].

<sup>16</sup>All evidence on ultra-processed food and long-term health outcomes is necessarily observational, since randomising participants to decades of dietary exposure is not ethically feasible. Lane et al. report the highest available tier of evidence synthesis for this question; their “convincing” (Class I) and “highly suggestive” (Class II) grades reflect the strength of association within that framework, not experimental proof of causation. The underlying GRADE quality ratings were predominantly low or very low (only 4 of 45 pooled analyses reached moderate; none reached high). Under GRADE methodology, observational studies are rated “low” quality by default, so these ratings reflect inherent limits of the study design rather than weakness in the observed associations.

<sup>17</sup>Nutrient-decline estimates derived from historical food-composition tables are contested. A critical review by Marles [104, Sections 3–4, 6–7] argues that changes in analytical methods, sampling protocols, and cultivar selection across editions render direct comparisons unreliable. However, Marles concedes that the archived-sample evidence of Fan et al. avoids these methodological limitations and that the dilution effect from high-yield cultivar selection is a real phenomenon, though he argues that its magnitude is too small to have a significant impact on consumer nutrition and that increased yield outweighs the dilution cost. The estimates from Davis et al. and Mayer et al. should therefore be read as indicative of direction and approximate magnitude, not as precise measurements.

Beyond nutrient decline, the narrowing of crop diversity documented above has also homogenised food supplies at a global scale. An analysis of national food supply data from 152 countries over 50 years found that national food supplies converged significantly towards a common pattern dominated by a few staple cereals and oil crops (notably wheat, rice, maize, soybean, and palm oil), even as the number of available crop commodities increased. The convergence was in proportional composition rather than in crop availability, with the most compositionally distinct regions (such as East and Southeast Asia and sub-Saharan Africa) experiencing the most rapid shift [107, Figure 3]. Modelling by Springmann et al. shows that current dietary trends (projected increases in consumption of animal products, sugar, and oils driven by population and income growth) could, absent technological changes and dedicated mitigation measures, push multiple planetary boundaries<sup>18</sup> beyond safe limits by 2050, including those for climate change, cropland use, freshwater use, and nitrogen and phosphorus pollution [108, p. 522]. This reinforces the conclusion, noted in Section 2.8, that no single intervention is sufficient and that systemic reform of the food system is required.

## 2.12 A systemic crisis

The problems documented in this section are not a collection of separable inefficiencies that could be corrected one by one. They form a system whose harms are structural and mutually reinforcing. Monoculture concentrates pest and disease pressure, which drives pesticide use, which suppresses the natural enemies and pollinators that would otherwise hold pests in check. Demand for uniform, high-yielding varieties concentrates the seed and agrochemical markets, and that concentration entrenches the input-intensive model while foreclosing the alternatives that might replace it. Soil degradation deepens dependence on synthetic fertilisers, whose losses reappear as eutrophied waters and atmospheric nitrous oxide. Each harm sustains the conditions that produce the others, and the market structure above them rewards precisely the practices that cause them.

Scale is the second defining feature. These are planetary effects rather than local externalities, measured in gigatonnes of greenhouse gas, millions of hectares of cleared forest, and the approach or breach of several planetary boundaries. On its current trajectory the food system alone would exhaust the remaining carbon budget for 1.5°C even if all fossil fuel combustion ceased immediately. At the same time it has left billions of people dependent for their daily nutrition on a handful of crops, firms, and exporting nations, a dependency whose fragility is revealed each time a drought, a conflict, or a closed shipping lane interrupts supply.

This structural character has a direct implication for any proposed remedy. A technology that raises yields or trims inputs while operating inside the same concentrated, proprietary, scale-driven system will tend to reproduce the dependencies that define it, as every productivity gain channelled through the agricultural treadmill has done before. No single intervention is proportionate to a crisis of this scale, and the evidence above points to none; meaningful reform will require change across diets, land use, supply chains, and the structure of the markets themselves. What the analysis does yield is a set of requirements that any genuine contribution must satisfy: it should be architecturally independent of proprietary input chains, it should reduce rather than relocate the land, water, and chemical demands that drive the ecological harm, and it

---

<sup>18</sup>Biophysical thresholds beyond which Earth-system processes risk destabilisation.

should distribute the means of production rather than concentrate them further. The remainder of this paper develops one component that aims to meet these requirements, an open-source and decentralised approach to controlled-environment production directed not at replacing field agriculture wholesale but at the segment where its advantages are clearest, and it assesses how far such a contribution can realistically reach.

## References

- [1] Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, “Global assessment report of the intergovernmental science-policy platform on biodiversity and ecosystem services,” IPBES Secretariat, Bonn, Germany, Tech. Rep., 2019, p. 1144. DOI: 10.5281/zenodo.3831673.
- [2] European Commission: Directorate-General for Financial Stability, Financial Services and Capital Markets Union, IIASA & Trinomics, *Study for a methodological framework and assessment of potential financial risks associated with biodiversity loss and ecosystem degradation – Final report*. Publications Office of the European Union, 2024. DOI: 10.2874/30768. [Online]. Available: <https://data.europa.eu/doi/10.2874/30768>.
- [3] L. Cabernard *et al.*, “Biodiversity impacts of recent land-use change driven by increases in agri-food imports,” *Nature Sustainability*, 2024. DOI: 10.1038/s41893-024-01433-4.
- [4] Food and Agriculture Organization of the United Nations, “Land statistics 2001–2023 – global, regional and country trends,” FAOSTAT Analytical Briefs, Rome, Tech. Rep. No. 107, 2025, FAO. DOI: 10.4060/cd5765en. [Online]. Available: <https://openknowledge.fao.org/handle/20.500.14283/cd5765en>.
- [5] E. S. Cassidy, P. C. West, J. S. Gerber, and J. A. Foley, “Redefining agricultural yields: From tonnes to people nourished per hectare,” *Environmental Research Letters*, vol. 8, no. 3, p. 034015, 2013. DOI: 10.1088/1748-9326/8/3/034015.
- [6] Food and Agriculture Organization of the United Nations, “The state of food and agriculture 2025 – addressing land degradation across landholding scales,” FAO, Rome, Tech. Rep., 2025, FAO publication. DOI: 10.4060/cd7067en. [Online]. Available: <https://doi.org/10.4060/cd7067en>.
- [7] D. Lawrence, M. Coe, W. Walker, L. Verchot, and K. Vandecar, “The unseen effects of deforestation: Biophysical effects on climate,” *Front. For. Glob. Change*, vol. 5, Mar. 2022. DOI: 10.3389/ffgc.2022.756115. [Online]. Available: <https://www.frontiersin.org/journals/forests-and-global-change/articles/10.3389/ffgc.2022.756115/full>.
- [8] F. Pendrill, T. A. Gardner, P. Meyfroidt, U. M. Persson, *et al.*, “Disentangling the numbers behind agriculture-driven tropical deforestation,” *Science*, vol. 377, no. 6611, eabm9267, 2022. DOI: 10.1126/science.abm9267.
- [9] B. J. Cardinale, J. E. Duffy, A. Gonzalez, *et al.*, “Biodiversity loss and its impact on humanity,” *Nature*, vol. 486, no. 7401, pp. 59–67, 2012. DOI: 10.1038/nature11148.
- [10] P. H. Raven and D. L. Wagner, “Agricultural intensification and climate change are rapidly decreasing insect biodiversity,” *Proceedings of the National Academy of Sciences*, vol. 118, no. 2, 2021. DOI: 10.1073/pnas.2002548117.
- [11] T. G. Benton, C. Bieg, H. Harwatt, R. Pudasaini, and L. Wellesley, “Food system impacts on biodiversity loss: Three levers for food system transformation in support of nature,” Royal Institute of International Affairs, 2021. [Online]. Available: <https://www.chathamhouse.org/2021/02/food-system-impacts-biodiversity-loss>.

- [12] C. A. Hallmann, M. Sorg, E. Jongejans, *et al.*, “More than 75 percent decline over 27 years in total flying insect biomass in protected areas,” *PLOS ONE*, vol. 12, no. 10, e0185809, 2017. DOI: 10.1371/journal.pone.0185809.
- [13] D. L. Wagner, E. M. Grames, M. L. Forister, M. R. Berenbaum, and D. Stopak, “Insect decline in the Anthropocene: Death by a thousand cuts,” *Proceedings of the National Academy of Sciences*, vol. 118, no. 2, e2023989118, 2021. DOI: 10.1073/pnas.2023989118.
- [14] R. van Klink, D. E. Bowler, K. B. Gongalsky, M. Shen, S. R. Swengel, and J. M. Chase, “Disproportionate declines of formerly abundant species underlie insect loss,” *Nature*, vol. 628, pp. 359–364, 2024. DOI: 10.1038/s41586-023-06861-4.
- [15] C. L. Outhwaite, P. McCann, and T. Newbold, “Agriculture and climate change are reshaping insect biodiversity worldwide,” *Nature*, vol. 605, pp. 97–102, 2022. DOI: 10.1038/s41586-022-04644-x.
- [16] D. M. Salcido, M. L. Forister, H. Garcia Lopez, and L. A. Dyer, “Loss of dominant caterpillar genera in a protected tropical forest,” *Scientific Reports*, vol. 10, p. 422, 2020. DOI: 10.1038/s41598-019-57226-9.
- [17] S. Rigal, V. Dakos, H. Alonso, *et al.*, “Farmland practices are driving bird population decline across Europe,” *Proceedings of the National Academy of Sciences*, vol. 120, no. 21, e2216573120, 2023. DOI: 10.1073/pnas.2216573120.
- [18] D. Goulson, E. Nicholls, C. Botías, and E. L. Rotheray, “Bee declines driven by combined stress from parasites, pesticides, and lack of flowers,” *Science*, vol. 347, no. 6229, p. 1255957, 2015. DOI: 10.1126/science.1255957.
- [19] Food and Agriculture Organization of the United Nations, *Revised World Soil Charter*, en. Rome, Italy: FAO, 2015. [Online]. Available: <https://openknowledge.fao.org/server/api/core/bitstreams/65be618a-f5e8-4f67-9cbb-d77f1f5cd31e/content>.
- [20] D. R. Montgomery, “Soil erosion and agricultural sustainability,” *Proceedings of the National Academy of Sciences*, vol. 104, no. 33, pp. 13268–13272, 2007. DOI: 10.1073/pnas.0611508104.
- [21] R. Lal, “Soil carbon sequestration to mitigate climate change,” *Geoderma*, vol. 123, no. 1–2, pp. 1–22, 2004. DOI: 10.1016/j.geoderma.2004.01.032.
- [22] IPCC, “Climate change and land: An IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems,” Intergovernmental Panel on Climate Change, Cambridge, UK and New York, NY, USA, Tech. Rep., 2019. DOI: 10.1017/9781009157988. [Online]. Available: <https://www.ipcc.ch/srccl/>.
- [23] D. Beillouin *et al.*, “A global meta-analysis of soil organic carbon in the Anthropocene,” *Nature Communications*, vol. 14, p. 3700, 2023. DOI: 10.1038/s41467-023-39338-z.
- [24] M. F. Chislock, E. Doster, R. A. Zitomer, and A. E. Wilson, “Eutrophication: Causes, Consequences, and Controls in Aquatic Ecosystems,” *Nature Education Knowledge*, vol. 4, no. 4, p. 10, 2013. [Online]. Available: <https://www.nature.com/scitable/knowledge/library/eutrophication-causes-consequences-and-controls-in-aquatic-102364466>.



- [25] S. R. Carpenter, N. F. Caraco, D. L. Correll, R. W. Howarth, A. N. Sharpley, and V. H. Smith, “Nonpoint pollution of surface waters with phosphorus and nitrogen,” *Issues in Ecology*, no. 3, 1998. [Online]. Available: <https://esa.org/wp-content/uploads/2013/03/issue3.pdf>.
- [26] R. J. Diaz and R. Rosenberg, “Spreading dead zones and consequences for marine ecosystems,” *Science*, vol. 321, no. 5891, pp. 926–929, 2008. DOI: 10.1126/science.1156401.
- [27] D. Breitburg, L. A. Levin, A. Oschlies, *et al.*, “Declining oxygen in the global ocean and coastal waters,” *Science*, vol. 359, no. 6371, eaam7240, 2018. DOI: 10.1126/science.aam7240.
- [28] D. Coskun, D. T. Britto, W. Shi, and H. J. Kronzucker, “Nitrogen transformations in modern agriculture and the role of biological nitrification inhibition,” *Nature Plants*, vol. 3, p. 17074, 2017. DOI: 10.1038/nplants.2017.74.
- [29] P. Forster, T. Storelvmo, K. Armour, *et al.*, “The earth’s energy budget, climate feedbacks, and climate sensitivity,” in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, V. Masson-Delmotte, P. Zhai, A. Pirani, *et al.*, Eds., Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press, 2021, ch. 7, pp. 923–1054. DOI: 10.1017/9781009157896.009.
- [30] H. Tian, N. Pan, R. L. Thompson, *et al.*, “Global nitrous oxide budget (1980–2020),” *Earth System Science Data*, vol. 16, no. 6, pp. 2543–2604, 2024. DOI: 10.5194/essd-16-2543-2024.
- [31] I. Shcherbak, N. Millar, and G. P. Robertson, “Global metaanalysis of the nonlinear response of soil nitrous oxide (n<sub>2</sub>o) emissions to fertilizer nitrogen,” *Proceedings of the National Academy of Sciences*, vol. 111, no. 25, pp. 9199–9204, 2014. DOI: 10.1073/pnas.1322434111. eprint: <https://www.pnas.org/doi/pdf/10.1073/pnas.1322434111>. [Online]. Available: <https://www.pnas.org/doi/abs/10.1073/pnas.1322434111>.
- [32] F. H. M. Tang, M. Lenzen, A. McBratney, and F. Maggi, “Risk of pesticide pollution at the global scale,” *Nature Geoscience*, vol. 14, pp. 206–210, Apr. 2021. DOI: 10.1038/s41561-021-00712-5.
- [33] L. Beaumelle, L. Tison, N. Eisenhauer, *et al.*, “Pesticide effects on soil fauna communities—A meta-analysis,” *Journal of Applied Ecology*, vol. 60, no. 7, pp. 1239–1253, 2023. DOI: 10.1111/1365-2664.14437.
- [34] M. A. Aizen, S. Aguiar, J. C. Biesmeijer, *et al.*, “Global agricultural productivity is threatened by increasing pollinator dependence without a parallel increase in crop diversification,” *Global Change Biology*, vol. 25, no. 10, pp. 3516–3527, 2019. DOI: 10.1111/gcb.14736.
- [35] C. Shekhar, R. Khosya, K. Thakur, *et al.*, “A systematic review of pesticide exposure, associated risks, and long-term human health impacts,” *Toxicology Reports*, vol. 13, p. 101840, 2024, ISSN: 2214-7500. DOI: 10.1016/j.toxrep.2024.101840. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S2214750024002233>.
- [36] H. Cavalier, L. Trasande, and M. Porta, “Exposures to pesticides and risk of cancer: Evaluation of recent epidemiological evidence in humans and paths forward,” *International Journal of Cancer*, vol. 152, no. 5, pp. 879–912, 2023. DOI: 10.1002/ijc.34300.

- [37] S. R. Belmain, Y. Tembo, A. G. Mkindi, S. E. J. Arnold, and P. C. Stevenson, “Elements of agroecological pest and disease management,” *Elementa: Science of the Anthropocene*, vol. 10, no. 1, p. 00099, 2022. DOI: 10.1525/elementa.2021.00099.
- [38] R. M. Anderson and R. M. May, “The invasion, persistence and spread of infectious diseases within animal and plant communities,” *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, vol. 314, no. 1167, pp. 533–570, 1986. DOI: 10.1098/rstb.1986.0072.
- [39] C. C. Mundt, “Use of multiline cultivars and cultivar mixtures for disease management,” *Annual Review of Phytopathology*, vol. 40, pp. 381–410, 2002. DOI: 10.1146/annurev.phyto.40.011402.113723.
- [40] Y. Zhu, H. Chen, J. Fan, *et al.*, “Genetic diversity and disease control in rice,” *Nature*, vol. 406, pp. 718–722, 2000. DOI: 10.1038/35021046.
- [41] A. K. Gibson and A. E. Nguyen, “Does genetic diversity protect host populations from parasites? A meta-analysis across natural and agricultural systems,” *Evolution Letters*, vol. 5, no. 1, pp. 16–32, 2021. DOI: 10.1002/evl3.206.
- [42] D. A. Andow, “Vegetational diversity and arthropod population response,” *Annual Review of Entomology*, vol. 36, no. 1, pp. 561–586, 1991. DOI: 10.1146/annurev.en.36.010191.003021.
- [43] R. B. Root, “Organization of a plant-arthropod association in simple and diverse habitats: The fauna of collards (*Brassica oleracea*),” *Ecological Monographs*, vol. 43, no. 1, pp. 95–124, 1973. DOI: 10.2307/1942161.
- [44] N.-F. Wan, Y.-Q. Wang, L. Fu, *et al.*, “Global evidence that plant diversity suppresses pests and promotes plant performance and crop production,” *Nature Ecology & Evolution*, 2025. DOI: 10.1038/s41559-025-02964-5.
- [45] J. F. Tooker and S. D. Frank, “Genotypically diverse cultivar mixtures for insect pest management and increased crop yields,” *Journal of Applied Ecology*, vol. 49, no. 5, pp. 974–985, 2012. DOI: 10.1111/j.1365-2664.2012.02173.x.
- [46] National Research Council, *Genetic Vulnerability of Major Crops*. Washington, DC: The National Academies Press, 1972. DOI: 10.17226/27600.
- [47] L. A. Tatum, “The southern corn leaf blight epidemic,” *Science*, vol. 171, no. 3976, pp. 1113–1116, 1971. DOI: 10.1126/science.171.3976.1113.
- [48] H. A. Bruns, “Southern corn leaf blight: A story worth retelling,” *Agronomy Journal*, vol. 109, no. 4, pp. 1218–1224, 2017. DOI: 10.2134/agronj2017.01.0006.
- [49] M. A. Mikel and J. W. Dudley, “Evolution of North American dent corn from public to proprietary germplasm,” *Crop Science*, vol. 46, no. 3, pp. 1193–1205, 2006. DOI: 10.2135/cropsci2005.10-0371.
- [50] A. Drenth and G. Kema, “The vulnerability of bananas to globally emerging disease threats,” *Phytopathology*, vol. 111, no. 12, pp. 2146–2161, 2021, PMID: 34231377. DOI: 10.1094/PHYTO-07-20-0311-RVW.

- [51] T. Munhoz, J. Vargas, L. Teixeira, C. Staver, and M. Dita, “Fusarium Tropical Race 4 in Latin America and the Caribbean: Status and global research advances towards disease management,” *Frontiers in Plant Science*, vol. 15, p. 1397617, 2024. DOI: 10.3389/fpls.2024.1397617.
- [52] R. C. Ploetz, “Fusarium wilt of banana is caused by several pathogens referred to as *Fusarium oxysporum* f. sp. *cubense*,” *Phytopathology*, vol. 96, no. 6, pp. 653–656, 2006. DOI: 10.1094/PHYTO-96-0653.
- [53] J. M. Roberts, L. C. Carvalhais, C. O’Dwyer, V. A. Rincón-Flórez, and A. Drenth, “Diagnostics of Fusarium wilt in banana: Current status and challenges,” *Plant Pathology*, vol. 73, no. 4, pp. 760–776, 2024. DOI: 10.1111/ppa.13863.
- [54] P. Talhinhos, D. Batista, I. Diniz, *et al.*, “The coffee leaf rust pathogen *Hemileia vastatrix*: One and a half centuries around the tropics,” *Molecular Plant Pathology*, vol. 18, no. 8, pp. 1039–1051, 2017. DOI: 10.1111/mpp.12512.
- [55] R. P. Singh, P. K. Singh, J. Rutkoski, *et al.*, “Disease impact on wheat yield potential and prospects of genetic control,” *Annual Review of Phytopathology*, vol. 54, pp. 303–322, 2016. DOI: 10.1146/annurev-phyto-080615-095835.
- [56] Z. H. Pervaiz, J. Iqbal, Q. Zhang, D. Chen, H. Wei, and M. Saleem, “Continuous cropping alters multiple biotic and abiotic indicators of soil health,” *Soil Systems*, vol. 4, no. 4, p. 59, 2020. DOI: 10.3390/soilsystems4040059. [Online]. Available: <https://doi.org/10.3390/soilsystems4040059>.
- [57] C. Li, L. Shi, K. Wang, *et al.*, “Crop rotation differentially increases soil bacterial and fungal diversities in global croplands: A meta-analysis,” *Nature Communications*, vol. 16, p. 11686, 2025. DOI: 10.1038/s41467-025-66823-4.
- [58] P. Li, J. Liu, M. Saleem, *et al.*, “Reduced chemodiversity suppresses rhizosphere microbiome functioning in the mono-cropped agroecosystems,” *Microbiome*, vol. 10, p. 108, 2022. DOI: 10.1186/s40168-022-01287-y.
- [59] Y. Zhou, Z. Yang, J. Liu, *et al.*, “Crop rotation and native microbiome inoculation restore soil capacity to suppress a root disease,” *Nature Communications*, vol. 14, no. 1, p. 8126, 2023. DOI: 10.1038/s41467-023-43926-4.
- [60] Q. Liu, Y. Zhao, T. Li, L. Chen, Y. Chen, and P. Sui, “Changes in soil microbial biomass, diversity, and activity with crop rotation in cropping systems: A global synthesis,” *Applied Soil Ecology*, vol. 186, p. 104815, 2023. DOI: 10.1016/j.apsoil.2023.104815.
- [61] M. D. McDaniel, L. K. Tiemann, and A. S. Grandy, “Does agricultural crop diversity enhance soil microbial biomass and organic matter dynamics? A meta-analysis,” *Ecological Applications*, vol. 24, no. 3, pp. 560–570, 2014. DOI: 10.1890/13-0616.1.
- [62] J. M. Tisdall and J. M. Oades, “Organic matter and water-stable aggregates in soils,” *Journal of Soil Science*, vol. 33, no. 2, pp. 141–163, 1982. DOI: 10.1111/j.1365-2389.1982.tb01755.x.

- [63] F. Zheng, X. Liu, W. Ding, X. Song, S. Li, and X. Wu, "Positive effects of crop rotation on soil aggregation and associated organic carbon are mainly controlled by climate and initial soil carbon content: A meta-analysis," *Agriculture, Ecosystems & Environment*, vol. 355, p. 108600, 2023. DOI: 10.1016/j.agee.2023.108600.
- [64] S. Mudare, J. Jing, D. Makowski, *et al.*, "Crop rotations synergize yield, nutrition, and revenue: A meta-analysis," *Nature Communications*, vol. 16, p. 9552, 2025. DOI: 10.1038/s41467-025-64567-9.
- [65] D. Renard and D. Tilman, "National food production stabilized by crop diversity," *Nature*, vol. 571, no. 7764, pp. 257–260, 2019. DOI: 10.1038/s41586-019-1316-y.
- [66] T. M. Bowles, M. Mooshammer, Y. Socolar, *et al.*, "Long-term evidence shows that crop-rotation diversification increases agricultural resilience to adverse growing conditions in North America," *One Earth*, vol. 2, no. 3, pp. 284–293, 2020. DOI: 10.1016/j.oneear.2020.02.007.
- [67] G. Tamburini, R. Bommarco, T. C. Wanger, *et al.*, "Agricultural diversification promotes multiple ecosystem services without compromising yield," *Science Advances*, vol. 6, no. 45, eaba1715, 2020. DOI: 10.1126/sciadv.aba1715.
- [68] S. Mukhovi and J. Jacobi, "Can monocultures be resilient? assessment of buffer capacity in two agroindustrial cropping systems in africa and south america," *Agriculture & Food Security*, vol. 11, no. 19, 2022. DOI: 10.1186/s40066-022-00356-7.
- [69] L. Alletto, A. Vandewalle, and P. Debaeke, "Crop diversification improves cropping system sustainability: An 8-year on-farm experiment in south-western france," *Agricultural Systems*, vol. 200, p. 103433, 2022. DOI: 10.1016/j.agsy.2022.103433.
- [70] ETC Group and GRAIN, "Top 10 agribusiness giants: Corporate concentration in food & farming in 2025," ETC Group and GRAIN, Tech. Rep., 2025. [Online]. Available: <https://www.etcgroup.org/content/top-10-agribusiness-giants>.
- [71] W. W. Cochrane, *Farm Prices: Myth and Reality*. Minneapolis: University of Minnesota Press, 1958.
- [72] G. Vanloqueren and P. V. Baret, "How agricultural research systems shape a technological regime that develops genetic engineering but locks out agroecological innovations," *Research Policy*, vol. 38, no. 6, pp. 971–983, 2009. DOI: 10.1016/j.respol.2009.02.008.
- [73] P. H. Howard, "Visualizing consolidation in the global seed industry: 1996–2008," *Sustainability*, vol. 1, no. 4, pp. 1266–1287, 2009. DOI: 10.3390/su1041266.
- [74] M. Mascarenhas and L. Busch, "Seeds of change: Intellectual property rights, genetically modified soybeans and seed saving in the united states," *Sociologia Ruralis*, vol. 46, no. 2, pp. 122–138, 2006. DOI: 10.1111/j.1467-9523.2006.00406.x.
- [75] L. L. Glenna and D. R. Cahoy, "Agribusiness concentration, intellectual property, and the prospects for rural economic benefits from the emerging biofuel economy," *Southern Rural Sociology*, vol. 24, no. 2, pp. 111–129, 2009, Also archived as *Journal of Rural Social Sciences* 24(2), Art. 6. [Online]. Available: <https://egrove.olemiss.edu/jrss/vol24/iss2/6>.

- [76] K. Deconinck, “Concentration in seed and biotech markets: Extent, causes, and impacts,” *Annual Review of Resource Economics*, vol. 12, pp. 129–147, 2020. DOI: 10.1146/annurev-resource-102319-100751.
- [77] L. Keenan, T. Monteath, and D. Wójcik, “Hungry for power: Financialization and the concentration of corporate control in the global food system,” *Geoforum*, vol. 147, p. 103 909, 2023. DOI: 10.1016/j.geoforum.2023.103909.
- [78] J. Clapp, “The rise of financial investment and common ownership in global agrifood firms,” *Review of International Political Economy*, vol. 26, no. 4, pp. 604–629, 2019. DOI: 10.1080/09692290.2019.1597755.
- [79] A. Puy, S. N. Linga, N. Wei, *et al.*, “Widely cited global irrigation statistics lack empirical support,” *PNAS Nexus*, vol. 4, no. 11, pgaf323, 2025. DOI: 10.1093/pnasnexus/pgaf323.
- [80] C. Dalin, Y. Wada, T. Kastner, and M. J. Puma, “Groundwater depletion embedded in international food trade,” *Nature*, vol. 543, pp. 700–704, 2017. DOI: 10.1038/nature21403.
- [81] X. Kuang, J. Liu, B. R. Scanlon, *et al.*, “The changing nature of groundwater in the global water cycle,” *Science*, vol. 383, no. 6686, eadf0630, 2024. DOI: 10.1126/science.adf0630.
- [82] S. Jasechko, H. Seybold, D. Perrone, *et al.*, “Rapid groundwater decline and some cases of recovery in aquifers globally,” *Nature*, vol. 625, no. 7996, pp. 715–721, 2024. DOI: 10.1038/s41586-023-06879-8.
- [83] D. Long, Y. Xu, Y. Cui, *et al.*, “Unprecedented large-scale aquifer recovery through human intervention,” *Nature Communications*, vol. 16, p. 7296, 2025. DOI: 10.1038/s41467-025-62719-5.
- [84] FAO, “Greenhouse gas emissions from agrifood systems – global, regional and country trends, 2000–2022,” Food and Agriculture Organization of the United Nations, Rome, FAOSTAT Analytical Brief Series 94, 2024. [Online]. Available: <https://openknowledge.fao.org/handle/20.500.14283/cd3167en>.
- [85] M. Crippa, E. Solazzo, D. Guizzardi, F. Monforti-Ferrario, F. Tubiello, and A. Leip, “Food systems are responsible for a third of global anthropogenic GHG emissions,” *Nature Food*, vol. 2, no. 3, pp. 198–209, 2021. DOI: 10.1038/s43016-021-00225-9.
- [86] F. N. Tubiello, K. Karl, A. Flammini, *et al.*, “Pre- and post-production processes increasingly dominate greenhouse gas emissions from agri-food systems,” *Earth System Science Data*, vol. 14, pp. 1795–1809, 2022. DOI: 10.5194/essd-14-1795-2022.
- [87] X. Xu, P. Sharma, S. Shu, *et al.*, “Global greenhouse gas emissions from animal-based foods are twice those of plant-based foods,” *Nature Food*, vol. 2, pp. 724–732, 2021. DOI: 10.1038/s43016-021-00358-x.
- [88] J. Poore and T. Nemecek, “Reducing food’s environmental impacts through producers and consumers,” *Science*, vol. 360, no. 6392, pp. 987–992, 2018. DOI: 10.1126/science.aaq0216.
- [89] M. A. Clark, N. G. G. Domingo, K. Colgan, *et al.*, “Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets,” *Science*, vol. 370, pp. 705–708, 2020. DOI: 10.1126/science.aba7357.

- [90] A. Flammini, H. Adzmir, R. Pattison, K. Karl, Y. Allouche, and F. N. Tubiello, “Greenhouse gas emissions from cold chains in agrifood systems,” *Sustainability*, vol. 16, no. 21, p. 9184, 2024. DOI: 10.3390/su16219184.
- [91] United States Environmental Protection Agency, “Quantifying methane emissions from landfilled food waste,” US EPA, EPA-600-R-23-064, 2023. [Online]. Available: <https://www.epa.gov/land-research/quantifying-methane-emissions-landfilled-food-waste>.
- [92] J. Clapp, “Concentration and crises: Exploring the deep roots of vulnerability in the global industrial food system,” *Journal of Peasant Studies*, vol. 50, no. 1, pp. 1–25, 2023. DOI: 10.1080/03066150.2022.2129013.
- [93] USDA Foreign Agricultural Service, “Grain: World markets and trade,” United States Department of Agriculture, Tech. Rep., Aug. 2025. [Online]. Available: <https://apps.fas.usda.gov/psdonline/circulars/grain.pdf>.
- [94] J. Clapp, “The problem with growing corporate concentration and power in the global food system,” *Nature Food*, vol. 2, no. 6, pp. 404–408, 2021. DOI: 10.1038/s43016-021-00297-7.
- [95] UNCTAD, “Strait of Hormuz disruptions: Implications for global trade and development,” United Nations Conference on Trade and Development, Geneva, Tech. Rep. UNCTAD/OSG/TT/INF/2026/1, 2026. [Online]. Available: <https://unctad.org/publication/strait-hormuz-disruptions-implications-global-trade-and-development>.
- [96] World Bank Group, “Commodity markets outlook,” World Bank Group, Washington, DC, Tech. Rep., Apr. 2026.
- [97] FAO. “Strait of Hormuz crisis: Fertilizer scarcity will affect next harvests and food supplies, FAO warns.” Statement by Director-General QU Dongyu at the MED9++ Ministerial Meeting, Rome, Food and Agriculture Organization of the United Nations. (May 2026), [Online]. Available: <https://www.fao.org/newsroom/detail/strait-of-hormuz-crisis--fertilizer-scarcity-will-affect-next-harvests-and-food-supplies--fao-warns/en>.
- [98] FAO, “The state of food and agriculture 2025 — addressing land degradation across landholding scales,” Food and Agriculture Organization of the United Nations, Rome, Tech. Rep., 2025. DOI: 10.4060/cd7067en.
- [99] C. A. Monteiro, M. L. C. Louzada, E. Steele-Martinez, G. Cannon, G. C. Andrade, P. Baker, *et al.*, “Ultra-processed foods and human health: The main thesis and the evidence,” *The Lancet*, vol. 406, pp. 2667–2684, 2025. DOI: 10.1016/S0140-6736(25)01565-X.
- [100] C. A. Monteiro, J.-C. Moubarac, G. Cannon, S. W. Ng, and B. Popkin, “Ultra-processed products are becoming dominant in the global food system,” *Obesity Reviews*, vol. 14, no. S2, pp. 21–28, 2013. DOI: 10.1111/obr.12107.
- [101] M. M. Lane, E. Gamage, S. Du, *et al.*, “Ultra-processed food exposure and adverse health outcomes: Umbrella review of epidemiological meta-analyses,” *BMJ*, vol. 384, e077310, 2024. DOI: 10.1136/bmj-2023-077310.

- [102] M.-S. Fan, F.-J. Zhao, S. J. Fairweather-Tait, P. R. Poulton, S. J. Dunham, and S. P. McGrath, “Evidence of decreasing mineral density in wheat grain over the last 160 years,” *Journal of Trace Elements in Medicine and Biology*, vol. 22, no. 4, pp. 315–324, 2008. DOI: 10.1016/j.jtemb.2008.07.002.
- [103] D. R. Davis, M. D. Epp, and H. D. Riordan, “Changes in USDA food composition data for 43 garden crops, 1950 to 1999,” *Journal of the American College of Nutrition*, vol. 23, no. 6, pp. 669–682, 2004. DOI: 10.1080/07315724.2004.10719409.
- [104] R. J. Marles, “Mineral nutrient composition of vegetables, fruits and grains: The context of reports of apparent historical declines,” *Journal of Food Composition and Analysis*, vol. 56, pp. 93–103, 2017. DOI: 10.1016/j.jfca.2016.11.012.
- [105] A.-M. B. Mayer, L. Trenchard, and F. Rayns, “Historical changes in the mineral content of fruit and vegetables in the UK from 1940 to 2019: A concern for human nutrition and agriculture,” *International Journal of Food Sciences and Nutrition*, vol. 73, no. 3, pp. 315–326, 2022. DOI: 10.1080/09637486.2021.1981831.
- [106] R. M. Welch and R. D. Graham, “Breeding for micronutrients in staple food crops from a human nutrition perspective,” *Journal of Experimental Botany*, vol. 55, no. 396, pp. 353–364, 2004. DOI: 10.1093/jxb/erh064.
- [107] C. K. Khoury, A. D. Bjorkman, H. Dempewolf, *et al.*, “Increasing homogeneity in global food supplies and the implications for food security,” *Proceedings of the National Academy of Sciences*, vol. 111, no. 11, pp. 4001–4006, 2014. DOI: 10.1073/pnas.1313490111.
- [108] M. Springmann *et al.*, “Options for keeping the food system within environmental limits,” *Nature*, vol. 562, pp. 519–525, 2018. DOI: 10.1038/s41586-018-0594-0.