

Decentralizing Agriculture

A shift in the industrial paradigm of food production

Marco Zolla Salazar

April 16, 2026

Outline

- 1 The many problems of industrial agriculture
- 2 Research methodology
- 3 Background
- 4 Proposed solution
- 5 References

Land use and biodiversity loss

- Cropping and animal husbandry cover over one third of the terrestrial land surface; agriculture drives roughly 90% of global deforestation [1, p. XVI], [2, p. 4].
- More than 90% of biodiversity impacts from land-use change are attributable to agriculture — crops alone account for 72% [3].
- A 27-year study found a 76% decline in flying insect biomass; European farmland birds fell 57% over 37 years [4], [5].

Soil degradation

- Conventional tillage erodes soil one to two orders of magnitude faster than it forms — “fundamentally unsustainable over civilizational timescales” [6].
- Cropland soils have lost 20–60% of their pre-cultivation organic carbon [7, TS.4].
- A meta-analysis of 25,000+ studies confirms 16–25% SOC loss following conversion of grassland or forest to cropland [8].

Chemical inputs: pesticides and fertilizers

- 64% of global agricultural land is at risk of pesticide pollution; 31% at high risk [9, p. 206].
- Fertilizer runoff has driven the spread of coastal dead zones to over 500 documented systems worldwide [10].
- Agricultural N_2O emissions have grown 40% since 1980, exceeding all CMIP6 emission scenarios [11, Section 4.2.1].

Monoculture and structural lock-in

- Diversified rice plantings suffered 94% less rice blast and yielded 89% more per hill than monoculture [12, pp. 718–721].
- Global banana production hangs on a single cultivar (Cavendish); its predecessor was wiped out by Fusarium wilt — and is now under threat from a new strain [13, pp. 2147–2150].
- Four firms control 56% of the global seed market and 61% of pesticides — agronomic uniformity is a structural outcome of input-market concentration [14], [15].

Freshwater depletion and greenhouse gas emissions

- Agriculture accounts for 70–85% of global freshwater withdrawals [16].
- Five crops drove most of the 22% rise in global groundwater depletion between 2000 and 2010 — much of it from non-renewable aquifers [17].
- The agrifood system emits ~30% of all anthropogenic GHGs (16.2 Gt CO₂eq); on a business-as-usual path, food alone could exhaust the 1.5°C carbon budget [18], [19].

Supply-chain externalities: transport, storage, food waste

- 25–30% of food produced is lost or wasted; food loss and waste alone account for 8–10% of anthropogenic GHGs [7, Section 5.5.2.5].
- Cold-chain refrigeration emits $\sim 10\%$ of all agrifood GHGs and has more than doubled since 2000 [20, Section 3].
- Food waste is responsible for an estimated 58% of US landfill methane, with 61% escaping uncaptured [21, Table 3].

Concentration and dependency at every level

- **Country level** — 7 countries plus the EU produce 88% of world wheat exports; most nations depend on a handful of suppliers [22, p. 16].
- **Corporate level** — four firms dominate the global seed (56%), pesticide (61%), and machinery (43%) markets [14].
- **Community level** — 0.1% of farms operate over half of all agricultural land; the 85% of farms under 2 ha hold just 9% [23, Chapter 3].

Suppression of dietary diversity and nutritional decline

- Industrial systems funnel commodity outputs into ultra-processed foods, linked to higher cardiovascular mortality, type-2 diabetes, and obesity (umbrella review, ~10M participants) [24, Figures 2–3].
- Wheat zinc, copper, and magnesium concentrations declined 20–49% after the introduction of high-yield cultivars in the 1960s [25, Tables 2, 4].
- Nutritional decline tracks the narrowing of crop genetic diversity that monoculture demands [26].

An iterative, source-grounded verification pipeline

Three custom slash commands form a single research pipeline:

- `/verify-paper` — paper-wide audit; one subagent per section dispatched in parallel; produces a master corrections table.
- `/verify-section` — interactive, claim-by-claim verification of citations, narrative, and structure for one section.
- `/expand-section` — gap-filling pass; identifies missing evidence and surfaces cross-section relevance.
- All findings persist in `sources/corrections.md` as stable IDed items, working across multiple sessions.

Per-citation verification protocol

One subagent per source file, reading both:

- OCR text — statistics, quotes, exact wording.
- Original PDF — figures, tables, layout.

Each claim is checked for:

- Factual and temporal accuracy
- Primary vs. secondary attribution
- Scope and caveats preservation
- Disconfirming evidence in the surrounding text
- Bibliography metadata (author, year, DOI)

The orchestrator independently spot-checks every non-OK verdict before any edit; pinpoint locators (`\cite[Section X]{key}`) are required on every modified citation.

Standing principles

- **No sycophancy** — ideas are evaluated against the thesis before agreement; corrections are preferred to retractions.
- **Primary sources only** — every secondary citation is flagged to the user, never filtered out by the assistant.
- **Reproducibility** — every claim must be traceable to a source on disk; no verification from model memory.
- **Failure mode avoided** — confident citation of things the source does not actually say.


Case study: The Netherlands

- “Dutch greenhouses aren’t your run-of-the-mill structures - they’re technological marvels.” [27]
- The Netherlands, despite its small land area, is the second-largest agricultural exporter in the world by value [28]

Objectives

- Agriculture decentralization
- Optimize production efficiency
- Create a scalable, automatic and widely available solution

Architecture overview



Optimization through collaboration

- Open source hardware, software, models and data
- Guides for personal and commercial use
- Seed sharing communities and markets

Bibliography I

- [1] Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, “Global assessment report of the intergovernmental science-policy platform on biodiversity and ecosystem services,” IPBES Secretariat, Bonn, Germany, Tech. Rep., 2019, IPBES Analytical Brief, No. 107, p. 1144. DOI: 10.5281/zenodo.3831673.
- [2] Food and Agriculture Organization of the United Nations, “The state of food and agriculture 2025 – addressing land degradation across landholding scales,” FAO, Rome, Tech. Rep., 2025, FAO publication. DOI: 10.4060/cd7067en. [Online]. Available: <https://doi.org/10.4060/cd7067en>.
- [3] L. Cabernard *et al.*, “Biodiversity impacts of recent land-use change driven by increases in agri-food imports,” *Nature Sustainability*, 2024. DOI: 10.1038/s41893-024-01433-4.

Bibliography II

- [4] C. A. Hallmann, M. Sorg, E. Jongejans, *et al.*, “More than 75 percent decline over 27 years in total flying insect biomass in protected areas,” *PLOS ONE*, vol. 12, no. 10, e0185809, 2017. DOI: 10.1371/journal.pone.0185809.
- [5] S. Rigal, V. Dakos, H. Alonso, *et al.*, “Farmland practices are driving bird population decline across Europe,” *Proceedings of the National Academy of Sciences*, vol. 120, no. 21, e2216573120, 2023. DOI: 10.1073/pnas.2216573120.
- [6] D. R. Montgomery, “Soil erosion and agricultural sustainability,” *Proceedings of the National Academy of Sciences*, vol. 104, no. 33, pp. 13 268–13 272, 2007. DOI: 10.1073/pnas.0611508104.

Bibliography III

- [7] IPCC, “Climate change and land: An IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems,” Intergovernmental Panel on Climate Change, Cambridge, UK and New York, NY, USA, Tech. Rep., 2019. DOI: 10.1017/9781009157988. [Online]. Available: <https://www.ipcc.ch/srccl/>.
- [8] D. Beillouin *et al.*, “A global meta-analysis of soil organic carbon in the Anthropocene,” *Nature Communications*, vol. 14, p. 3700, 2023. DOI: 10.1038/s41467-023-39338-z.
- [9] F. H. M. Tang, M. Lenzen, A. McBratney, and F. Maggi, “Risk of pesticide pollution at the global scale,” *Nature Geoscience*, vol. 14, pp. 206–210, Apr. 2021. DOI: 10.1038/s41561-021-00712-5.

Bibliography IV

- [10] D. Breitburg, L. A. Levin, A. Oschlies, *et al.*, “Declining oxygen in the global ocean and coastal waters,” *Science*, vol. 359, no. 6371, eaam7240, 2018. DOI: 10.1126/science.aam7240.
- [11] H. Tian, N. Pan, R. L. Thompson, *et al.*, “Global nitrous oxide budget (1980–2020),” *Earth System Science Data*, vol. 16, no. 6, pp. 2543–2604, 2024. DOI: 10.5194/essd-16-2543-2024.
- [12] Y. Zhu, H. Chen, J. Fan, *et al.*, “Genetic diversity and disease control in rice,” *Nature*, vol. 406, pp. 718–722, 2000. DOI: 10.1038/35021046.

Bibliography V

- [13] A. Drenth and G. Kema, “The vulnerability of bananas to globally emerging disease threats,” *Phytopathology*, vol. 111, no. 12, pp. 2146–2161, 2021, PMID: 34231377. DOI: 10.1094/PHYTO-07-20-0311-RVW.
- [14] ETC Group and GRAIN, “Top 10 agribusiness giants: Corporate concentration in food & farming in 2025,” ETC Group and GRAIN, Tech. Rep., 2025. [Online]. Available: <https://www.etcgroup.org/content/top-10-agribusiness-giants>.
- [15] P. H. Howard, “Visualizing consolidation in the global seed industry: 1996–2008,” *Sustainability*, vol. 1, no. 4, pp. 1266–1287, 2009. DOI: 10.3390/su1041266.

Bibliography VI

- [16] A. Puy, S. N. Linga, N. Wei, *et al.*, “Widely cited global irrigation statistics lack empirical support,” *PNAS Nexus*, vol. 4, no. 11, pgaf323, 2025. DOI: [10.1093/pnasnexus/pgaf323](https://doi.org/10.1093/pnasnexus/pgaf323).
- [17] C. Dalin, Y. Wada, T. Kastner, and M. J. Puma, “Groundwater depletion embedded in international food trade,” *Nature*, vol. 543, pp. 700–704, 2017. DOI: [10.1038/nature21403](https://doi.org/10.1038/nature21403).
- [18] FAO, “Greenhouse gas emissions from agrifood systems – global, regional and country trends, 2000–2022,” Food and Agriculture Organization of the United Nations, Rome, FAOSTAT Analytical Brief Series 94, 2024. [Online]. Available: <https://openknowledge.fao.org/handle/20.500.14283/cd3167en>.

Bibliography VII

- [19] M. A. Clark, N. G. G. Domingo, K. Colgan, *et al.*, "Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets," *Science*, vol. 370, pp. 705–708, 2020. DOI: [10.1126/science.aba7357](https://doi.org/10.1126/science.aba7357).
- [20] A. Flammini, H. Adzmir, R. Pattison, K. Karl, Y. Allouche, and F. N. Tubiello, "Greenhouse gas emissions from cold chains in agrifood systems," *Sustainability*, vol. 16, no. 21, p. 9184, 2024. DOI: [10.3390/su16219184](https://doi.org/10.3390/su16219184).
- [21] United States Environmental Protection Agency, "Quantifying methane emissions from landfilled food waste," US EPA, EPA-600-R-23-064, 2023. [Online]. Available: <https://www.epa.gov/land-research/quantifying-methane-emissions-landfilled-food-waste>.

Bibliography X

- [27] H. B. School, “High-tech greenhouse innovations in the netherlands,” , 2024, Accessed: 2026-03-10. [Online]. Available:
<https://www.hbs.edu/environment/blog/post/IFC-2024-Greenhouses>.
- [28] International Trade Administration, U.S. Department of Commerce, *Netherlands – agriculture*, Accessed: 2026-03-21. States: “The Netherlands is the world’s second largest exporter of agricultural products after the United States.”, 2026. [Online]. Available:
<https://www.trade.gov/country-commercial-guides/netherlands-agriculture>.