


# 3

## Atoms

**M**ATTER is everywhere around you, from the water you drink to the air you inhale. Matter is made of elements and elements are made of atoms. Even the atoms of an element can be different, having a distinct number of protons and neutrons. This chapter covers the principles of atomic structure. You will learn what makes an atom and will be able to quantify the particles that make atoms. Perhaps more importantly, you will also learn about the periodic table of the elements and the different types of chemical formulas.



### 3.1 The periodic table

The periodic table is a chart containing all known elements arranged in increasing number of electrons per atom in a way that elements with similar chemical and physical properties are located together. The periodic table contains all existing elements—some of them are synthetic others are natural—that form the matter arranged in columns and rows. Every element has a different name accompanied by a symbol that represents their name. The tabular arrangement of elements in the form of rows and columns allow further classification of the elements according to their properties. This section will cover the different features of the periodic table.

*Elements and Symbols* Elements cannot be broken down into simpler substances. For example aluminum is an element only made of aluminum atoms and if you analyze the composition of a piece of this metals you would only find aluminium atoms. Chemical symbols are one- or two-letter abbreviations that represent the names of the elements. Only the first letter is capitalized and if a second letter exist in the element's name, the second letter should be lowercase. For example, the chemical symbol for aluminum is Al, written as capital A and lowercase l.

#### Sample Problem 1

Give the symbol or name the following elements: C, Oxygen, N, Phosphorus, Au, Iron, Na and Iodine.

#### **SOLUTION**

The name of the element with symbol C is carbon, whereas the chemical symbol of oxygen is O. Similarly, N stands for nitrogen, whereas the chemical symbol of Phosphorus is P. The chemical symbol of Au is Gold. The chemical symbol of Iron is Fe and the chemical symbol of Iodine is I.

#### **STUDY CHECK**

Give the symbol or name the following elements: Ni.

*Periods and groups* The periodic table contains all elements arranged in rows and columns. The horizontal rows are called *periods* and the vertical columns are called *groups or families*. For example, the first period contains hydrogen (H) and helium (He), whereas the second group contains Beryllium (Be), Magnesium (Mg), Calcium (Ca), Strontium (Sr), Barium (Ba) and Radium (Ra). There are seven periods (periods 1-7) and 18 groups. Some of the groups are labeled with an A (e.g. group 8A) whereas others are labeled with a B (e.g. group 8B). The groups numbers can be found written with roman numbers and a letters (A or B) or with a more modern group numbering of 1-18 going across the periodic table. For example, the group 2 (Mg-Ra) can also be called IIA, and the group 13 (B-Ti) is also known as IIIA.

*Properties in the periodic table* The physical and chemical properties of some elements of the table are similar, and these similarities led to the organization of the periodic table. Elements in the same group share properties and for example, oxygen and sulfur have similar properties: both are reactive elements. Differently, the properties across periods change going from metals to nonmetals. For example, the properties of Li and Ne are very different and lithium is a reactive metal whereas neon is a nonreactive gas.

*Metals, Nonmetals, and Metalloids* Overall, the elements of the periodic table can be classify as metals, nonmetals, and metalloids. Metals are those elements on the left of the table and nonmetals are the elements on the right of the table. The elements between metals and nonmetals are called metalloids and include only B, Si, Ge, As, Sb, Te, Po, and At. Metals are shiny solids and usually melt at higher temperatures. Some examples of metals are Gold (Au) or Iron (Fe). Nonmetals are often poor conductors of heat and electricity with low melting points. They also tend to be matt (non-shinny), malleable, or ductile. Some examples of nonmetals are Carbon (C) or Nitrogen (N). Metalloids are elements that share some properties with metals and others with the nonmetals. For example, they are better conductors of heat and electricity than the nonmetals, but not as good conductors as the metals. Metalloids are semiconductors because they can act as both conductors and insulators under certain conditions. An example of metalloids is Silicon (Si) that should not be confused from silicone, a chemical employed in prosthetics.

#### Sample Problem 2

Answer the following questions: (a) Give the group and period of the following elements, and give the name: Ca, Ir, and C. (b) Classify as alkali metal, alkali earth metal, transition metal, halogen or noble gas, and give the name: Mg, Li, Co, He, F. (c) Classify as metal, nonmetal or metalloid, and give the name: Ba, N, Si.

#### **SOLUTION**

(a)The period and group of Ca (Calcium) is 2 (2A) and 4, respectively. The period and group of Ir (Iridium) is 9 (8B) and 6, respectively. The period and group of C (Carbon) is 14 (IVA) and 2, respectively. (b) Mg (Magnesium) is an alkali earth metal, whereas Li (Lithium) is a alkali metal. Co (Cobalt) is a transition metal. He (Helium) is a noble gas. F (Fluorine) is an halogen. (c) Ba (Barium) is a metal. N (Nitrogen) is a nonmetal. Si (Silicon) is a metalloid.

### STUDY CHECK

Answer the following questions: (a) Give the group and period of the following elements, and give the name: Cl. (b) Classify as alkali metal, alkali earth metal, transition metal, halogen or noble gas, and give the name: Ne. (c) Classify as metal, nonmetal or metalloid, and give the name: W.

*Classification of elements in terms of groups* Some of the groups in the periodic table have specific names such as alkali metals, alkaline earth metals, transition metals, chalcogens, halogens or noble gases. Alkali metals are the group 1A elements: lithium (Li), sodium (Na), potassium (K), rubidium (Rb), cesium (Cs), and francium (Fr). Alkali elements are soft and shiny metals, and they are also good conductors of heat and electricity, with low melting points. Alkali earth metals are the group 2A (2) elements: beryllium (Be), magnesium (Mg), calcium (Ca), strontium (Sr), barium (Ba), and radium (Ra). Transition metals are the elements from group 3 to 12 and they are located in the middle of the table. Chalcogens are the group 6A (16) elements: oxygen (O), sulfur (S), selenium (Se), tellurium (Te), and polonium (Po). Halogens are the group 7A (17) elements: fluorine (F), chlorine (Cl), bromine (Br), iodine (I), and astatine (At). Halogens are very reactive elements. Finally, noble gases are the group 8A (18) elements: helium (He), neon (Ne), argon (Ar), krypton (Kr), xenon (Xe), and radon (Rn). They are inert and rarely combine with other elements in the periodic table, like a noble family: have you ever meet a royal?

*How to classify Hydrogen* At first sight, hydrogen (H) may seem to be put in the wrong spot at the periodic table. Although it is located at the top of Group 1A (1), it is not an alkali metal, as it has very different properties. Thus hydrogen does not belong to the alkali metals, being nonmetal.


1 IA																		2 IIA																		13 IIIA																		14 IVA		15 VA		16 VIA		17 VIIA		18 VIIIA																																																																																																																																																																																																																																																																					
1 1.0079 H Hydrogen																		2 4.0026 He Helium																																																																																																																																																																																																																																																																																																																	
3 6.941 Li Lithium																		4 9.0122 Be Beryllium																																																																																																																																																																																																																																																																																																																	
11 22.990 Na Sodium																		12 24.305 Mg Magnesium																																																																																																																																																																																																																																																																																																																	
19 39.098 K Potassium																		20 40.078 Ca Calcium																		21 44.956 Sc Scandium																		22 47.867 Ti Titanium																		23 50.942 V Vanadium																		24 51.996 Cr Chromium																		25 54.938 Mn Manganese																		26 55.845 Fe Iron																		27 58.933 Co Cobalt																		28 58.693 Ni Nickel																		29 63.546 Cu Copper																		30 65.39 Zn Zinc																		31 69.723 Ga Gallium																		32 72.64 Ge Germanium																		33 74.922 As Arsenic																		34 78.96 Se Selenium																		35 79.904 Br Bromine																		36 83.8 Kr Krypton																	
37 85.468 Rb Rubidium																		38 87.62 Sr Strontium																		39 88.906 Y Yttrium																		40 91.224 Zr Zirconium																		41 92.906 Nb Niobium																		42 95.94 Mo Molybdenum																		43 96 Tc Technetium																		44 101.07 Ru Ruthenium																		45 102.91 Rh Rhodium																		46 106.42 Pd Palladium																		47 107.87 Ag Silver																		48 112.41 Cd Cadmium																		49 114.82 In Indium																		50 118.71 Sn Tin																		51 121.76 Sb Antimony																		52 127.6 Te Tellurium																		53 126.9 I Iodine																		54 131.29 Xe Xenon																	
55 132.91 Cs Cesium																		56 137.33 Ba Barium																		57-71 La-Lu Lanthanide																		72 178.49 Hf Hafnium																		73 180.95 Ta Tantalum																		74 183.84 W Tungsten																		75 186.21 Re Rhenium																		76 190.23 Os Osmium																		77 192.22 Ir Iridium																		78 195.08 Pt Platinum																		79 196.97 Au Gold																		80 200.59 Hg Mercury																		81 204.38 Tl Thallium																		82 207.2 Pb Lead																		83 208.98 Bi Bismuth																		84 209 Po Polonium																		85 210 At Astatine																		86 222 Rn Radon																	
87 223 Fr Francium																		88 226 Ra Radium																		89-103 Ac-Lr Actinide																		104 261 Rf Rutherfordium																		105 262 Db Dubnium																		106 266 Sg Seaborgium																		107 264 Bh Bohrium																		108 Hs Hassium																		109 Mt Meitnerium																		110 Ds Darmstadtium																		111 Rg Roentgenium																		112 Uub Ununbium																		113 Uut Ununtrium																		114 Uuq Ununquadium																		115 Uup Ununpentium																		116 Uuh Ununhexium																		117 Uus Ununseptium																		118 Uuo Ununoctium																	
57 138.91 La Lanthanum																		58 140.12 Ce Cerium																		59 140.91 Pr Praseodymium																		60 144.24 Nd Neodymium																		61 145 Pm Promethium																		62 150.36 Sm Samarium																		63 151.96 Eu Europium																		64 157.25 Gd Gadolinium																		65 158.93 Tb Terbium																		66 162.50 Dy Dysprosium																		67 164.93 Ho Holmium																		68 167.26 Er Erbium																		69 168.93 Tm Thulium																		70 173.04 Yb Ytterbium																		71 174.97 Lu Lutetium																																																																							
89 227 Ac Actinium																		90 232.04 Th Thorium																		91 231.04 Pa Protactinium																		92 238.03 U Uranium																		93 237 Np Neptunium																		94 244 Pu Plutonium																		95 243 Am Americium																		96 247 Cm Curium																		97 247 Bk Berkelium																		98 251 Cf Californium																		99 252 Es Einsteinium																		100 257 Fm Fermium																		101 258 Md Mendelevium																		102 259 No Nobelium																		103 262 Lr Lawrencium																																																																							

Figure 1.1 The periodic table of the elements

## 3.2 Early experiments of the atom

Scientists wondered about the nature of the atom and its structure for years. In a series of experiments carried in the late nineteenth century, scientists such as J.J. Thomson, Henri Becquerel, or Ernest Rutherford gained insight into the nature and structure of the atom. These remarkable scientists and these creative experiments helped shape the view of the atom that we have nowadays.

▼ A cathode tube



© www.wallpaperflare.com

▼ Millikan apparatus




© www.wallpaperflare.com

▼ A plum pudding, with the electrons represented by the raisins



© wikipedia



**Figure 1.2** Cathode-ray tube made of two electrodes and a partially evacuated gas tube. The electrons, generated on the negatively charged electrode names cathode, excite the gas in the tube generating glow.

*Charge to mass ratio of an electron* The English researcher J.J. Thomson investigated electric discharges in partially evacuated tubes—tubes in which the air has been partially removed (See Figure 1.2); these tubes, made of a positive and negative electrode, are the base of old-fashion, bulky televisions. Thomson found that rays emanated from the negative electrode when applying high-voltage to these tubes. These rays were names cathodic rays as the negative electrode of the tube is called the cathode. Thomson also found that cathodic rays were repelled from the negative pole of an electric field. Hence, these rays were postulated to be a stream of negatively charged particles, now known as electrons. By studying the deflection of these rays by an electric field, Thomson was able to calculate the charge-to-mass ratio of an electron:

$$\frac{e}{m_e} = -1.76 \times 10^8 C/g \quad (3.1)$$

where:

$e$  is the charge of an electron

$m_e$  is the mass of an electron


Overall, the biggest of Thomson's discoveries was that all atoms are made of negatively charged particles. As atoms are charge-neutral, they are also made of positively charged particles. These observations led to a new atomic model, the *plum pudding model*, that envisioned atoms as a diffuse cloud of positive charges with negative electrons embedded in it. The name plum pudding comes from an English dessert that contains a raising spread on it. A different scientists, Robert Millikan, revealed the magnitude of the electric charge of the electron. Millikan used an apparatus that dispersed charged oil droplets falling under the influence of an electric field. Given the applied voltage and the droplet mass, Millikan was able to calculate the droplet charge. He found that the oil drop charge was always a whole-number time the electron charge,

$$e = 1.60 \times 10^{-19} C \quad (3.2)$$

where a Coulomb is a unit of charge. With the value of the charge-to-mass ratio of an electron and the electron charge, Millikan was also able to calculate the mass of an electron,

$$m_e = 9.11 \times 10^{-31} \text{ kg.} \quad (3.3)$$

*The atom nucleus* Ernest Rutherford carried further experiments to validate the plum pudding model of the atom. He exposed a thin sheet of metal foil to  $\alpha$  particles known to be massive and positively charged particles. According to the plum pudding model, the bulky  $\alpha$  particles should have crashed through the thin foil and traverse through without being deflected. However, the results did not corroborate his expectations. Indeed, some particles traversed the film, whereas other were slightly deflected and some were strongly deflected at large angles. These observations did not corroborate the plum pudding model. However, they contributed to the creation of the modern atomic model in which a large number of positive charges were concentrated at a point—called the nucleus—instead of being spread whereas the electrons move around the nucleus at large distances from it. The figure below represents the experiment carried out by Rutherford in which alpha particles were scattered on a thin foil made of gold.




**Figure 1.3** Rutherford's scattering: the elastic scattering of charged particles by the Coulomb interaction after a particle beam passed through a thin gold foil obstruction.

*Radiation* In the last part of the nineteenth century, scientists came to the discovery that some materials were able to produce high-energy radiation. Among the scientist working in this field, Henri Becquerel discovered that pitchblende, a mineral containing uranium, was able to produce an image on a photographic plate in the absence of light. In the early twentieth century, three different types of radiation were discovered: alpha radiation, beta radiation, and gamma radiation. Further studies revealed that gamma radiation was made of gamma particles, high-energy radiation, whereas beta radiation was made of high-energy electrons. Alpha particles were found to be positively charged, with a charge twice the charge of an electron and a mass 7300 times that of the electron. These particles were indeed Helium nucleus, resulting from removing electrons from atoms of Helium.

### 3.3 The atom

Atoms are the smallest piece of an element that retains its characteristics. They are the building blocks of matter. This section covers the structure of the atom. You will learn how

to calculate the number of subatomic particles that made an atom and how to differentiate atoms of an element—all atoms of an element are not equal.




**Figure 1.4** Three models of the atom. From left to right: the plum pudding model, the updated plum pudding model according to Rutherford's observations, (two right images) the modern atomic model.  $Z$  is the atomic number of the atom.

*Atomic Structure* Atoms contain three atomic particles: the proton, neutron, and electron. Protons have positive charge (+), whereas electrons carry negative charge (-). Neutrons on the other hand are neutral, and they have no electrical charge. Protons and neutrons are located in the core of the atom, which is called the nucleus, and account for the mass of the atom. Electrons are delocalized in the exterior part of the atoms. They are not necessarily located in a specific spot and their existence spreads in the area next to the nucleus. When an atom is neutral it has no charge and the number of electrons and protons are the same. Some atoms have positive charge, resulting from removing electrons, and we call these cations. Others—called anions—can have negative charge as a result of accepting a negatively charged electron.

*Atomic and mass number* Elements are made of atoms, and each atom of an element is characterized by a atomic number ( $Z$ ) and a mass number ( $A$ ). The atomic number ( $Z$ ) of an element indicates the number of protons of an atom. This number can be easily located in the periodic table. The mass number ( $A$ ) of an element indicates the combined number of protons and neutrons. Mass numbers are nowhere located in the periodic table as different atoms of the same element can have different mass numbers. Both  $A$  and  $Z$  for an atom  $X$  are indicated in the following form called isotope notation:

$${}^A_ZX$$

As an example, the notation  ${}^{24}_{12}\text{Mg}$  means that the atomic number of Mg is  $Z=12$  and the mass number is  $A=24$ . Remember that the atomic number can be found in the periodic table whereas the mass number  $A$  is not on the table. By means of the isotope notation, one can quickly identify the number of protons, neutrons and electrons in an atom. As the atomic number is always indicated on the bottom part (e.g. Mg has 12 electrons). At the same time, the number of electrons and protons in a neutral atom is the same—neutral means an atom without a charge. The number of neutrons of an isotope can be computed by subtracting the atomic number from the mass number.



## Sample Problem 3

Calculate the number of protons, neutrons and electrons of the following atoms:

- (a)  ${}^{27}_{12}\text{Mg}$                       (b)  ${}^{22}_{10}\text{Ne}$                       (c)  ${}^{20}_{10}\text{Ne}$

**SOLUTION**

(a)  ${}^{27}_{12}\text{Mg}$  has 12 electrons ( $Z=12$ ) and 12 protons as well (the number of electrons and protons are the same if the atom is neutral), and 15 neutrons, as  $27-12=15$ .

(b)  ${}^{22}_{10}\text{Ne}$  has 10 electrons and 10 protons, and 12 neutrons. (c)  ${}^{20}_{10}\text{Ne}$  has 10 electrons and 10 protons, and 10 neutrons as well.

**STUDY CHECK**

Calculate the number of protons, neutrons and electrons of the following atoms:

- (a)  ${}^{32}_{16}\text{S}$  (b)  ${}^{34}_{16}\text{S}$  (c)  ${}^{36}_{16}\text{S}$

**Isotopes** All atoms of the same element are not the same. Some are heavier than others. Isotopes are atoms of the same element with different numbers of neutrons and therefore with different mass number but with the same atomic number. For example:  ${}^{24}_{12}\text{Mg}$ ,  ${}^{25}_{12}\text{Mg}$  and  ${}^{26}_{12}\text{Mg}$  are three isotopes of Mg.  ${}^{27}_{12}\text{Mg}$  is heavier than  ${}^{24}_{12}\text{Mg}$  as it contains more neutrons and protons in the nucleus. Each of the isotopes has a specific abundance, as some isotopes are more abundant than others. For example, the abundance of  ${}^{24}_{12}\text{Mg}$  is 79%, and the abundance of  ${}^{25}_{12}\text{Mg}$  and  ${}^{26}_{12}\text{Mg}$  is 10% and 11%, respectively. This means,  ${}^{24}_{12}\text{Mg}$  is more abundant than for example  ${}^{26}_{12}\text{Mg}$ .

**Average atomic mass** The average atomic mass represents the mass of the atoms of an element and results from all existing isotopes taking into account their abundance. The units of atomic mass are called *amu*, which stands for atomic mass units. This value can be simply found at any periodic table. Using the periodic table provided in this manual Figure 1.1, you can find the atomic mass of each element on top of the symbol at the right side. For example, the atomic mass of oxygen (O) is 15.999 amu and the atomic mass of nitrogen (N) is 14.007 amu. As atoms are made of numerous isotopes—this means different atoms of the same element but with different number of neutrons and hence different weight—the atomic mass found in the periodic table is an average that results from including the mass of the different isotopes and their abundance. That is you need to do an average of the mass of each isotope using values of abundance. In other words: the *average atomic mass* of an element—herein called simply atomic mass—, expressed in *amu* (atomic mass units), is the weighted average of the masses of the individual isotopes of the element. For an element with  $n$  isotopes with different masses ( $A_1, A_2, \dots, A_n$ ) and different fractional abundances for each isotope ( $f_1, f_2, \dots, f_n$ ), the atomic mass is given by

$$\text{Atomic mass} = \sum_{i=1}^n A_i \cdot f_i = A_1 \cdot f_1 + A_2 \cdot f_2 + \cdots + A_n \cdot f_n$$

Also, note that when adding the fractional abundances of all isotopes, one should obtain a value of one:

$$\sum_{i=1}^n f_i = f_1 + f_2 + \cdots + f_n = 1$$

## Sample Problem 4

Naturally occurring copper (Cu) consists of 69.17%  ${}^{63}\text{Cu}$  and 30.83%  ${}^{65}\text{Cu}$ . The mass of  ${}^{63}\text{Cu}$  is 62.939598 amu, and the mass of  ${}^{65}\text{Cu}$  is 64.927793 amu.



What is the atomic mass of copper?

### SOLUTION

The weighted average is the sum of the mass of each isotope times its fractional abundance. We have that the isotope  $^{63}\text{Cu}$  has a mass of 62.939598 amu and an abundance of 69.17%, that is the same as 0.6917. At the same time, the isotope  $^{65}\text{Cu}$  has a mass of 64.927793 amu and an abundance of 0.3083. After adding both contributions, we have:

$$62.939598 \text{ amu} \times \frac{69.17}{100} + 64.927793 \text{ amu} \times \frac{30.83}{100} = 63.55 \text{ amu}$$

### STUDY CHECK

Lithium is made up of two isotopes, Li-7 (7.016003 amu) and Li-6 (6.015121 amu). Calculate the percent abundance of each isotope knowing that lithium's atomic weight is 6.94 amu.

**Table 1.1 Isotope abundance of some elements**

Element	Isotope	% Abundance	Element	Isotope	% Abundance
Hydrogen	$^1\text{H}$	99.9885%	Silicon	$^{28}\text{Si}$	92.2297%
	$^2\text{H}$	0.0115%		$^{29}\text{Si}$	4.6832%
Helium	$^3\text{He}$	0.000137%		$^{30}\text{Si}$	3.0872%
	$^4\text{He}$	99.999863%	Sulfur	$^{32}\text{S}$	94.93%
Lithium	$^6\text{Li}$	7.59%		$^{33}\text{S}$	0.76%
	$^7\text{Li}$	92.41%		$^{34}\text{S}$	4.29%
Boron	$^{10}\text{B}$	19.9%	Chlorine	$^{36}\text{S}$	0.02%
	$^{11}\text{B}$	80.1%		$^{35}\text{Cl}$	75.78%
Carbon	$^{12}\text{C}$	98.93%		$^{37}\text{Cl}$	24.22%
	$^{13}\text{C}$	1.07%	Argon	$^{36}\text{Ar}$	0.3365%
Nitrogen	$^{14}\text{N}$	99.632%		$^{38}\text{Ar}$	0.0632%
	$^{15}\text{N}$	0.368%		$^{40}\text{Ar}$	99.6003%
Oxygen	$^{16}\text{N}$	99.757%	Potassium	$^{39}\text{K}$	93.2581%
	$^{17}\text{O}$	0.038%		$^{40}\text{K}$	0.0117%
	$^{16}\text{O}$	0.205%		$^{41}\text{K}$	6.7302%

## 3.4 An introduction to molecules

The periodic table contains all elements in nature. At the same time, elements combine to form molecules. For example, in the air there are traces of Argon—this is an element—and also water, a molecule ( $\text{H}_2\text{O}$ ) that results from the combination of two elements, hydrogen (H) and oxygen (O). This section will first introduce some of the properties of molecules, without paying attention to their chemical names that will be covered in the following chapters.

*Molecular weight* Here are two examples of molecules: molecular oxygen  $\text{O}_2$  and molecular nitrogen  $\text{N}_2$ . How do we interpret these formulas? The subscript "2" indicates that each molecule contains two atoms. For example, a  $\text{O}_2$  molecule is made of two oxygen atoms O. At the same time, the weight of a set of molecules is called the molecular weight (MW). However, you will find different terms to refer to the same property such as: molecular mass, molar mass, or formula unit mass. All these terms



indeed mean the weight a large set of molecules. We can calculate the MW by adding the weight of each atom that form the molecule taking into the account the number of atoms of each element present in the molecule. The units of molecular weight are the same as the units of atomic weight: amu, atomic mass units.

#### Sample Problem 5

Calculate: (a) The atomic weight of O; (b) the molecular mass of molecular oxygen, O<sub>2</sub>

#### SOLUTION

- (a) According to the periodic table the atomic weight (AW) of O is 15.999 amu.  
 (b) The molar mass of O<sub>2</sub> is the result of adding the atomic masses of 2O atoms, that is 31.998 amu, close to 32 amu.

#### STUDY CHECK

Calculate the molar mass of water H<sub>2</sub>O and ammonia, NH<sub>3</sub>

*Mass percent composition of a compound* Look at these two molecules: C<sub>2</sub>H<sub>2</sub> and C<sub>2</sub>H<sub>6</sub>. They contain different amounts of hydrogen. We quantify the amount of an element in a compound by means of the mass % composition. The mass % of an element in a compound is the mass of the element with respect to the molecular weight of the molecule in percent form. Mind that you have to take into account the molecular indexes in the compound as C<sub>2</sub>H<sub>2</sub> is made of 2H and C<sub>2</sub>H<sub>6</sub> is made of 6H. For example, given that the molar mass of C<sub>2</sub>H<sub>2</sub> is 26 amu, the mass % of hydrogen in C<sub>2</sub>H<sub>2</sub> would be:

$$\%_H \text{ in C}_2\text{H}_2 = \frac{2 \cdot AW(H)}{MW(\text{C}_2\text{H}_2)} \times 100 = \frac{2 \cdot 1}{26} \times 100 = 7.7\%$$

Similarly, the mass % of C would be:

$$\%_C \text{ in C}_2\text{H}_2 = \frac{2 \cdot AW(C)}{MW(\text{C}_2\text{H}_2)} \times 100 = \frac{2 \cdot 12}{26} \times 100 = 92.3\%$$

By adding the mass % of all elements in a molecule we should obtain 100.

$$\%_H \text{ in C}_2\text{H}_2 + \%_C \text{ in C}_2\text{H}_2 = 100$$

#### Sample Problem 6

Calculate the mass % composition for each element of glucose, C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>.

#### SOLUTION

We first need the molecular weight of glucose, C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>, that is: 6 · 12 + 12 · 1 + 6 · 16 = 180 amu. Now we can calculate the mass percent of carbon, hydrogen and oxygen:

$$\%_C \text{ in C}_6\text{H}_{12}\text{O}_6 = \frac{6 \cdot 12}{180} \times 100 = 40\%$$

$$\%_H \text{ in C}_6\text{H}_{12}\text{O}_6 = \frac{12 \cdot 1}{180} \times 100 = 6.6\%$$

By subtraction, we have that %<sub>C</sub> in C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> = 53.4.

#### STUDY CHECK

Urea CO(NH<sub>2</sub>)<sub>2</sub> is a colorless crystalline compound excreted in urine, product of protein metabolism in mammals. Calculate the mass % composition for each element of urea.

### 3.5 Empirical and molecular formula of a chemical

There are two different types of formulas: molecular formulas and empirical formulas. Empirical formula (EFs) are simplified formulas resulting from an experiment, whereas molecular formulas (MFs) are exact formulas of molecules. For example: the molecular formula of hydrogen peroxide, a mild antiseptic used on the skin to prevent infection of minor cuts, is  $\text{H}_2\text{O}_2$  as the hydrogen peroxide molecule is made of two oxygen and two hydrogen atoms. Differently, the empirical formula of the same chemical is  $\text{HO}$ , being this the result of the simplification of  $\text{H}_2\text{O}_2$ . One can obtain empirical formulas simply by dividing the molecular formula by the smallest integer number, of course, given you know the molecular formula. The word empirical means "from an experiment", and the use of empirical formulas comes from the fact that the formulas of all chemicals actually come from experiments, and from experiments one normally can only obtain ratios of atomic composition.

#### Sample Problem 7

From the following formulas identify the empirical and molecular formulas:  $\text{P}_4\text{O}_{10}$ ,  $\text{C}_3\text{H}_6\text{O}$ ,  $\text{N}_2\text{O}_4$  and  $\text{C}_5\text{H}_{11}$ .

#### **SOLUTION**

Empirical formulas are simplified versions of molecular formulas. For example,  $\text{C}_3\text{H}_6\text{O}$  and  $\text{C}_5\text{H}_{11}$  are empirical formulas. Differently,  $\text{P}_4\text{O}_{10}$  and  $\text{N}_2\text{O}_4$  are molecular formulas.

#### **STUDY CHECK**

Given the following molecular formulas, obtain the corresponding empirical formula:  $\text{P}_4\text{O}_{10}$ ,  $\text{N}_2\text{O}_4$  and  $\text{C}_6\text{H}_{18}\text{O}_3$ .

#### *Molecular weight of empirical formulas and molecular formulas*

The molecular weight of an empirical formula and its corresponding molecular formula are related by the following formula:

$$n = \frac{MW_{MF}}{MW_{EF}}$$

where:

$MW_{EF}$  is the molecular weight of the empirical formula

$MW_{MF}$  is the molecular weight of the molecular formula

$n$  is a integer number such as 1, 2, 3...

Understanding the formula above is simple. On one hand, the MW of a molecular formula  $\text{H}_2\text{O}_2$  that is 34 amu. On the other hand, the molecular weight of the empirical formula of the same chemical  $\text{HO}$  is 17 amu. If we do  $34/17$  we would obtain 2, as we need to multiply  $\text{HO}$  by two in order to obtain  $\text{H}_2\text{O}_2$ . As a final note, mind that empirical formulas are just simplified formulas. So when we think about the molecular weight of a chemical we normally have molecular formula in mind. Let us work on an example:

## Sample Problem 8

Given that the empirical formula of dichloromethane is  $\text{ClCH}_2$  and the molecular weight of the chemical is 98 amu, calculate the molecular formula of dichloromethane.

**SOLUTION**

Given the empirical formula of dichloromethane one can think of many different molecular formulas, for example:  $\text{Cl}_3\text{C}_3\text{H}_6$  or  $\text{Cl}_2\text{C}_2\text{H}_4$ . From these, and many other, there is only one real molecular formula. How do we calculate the real molecular formula? By comparing the MW of the molecular and empirical formula we can figure out the number of times we need to multiply the MF to obtain the EF. We know the MW is 98 amu. Using the EF we can also calculate a MW:  $35 + 12 + 2 \cdot 1 = 49$  amu. If we compare both numbers using the formula:

$$n = \frac{MW_{MF}}{MW_{EF}}$$

we have:  $n = 98/49$  and solving we have  $n = 2$ . Therefore the MF is:  $\text{Cl}_2\text{C}_2\text{H}_4$ .

**STUDY CHECK**

The empirical formula of dinitrogen tetroxide, a red-brown liquid with an unpleasant chemical odor, is  $\text{NO}_2$  and the molecular weight of the chemical is 92 amu. Calculate the molecular formula of dinitrogen tetroxide.

### 3.6 Determining empirical formulas

We said that the formula of a chemical that takes into account the correct number of atoms in a molecule is the molecular formula and therefore the real molecular weight of a chemical comes from these formulas. Empirical formulas are obtained from experiments in which a chemical is fragmented and analyzed so that the elements in the molecule and the mass percentage of each element is determined. Molecular formulas are obtained by using the molecular weight of the chemical and the empirical formula. Mind that the formula of a chemical that takes into account the correct number of atoms in a molecule is the molecular formula and therefore the real molecular weight of a chemical comes from these formulas. Let us work on an example in order to learn the procedure of obtaining molecular formulas.

*Calculating molecular formulas* By means of an experiment, we want to calculate the empirical formula of a chemical given that the chemical contains 2.8 g of nitrogen and 6.4 g of oxygen. In order to calculate the EF we will set up a table like the one presented below.

Empirical Formula Calculation		
	N	O
Grams	2.8g	6.4g
AW	14	16
Grams/AW	0.2	0.4
∇• by smallest	1	2
Formula	$\text{N}_1\text{O}_2=\text{NO}_2$	

In each column we will add each of the elements that form the molecule. In the first row we will include the grams of each element (sometimes this information is given in terms of mass %), in the second we will divide the grams of each element by its atomic weight ( $\text{AW}(\text{N})=14\text{amu}$ ,  $\text{AW}(\text{O})=16\text{amu}$ ). Among all numbers of the second row (in this example 0.2 and 0.4), we will select the smallest number (0.2). Once we have the smallest, we will divide all numbers by the smallest and that will give us round numbers (1 and 2); these will be the numbers in an empirical formula:  $\text{NO}_2$ .

#### Sample Problem 9

The mass percentage composition of a compound is: 18.59% O, 37.25% S, and 44.16% F. Calculate its empirical formula.

#### **SOLUTION**

We will set up the the molecular formula table, knowing that the percentage are mass percentages, that is the mass of each element in the chemical, hence they should go in the grams row. Also the atomic weights of O, S and F are 16, 32 and 19 amu.

Empirical Formula Calculation			
	O	S	F
Grams	0.1859g	0.3725g	0.4416g
AW	16	32	19
Grams/AW	0.0116	0.0116	0.0232
∇• by smallest	1	1	2
Formula	$\text{OSF}_2$		

#### **STUDY CHECK**

What is the empirical formula of a compound if a sample contains 10.28 g of C, 1.71 H and 12.71 g of oxygen?