

© PxFuel

Ch. 8. Reactions in solution

HE most common reactions happen in solution and involve cations and anions. Think, for example, when you add sugar to your coffee or how metal rusts when it gets wet. The first example involves a dissolution reaction, whereas the second is a redox reaction in which electrons are exchanged. Overall there are three main different types of reactions happening in solutions. It is not only important to be able to differentiate these reactions, but it is also important to understand the nuances between acid-base reactions, precipitation reactions, and redox reactions. It is also critical to understand the properties of solutions and learn how to quantify and compare the amount of solute in a solution. On one hand, most of you will be surprised to know that water does not conduct electricity. This is because pure water is a weak electrolyte. On the other hand, the importance of electrolytes is well-known among the sports community. If you have ever played a sport, you have probably chugged a sports drink. These are electrolyte solutions. However, few know the specifics of their function. Electrolytes are salts that conduct electricity in water by separating them into positive and negative ions. To understand the properties of solutions it is important to identify and differentiate the different types of electrolytes.

8.1 Solutions and composition

Solutions are homogeneous mixtures of two components. The state of the matter of both components of the mixture or their polarity affects the formation of a solution. For example, a solution will not result from mixing oil and water as they have different polar characteristics and it will form from mixing table salt and water as both are polar chemicals. At the same time, the more solute you add to a solution the more concentrated the solution will be. This section covers polarity and the composition of solutions.

What makes a solution? Solutions are homogeneous mixtures of a solute and a solvent. Homogeneous means that if you look at the mixture you will not be able to differentiate both components and you will only see it as a whole. In a solution, the solute is the component of the mixture in less amount, whereas the solvent is the component in a larger amount. Think about mixing sugar with water. Sugar is sweet and water tasteless. When you mix both, you form a solution of sugar (solute) in water (solvent) and you will not see sugar in the solution as it is dissolved. In this particular example, sugar will be the solute in the solution, as the sugar is in less amount than water. Is important to remember that a solution is a result or mixing a solute and a solvent:

Solution = Solute + Solvent

Sample Problem

Identify the solute and the solvent in the following mixtures: (a) 10g of $H_2O(1)$ mixed with 3g of KCl(s); (b) 10g Cu(s) mixed with 3g Zn(s)

SOLUTION

(a) Potassium chloride is in less amount and hence will be the solute. Water, in larger quantity, will be the solvent. This is an example of an aquose-based solution. (b) Zn is in less amount and hence will be the solute, and Cu the solvent. This is a solid solution.

STUDY CHECK

Identify solute and solvent in the following mixture: (a) 10g of $H_2O(l)$ and 20g of $CH_3OH(l)$; (b) 1g of $I_2(s)$ and 100g of $CH_3CH_2OH(l)$

Types of solutions You can prepare different types of solutions by mixing a solid and a liquid, like when you mix sugar and water, or salt and water. You can create solutions as well by mixing two liquids or two solids. Examples are vinegar—a liquid solution of acetic acid (liquid) in water (liquid)—or steel—a solid solution that contains iron and carbon, both solids.




Figure 8.1 A solution results from dissolving a solute into a solvent

Empirical rules of polarity The affinity between two chemicals is related to a concept called polarity. Molecules contain electrons and depending on the electron distribution within the molecules, molecules can be polar or non-polar. Molecules with an even electron distribution are non-polar as they have no permanent dipole moment. An example of this is H₂ molecule. Differently, HF is a polar molecule, as F concentrates more on the electron density of the molecule than H. The polar nature of substances—with a permanent dipole moment—is related to miscibility and molecules with similar polar character will mingle and mix creating a single visible phase. As an example, water (H₂O, polar) and methanol (CH₃OH, polar) will mix together. Differently, water (polar) and oil (non-polar) are immiscible due to their different polar nature and they will not mix. Even if the rules or polarity are based on

the nature and structure of the molecule, one can use very simple empirical rules to classify molecules as polar or non-polar. These rules work in general well for the case of diatomic and very large molecules:

- Diatomic molecules made of the same element (e.g. H₂) are non-polar.
- Diatomic molecules made of different elements (e.g. HI) are polar.
- Poliatomic molecules (with more than four atoms) made of C and H (e.g. CH₄) are in general non-polar.
- Poliatomic molecules (with more than four atoms) containing C, H, and a different atom (e.g. CH₃F) are in general polar.

Sample Problem 2

Classify the following molecules as polar or nonpolar: H₂, HCl, CH₃CH₃, and CH₃CH₂Cl.

SOLUTION

H₂ is a non-polar molecule, being a diatomic molecule containing two atoms of the same element. Differently HCl is polar. CH₃CH₃ is a non-polar poliatomic molecule made of C and H atoms, whereas CH₃CH₂Cl is polar.

STUDY CHECK

Classify the following molecules as polar or nonpolar: HF, Cl_2 , C_2H_4 , and $\text{C}_2\text{H}_3\text{Cl}$.

Mixing and polarity A solution is formed when both the solute and the solvent mix. However, they will only mix if they have the same polarity. As an example, water (H_2O) is a polar molecule, and methanol (CH_3-OH) is too. Hence they will both mix and form a solution. If the elements of a mixture have different polarity they will not mix. An example is benzene $(C_6H_6$, nonpolar) and water, or for example oil (nonpolar) and water (polar).

Sample Problem 3

Use polarity arguments to indicate if the following substances will mix: (a) $H_2O_{(g)}$ and $CH_{4(g)}$; (b) $H_2O_{(g)}$ and $HCl_{(g)}$

SOLUTION

(a) Water and methane (CH_4) will not mix, as water is a polar molecule and CH_4 (methane) is nonpolar. (b) They will mix as HCl is a polar molecule and so is water.

STUDY CHECK

Use polarity arguments to indicate if the following substances will mix: (a) $H_2O_{(l)}$ and $CH_3Cl_{(l)}$; (b) $CH_3Cl_{(l)}$ and $CCl_{4(l)}$

Concentration of solutions The concentration of a solution refers to the amount of solute with respect to the amount of solution. The larger concentration the larger the number of solute particles with respect to the particles of solvent. Concentration is one of the most important properties of a solution as it affects the physical properties of a solution such as the freezing and boiling point. There are many different concentration units, such as molarity, mass percent concentration, or volume percent concentration. All these different units overall express the ratio between the particles—mass or volume—of solute and solvent.

Table 8.1 Polarity and mixing			
Solvent	Solute	Mixing?	
Polar	Polar	Yes	
Polar	Nonpolar	No	
Nonpolar	polar	No	
Nonpolar	Nonpolar	Yes	

Mass percent concentration The mass percent (m/m) is the amount of solute in grams per gram of solution in percent form

$$\left[m/m = \frac{\text{g of solute}}{\text{g of solution}} \times 100 \right]$$

Sample Problem 4

A NaCl solution is prepared by mixing 4g of NaCl with 50g of H_2O . Calculate the percent (m/m) of the solution.

SOLUTION

We need the grams of solute and the grams of solution. The grams of solute are given (4g of NaCl), whereas the grams of solution result from adding the grams of solvent and solute: 54 g of solution. Using the formula for the percent (m/m), he have:

$$m/m = \frac{4 \text{ g of solute}}{54 \text{ g of solution}} \times 100 = 7.4\%$$

This means that by mixing 4g of NaCl with 50g of H_2O you prepare a 7.4% (m/m) solution.

STUDY CHECK

A KCl solution is prepared by mixing 5g of KCl with 200g of H_2O . Calculate the percent (m/m) of the solution.

Volume percent concentration The volume percent concentration (v/v) is the volume of solute per volume of solution in percent form

$$v/v = \frac{\text{volume of solute}}{\text{volume of solution}} \times 100$$

Mass/volume percent concentration The mass/volume percent concentration (m/v) is the mass of solute per mL of solution in percent form.

$$\left[m/v = \frac{\text{g of solute}}{\text{mL of solution}} \times 100 \right]$$

 ${\it Molarity concentration}$ The molarity (M) is the moles of solute per L of solution.

$$M = \frac{\text{moles of solute}}{\text{L of solution}}$$

Sample Problem 5

A NaCl solution is prepared by mixing 4g of NaCl (MW=58.4g/mol) with 50 g of water until a final volume of 52mL of solution. Calculate: (a) the mass percent (m/m) concentration; (b) the molarity.

SOLUTION

(a) to calculate the mass percent (m/m) we just need the grams of solute and the grams of solution—that is four plus fifty. Both numbers are already given:

$$m/m = \frac{\text{g of solute}}{\text{g of solution}} \times 100 = \frac{4 \text{ g of solute}}{54 \text{ g of solution}} \times 100 = 9.2\%$$

(b) To calculate molarity we need the moles of solute and the liters of solution.

$$M = \frac{\text{moles of solute}}{\text{L of solution}} = \frac{0.068 \text{ moles of solute}}{5.2 \times 10^{-2} \text{L of solution}} = 1.31 M$$

STUDY CHECK

(a) A solution is prepared by mixing 8g of NaCl (MW=74g/mol) with 250mL of H₂O. Calculate the molarity; (b) A KCl solution is prepared by mixing 45g of KCl with 200g of H₂O. Calculate the percent (m/m) of the solution.

Concentration units as conversion factors Each of the different concentration units—molarity, mass percent, volume percent, mass/volume percent—can be used in a conversion factor form. For example, if the molarity of a solution is 3M, this means that in the solution there are 3 moles of solute in every liter of solution.

Similarly, if the mass percent of a solution is 5% this means that there are 5 grams of solute for every 100 grams of solution. We often use concentration units as conversion factors when we need to transform between one unit on top (bottom) of the conversion factor and the unit on the bottom (top).

Sample Problem 6

How much volume of a 4M solution do you need to provide 5 moles of solute.

We will use the conversion factor of Molarity using the volume on top and the moles on the bottom in order to cancel the units:

$$5\underline{\text{moles of solute}} \times \frac{1 \text{ L of solution}}{4\underline{\text{moles of solute}}} = 1.25 \text{L}$$

This means that 1.25L of a 4M solution will provide 5 moles of solute.

STUDY CHECK

How many grams of a 6% (m/m) solution do you need to provide 5 grams of solute.

Dilution Dilution is the process of preparing a diluted solution from a more concentrated solution. Solutions are often stored in a stock room in concentrated form. These stocks should be diluted before use. To dilute a solution we need to take a certain amount of the concentrated solution and add water. When adding water, the number of moles of solute does not change, and the concentration always decreases. We have a concentrated solution (c_1) and we need to prepare a certain volume (V_2) of a more diluted solution (c_2) . The question is how much volume of the concentrated solution (V_1) we need to take. To answer this we should use the following formula:

$$\boxed{c_1 \cdot V_1 = c_2 \cdot V_2}$$

Sample Problem 7

How many liters of a 3M NaCl solution are required to prepare 2L of a 1M diluted NaCl solution.

SOLUTION

We have a concentrated solution of 3M molarity and we want to prepare a more dilute solution. In particular 2L of a 1M. Hence: $c_1=3$ and $c_2=1M$ and $V_2=2L$. Using the dilution formula:

$$3M \cdot V_1 = 1M \cdot 2L$$

Solving for V_1 we have a volume of 0.66L.

STUDY CHECK

How many liters of a 5M NaCl solution are required to prepare 3L of a 3M diluted NaCl solution.

Molarity calculation involving grams As molarity is just the ratio between the moles of solute and the litters of the solution, we can use this property to calculate the mass of solute contained in a certain volume of solution. We will proceed by first calculating the number of moles in that volume to then convert moles into grams.

8.2 Electrolytes and insoluble compounds

On one hand, electrolytes are compounds that conduct electricity once dissolved in water. Differently, nonelectrolytes are compounds that do not conduct electricity once dissolved in water. On the other hand, insoluble compounds are not soluble in water, whereas soluble compounds can be dissolved in water. This section covers the properties of electrolytes and insoluble (and soluble) compounds. At the end of this section, you should be able to classify a chemical in terms of its electrolyte type and solubility character.

Solubility formula Solubility (s) is the grams of a solute per 100 g of solvent:

$$s = \frac{\text{g of solute}}{100 \text{ g of solvent}}$$

A saturated solution can be achieved when you fit the maximum amount of solute in the solvent. If you continue adding solute to a saturated solution it will precipitate and a solid will form.

Soluble and insoluble salts Soluble compounds dissolve in water, whereas insoluble compounds do not. For example, barium chromate $(BaCrO_{4(s)})$ is an insoluble salt. How do we know that? The Table below will help you predict the solubility of a salt. To do this, you need to start by assessing the right ion (the anion, CrO_4^{2-}) located in the left column of the Table below. After that, you need to assess the left ion (the cation, Ba^{2+}) located in the right column. If you follow this, you will see that chromate is insoluble and barium is not part of any exception. Let us predict for example the soluble/insoluble nature of $CaSO_4$, calcium sulfate. We start by looking for SO_4^{2-} in the left column to find out is soluble. Next, we continue in the same line as SO_4^{2-} and look for the ion in the left Ca^{2+} . In conclusion, even when SO_4^{2-} is soluble,

when combined with Ca²⁺, we have that CaSO₄ is insoluble, and overall CaSO_{4(s)} is insoluble.

Predict the soluble/insoluble nature of the following compounds: (a) K₂CO₃,

(b) NaNO₃ and (c) Ca(OH)₂.

SOLUTION

(a) K₂CO_{3(aq)} is soluble, as CO₃⁻² is insoluble but when combined with K⁺ the salt becomes soluble. (b) All nitrates are soluble without exceptions. (c) $Ca(OH)_{2(aq)}$ is soluble.

STUDY CHECK

Predict the soluble/insoluble nature of the following compounds: (a) Li₃PO₄

(b) Na₂S (c) AgCl

Table 8.2 Soluble and insoluble compounds			
Ions that form soluble compounds	except when combined with		
Group I ions (Na ⁺ , Li ⁺ , K ⁺ , etc) Ammonium (NH ₄ ⁺) Nitrate (NO ₃ ⁻) Acetate (CH ₃ COO ⁻) Hydrogen carbonate (HCO ₃ ⁻) Chlorate (ClO ₃ ⁻) Halide (F ⁻ , Cl ⁻ , Br ⁻) Sulfate (SO ₄ ²⁻)	no exceptions no exceptions no exceptions no exceptions no exceptions no exceptions Pb ²⁺ , Ag ⁺ and Hg ₂ ²⁺ Ag ⁺ , Ca ²⁺ , Sr ²⁺ , Ba ²⁺ , Hg ₂ ²⁺ and Pb ²⁺		
Ions that form <i>insoluble</i> compounds	except when combined with		
Carbonates (CO_3^{2-}) Chromates (CrO_4^{2-}) Phosphates (PO_4^{3-}) Sulfides (S^{2-}) Hydroxides (OH^-)	group I ions (Na ⁺ , Li ⁺ , K ⁺ , etc) or ammonium (NH ₄ ⁺) group I ions (Na ⁺ , Li ⁺ , K ⁺ , etc) or Ca ²⁺ , Mg ²⁺ or ammonium (NH ₄ ⁺) group I ions (Na ⁺ , Li ⁺ , K ⁺ , etc) or ammonium (NH ₄ ⁺) group I ions (Na ⁺ , Li ⁺ , K ⁺ , etc) or Ca ²⁺ , Mg ²⁺ , Sr ²⁺ or ammonium (NH ₄ ⁺)		

Strong electrolytes Strong electrolytes completely dissociate in water. Hence, in a solution of a strong electrolyte, you will only have ions and never molecules. Strong electrolytes are typically ionic compounds such as MgCl₂ or NaCl (table salt). We represent the dissociation of a strong electrolyte with a single arrow, meaning that the reaction proceeds to completion, and for the example below, in the solution, we will only have ions $(Mg^{2+}_{(aq)} + 2 Cl^{-}_{(aq)})$ and not molecules $(MgCl_{2(s)})$:

$$MgCl_{2(s)} \xrightarrow{H_2O} Mg^{2+}_{(aq)} + 2\,Cl^-_{(aq)}.$$

Weak electrolytes Weak electrolytes partially dissociate in water, and this is indicated using a chemical reaction with a double arrow. Hence in a solution of a weak electrolyte, you will have ions as well as molecules at the same time. Examples of weak electrolytes are hydrofluoric acid, water, ammonia, or acetic acid. The dissociation of hydrochloric acid (HF) proceeds as:

$$HF_{(l)} \stackrel{H_2O}{\longleftrightarrow} H_{(aq)}^+ + F_{(aq)}^-$$

Acetic acid (CH₃COOH) is an important weak electrolyte and its dissociation proceeds somehow in a peculiar way:

$$CH_3COOH_{(l)} \stackrel{H_2O}{\longleftarrow} CH_3COO_{(aq)}^- + H_{(aq)}^+$$

Nonelectrolytes Nonelectrolytes do not dissociate in water. Hence a solution of a nonelectrolyte will only contain molecules and not ions. Examples of nonelectrolytes are carbon-based chemicals such as methanol, ethanol, urea, or sucrose. The dissociation of urea for example CH_4N_2O proceeds as:

$$CH_4N_2O_{(s)} \xrightarrow{H_2O} CH_4N_2O_{(aq)}$$

Identify the electrolyte character of a chemical You can use the Table below to identify the electrolyte character of a chemical. Ionic compounds are in general strong electrolytes, and most acids are as well. There are four important weak electrolytes: water, acetic acid, ammonia, and hydrofluoric acid. Covalent compounds are in general nonelectrolytes. Organic compounds, compounds based on carbon atoms (e.g. $C_{12}H_{22}O_{11}$) are in general nonelectrolytes.

Table 8.3 Different types of electrolytess			
Electrolyte Type	Dissociation	Particles in solution	Examples
Strong	Fully	Mostly ions	Ionic Compounds and most acids and bases (hydroxides): NaCl, NaOH,
			HCl, MgCl ₂ , H ₂ SO ₄ , etc
Weak	Partially	Ions & molecules	NH ₃ , CH ₃ COOH (acetic acid), HF, H ₂ O
Nonelectrolytes	No	molecules	Most covalent compounds: CH ₃ OH (methanol),
			CH ₃ CH ₂ OH (ethanol), C ₁₂ H ₂₂ O ₁₁ (sucrose), CH ₄ NO ₂ (urea)

Breaking down electrolytes into ions Electrolytes—in particular strong electrolytes—dissociate producing ions. This way, a solution of for example NaCl does not contain NaCl molecules but $\mathrm{Na}^+_{(\mathrm{aq})}$ cations and $\mathrm{Cl}^-_{(\mathrm{aq})}$ anions. Hence it is important to correctly break down electrolytes into ions. To do this, you need to revert the combination of ions that produce a given chemical while making sure the charges are balanced. For example, let us beak magnesium chloride $\mathrm{MgCl}_{2(\mathrm{aq})}$ into ions. This is a strong electrolyte formed by magnesium cations and chloride anions. The valence of magnesium is +II and the valence of chlorine is -I. The MgCl_2 formula also tells us we have one magnesium and two chlorines. The overall process is:

$$MgCl_{2(aq)} \longrightarrow Mg^{2+}_{(aq)} + 2Cl^{-}_{(aq)}$$

Another example, magnesium nitrate $Mg(NO_3)_2$. This strong electrolyte—as this is an ionic salt—is made of magnesium with valence +II and nitrate with valence -I. The formula indicated we have one $Mg_{(aq)}^{2+}$ and two $NO_3^{-}_{(aq)}$. Hence:

$$Mg(NO_3)_{2(aq)} \longrightarrow Mg^{2+}_{(aq)} + 2NO_3^{-}_{(aq)}$$

Sample Problem 9

For the following chemicals indicate whether you will have in the solution (a) only ions, (b) ions and some molecules, or (c) molecules: NH_3 , KOH, and $C_{12}H_{22}O_{11}$.

SOLUTION

Ammonia (NH₃) is a weak electrolyte and a solution of ammonia will contain ions and well as ammonia molecules. Differently KOH is a strong electrolyte

and in solution you would find only ions (K^+ and OH^-). Sucrose ($C_{12}H_{22}O_{11}$) is a nonelectrolyte and in solution you will find molecules.

♥ STUDY CHECK

For the following chemicals indicate whether you will have in the solution only ions, ions and some molecules, or molecules: (a) H₂SO₄, HNO₃ and CH₃OH.

Sample Problem 10

Break down the following chemicals into ions, if possible:

Chemical	Particles in solution
$K_2CrO_{4(aq)}$	
$Ba(NO_3)_{2(aq)}$	
$BaCrO_{4(s)}$	
$KNO_{3(aq)}$	

SOLUTION

We can only break down into ions ionic compounds and oxosalts that are not solid. From the list of chemicals in the example, we will not be able to break down $BaCrO_{4(s)}$ into ions as it is a solid. From the other chemicals, $K_2CrO_{4(aq)}$ is named potassium chromate and contains $2K_{(aq)}^+$ and $CrO_4^{2-}_{(aq)}$ ions. Barium nitrate– $Ba(NO_3)_{2(aq)}$ –will produce $Ba_{(aq)}^{2+}$ and $2NO_3^-_{(aq)}$. Finally, potassium nitrate– $KNO_{3(aq)}$ –will produce $K_{(aq)}^+$ and $NO_3^-_{(aq)}$. In the table:

Chemical	Particles in solution
K ₂ CrO _{4(aq)}	$2K_{(aq)}^{+} + CrO_4^{2-}{}_{(aq)}$
$Ba(NO_3)_{2(aq)}$	$Ba_{(aq)}^{2+} + 2NO_3^{-}_{(aq)}$
$BaCrO_{4(s)}$	$BaCrO_{4(s)}$
$KNO_{3(aq)}$	$K_{(aq)}^+ + NO_3^{(aq)}$

STUDY CHECK

Break down the following chemicals into ions, if possible: $H_2O_{(l)}$, $NH_{3(l)}$, $AgNO_{3(aq)}$.

8.3 An introduction to reactions in solution

There are three different reactions in solution: acid-base reactions, precipitation reactions, and redox reactions. The key to identifying acid-base reactions is in the reactants, as an acid-base reaction results from the reaction between an acid and a base. Precipitation reactions are reactions that produce a precipitate. Hence, the key to identifying a precipitation reaction is in the products. Precipitation reactions always contain a solid as a product. Redox reactions contain two elements with different redox numbers in the reactants and products. The key to identifying redox reactions is to be able to spot elements with different oxidation states, for example: Cu and Cu^{2+} or H^+ and H_2 . In the following, we will describe more about the three different types of reactions in solution. The goal of this section is for you to be able to identify each type.

Acid-base reactions Acid-base reactions result from the reaction of an acid with a base. Both produce water and another chemical. An example is:

$$HBr_{(aq)} + KOH_{(aq)} \longrightarrow KBr_{(aq)} + H_2O_{(l)}$$
 (acid-base reaction)

Hydrobromic acid (HBr) is an acid and potassium hydroxide (KOH) is a base. The result of an acid-base reaction is always water and an ionic compound, in this case KBr.

Precipitation reactions Precipitation reactions result in an insoluble chemical, that is, results in a solid chemical. An example would be:

 $K_2CrO_{4(aq)} + Ba(NO_3)_{2(aq)} \longrightarrow BaCrO_{4(s)} \downarrow + 2 \, KNO_{3(aq)}$ (precipitation reaction) The chemical $BaCrO_{4(s)}$ is a solid that precipitates in the solution, hence the name of the type of reaction. The symbol on $BaCrO_{4(s)} \downarrow$ represents the precipitation process. The solubility of a given solute such as $BaCrO_{4(s)}$ is the amount of solute (in grams) that can be dissolved in a given mass of solvent (in particular 100 g of solvent). A solute with a low solubility will be hard to dissolve. Think about cacao and water. The solubility of cacao is low and hence by simply adding cacao powder to water, you will not be able to make a solution. However, solubility depends on the solute and solvent combination, but also on the temperature, and by warming up a solvent you can increase solubility and fit more solute in the same amount of solvent. This section covers different aspects of solubility.

Redox reactions Redox reactions are different than acid-base or precipitation reactions. They contain the same chemical element in two different states resulting from the loss or win of electrons. Look for the example:

$$2 \operatorname{Al}_{(s)} + 3 \operatorname{Cu}_{(aq)}^{+2} \longrightarrow 2 \operatorname{Al}_{(aq)}^{+3} + 3 \operatorname{Cu}_{(s)} (\text{redox reaction})$$

We have that neither $Al_{(s)}$ or $Cu_{(aq)}^{+2}$ are an acid or a base, therefore this is not an acid-base reaction. Also, there is no insoluble product, hence this is not a precipitation reaction. Indeed, this is a redox reaction, as we have Al in two different states: as metallic $Al_{(s)}$ and as ionic $Al_{(aq)}^{+3}$, which result from the loss of three electrons. Therefore in a redox reaction, there are always elements in the chemicals that lose electrons. In redox reactions, there is also an element that wins electrons. For example, $Cu_{(s)}$ and $Cu_{(aq)}^{+2}$ have different redox number. In particular, $Cu_{(aq)}^{+2}$ is the result of removing three electrons from $Cu_{(s)}$. At this point, we have that this reaction is redox as it contains an element that gains electrons and an element that loses electrons. Sometimes, the redox state of the elements is not that obvious. Look at this example:

$$Fe_{(s)} + CuSO_{4(aq)} \longrightarrow FeSO_{4(aq)} + Cu_{(s)}$$
 (redox reaction)

This is a redox reaction as you can find iron and copper in two states, metallic and also ionic. Therefore, these two metals have two different redox numbers in the reaction.

Sample Problem 11

Classify the following reactions as acid-base, redox or precipitation.

(a)
$$Fe_{(s)} + Cu_{(aq)}^{+2} \longrightarrow Fe_{(aq)}^{+2} + Cu_{(s)}$$

$$(b) \ AgNO_{3(aq)} + NaCl_{(aq)} \longrightarrow AgCl_{(s)} + NaNO_{3(aq)} \\$$

(c)
$$2 \text{HCl}_{(aq)} + \text{Ca}(\text{OH})_{2(aq)} \longrightarrow \text{CaCl}_{2(aq)} + 2 \text{H}_2\text{O}_{(l)}$$

SOLUTION

The first reaction is a redox reaction. This is because we can find two different oxidation states for Cu and also for Fe. That means one of these elements lost

electrons and the other won electrons. The second reaction is a precipitation reaction as it produces a solid. The last reaction is an acid base, as the reactants are an acid and a base.

STUDY CHECK

Classify the following reactions as acid-base, redox or precipitation.

- (a) $HNO_{2(aq)} + NaOH_{(aq)} \longrightarrow NaNO_{2(aq)} + H_2O_{(l)}$
- $\text{(b) } 2\,Na_{(s)} + Cl_{2(g)} \longrightarrow 2\,NaCl_{(s)}$
- (c) $MgCl_{2(aq)} + 2 AgNO_{3(aq)} \longrightarrow 2 AgCl_{(s)} + Mg(NO_3)_{2(aq)}$

CHAPTER 8

SOLUTIONS

- **8.1** A solution is prepared by mixing 4 g of $C_6H_{6(1)}$ and 10 g of $CCl_{4(1)}$. Indicate the true statement: (a) C_6H_6 is the solute (b) CCl₄ is the solute (c) Both chemicals do not mix (d) The mixture is not a solution
- **8.2** A solution is prepared by mixing 5 g of $Au_{(s)}$ and 2 g of $Cu_{(s)}$. Indicate the true statement: (a) Au is the solute (b) Cu is the solute (c) Both elements do not mix (d) The mixture is not a solution
- **8.3** Classify the following molecules as polar or nonpolar: (a) $C_6H_{14(1)}$ (b) $CH_3CH_2CH_2OH_{(1)}$ (c) $C_5H_{10(1)}$ (d) $C_6H_5CH_{3(1)}$ (e) $CH_3CH_2OH_{(1)}$ (f) $C_6H_5NH_{2(1)}$
- **8.4** Classify the following molecules as polar or nonpolar: (a) $H_2O_{(1)}$ (b) $C_5H_{12(1)}$ (c) $CH_3COOH_{(1)}$ (d) $CH_3OH_{(1)}$
- **8.5** Classify the following molecules as polar or nonpolar: (a) $H_2O_{(1)}$ (b) $C_5H_{12(1)}$ (c) $CH_3COOH_{(1)}$ (d) $CH_3OH_{(1)}$
- **8.6** Which chemicals from the following list will mix with $H_2O_{(1)}$: (a) $NH_{3(1)}$ (b) $C_5H_{12(1)}$ (c) $C_6H_{14(1)}$
- **8.7** Which chemicals from the following list will mix with $C_5H_{12(1)}$: (a) $H_2O_{(1)}$ (b) $C_6H_{14(1)}$ (c) $CH_3COOH_{(1)}$

ELECTROLYTES AND INSOLUBLE COMPOUNDS

8.8 Indicate whether solutions of the following chemicals will have ions (I), ions and molecules (I+M), or just molecules (M):

Chemical	I	I+M	M
NaCl			
HCl			
CaCl ₂			
H_2O			
NO ₂			

8.9 Indicate the soluble/insoluble character of the following compounds:

Chemical	Soluble	Insoluble
$AgNO_3$		
AgBr		
CaCO ₃		
Na_2CO_3		

8.10 Indicate the soluble/insoluble character of the following compounds:

Chemical	Soluble	Insoluble
NaCH ₃ COO		
NaHCO ₃		
Ag_2SO_4		
NaCrO ₄		
CaS		

CONCENTRATION UNITS

- **8.11** Order the following solution from more to less concentrated: (a) A vanillin solution $(1.3 \times 10^{-1} \text{M})$ (b) An adrenaline solution $(1.1 \times 10^3 \text{M})$ (c) Vinegar, an acetic acid solution $(1.2 \times 10^0 \text{M})$
- **8.12** Sodium hydroxide NaOH, a very strong base, is a chemical used in drain cleaners. A drain cleaning solution is prepared by mixing 25g of NaOH in 250g of water. Calculate the mass percent of solute.
- **8.13** A solution is prepared by mixing 15g of KCl in 50g of water. Calculate the mass percent of solute.
- **8.14** Alcohol-hydroxide is a mixture of a base with an organic alcohol, employed to clean glass. An alcoholhydroxide mixture is prepared by mixing 60g of NaOH with 500g of ethanol. Calculate: (a) the mass percent of solvent. (b) the mass percent of solute.
- **8.15** Vinegar is not a pure chemical, it is indeed a (m/m) 5% acetic acid solution. How many grams of acetic acid are there in 2g of vinegar.

- **8.16** Vanilla extract is a solution vanillin in ethanol. A vanilla solution is made by mixing 15 mL of pure vanillin and 50mL of ethanol.
- **8.17** An HCl solution is prepared by mixing 4 moles of HCl with water until reaching a volume of 250mL. Calculate the molarity of the solution.
- **8.18** How many mL of a 3M KCl solution contains 0.06 moles of KCl.
- **8.19** How many mL of a 4M NaCl (MW=58g/mol) solution contains 5 grams of NaCl.
- **8.20** How many grams of solute are there in 100mL of a 0.01M HNO₃ (MW=63g/mol) solution.
- **8.21** How many mL of a 0.001M Ca(OH)₂ (MW=74g/mol) solution can be prepared from 5 mg of Ca(OH)₂.
- **8.22** What is the final volume when 50mL of a 2M NaCl solution is diluted to a 1M.
- **8.23** What is the concentration of a solution prepared when 100mL a 4% HCl solution is diluted to a final volume of 500mL.
- **8.24** Describe how to prepare 50mL of a 0.5M H₂SO₄ solution, starting with a 1M stock H₂SO₄ solution.

PRECIPITATION AND ACID-BASE REACTIONS

- **8.25** Classify the following reaction as acid-base or precipitation:
- (a) $NaCl_{(aq)} + AgNO_{3}$ (aq) $\longrightarrow NaNO_{3(aq)} + AgCl_{(s)} \downarrow$
- (b) $H_2SO_{4(aq)} + 2 NaOH_{(aq)} \longrightarrow 2 H_2O_{(1)} + Na_2SO_{4(aq)}$
- $\begin{array}{ccc} \text{(c)} & 2\,Na_3PO_{4(aq)} + 3\,CaCl_{2(aq)} \longrightarrow \\ & 6\,NaCl_{(aq)} + Ca_3(PO_4)_{2(s)} \downarrow \end{array}$
- **8.26** Classify the following reaction as acid-base or precipitation:
- (a) $H_2SO_{4(aq)} + Ba(OH)_{2(aq)} \longrightarrow BaSO_{4(aq)} + 2 H_2O_{(l)}$
- $(b) \ \ Na_{2}CO_{3(aq)} + BaCl_{2(aq)} \longrightarrow BaCO_{3(s)} + 2\,NaCl_{(aq)}$
- (c) $4 \text{ Fe} + 3 \text{ O}_2 \longrightarrow 2 \text{ Fe}_2 \text{O}_3$

Answers 8.1 C_6H_6 is the solute **8.2** C_6H_6 is the solute **8.3** (a) $C_6H_{14(1)}$ (nonpolar) (b) $CH_3CH_2CH_2OH_{(1)}$ $(polar) \ \ (c) \ \ C_5H_{10(l)} \ \ (nonpolar) \ \ (d) \ \ C_6H_5CH_{3(l)} \ \ (nonpolar) \ \ (e) \ \ CH_3CH_2OH_{(l)} \ \ (polar) \ \ (f) \ \ C_6H_5NH_{2(l)} \ \ (polar)$ **8.4** (a) $H_2O_{(1)}$ (polar) (b) $C_5H_{12(1)}$ (nonpolar) (c) $CH_3COOH_{(1)}$ (polar) (d) $CH_3OH_{(1)}$ (polar) **8.5** (a) $H_2O_{(1)}$ (polar) (b) $C_5H_{12(l)}$ (nonpolar) (c) $CH_3COOH_{(l)}$ (polar) (d) $CH_3OH_{(l)}$ (polar) **8.6** $NH_{3(l)}$ (polar) **8.7** $C_6H_{14(l)}$ (nonpolar) **8.8** NaCl(I); HCl (I); CaCl₂ (I); H₂O (M+I); CO₂ (M) **8.9** AgNO₃(Soluble); AgBr (Insol $uble); \quad CaCO_3(Soluble); Na_2CO_3(Soluble) \\ \quad \textbf{8.10} \ NaCH_3COO(Soluble); \quad NaHCO_3(Soluble); \quad Ag_2SO_4(Insoluble); \\ \quad Ag_2SO_4(Insoluble)$ NaCrO₄(Soluble); CaS(Insoluble); **8.11** adrenaline winegar vanillin **8.12** 9% **8.13** 23% **8.14** (a) 89% (b) 11% 8.15 0.1 g 8.16 23% 8.17 16M 8.18 20mL 8.19 21mL 8.20 0.062g 8.21 67.56mL 8.22 100 mL 8.23 0.8 % **8.24** 25mL **8.25** (a) Precipitation (b) Acid-base (c) Precipitation **8.26** (a) Acid-base (b) Precipitation (c) redox