

1

The periodic table

MATTER is everywhere around you, from the water you drink to the air you inhale. Matter is made of elements and elements are made of atoms. Even the atoms of an element can be different, having a distinct number of protons and neutrons. This chapter covers the principles of atomic structure. You will learn what makes an atom and will be able to quantify the particles that make atoms. Perhaps more importantly, you will also learn about the periodic table of the elements and the different types of chemical formulas.

1.1 The periodic table

The periodic table is a chart containing all known elements arranged in increasing number of electrons per atom in a way that elements with similar chemical and physical properties are located together. The periodic table contains all existing elements—some of them are synthetic others are natural—that form the matter arranged in columns and rows. Every element has a different name accompanied by a symbol that represents their name. The tabular arrangement of elements in the form of rows and columns allow further classification of the elements according to their properties. This section will cover the different features of the periodic table.

Elements and Symbols Elements cannot be broken down into simpler substances. For example aluminum is an element only made of aluminum atoms and if you analyze the composition of a piece of this metals you would only find aluminium atoms. Chemical symbols are one- or two-letter abbreviations that represent the names of the elements. Only the first letter is capitalized and if a second letter exist in the element's name, the second letter should be lowercase. For example, the chemical symbol for aluminum is Al, written as capital A and lowercase l.


Sample Problem 1

Give the symbol or name the following elements: C, Oxygen, N, Phosphorus, Au, Iron, Na and Iodine.

SOLUTION

The name of the element with symbol C is carbon, whereas the chemical symbol of oxygen is O. Similarly, N stands for nitrogen, whereas the chemical symbol of Phosphorus is P. The chemical symbol of Au is Gold. The chemical symbol of Iron is Fe and the chemical symbol of Iodine is I.

STUDY CHECK



GOALS

- 1 Navigate the periodic table
- 2 Calculate average atomic masses, number of electrons, protons and neutrons in an atom
- 3 Calculate simple molecular weights
- 4 Calculate molecular formulas from empirical formulas

Discussion: Why having a periodic table? What is the basic information all periodic tables provide? Do you know any other periodic table design?

“Nothing exists except atoms and empty space; everything else is opinion.”
Democritus

Give the symbol or name the following elements: Ni.

Periods and groups The periodic table contains all elements arranged in rows and columns. The horizontal rows are called *periods* and the vertical columns are called *groups or families*. For example, the first period contains hydrogen (H) and helium (He), whereas the second group contains Beryllium (Be), Magnesium (Mg), Calcium (Ca), Strontium (Sr), Barium (Ba) and Radium (Ra). There are seven periods (periods 1-7) and 18 groups. Some of the groups are labeled with an A (e.g. group 8A) whereas others are labeled with a B (e.g. group 8B). The groups numbers can be found written with roman numbers and a letters (A or B) or with a more modern group numbering of 1-18 going across the periodic table. For example, the group 2 (Mg-Ra) can also be called IIA, and the group 13 (B-Ti) is also known as IIIA.

Properties in the periodic table The physical and chemical properties of some elements of the table are similar, and these similarities led to the organization of the periodic table. Elements in the same group share properties and for example, oxygen and sulfur have similar properties: both are reactive elements. Differently, the properties across periods change going from metals to nonmetals. For example, the properties of Li and Ne are very different and lithium is a reactive metal whereas neon is a nonreactive gas.

Metals, Nonmetals, and Metalloids Overall, the elements of the periodic table can be classify as metals, nonmetals, and metalloids. Metals are those elements on the left of the table and nonmetals are the elements on the right of the table. The elements between metals and nonmetals are called metalloids and include only B, Si, Ge, As, Sb, Te, Po, and At. Metals are shiny solids and usually melt at higher temperatures. Some examples of metals are Gold (Au) or Iron (Fe). Nonmetals are often poor conductors of heat and electricity with low melting points. They also tend to be matt (non-shinny), malleable, or ductile. Some examples of nonmetals are Carbon (C) or Nitrogen (N). Metalloids are elements that share some properties with metals and others with the nonmetals. For example, they are better conductors of heat and electricity than the nonmetals, but not as good conductors as the metals. Metalloids are semiconductors because they can act as both conductors and insulators under certain conditions. An example of metalloids is Silicon (Si) that should not be confused from silicone, a chemical employed in prosthetics.

Sample Problem 2

Answer the following questions: (a) Give the group and period of the following elements, and give the name: Ca, Ir, and C. (b) Classify as alkali metal, alkali earth metal, transition metal, halogen or noble gas, and give the name: Mg, Li, Co, He, F. (c) Classify as metal, nonmetal or metalloid, and give the name: Ba, N, Si.

SOLUTION

(a) The period and group of Ca (Calcium) is 2 (2A) and 4, respectively. The period and group of Ir (Iridium) is 9 (8B) and 6, respectively. The period and group of C (Carbon) is 14 (IVA) and 2, respectively. (b) Mg (Magnesium) is an alkali earth metal, whereas Li (Lithium) is a alkali metal. Co (Cobalt) is a transition metal. He (Helium) is a noble gas. F (Fluorine) is an halogen. (c) Ba (Barium) is a metal. N (Nitrogen) is a nonmetal. Si (Silicon) is a metalloid.

STUDY CHECK

Answer the following questions: (a) Give the group and period of the following elements, and give the name: Cl. (b) Classify as alkali metal, alkali earth metal, transition metal, halogen or noble gas, and give the name: Ne. (c) Classify as metal, nonmetal or metalloid, and give the name: W.

60

Classification of elements in terms of groups Some of the groups in the periodic table have specific names such as alkali metals, alkaline earth metals, transition metals, chalcogens, halogens or noble gases. Alkali metals are the group 1A elements: lithium (Li), sodium (Na), potassium (K), rubidium (Rb), cesium (Cs), and francium (Fr). Alkali elements are soft and shiny metals, and they are also good conductors of heat and electricity, with low melting points. Alkali earth metals are the group 2A (2) elements: beryllium (Be), magnesium (Mg), calcium (Ca), strontium (Sr), barium (Ba), and radium (Ra). Transition metals are the elements from group 3 to 12 and they are located in the middle of the table. Chalcogens are the group 6A (16) elements: oxygen (O), sulfur (S), selenium (Se), tellurium (Te), and polonium (Po). Halogens are the group 7A (17) elements: fluorine (F), chlorine (Cl), bromine (Br), iodine (I), and astatine (At). Halogens are very reactive elements. Finally, noble gases are the group 8A (18) elements: helium (He), neon (Ne), argon (Ar), krypton (Kr), xenon (Xe), and radon (Rn). They are inert and rarely combine with other elements in the periodic table, like a noble family: have you ever meet a royal?

65

70


75

How to classify Hydrogen At first sight, hydrogen (H) may seem to be put in the wrong spot at the periodic table. Although it is located at the top of Group 1A (1), it is not an alkali metal, as it has very different properties. Thus hydrogen does not belong to the alkali metals, being nonmetal.

80

1 IA																		18 VIIIA																	
1 1 H Hydrogen																		2 2 He Helium																	
2 2 Li Lithium																		3 3 Be Beryllium																	
3 3 Na Sodium																		4 4 Mg Magnesium																	
4 4 K Potassium																		5 5 Ca Calcium																	
5 5 Rb Rubidium																		6 6 Sr Strontium																	
6 6 Cs Cesium																		7 7 Ba Barium																	
7 7 Fr Francium																		8 8 Ra Radium																	
8 8 Ac-Lr Actinide																		9 9 La-Lu Lanthanide																	
9 9 Pr Praseodymium																		10 10 Nd Neodymium																	
10 10 Ce Cerium																		11 11 Pm Promethium																	
11 11 Eu Europium																		12 12 Gd Gadolinium																	
12 12 Tb Terbium																		13 13 Dy Dysprosium																	
13 13 Ho Holmium																		14 14 Er Erbium																	
14 14 Tm Thulium																		15 15 Yb Ytterbium																	
15 15 Lu Lutetium																		16 16 La Lanthanum																	
16 16 Ce Cerium																		17 17 Pr Praseodymium																	
17 17 Nd Neodymium																		18 18 Pm Promethium																	
18 18 Eu Europium																		19 19 Gd Gadolinium																	
19 19 Tb Terbium																		20 20 Dy Dysprosium																	
20 20 Ho Holmium																		21 21 Er Erbium																	
21 21 Tm Thulium																		22 22 Yb Ytterbium																	
22 22 Lu Lutetium																		23 23 La Lanthanum																	
23 23 Ce Cerium																		24 24 Pr Praseodymium																	
24 24 Nd Neodymium																		25 25 Pm Promethium																	
25 25 Eu Europium																		26 26 Gd Gadolinium																	
26 26 Tb Terbium																		27 27 Dy Dysprosium																	
27 27 Ho Holmium																		28 28 Er Erbium																	
28 28 Tm Thulium																		29 29 Yb Ytterbium																	
29 29 Lu Lutetium																		30 30 La Lanthanum																	
30 30 Ce Cerium																		31 31 Pr Praseodymium																	
31 31 Nd Neodymium																		32 32 Pm Promethium																	
32 32 Eu Europium																		33 33 Gd Gadolinium																	
33 33 Tb Terbium																		34 34 Dy Dysprosium																	
34 34 Ho Holmium																		35 35 Er Erbium																	
35 35 Tm Thulium																		36 36 Yb Ytterbium																	
36 36 Lu Lutetium																		37 37 La Lanthanum																	
37 37 Ce Cerium																		38 38 Pr Praseodymium																	
38 38 Nd Neodymium																		39 39 Pm Promethium																	
39 39 Eu Europium																		40 40 Gd Gadolinium																	
40 40 Tb Terbium																		41 41 Dy Dysprosium																	
41 41 Ho Holmium																		42 42 Er Erbium																	
42 42 Tm Thulium																		43 43 Yb Ytterbium																	
43 43 Lu Lutetium																		44 44 La Lanthanum																	
44 44 Ce Cerium																		45 45 Pr Praseodymium																	
45 45 Nd Neodymium																		46 46 Pm Promethium																	
46 46 Eu Europium																		47 47 Gd Gadolinium																	
47 47 Tb Terbium																		48 48 Dy Dysprosium																	
48 48 Ho Holmium																		49 49 Er Erbium																	
49 49 Tm Thulium																		50 50 Yb Ytterbium																	
50 50 Lu Lutetium																		51 51 La Lanthanum																	
51 51 Ce Cerium																		52 52 Pr Praseodymium																	
52 52 Nd Neodymium																		53 53 Pm Promethium																	
53 53 Eu Europium																		54 54 Gd Gadolinium																	
54 54 Tb Terbium																		55 55 Dy Dysprosium																	
55 55 Ho Holmium																		56 56 Er Erbium																	
56 56 Tm Thulium																		57 57 Yb Ytterbium																	
57 57 Lu Lutetium																		58 58 La Lanthanum																	
58 58 Ce Cerium																		59 59 Pr Praseodymium																	
59 59 Nd Neodymium																		60 60 Pm Promethium																	
60 60 Eu Europium																		61 61 Gd Gadolinium																	
61 61 Tb Terbium																		62 62 Dy Dysprosium																	
62 62 Ho Holmium																		63 63 Er Erbium																	
63 63 Tm Thulium																		64 64 Yb Ytterbium																	
64 64 Lu Lutetium																		65 65 La Lanthanum																	
65 65 Ce Cerium																		66 66 Pr Praseodymium																	
66 66 Nd Neodymium																		67 67 Pm Promethium																	
67 67 Eu Europium																		68 68 Gd Gadolinium																	
68 68 Tb Terbium																		69 69 Dy Dysprosium																	
69 69 Ho Holmium																		70 70 Er Erbium																	
70 70 Tm Thulium																		71 71 Yb Ytterbium																	
71 71 Lu Lutetium																		72 72 La Lanthanum																	
72 72 Ce Cerium																		73 73 Pr Praseodymium																	
73 73 Nd Neodymium																		74 74 Pm Promethium																	
74 74 Eu Europium																		75 75 Gd Gadolinium																	
75 75 Tb Terbium																		76 76 Dy Dysprosium																	
76 76 Ho Holmium																		77 77 Er Erbium																	
77 77 Tm Thulium																		78 78 Yb Ytterbium																	
78 78 Lu Lutetium																		79 79 La Lanthanum																	
79 79 Ce Cerium																		80 80 Pr Praseodymium																	
80 80 Nd Neodymium																		81 81 Pm Promethium																	
81 81 Eu Europium																		82 82 Gd Gadolinium																	
82 82 Tb Terbium																		83 83 Dy Dysprosium																	
83 83 Ho Holmium																		84 84 Er Erbium																	
84 84 Tm Thulium																		85 85 Yb Ytterbium																	
85 85 Lu Lutetium																		86 86 La Lanthanum																	
86 86 Ce Cerium																		87 87 Pr Praseodymium																	
87 87 Nd Neodymium																		88 88 Pm Promethium																	
88 88 Eu Europium																		89 89 Gd Gadolinium																	
89 89 Tb Terbium																		90 90 Dy Dysprosium																	
90 90 Ho Holmium																		91 91 Er Erbium																	
91 91 Tm Thulium																		92 92 Yb Ytterbium																	
92 92 Lu Lutetium																		93 93 La Lanthanum																	
93 93 Ce Cerium																		94 94 Pr Praseodymium																	
94 94 Nd Neodymium																		95 95 Pm Promethium																	
95 95 Eu Europium																		96 96 Gd Gadolinium																	
96 96 Tb Terbium																		97 97 Dy Dysprosium																	
97 97 Ho Holmium																		98 98 Er Erbium																	
98 98 Tm Thulium																		99 99 Yb Ytterbium																	
99 99 Lu Lutetium																		100 100 La Lanthanum																	
100 100 Ce Cerium																		101 101 Pr Praseodymium																	
101 101 Nd Neodymium																		102 102 Pm Promethium																	
102 102 Eu Europium																		103 103 Gd Gadolinium																	
103 103 Tb Terbium																		104 104 Dy Dysprosium																	
104 104 Ho Holmium																		105 105 Er Erbium																	
105 105 Tm Thulium																		106 106 Yb Ytterbium																	
106 106 Lu Lutetium																		107 107 La Lanthanum																	
107 107 Ce Cerium																		108 108 Pr Praseodymium																	
108 108 Nd Neodymium																		109 109 Pm Promethium																	
109 109 Eu Europium																		110 110 Gd Gadolinium																	
110 110 Tb Terbium																		111 111 Dy Dysprosium																	
111 111 Ho Holmium																		112 112 Er Erbium																	
112 112 Tm Thulium																		113 113 Yb Ytterbium																	
113 113 Lu Lutetium																		114 114 La Lanthanum																	
114 114 Ce Cerium																		115 115 Pr Praseodymium																	
115 115 Nd Neodymium																		116 116 Pm Promethium																	
116 116 Eu Europium																		117 117 Gd Gadolinium																	
117 117 Tb Terbium																		118 118 Dy Dysprosium																	
118 118 Ho Holmium																		119 119 Er Erbium																	
119 119 Tm Thulium																		120 120 Yb Ytterbium																	
120 120 Lu Lutetium																		121 121 La Lanthanum																	
121 121 Ce Cerium																		122 122 Pr Praseodymium																	
122 122 Nd Neodymium																		123 123 Pm Promethium																	
123 123 Eu Europium																		124 124 Gd Gadolinium																	
124 124 Tb Terbium																		125 125 Dy Dysprosium																	
125 125 Ho Holmium																		126 126 Er Erbium																	
126 126 Tm Thulium																		127 127 Yb Ytterbium																	
127 127 Lu Lutetium																		128 128 La Lanthanum																	
128 128 Ce Cerium																		129 129 Pr Praseodymium																	
129 129 Nd Neodymium																		130 130 Pm Promethium																	
130 130 Eu Europium																		131 131 Gd Gadolinium																	
131 131 Tb Terbium																		132 132 Dy Dysprosium																	
132 132 Ho Holmium																		133 133 Er Erbium																	
133 133 Tm Thulium																		134 134 Yb Ytterbium																	
134 134 Lu Lutetium																		135 135 La Lanthanum																	
135 135 Ce Cerium																		136 136 Pr Praseodymium																	
136 136 Nd Neodymium																		137 137 Pm Promethium																	
137 137 Eu Europium																		138 138 Gd Gadolinium																	
138 138 Tb Terbium																		139 139 Dy Dysprosium																	
139 139 Ho Holmium																		140 140 Er Erbium																	
140 140 Tm Thulium																		141 141 Yb Ytterbium																	
141 141 Lu Lutetium																		142 142 La Lanthanum																	
142 142 Ce Cerium																		143 143 Pr Praseodymium																	
143 143 Nd Neodymium																		144 144 Pm Promethium																	
144 144 Eu Europium																		145 145 Gd Gadolinium																	
145 145 Tb Terbium																		146 146 Dy Dysprosium																	
146 146 Ho Holmium																		147 147 Er Erbium																	
147 147 Tm Thulium																		148 148 Yb Ytterbium																	
148 148 Lu Lutetium																		149 149 La Lanthanum																	
149 149 Ce Cerium																		150 150 Pr Praseodymium																	
150 150 Nd Neodymium																		151 151 Pm Promethium																	
151 151 Eu Europium																		152 152 Gd Gadolinium																	
152 152 Tb Terbium																		153 153 Dy Dysprosium																	
153 153 Ho Holmium																		154 154 Er Erbium																	
154 154 Tm Thulium																		155 155 Yb Ytterbium																	
155 155 Lu Lutetium																		156 156 La Lanthanum																	
156 156 Ce Cerium																		157 157 Pr Praseodymium																	
157 157 Nd Neodymium																		158 158 Pm Promethium																	
158 158 Eu Europium																		159 159 Gd Gadolinium																	
159 159 Tb Terbium																		160 160 Dy Dysprosium																	
160 160 Ho Holmium																		161 161 Er Erbium																	
161 161 Tm Thulium																		162 162 Yb Ytterbium																	
162 162 Lu Lutetium																		163 163 La Lanthanum																	
163 163 Ce Cerium																		164 164 Pr Praseodymium																	
164 164 Nd Neodymium																		165 165 Pm Promethium																	
165 165 Eu Europium																		166 166 Gd Gadolinium																	
166 166 Tb Terbium																		167 167 Dy Dysprosium																	
167 167 Ho Holmium																		168 168 Er Erbium																	
168 168 Tm Thulium																		169 169 Yb Ytterbium																	
169 169 Lu Lutetium																		170 170 La Lanthanum																	
170 170 Ce Cerium																		171 171 Pr Praseodymium																	
171 171 Nd Neodymium																		172 172 Pm Promethium																	
172 172 Eu Europium																		173 173 Gd Gadolinium																	
173 173 Tb Terbium																		174 174 Dy Dysprosium																	
174 174 Ho Holmium																		175 175 Er Erbium																	
175 175 Tm Thulium																		176 176 Yb Ytterbium																	
176 176 Lu Lutetium																		177 177 La Lanthanum																	
177 177 Ce Cerium																		178 178 Pr Praseodymium																	
178 178 Nd Neodymium																		179 179 Pm Promethium																	
179 179 Eu Europium																		180 180 Gd Gadolinium																	
180 180 Tb Terbium																		181 181 Dy Dysprosium																	
181 181 Ho Holmium																		182 182 Er Erbium																	
182 182 Tm Thulium																		183 183 Yb Ytterbium																	
183 183 Lu Lutetium																		184 184 La Lanthanum																	
184 184 Ce Cerium																		185 185 Pr Praseodymium																	
185 185 Nd Neodymium																		186 186 Pm Promethium																	
186 186 Eu Europium																		187 187 Gd Gadolinium																	
187 187 Tb Terbium																		188 188 Dy Dysprosium																	
188 188 Ho Holmium																		189 189 Er Erbium																	
189 189 Tm Thulium																		190 190 Yb Ytterbium																	
190 190 Lu Lutetium																		191 191 La Lanthanum																	
191 191 Ce Cerium																		192 192 Pr Praseodymium																	
192 192 Nd Neodymium																		193 193 Pm Promethium																	
193 193 Eu Europium																		194 194 Gd Gadolinium																	
194 194 Tb Terbium																		195 195 Dy Dysprosium																	
195 195 Ho Holmium																		196 196 Er Erbium																	
196 196 Tm Thulium																		197 197 Yb Ytterbium																	
197 197 Lu Lutetium																		198 198 La Lanthanum																	
198 198 Ce Cerium																		199 199 Pr Praseodymium																	
199 199 Nd Neodymium																		200 200 Pm Promethium																	
200 200 Eu Europium																		201 201 Gd Gadolinium																	
201 201 Tb Terbium																		202 202 Dy Dysprosium																	
202 202 Ho Holmium																		203 203 Er Erbium																	
203 203 Tm Thulium																		204 204 Yb Ytterbium																	
204 204 Lu Lutetium																		205 205 La Lanthanum																	
205 205 Ce Cerium																		206 206 Pr Praseodymium																	
206 206 Nd Neodymium																		207 207 Pm Promethium																	
207 207 Eu Europium																		208 208 Gd Gadolinium																	
208 208 Tb Terbium																		209 209 Dy Dysprosium																	
209 209 Ho Holmium																		210 210 Er Erbium																	
210 210 Tm Thulium																		211 211 Yb Ytterbium																	
211 211 Lu Lutetium																		212 212 La Lanthanum																	
212 212 Ce Cerium																		213 213 Pr Praseodymium																	
213 213 Nd Neodymium																		214 214 Pm Promethium																	
214 214 Eu Europium																		215 215 Gd Gadolinium																	
215 215 Tb Terbium																		216 216 Dy Dysprosium																	
216 216 Ho Holmium																		217 217 Er Erbium																	
217 217 Tm Thulium																		218 218 Yb Ytterbium																	
218 218 Lu Lutetium																		219 219 La Lanthanum																	
219 219 Ce Cerium																		220 220 Pr Praseodymium																	
220 220 Nd Neodymium																		221 221 Pm Promethium																	
221 221 Eu Europium																		222 222 Gd Gadolinium																	
222 222 Tb Terbium																		223 223 Dy Dysprosium																	
223 223 Ho Holmium																		224 224 Er Erbium																	
224 224 Tm Thulium																		225 225 Yb Ytterbium																	
225 225 Lu Lutetium</																																			

▼A cathode tube



© www.wallpaperflare.com

▼Millikan apparatus



© www.wallpaperflare.com

▼A plum pudding, with the electrons represented by the raisins



© wikipedia

1.2 Early experiments of the atom

Scientists wondered about the nature of the atom and its structure for years. In a series of experiments carried in the late nineteenth century, scientists such as J.J. Thomson, Henri Becquerel, or Ernest Rutherford gained insight into the nature and structure of the atom. These remarkable scientists and these creative experiments helped shape the view of the atom that we have nowadays.




Figure ?? Cathode-ray tube made of two electrodes and a partially evacuated gas tube. The electrons, generated on the negatively charged electrode names cathode, excite the gas in the tube generating glow.

Charge to mass ratio of an electron The English researcher J.J. Thomson investigated electric discharges in partially evacuated tubes—tubes in which the air has been partially removed (See Figure ??); these tubes, made of a positive and negative electrode, are the base of old-fashion, bulky televisions. Thomson found that rays emanated from the negative electrode when applying high-voltage to these tubes. These rays were names cathodic rays as the negative electrode of the tube is called the cathode. Thomson also found that cathodic rays were repelled from the negative pole of an electric field. Hence, these rays were postulated to be a stream of negatively charged particles, now known as electrons. By studying the deflection of these rays by an electric field, Thomson was able to calculate the charge-to-mass ratio of an electron:

$$\frac{e}{m_e} = -1.76 \times 10^8 C/g \tag{1.1}$$

where:

e is the charge of an electron

m_e is the mass of an electron

Overall, the biggest of Thomson’s discoveries was that all atoms are made of negatively charged particles. As atoms are charge-neutral, they are also made of positively charged particles. These observations led to a new atomic model, the *plum pudding model*, that envisioned atoms as a diffuse cloud of positive charges with negative electrons embedded in it. The name plum pudding comes from an English dessert that contains a raising spread on it. A different scienticts, Robert Millikan, revealed the magnitude of the electric charge of the electron. Millikan used an apparatus that dispersed charged oil droplets falling under the influence of an electric field. Given the applied voltage and the droplet mass, Millikan was able to calculate the droplet charge. He found that the oil drop charge was always a whole-number time the electron charge,

$$e = 1.60 \times 10^{-19} C \tag{1.2}$$

where a Coulomb is a unit of charge. With the value of the charge-to-mass ratio of an electron and the electron charge, Millikan was also able to calculate the mass of an electron,

$$m_e = 9.11 \times 10^{-31} \text{ kg.} \quad (1.3)$$

The atom nucleus Ernest Rutherford carried further experiments to validate the plum pudding model of the atom. He exposed a thin sheet of metal foil to α particles known to be massive and positively charged particles. According to the plum pudding model, the bulky α particles should have crashed through the thin foil and traverse through without being deflected. However, the results did not corroborate his expectations. Indeed, some particles traversed the film, whereas other were slightly deflected and some were strongly deflected at large angles. These observations did not corroborate the plum pudding model. However, they contributed to the creation of the modern atomic model in which a large number of positive charges were concentrated at a point—called the nucleus—instead of being spread whereas the electrons move around the nucleus at large distances from it. The figure below represents the experiment carried out by Rutherford in which alpha particles were scattered on a thin foil made of gold.




Figure ?? Rutherford's scattering: the elastic scattering of charged particles by the Coulomb interaction after a particle beam passed through a thin gold foil obstruction.

Radiation In the last part of the nineteenth century, scientists came to the discovery that some materials were able to produce high-energy radiation. Among the scientist working in this field, Henri Becquerel discovered that pitchblende, a mineral containing uranium, was able to produce an image on a photographic plate in the absence of light. In the early twentieth century, three different types of radiation were discovered: alpha radiation, beta radiation, and gamma radiation. Further studies revealed that gamma radiation was made of gamma particles, high-energy radiation, whereas beta radiation was made of high-energy electrons. Alpha particles were found to be positively charged, with a charge twice the charge of an electron and a mass 7300 times that of the electron. These particles were indeed Helium nuclei, resulting from removing electrons from atoms of Helium.

Atoms are the smallest piece of an element that retains its characteristics. They are the building blocks of matter. This section covers the structure of the atom. You will learn how to calculate the number of subatomic particles that made an atom and how to differentiate atoms of an element—all atoms of an elements are not equal.

145




Figure ?? Three models of the atom. From left to right: the plum puding model, the updated plum pudding model according to Rutherford's observations, (two right images) the modern atomic model. Z is the atomic number of the atom.

Atomic Structure Atoms contain three atomic particles: the proton, neutron, and electron. Protons have positive charge (+), whereas electrons carry negative charge (-). Neutrons on the other hand are neutral, and they have no electrical charge. Protons and neutrons are located in the core of the atom, which is called the nucleus, and account for the mass of the atom. Electrons are delocalized in the exterior part of the atoms. They are not necessarily located in a specific spot and their existence spreads in the area next to the nucleus. When an atoms is neutral it has no charge and the number of electrons and protons are the same. Some atoms have positive charge, resulting of removing electrons, and we call these cations. Others—called anions—can have negative charge as a result of accepting a negatively charged electron.

150

155

Atomic and mass number Elements are made of atoms, and each atom of an element is characterized by a atomic number (Z) and a mass number (A). The atomic number (Z) of an element indicates the number of electrons of an atom. This number can be easily located in the periodic table. The mass number (A) of an element indicates the combined number of protons and neutrons. Mass numbers are nowhere located in the periodic table as different atoms of the same element can have different mass numbers. Both A and Z for an atom X are indicated in the following form called isotope notation:


160

165

A_ZX

As an example, the notation ${}^{24}_{12}\text{Mg}$ means that the atomic number of Mg is $Z=12$ and the mass number is $A=24$. Remember that the atomic number can be found in the periodic table whereas the mass number A is not on the table. By means of the isotope notation, one can quickly identify the number of protons, neutrons and electrons in an atom. As the atomic number is always indicated on the bottom part (e.g. Mg has 12 electrons). At the same time, the number of electrons and protons in a neutral atom is the same—neutral means an atom without a charge. The number of neutrons of an isotope can be computed by subtracting the atomic number from the mass number.

170



175

Sample Problem 3

Calculate the number of protons, neutrons and electrons of the following atoms:

- (a) ${}^{27}_{12}\text{Mg}$ (b) ${}^{22}_{10}\text{Ne}$ (c) ${}^{20}_{10}\text{Ne}$

SOLUTION

(a) ${}^{27}_{12}\text{Mg}$ has 12 electrons ($Z=12$) and 12 protons as well (the number of electrons and protons are the same if the atom is neutral), and 15 neutrons, as $27-12=15$.

(b) ${}^{22}_{10}\text{Ne}$ has 10 electrons and 10 protons, and 12 neutrons. (c) ${}^{20}_{10}\text{Ne}$ has 10 electrons and 10 protons, and 10 neutrons as well.

STUDY CHECK

Calculate the number of protons, neutrons and electrons of the following atoms:

- (a) ${}^{32}_{16}\text{S}$ (b) ${}^{34}_{16}\text{S}$ (c) ${}^{36}_{16}\text{S}$

Isotopes All atoms of the same element are not the same. Some are heavier than others. Isotopes are atoms of the same element with different numbers of neutrons and therefore with different mass number but with the same atomic number. For example: ${}^{24}_{12}\text{Mg}$, ${}^{25}_{12}\text{Mg}$ and ${}^{26}_{12}\text{Mg}$ are three isotopes of Mg. ${}^{27}_{12}\text{Mg}$ is heavier than ${}^{24}_{12}\text{Mg}$ as it contains more neutrons and protons in the nucleus. Each of the isotopes has a specific abundance, as some isotopes are more abundant than others. For example, the abundance of ${}^{24}_{12}\text{Mg}$ is 79%, and the abundance of ${}^{25}_{12}\text{Mg}$ and ${}^{26}_{12}\text{Mg}$ is 10% and 11%, respectively. This means, ${}^{24}_{12}\text{Mg}$ is more abundant than for example ${}^{26}_{12}\text{Mg}$.

Average atomic mass The average atomic mass represents the mass of the atoms of an element and results from all existing isotopes taking into account their abundance. The units of atomic mass are called *amu*, which stands for atomic mass units. This value can be simply found at any periodic table. Using the periodic table provided in this manual Figure ??, you can find the atomic mass of each element on top of the symbol at the right side. For example, the atomic mass of oxygen (O) is 15.999 amu and the atomic mass of nitrogen (N) is 14.007 amu. As atoms are made of numerous isotopes—this means different atoms of the same element but with different number of neutrons and hence different weight—the atomic mass found in the periodic table is an average that results from including the mass of the different isotopes and their abundance. That is you need to do an average of the mass of each isotope using values of abundance. In other words: the *average atomic mass* of an element—herein called simply atomic mass—, expressed in *amu* (atomic mass units), is the weighted average of the masses of the individual isotopes of the element. For an element with n isotopes with different masses (A_1, A_2, \dots, A_n) and different fractional abundances for each isotope (f_1, f_2, \dots, f_n), the atomic mass is given by

$$\text{Atomic mass} = \sum_{i=1}^n A_i \cdot f_i = A_1 \cdot f_1 + A_2 \cdot f_2 + \dots + A_n \cdot f_n$$

Also, note that when adding the fractional abundances of all isotopes, one should obtain a value of one:

$$\sum_{i=1}^n f_i = f_1 + f_2 + \dots + f_n = 1$$

Sample Problem 4

Naturally occurring copper (Cu) consists of 69.17% ${}^{63}\text{Cu}$ and 30.83% ${}^{65}\text{Cu}$. The mass of ${}^{63}\text{Cu}$ is 62.939598 amu, and the mass of ${}^{65}\text{Cu}$ is 64.927793 amu.

What is the atomic mass of copper?

SOLUTION

The weighted average is the sum of the mass of each isotope times its fractional abundance. We have that the isotope ^{63}Cu has a mass of 62.939598 amu and an abundance of 69.17%, that is the same as 0.6917. At the same time, the isotope ^{65}Cu has a mass of 64.927793 amu and an abundance of 0.3083. After adding both contributions, we have:

$$62.939598 \text{ amu} \times \frac{69.17}{100} + 64.927793 \text{ amu} \times \frac{30.83}{100} = 63.55 \text{ amu}$$

STUDY CHECK

Lithium is made up of two isotopes, Li-7 (7.016003 amu) and Li-6 (6.015121 amu). Calculate the percent abundance of each isotope knowing that lithium's atomic weight is 6.94 amu.

Table ?? Isotope abundance of some elements

Element	Isotope	% Abundance	Element	Isotope	% Abundance
Hydrogen	^1H	99.9885%	Silicon	^{28}Si	92.2297%
	^2H	0.0115%		^{29}Si	4.6832%
Helium	^3He	0.000137%		^{30}Si	3.0872%
	^4He	99.999863%	Sulfur	^{32}S	94.93%
Lithium	^6Li	7.59%		^{33}S	0.76%
	^7Li	92.41%		^{34}S	4.29%
Boron	^{10}B	19.9%	Chlorine	^{36}S	0.02%
	^{11}B	80.1%		^{35}Cl	75.78%
Carbon	^{12}C	98.93%		^{37}Cl	24.22%
	^{13}C	1.07%	Argon	^{36}Ar	0.3365%
Nitrogen	^{14}N	99.632%		^{38}Ar	0.0632%
	^{15}N	0.368%		^{40}Ar	99.6003%
Oxygen	^{16}N	99.757%	Potassium	^{39}K	93.2581%
	^{17}O	0.038%		^{40}K	0.0117%
	^{16}O	0.205%		^{41}K	6.7302%

205

1.4 An introduction to molecules

The periodic table contains all elements in nature. At the same time, elements combine to form molecules. For example, in the air there are traces of Argon—this is an element—and also water, a molecule (H_2O) that results from the combination of two elements, hydrogen (H) and oxygen (O). This section will first introduce some of the properties of molecules, without paying attention to their chemical names that will be covered in the following chapters.

210

Molecular weight Here are two examples of molecules: molecular oxygen O_2 and molecular nitrogen N_2 . How do we interpret these formulas? The subscript "2" indicates that each molecule contains two atoms. For example, a O_2 molecule is made of two oxygen atoms O. At the same time, the weight of a set of molecules is called the molecular weight (MW). However, you will find different terms to refer to the same property such as: molecular mass, molar mass, or formula unit mass. All these terms

215

indeed mean the weight a large set of molecules. We can calculate the MW by adding the weight of each atom that form the molecule taking into the account the number of atoms of each element present in the molecule. The units of molecular weight are the same as the units of atomic weight: amu, atomic mass units.

Sample Problem 5

Calculate: (a) The atomic weight of O; (b) the molecular mass of molecular oxygen, O₂

SOLUTION

- (a) According to the periodic table the atomic weight (AW) of O is 15.999 amu.
 (b) The molar mass of O₂ is the result of adding the atomic masses of 2O atoms, that is 31.998 amu, close to 32 amu.

STUDY CHECK

Calculate the molar mass of water H₂O and ammonia, NH₃

Mass percent composition of a compound Look at these two molecules: C₂H₂ and C₂H₆. They contain different amounts of hydrogen. We quantify the amount of an element in a compound by means of the mass % composition. The mass % of an element in a compound is the mass of the element with respect to the molecular weight of the molecule in percent form. Mind that you have to take into account the molecular indexes in the compound as C₂H₂ is made of 2H and C₂H₆ is made of 6H. For example, given that the molar mass of C₂H₂ is 26 amu, the mass % of hydrogen in C₂H₂ would be:

$$\%_H \text{ in C}_2\text{H}_2 = \frac{2 \cdot AW(H)}{MW(\text{C}_2\text{H}_2)} \times 100 = \frac{2 \cdot 1}{26} \times 100 = 7.7\%$$

Similarly, the mass % of C would be:

$$\%_C \text{ in C}_2\text{H}_2 = \frac{2 \cdot AW(C)}{MW(\text{C}_2\text{H}_2)} \times 100 = \frac{2 \cdot 12}{26} \times 100 = 92.3\%$$

By adding the mass % of all elements in a molecule we should obtain 100.

$$\%_H \text{ in C}_2\text{H}_2 + \%_C \text{ in C}_2\text{H}_2 = 100$$

Sample Problem 6

Calculate the mass % composition for each element of glucose, C₆H₁₂O₆.

SOLUTION

We first need the molecular weight of glucose, C₆H₁₂O₆, that is: 6 · 12 + 12 · 1 + 6 · 16 = 180 amu. Now we can calculate the mass percent of carbon, hydrogen and oxygen:

$$\%_C \text{ in C}_6\text{H}_{12}\text{O}_6 = \frac{6 \cdot 12}{180} \times 100 = 40\%$$

$$\%_H \text{ in C}_6\text{H}_{12}\text{O}_6 = \frac{12 \cdot 1}{180} \times 100 = 6.6\%$$

By subtraction, we have that %_C in C₆H₁₂O₆ = 53.4.

STUDY CHECK

Ureas CO(NH₂)₂ is a colorless crystalline compound excreted in urine, product of protein metabolism in mammals. Calculate the mass % composition for each element of urea.

1.5 Empirical and molecular formula of a chemical

There are two different types of formulas: molecular formulas and empirical formulas. Empirical formula (EFs) are simplified formulas resulting from an experiment, whereas molecular formulas (MFs) are exact formulas of molecules. For example: the molecular formula of hydrogen peroxide, a mild antiseptic used on the skin to prevent infection of minor cuts, is H_2O_2 as the hydrogen peroxide molecule is made of two oxygen and two hydrogen atoms. Differently, the empirical formula of the same chemical is HO , being this the result of the simplification of H_2O_2 . One can obtain empirical formulas simply by dividing the molecular formula by the smallest integer number, of course, given you know the molecular formula. The word empirical means "from an experiment", and the use of empirical formulas comes from the fact that the formulas of all chemicals actually come from experiments, and from experiments one normally can only obtain ratios of atomic composition.

Sample Problem 7

From the following formulas identify the empirical and molecular formulas: P_4O_{10} , $\text{C}_3\text{H}_6\text{O}$, N_2O_4 and C_5H_{11} .

SOLUTION

Empirical formulas are simplified versions of molecular formulas. For example, $\text{C}_3\text{H}_6\text{O}$ and C_5H_{11} are empirical formulas. Differently, P_4O_{10} and N_2O_4 are molecular formulas.

STUDY CHECK

Given the following molecular formulas, obtain the corresponding empirical formula: P_4O_{10} , N_2O_4 and $\text{C}_6\text{H}_{18}\text{O}_3$.

Molecular weight of empirical formulas and molecular formulas

The molecular weight of an empirical formula and its corresponding molecular formula are related by the following formula:

$$n = \frac{MW_{MF}}{MW_{EF}}$$

where:

MW_{EF} is the molecular weight of the empirical formula

MW_{MF} is the molecular weight of the molecular formula

n is a integer number such as 1, 2, 3...

Understanding the formula above is simple. On one hand, the MW of a molecular formula H_2O_2 that is 34 amu. On the other hand, the molecular weight of the empirical formula of the same chemical HO is 17 amu. If we do $34/17$ we would obtain 2, as we need to multiply HO by two in order to obtain H_2O_2 . As a final note, mind that empirical formulas are just simplified formulas. So when we think about the molecular weight of a chemical we normally have molecular formula in mind. Let us work on an example:

Sample Problem 8

Given that the empirical formula of dichloromethane is ClCH_2 and the molecular weight of the chemical is 98 amu, calculate the molecular formula of dichloromethane.

SOLUTION

Given the empirical formula of dichloromethane one can think of many different molecular formulas, for example: $\text{Cl}_3\text{C}_3\text{H}_6$ or $\text{Cl}_2\text{C}_2\text{H}_4$. From these, and many other, there is only one real molecular formula. How do we calculate the real molecular formula? By comparing the MW of the molecular and empirical formula we can figure out the number of times we need to multiply the MF to obtain the EF. We know the MW is 98 amu. Using the EF we can also calculate a MW: $35 + 12 + 2 \cdot 1 = 49$ amu. If we compare both numbers using the formula:

$$n = \frac{MW_{MF}}{MW_{EF}}$$

we have: $n = 98/49$ and solving we have $n = 2$. Therefore the MF is: $\text{Cl}_2\text{C}_2\text{H}_4$.

STUDY CHECK

The empirical formula of dinitrogen tetroxide, a red-brown liquid with an unpleasant chemical odor, is NO_2 and the molecular weight of the chemical is 92 amu. Calculate the molecular formula of dinitrogen tetroxide.

1.6 Determining empirical formulas

We said that the formula of a chemical that takes into account the correct number of atoms in a molecule is the molecular formula and therefore the real molecular weight of a chemical comes from these formulas. Empirical formulas are obtained from experiments in which a chemical is fragmented and analyzed so that the elements in the molecule and the mass percentage of each element is determined. Molecular formulas are obtained by using the molecular weight of the chemical and the empirical formula. Mind that the formula of a chemical that takes into account the correct number of atoms in a molecule is the molecular formula and therefore the real molecular weight of a chemical comes from these formulas. Let us work on an example in order to learn the procedure of obtaining molecular formulas.

Calculating molecular formulas By means of an experiment, we want to calculate the empirical formula of a chemical given that the chemical contains 2.8 g of nitrogen and 6.4 g of oxygen. In order to calculate the EF we will set up a table like the one presented below.

Empirical Formula Calculation		
	N	O
Grams	2.8g	6.4g
AW	14	16
Grams/AW	0.2	0.4
∇• by smallest	1	2
Formula	N ₁ O ₂ =NO ₂	

In each column we will add each of the elements that form the molecule. In the first row we will include the grams of each element (sometimes this information is given in terms of mass %), in the second we will divide the grams of each element by its atomic weight (AW(N)=14amu,AW(O)=16amu). Among all numbers of the second row (in this example 0.2 and 0.4), we will select the smallest number (0.2). Once we have the smallest, we will divide all numbers by the smallest and that will give us round numbers (1 and 2); these will be the numbers in an empirical formula: NO₂.

Sample Problem 9

The mass percentage composition of a compound is: 18.59% O, 37.25% S, and 44.16% F. Calculate its empirical formula.

SOLUTION

We will set up the the molecular formula table, knowing that the percentage are mass percentages, that is the mass of each element in the chemical, hence they should go in the grams row. Also the atomic weights of O, S and F are 16, 32 and 19 amu.

Empirical Formula Calculation			
	O	S	F
Grams	0.1859g	0.3725g	0.4416g
AW	16	32	19
Grams/AW	0.0116	0.0116	0.0232
∇• by smallest	1	1	2
Formula	OSF ₂		

STUDY CHECK

What is the empirical formula of a compound if a sample contains 10.28 g of C, 1.71 H and 12.71 g of oxygen?

CHAPTER 1

THE PERIODIC TABLE

1.1 Select from below the atomic symbol for the element Gold is: (a) Go (b) Au (c) G (d) Ca (e) Ol

1.2 The atomic symbol for aluminum is: (a) Al (b) Am (c) A (d) Sn (e) Ag

1.3 The atomic symbol for iron is: (a) Ir (b) Fs (c) Fe (d) In (e) Ir

1.4 Ca is the symbol for: (a) Carbon (b) Calcium (c) Cobalt (d) Copper (e) Cadmium

1.5 Which of the following elements is a metal? (a) Nitrogen (b) Lithium (c) Calcium (d) Iron (e) Iodine

1.6 Which of the following elements is an alkaline metal? (a) Nitrogen (b) Lithium (c) Calcium (d) Iron (e) Ruthenium

1.7 Which of the following elements is a nonmetal? (a) Nitrogen (b) Lithium (c) Calcium (d) Iron (e) Iodine

1.8 Which of the following elements is a halogen? (a) Nitrogen (b) Lithium (c) Calcium (d) Iron (e) Iodine

1.9 What is the symbol of the element in Period 4 and Group 2? (a) Be (b) Mg (c) Ca (d) C (e) Si

EARLY EXPERIMENTS OF THE ATOM

1.10 From the following scientist, J.J. Thomson, Robert Millikan, Henri Becquerel, and Ernest Rutherford, indicate who: (a) Discovered the element Uranium in a mineral (b) Scattered atoms on helium on a gold thin layer (c) Unsuccessfully validated the plum pudding model

1.11 From the following scientist, J.J. Thomson, Robert Millikan, Henri Becquerel, and Ernest Rutherford, indicate who: (a) Worked with cathodic rays (b) Calculate the charge-to-mass ratio of the electron (c) Worked with oil drops to calculate the electric charge of the electron

THE ATOM

1.12 In an atom, the nucleus contains: (a) an equal number of protons and electrons. (b) all the protons and neutrons (c) all the protons and electrons (d) only neutrons. (e) only protons.

1.13 The atomic number of an atom is equal to the number of: (a) nuclei (b) neutrons (c) neutrons plus protons (d) electrons plus protons (e) electrons

1.14 The mass number of an atom is equal to the number of: (a) nuclei (b) neutrons (c) neutrons plus protons (d) electrons plus protons (e) electrons

1.15 The mass number of an atom is equal to the number of: (a) electrons (b) neutrons (c) neutrons plus protons (d) protons

1.16 Consider a neutral atom with 30 protons and 34 neutrons. The atomic number of the element is: (a) 30 (b) 32 (c) 34 (d) 64 (e) 94

1.17 Consider a neutral atom with 30 protons and 34 neutrons. The mass number of the element is: (a) 30 (b) 32 (c) 34 (d) 64 (e) 94

1.18 The atomic mass of Ga is 69.72 amu. There are only two naturally occurring isotopes of gallium: ^{69}Ga , with a mass of 69.0 amu, and ^{71}Ga , with a mass of 71.0 amu. Calculate the natural abundance of the ^{69}Ga isotope.

1.19 Magnesium contains three different isotopes: magnesium-24 with an abundance of 79% and a mass of 23.9850423 amu, magnesium-25 with an abundance of 10% and a mass of 24.9858374 amu, and magnesium-26 with a mass of 25.9825937 amu. Calculate the abundance of magnesium-26 and the average atomic mass of a sample of magnesium.

AN INTRODUCTION TO MOLECULES

1.20 Calculate the molecular mass of the following molecule: CCl_2F_2

1.21 Calculate the molecular mass of the following molecule: C_4H_{10}

1.22 Calculate the molecular mass of the following molecule: $\text{C}_6\text{H}_{10}\text{O}_8$

EMPIRICAL AND MOLECULAR FORMULAS

1.23 What is the empirical formula of a compound if a sample of this compound contains 2.8 g of nitrogen and 3.2 g of oxygen?

1.24 What is the empirical formula and the molecular formula of a compound if a sample contains 3 g of C, 0.5 H and 4 g of oxygen? $\text{MW}=60\text{amu}$

1.25 What is the empirical and molecular formula of a compound with a percent composition of 49.47% C, 5.201% H, 28.84% N, and 16.48% O, if its molecular mass is 194.2 amu.

1.26 A 1.587 g sample of a compound containing N and O was analyzed finding a composition of 0.483 g of Nitrogen and 1.104 g of Oxygen. Calculate the empirical formula of the compound.

Answers 1.1 (b) 1.3 (c) 1.5 (d) 1.7 (a) 1.9 (c) 1.11 (a) J.J. Thomson (b) J.J. Thomson (c) Robert Millikan
1.13 (e) 1.15 (c) 1.17 (d) 1.19 24.3048 amu 1.21 58 amu 1.23 NO