

Abstract Algebra by Pinter, Chapter 18

Amir Taaki

Abstract

Chapter 18 on Ideals

Contents

1	A. Examples of Subrings	2
1.1	Q1	2
1.2	Q2	3
1.3	Q3	3
1.4	Q4	3
1.5	Q5	3
1.6	Q6	3
2	B. Examples of Ideals	3
2.1	Q1	3
2.1.1	$\{(n, n) : n \in \mathbb{Z}\}$	3
2.1.2	$\{(5n, 0) : n \in \mathbb{Z}\}$	3
2.1.3	$\{(n, m) : n + m \text{ is even}\}$	4
2.1.4	$\{(2n, 3m) : n, m \in \mathbb{Z}\}$	4
2.2	Q2	4
2.3	Q3	4
2.4	Q4	4
2.5	Q5	4
2.6	Q6	4
2.6.1	a	4
2.6.2	b	4
2.7	Q7	5
2.8	Q8	5
2.9	Q9	5
3	C. Elementary Properties of Subrings	5
3.1	Q1	5
3.2	Q2	5
3.3	Q3	5
3.4	Q4	5
3.5	Q5	5
3.6	Q6	5
3.7	Q7	5
3.7.1	a	5
3.7.2	b	6
3.8	Q8	6
3.9	Q9	6
4	D. Elementary Properties of Ideals	6
4.1	Q1	6
4.2	Q2	6
4.3	Q3	6
4.4	Q4	7
4.5	Q5	7
4.6	Q6	7

5	E. Examples of Homomorphisms	7
5.1	Q1	7
5.2	Q2	7
5.3	Q3	7
5.4	Q4	8
5.5	Q5	8
5.6	Q6	8
5.7	Q7	9
6	F. Elementary Properties of Homomorphisms	9
6.1	Q1	9
6.2	Q2	9
6.3	Q3	9
6.4	Q4	10
6.5	Q5	10
6.6	Q6	10
6.7	Q7	11
7	G. Examples of Isomorphisms	11
7.1	Q1	11
	7.1.1 Addition	11
	7.1.2 Multiplication	11
7.2	Q2	11
	7.2.1 Addition	11
	7.2.2 Multiplication	11
7.3	Q3	12
7.4	Q4	12
7.5	Q5	12
8	H. Further Properties of Ideals	12
8.1	Q1	12
8.2	Q2	12
8.3	Q3	13
8.4	Q4	13
8.5	Q5	13
9	I. Further Properties of Homomorphisms	13
9.1	Q1	13
9.2	Q2	14
9.3	Q3	14
9.4	Q4	14
9.5	Q5	14
10	J. A Ring of Endomorphisms	14
10.1	Q1	14
10.2	Q2	14
10.3	Q3	14
10.4	Q4	14
10.5	Q5	15
10.6	Q6	15

1 A. Examples of Subrings

1.1 Q1

$$\{x + \sqrt{3}y : x, y \in \mathbb{Z}\}$$

Closed wrt subtraction

$$(x + \sqrt{3}y)(v + \sqrt{3}w) = xv + \sqrt{3}(yv + xw) + 3yw \in \mathbb{R}$$

Thus it's a subring.

1.2 Q2

As before, it's closed under subtraction and multiplication.

1.3 Q3

$$\{x2^y : x, y \in \mathbb{Z}\}$$

Closed under multiplication because:

$$x_1 2^{y_1} \cdot x_2 2^{y_2} = (x_1 x_2) 2^{y_1 + y_2}$$

Also contains negatives since $x \in \mathbb{Z}$.

To show closure under addition is trivial for positive powers since

$$x2^y + v2^w = x2^{(y-w)}2^w + v2^w = (x2^{(y-w)} + v)2^w$$

Now for the negative case, assume $y > w$, hence $y - w$ is positive and the formulation still holds.

1.4 Q4

The sum and product of continuous functions are continuous.

1.5 Q5

The sum and product on any interval $[0, 1]$ also remains continuous, and hence also includes \mathcal{C}

1.6 Q6

Addition and negatives remain in $\mathcal{M}_2(\mathbb{R})$ as does multiplication

$$\begin{pmatrix} 0 & 0 \\ 0 & x \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & y \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & xy \end{pmatrix}$$

2 B. Examples of Ideals

2.1 Q1

Identify which of the following are ideals of $\mathbb{Z} \times \mathbb{Z}$

2.1.1 $\{(n, n) : n \in \mathbb{Z}\}$

$$\begin{aligned} (n, n) + (m, m) &= (m + n, m + n) \in I \\ -(n, n) &= (-n, -n) \in I \\ (n, n) \cdot (a, b) &= (na, nb) \notin I \end{aligned}$$

Not an ideal.

2.1.2 $\{(5n, 0) : n \in \mathbb{Z}\}$

$$\begin{aligned} (5m, 0) + (5n, 0) &= (5(m + n), 0) \in I \\ -(5n, 0) &= (5(-n), 0) \in I \\ (5n, 0) \cdot (a, b) &= (5(na), 0) \in I \end{aligned}$$

Is an ideal.

2.1.3 $\{(n, m) : n + m \text{ is even}\}$

$$\begin{aligned}(n_1, m_1) + (n_2, m_2) &= (n_1 + n_2, m_1 + m_2) \in I \\ -(n, m) &\in I \\ (n, m) \cdot (a, b) &= (na, mb)\end{aligned}$$

na is even and mb is even, so $na + mb$ is even so $(na, mb) \in I$.

Is an ideal.

2.1.4 $\{(2n, 3m) : n, m \in \mathbb{Z}\}$

$$\begin{aligned}(2n_1, 3m_1) + (2n_2, 3m_2) &= (2(n_1 + n_2), 3(m_1 + m_2)) \in I \\ -(2n, 3m) &= (2(-n), 3(-m)) \in I \\ (2n, 3m) \cdot (a, b) &= (2na, 3mb) \in I\end{aligned}$$

Is an ideal

2.2 Q2

List all the ideals of \mathbb{Z}_{12}

$\mathbb{Z}_{12} = \langle 1 \rangle$ and is cyclic. All subgroups are also cyclic.

- $\mathbb{Z}_{12} = \langle 1 \rangle = \langle 5 \rangle = \langle 7 \rangle = \langle 11 \rangle$ because $\gcd(m, 12) = 1$
- $\langle 4 \rangle = \langle 8 \rangle, \langle 4 \rangle = \{4, 8, 0\}$
- $\langle 3 \rangle = \langle 9 \rangle, \langle 3 \rangle = \{3, 6, 9, 0\}$
- $\langle 2 \rangle = \langle 10 \rangle, \langle 2 \rangle = \{2, 4, 6, 8, 10, 0\}$
- $\langle 6 \rangle = \{6, 0\}$
- $\langle 0 \rangle = \{0\}$

Let $m \in \bar{m} = \langle m \rangle$, then $\langle m \rangle = \{mj : j \in \mathbb{Z}_{12}\}$

Let $x \in \langle m \rangle$ and $y \in \mathbb{Z}_{12}$, since $x \in \langle m \rangle$, then $x = mj$ for some $j \in \mathbb{Z}_{12}$, thus $xy = m.jy$, thus $\langle m \rangle$ is an ideal of \mathbb{Z}_{12} .

Ideals are $\langle 0 \rangle, \langle 1 \rangle, \langle 2 \rangle, \langle 3 \rangle, \langle 4 \rangle, \langle 6 \rangle$

2.3 Q3

See previous exercise

2.4 Q4

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & x \end{pmatrix} = \begin{pmatrix} 0 & x \\ 0 & x \end{pmatrix} \notin \mathcal{M}_2(\mathbb{R})$$

2.5 Q5

The product of a continuous and non-continuous function are non-continuous, hence $\mathcal{C}(\mathbb{R})$ is not an ideal of $\mathcal{F}(\mathbb{R})$

2.6 Q6

2.6.1 a

Assume he means multiplication here.

$$f(x) \cdot g(x) = 0 \quad \forall x \in \mathbb{Q}$$

Thus $f \cdot g \in I$

2.6.2 b

Likewise $f(0)g(0) = 0g(0) = 0$, so $f \cdot g \in I$

2.7 Q7

Ideals of P_3 such that $AB = A \cap B \in I$. See also 17D5

$$P_3 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$$

Any subgroup must contain \emptyset .

$A + A = \emptyset$ so A is its own negative.

$\{\emptyset, \{a\}\}, \{\emptyset, \{b\}\}, \{\emptyset, \{c\}\}$ are all ideals since $\{a\}\{a, c\} = \{a\}$ and $\{a\}\{b, c\} = \emptyset$.

Likewise $\{\emptyset, \{a\}, \{c\}, \{a, c\}\}, \{\emptyset, \{a\}, \{b\}, \{a, b\}\}, \{\emptyset, \{b\}, \{c\}, \{b, c\}\}$ since $\{a, c\}\{b, c\} = \{c\}$

Lastly we have P_3 itself

2.8 Q8

Example of a non-ideal subring is $\{\emptyset, \{a, c\}\}$ which is closed under addition, negatives and multiplication.

2.9 Q9

$$A = \langle (1, 1) \rangle = \{(0, 0), (1, 1), (2, 2)\}$$

3 C. Elementary Properties of Subrings

3.1 Q1

Let $x \in B$ and since B is a ring then $0 \in B$, thus $0 - x = -x \in B$.

So B is closed wrt negatives and hence addition since $x - (-y) = x + y \in B$

3.2 Q2

As per part 1

3.3 Q3

A ring is a group under addition. Hence order of a subring divides ring by Lagrange.

3.4 Q4

A has no zero divisors, hence neither does $B \implies B$ is an integral domain.

3.5 Q5

B is a subring of field F . Let $b \in B, b \neq 0$, then $b^{-1} \in F$ (because F is a field and contains inverses). Every field is an integral domain, hence so is B .

3.6 Q6

F is a commutative ring with inverses and unity.

Since B is a subring, it also is commutative.

Since B also contains inverses and is closed wrt multiplication, it must contain 1_F .

Thus B is a field.

3.7 Q7

3.7.1 a

$$B = \langle 2 \rangle = \{0, 2, 4, \dots, 16\}$$

3.7.2 b

$$B = \langle 9 \rangle = \{0, 9\}$$

3.8 Q8

$$f(e) = e$$

$$f(x_1 + x_2) = f(x_1) + f(x_2)$$

$$f(x_1 x_2) = f(x_1) f(x_2)$$

But $\forall x \in B \quad f(x) = x$

$$x_1, x_2 \in B$$

$$x_1 + x_2 = f(x_1) + f(x_2) = f(x_1 + x_2)$$

Likewise for multiplication.

Since A is a ring $\forall -x \in A$ st $x + (-x) = e$ but $x \in B$

$$f(x) + f(-x) = f(e) = f(x + (-x)) = x + (-x)$$

Hence $-x \in B$ also.

3.9 Q9

$$ax = xa \quad bx = xb$$

$$(a + b)x = x(a + b)$$

So $a + b$ also is in the center.

$$(ab)x = axb = x(ab)$$

Finally $0x = 0 = x0$

$$-a \in A$$

$$-ax = -(ax) = -(xa) = -xa$$

By associativity.

4 D. Elementary Properties of Ideals

4.1 Q1

Explain why J is an ideal of A iff J is closed with respect to subtraction and J absorbs products in A .

$$0 - x = -x \in J$$

$$x - (-y) = x + y \in J$$

So J is closed wrt negatives and addition from the statement about subtraction.

4.2 Q2

If A is a ring with unity, prove that J is an ideal of A iff J is closed with respect to addition and J absorbs products in A .

Note that A is a ring with unity, and by definition must include -1 .

Then note that since J absorbs products, that $(-1) \cdot a = -a \in J$.

4.3 Q3

Prove that the intersection of any two ideals of A is an ideal of A .

1. Since $x, y \in I_j$, and I_j is an ideal, $x - y \in I_j, \forall j \in J$. Therefore $x - y \in \bigcap_{j \in J} I_j = I$.
2. Since $x \in I_j, rx \in I_j, \forall j \in J$. Therefore $rx \in I$.

4.4 Q4

Prove that J is an ideal of A and $1 \in J$, then $J = A$.

Since ideals absorb products, then if $1 \in J$, then since $a \cdot 1 = a \in J$, then $J = A$.

4.5 Q5

Prove that if J is an ideal of A and J contains an invertible element a of A , then $J = A$.

$$a \cdot a^{-1} = 1 \in J$$

By previous exercise $J = A$.

4.6 Q6

Explain why a field F can have no nontrivial ideals.

Every nonzero element of a field is invertible. Hence the only ideals are $\{0\}$ or F itself.

5 E. Examples of Homomorphisms

5.1 Q1

Let $f, g \in \mathcal{F}(\mathbb{R})$

$$\begin{aligned}\phi(f + g) &= (f + g)(0) = f(0) + g(0) = \phi(f) + \phi(g) \\ \phi(f \cdot g) &= (f \cdot g)(0) = f(0)g(0) = \phi(f)\phi(g)\end{aligned}$$

$$K = \{f \in \mathcal{F}(\mathbb{R}) : f(0) = 0\}$$

Range is $[-\infty, \infty]$.

5.2 Q2

$$h : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$$

$$h(x, y) = x$$

$$h(x_1 + x_2, y_1 + y_2) = x_1 + x_2 = h(x_1, y_1) + h(x_2, y_2)$$

$$h(x_1 x_2, y_1 y_2) = x_1 x_2 = h(x_1, y_1)h(x_2, y_2)$$

$$\begin{aligned}K &= \{x, y \in \mathbb{R} \times \mathbb{R} : h(x, y) = 0\} \\ &= \{(0, y) : y \in \mathbb{R}\}\end{aligned}$$

Range is $[-\infty, \infty]$

5.3 Q3

$$h : \mathbb{R} \rightarrow \mathcal{M}_2(\mathbb{R})$$

$$h(x) = \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix}$$

$$h(x + y) = \begin{pmatrix} x + y & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} y & 0 \\ 0 & 0 \end{pmatrix} = h(x) + h(y)$$

$$h(xy) = \begin{pmatrix} xy & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} y & 0 \\ 0 & 0 \end{pmatrix} = h(x)h(y)$$

$$K = \{0\}$$

Range is

$$\begin{pmatrix} \pm\infty & 0 \\ 0 & 0 \end{pmatrix}$$

5.4 Q4

$$h : \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{M}_2(\mathbb{R})$$

$$h(x, y) = \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix}$$

$$\begin{aligned} h(x_1 + x_2, y_1 + y_2) &= \begin{pmatrix} x_1 + x_2 & 0 \\ 0 & y_1 + y_2 \end{pmatrix} \\ &= \begin{pmatrix} x_1 & 0 \\ 0 & y_1 \end{pmatrix} + \begin{pmatrix} x_2 & 0 \\ 0 & y_2 \end{pmatrix} \\ &= h(x_1, y_1) + h(x_2, y_2) \end{aligned}$$

$$K = \{(0, 0)\}$$

Range is

$$\begin{pmatrix} \pm\infty & 0 \\ 0 & \pm\infty \end{pmatrix}$$

5.5 Q5

$$f : \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{M}_2(\mathbb{R})$$

$$f(x, y) = \begin{pmatrix} x & 0 \\ y & 0 \end{pmatrix}$$

$$\begin{aligned} f(x_1 + x_2, y_1 + y_2) &= \begin{pmatrix} x_1 + x_2 & 0 \\ y_1 + y_2 & 0 \end{pmatrix} \\ &= \begin{pmatrix} x_1 & 0 \\ y_1 & 0 \end{pmatrix} + \begin{pmatrix} x_2 & 0 \\ y_2 & 0 \end{pmatrix} \\ &= f(x_1, y_1) + f(x_2, y_2) \end{aligned}$$

$$f((x_1, y_1) \otimes (x_2, y_2)) = f(x_1 x_2, y_1 x_2) = \begin{pmatrix} x_1 x_2 & 0 \\ y_1 x_2 & 0 \end{pmatrix}$$

$$f(x_1, y_1) f(x_2, y_2) = \begin{pmatrix} x_1 & 0 \\ y_1 & 0 \end{pmatrix} \begin{pmatrix} x_2 & 0 \\ y_2 & 0 \end{pmatrix} = \begin{pmatrix} x_1 x_2 & 0 \\ y_1 x_2 & 0 \end{pmatrix}$$

$$K = \{(0, 0)\}$$

5.6 Q6

$$h : P_C \rightarrow P_C$$

$$h(A) = A \cap D$$

$$D \subseteq C$$

$$\begin{aligned} h(A + B) &= h((A - B) \cup (B - A)) \\ &= [(A - B) \cup (B - A)] \cap D \\ &= [(A - B) \cap D] \cup [(B - A) \cap D] \\ &= [A \cap D - B \cap D] \cup [B \cap D - A \cap D] \\ &= h(A) + h(B) \\ h(AB) &= h(A \cap B) = A \cap B \cap D \\ &= (A \cap D) \cap (B \cap D) \\ &= h(A)h(B) \end{aligned}$$

$$K = \{A \in P_C : A \cap D = \emptyset\}$$

Range is every subset of D .

5.7 Q7

Rules for ring homomorphisms:

$$f(a + b) = f(a) + f(b) \quad f(ab) = f(a)f(b)$$

$$f(0) = 0 \quad f(1_A) = 1_B$$

$$f(n) = f(1 + \cdots + 1) = f(1) + \cdots + f(1) = nf(1)$$

$$mf(1) = 0 \quad f(1)^2 = f(1)$$

Homomorphisms for $\phi_i : \mathbb{Z}_2 \rightarrow \mathbb{Z}_4$

$$\phi_e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

The other mappings do not work:

- $1 \rightarrow 1$ then $2f(1) \neq 0$
- $1 \rightarrow 2$ then $f(1)^2 \neq f(1)$
- $1 \rightarrow 3$ then $f(1)^2 \neq f(1)$

Homomorphisms for $\phi_i : \mathbb{Z}_2 \rightarrow \mathbb{Z}_4$

$$\phi_e = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\phi_a = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 4 & 2 \end{pmatrix}$$

The other mappings do not work:

- $1 \rightarrow 1$ then $f(1)^2 \neq f(1)$
- $1 \rightarrow 2$ then $f(1)^2 \neq f(1)$
- $1 \rightarrow 3$ then $3f(1) \neq 0$
- $1 \rightarrow 5$ then $3f(1) \neq 0$

6 F. Elementary Properties of Homomorphisms

6.1 Q1

Prove $f(A) = \{f(x) : x \in A\}$ is a subring of B .

Since f is a homomorphism, ring operations are obeyed in the homomorphism. For negatives we note that $f(0_A) = 0_B = 1_B + (-1_B)$ and every negative is expressible as $(-1_B) \cdot a$ where $a \in B$.

6.2 Q2

Prove the kernel of f is an ideal of A .

$$K = \{x \in A : f(x) = 0_B\}$$

From f being a homomorphism, we conclude K is a subring of A .

To show it's an ideal, for any $a \in A$ and $x \in K$, then $f(ax) = 0 = f(x)$. So K absorbs the product ax .

Thus the kernel of a homomorphism is an ideal of the input ring.

6.3 Q3

Prove $f(0) = 0$, and for every $a \in A$, $f(-a) = -f(a)$.

$$f(0) = f(0 + 0) = f(0) + f(0) \implies f(0) = 0$$

$$f(a + (-a)) = f(a) + f(-a) = f(0) = 0 = f(a) - f(a)$$

$$\implies f(-a) = -f(a)$$

6.4 Q4

Prove f is injective iff its kernel is equal to $\{0\}$.

$$f(x) = f(y) \iff f(y - x) = 0 \iff y - x \in K$$

Let $x \in K$

$$\implies f(x) = 0$$

$$\implies f(x) = f(0) \quad [\text{since } f(0) = 0]$$

$$\implies x = 0 \quad [\text{since } f \text{ is injective}]$$

It follows $K = \{0\}$

Thus f is injective $\implies K = \{0\}$

Now suppose $K = \{0\}$. Then

$$f(x) = f(y)$$

$$\implies f(x) - f(y) = 0$$

$$\implies f(x - y) = 0$$

$$\implies x - y \in K$$

$$\implies x - y = 0 \quad [\text{since } K = \{0\}]$$

$$\implies x = y$$

Hence f is injective.

Thus $K = \{0\} \implies f$ is injective.

Hence f is injective $\iff K = \{0\}$

6.5 Q5

If B is an integral domain, then either $f(1) = 1$ or $f(1) = 0$. If $f(1) = 0$ then $f(x) = 0$ for every $x \in A$. If $f(1) = 1$, the image of every invertible element of A is an invertible element of B .

Integral domain has the cancellation property such that $ab = ac \implies b = c$.

$$f(1) = f(1 \cdot 1) = f(1)f(1)$$

$$f(1) = f(1)f(1)$$

$$f(1) = 0 \text{ or } 1$$

$$\text{If } f(1) = 0 \text{ then } \forall a \in A, f(a) = f(1 \cdot a) = f(1)f(a) = 0$$

$$\text{If } f(1) = 1 \text{ and } \exists x, y \in A \text{ such that } xy = 1$$

$$f(xy) = f(x)f(y) \text{ where } f(y) = (f(x))^{-1}$$

6.6 Q6

Any homomorphic image of a commutative ring is a commutative ring. Any homomorphic image of a field is a field.

Let $a, b \in A$, then $f(a)f(b) = f(b)f(a)$ because $f(ab) = f(ba)$.

If A is a field, then $\forall x \in A, \exists x^{-1} \in A$. So by the last exercise, $f(x^{-1}) = (f(x))^{-1}$ and so the inverse of $f(x)$ is a member of B .

$$(f(x))^{-1} \in B$$

6.7 Q7

If the domain A of the homomorphism f is a field, and if the range of f has more than one element, then f is injective.

Since A is a field, the kernel of A is either $\{0\}$ or A itself.

But the range of f is more than one element, so the kernel of A cannot be A and must be $\{0\}$.

Since the kernel of f is $\{0\}$, then f is injective.

7 G. Examples of Isomorphisms

7.1 Q1

$$a \oplus b = a + b + 1$$

$$a \otimes b = ab + a + b$$

7.1.1 Addition

$$f(a + b) = a + b - 1$$

$$\begin{aligned} f(a) \oplus f(b) &= (a - 1) + (b - 1) - 1 \\ &= a + b - 1 \end{aligned}$$

7.1.2 Multiplication

$$f(ab) = ab - 1$$

$$\begin{aligned} f(a) \otimes f(b) &= (a - 1)(b - 1) + (a - 1) + (b - 1) \\ &= ab - b - a + 1 + a + b - 1 - 1 \\ &= ab - 1 \end{aligned}$$

7.2 Q2

$$\mathcal{J} = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} : a, b \in \mathbb{R} \right\}$$

$$f : \mathbb{C} \rightarrow \mathcal{J}$$

$$f(a + bi) = \begin{pmatrix} a & b \\ -b & a \end{pmatrix}$$

$$a + bi = c + di \implies f(a + bi) = f(c + di)$$

7.2.1 Addition

$$f((a + bi) + (c + di)) = \begin{pmatrix} a + c & b + d \\ -(b + d) & a + c \end{pmatrix}$$

$$f(a + bi) + f(c + di) = \begin{pmatrix} a + c & b + d \\ -(b + d) & a + c \end{pmatrix}$$

7.2.2 Multiplication

$$f((a + bi)(c + di)) = f((ac - bd) + (ad + bc)i)$$

$$= \begin{pmatrix} a + c & b + d \\ -(b + d) & a + c \end{pmatrix}$$

$$f(a + bi)f(c + di) = \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \begin{pmatrix} c & d \\ -d & c \end{pmatrix}$$

$$= \begin{pmatrix} a + c & b + d \\ -(b + d) & a + c \end{pmatrix}$$

7.3 Q3

$$A = \{(x, x) : x \in \mathbb{Z}\}$$

$$\forall x, y \in \mathbb{Z}, (x, x) \in A, (y, y) \in A, (x + y, x + y) \in A \text{ and } (xy, xy) \in A$$

Thus A is a subring of $\mathbb{Z} \times \mathbb{Z}$

The homomorphism $f : \mathbb{Z} \rightarrow A$ by $f(x) = (x, x)$ is isomorphic because it is one to one

$$f(x) = f(y) \implies x = y$$

and onto

$$\forall (x, x) \in A, \exists x \in \mathbb{Z} \text{ such that } f(x) = (x, x)$$

Thus

$$\{(x, x) : x \in \mathbb{Z}\} \cong \mathbb{Z}$$

7.4 Q4

Addition and negatives trivially remain inside the set.

$$\begin{pmatrix} 0 & 0 \\ 0 & x \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & y \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & xy \end{pmatrix}$$

Hence the set is a subring.

$$A = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & x \end{pmatrix} : x \in \mathbb{R} \right\}$$

Define $f : \mathbb{R} \rightarrow A$ by $f(x) = \begin{pmatrix} 0 & 0 \\ 0 & x \end{pmatrix}$, then f is an homomorphism from \mathbb{R} to A .

Hence $A \cong \mathbb{R}$

7.5 Q5

$$f : k\mathbb{Z} \rightarrow l\mathbb{Z}$$

$$f(k) = ln \text{ for some } n \neq 0$$

$$\begin{aligned} f(k^2) &= l^2 n^2 \\ &= f(k \cdot k) = f(k + \dots + k) = kf(k) \\ &= kln \\ k &= ln \end{aligned}$$

But $k \neq l$, so ln does not generate $l\mathbb{Z}$ and f is not an isomorphism.

8 H. Further Properties of Ideals

8.1 Q1

If $J \cap K = \{0\}$, then $jk = 0$ for every $j \in J$ and $k \in K$.

J and K are ideals, so for every $j \in J$ and $k \in K$, then $jk \in J$ and $jk \in K$, so $jk \in J \cap K$.

8.2 Q2

For any $a \in A$, $I_a = \{ax + j + k : x \in A, j \in J, k \in K\}$ is an ideal of A .

$\forall i \in I_a$, and $b \in A$, then $bi = b(ax + j + k) = a(bx) + bj + bk \in I_a$ since J and K are ideals and $bx \in A$.

8.3 Q3

The radical of J is the set $\text{rad } J = \{a \in A : a^n \in J \text{ for some } n \in \mathbb{Z}\}$. For any ideal J , $\text{rad } J$ is an ideal of A .

$$\begin{aligned} a^n \in J & \quad b^m \in J \\ (a+b)^{n+m} \in J & \quad [\text{see 17m3}] \end{aligned}$$

$x \in A$ and $a \in \text{rad } J$ then $(xa)^n = x^n a^n \in J$, so $xa \in \text{rad } J$.

$a, b \in \text{rad } J$, then $(a+b)^{m+n} \in J$ and so $a+b \in \text{rad } J$.

8.4 Q4

For any $a \in A$, $\{x \in A : ax = 0\}$ is an ideal (called the annihilator of a).

Furthermore, $\{x \in A : ax = 0 \text{ for every } a \in A\}$ is an ideal (called the annihilating ideal of A). If A is a ring with unity, its annihilating ideal is equal to $\{0\}$.

Let $b \in A$, then $bx \in \text{Ann}(a)$ because $ax = 0$ so $b(ax) = bxa = 0$.

Let $x, y \in \text{Ann}(a)$ then $a(x+y) = 0$ so $x+y \in \text{Ann}(a)$.

$$I = \{x \in A : ax = 0 \text{ for every } a \in A\}$$

If A is a ring with unity then $a = 1 \implies x = 0$ so $I = \{0\}$.

8.5 Q5

Show that $\{0\}$ and A are ideals of A . (They are trivial ideals; every other ideal of A is a proper ideal.) A proper J of A is called maximal if it is not strictly contained in any strictly larger proper ideal: that is if $J \subseteq K$, where K is an ideal containing some element not in J , then necessarily $K = A$.

Show the following is an example of a maximal ideal: in $\mathcal{F}(\mathbb{R})$, the ideal $J = \{f : f(0) = 0\}$.

$$g \in K \quad g(0) \neq 0 \quad g \notin J$$

$$\begin{aligned} h(x) &= g(x) - g(0) \in J \\ h(x) - g(x) &\in K \end{aligned}$$

Continuous function with a nonzero value is invertible.

$h(x) - g(x) = -g(0) \in K$ but $g(0) \neq 0$ so $-1/g(0) \in A$.

But since K is an ideal, that is

$$g(0) \cdot 1/g(0) \in K$$

but this equals 1, and $1 \in K$ so $K = A$ and is maximal.

9 I. Further Properties of Homomorphisms

9.1 Q1

If $f : A \rightarrow B$ is a homomorphism from A onto B with kernel K , and J is an ideal of A such that $K \subseteq J$, then $f(J)$ is an ideal of B .

f is onto $\exists x : f(x) = y$ so it's an ideal. Closed under addition and negatives and absorbs products.

See also [here](#)

9.2 Q2

If $f : A \rightarrow B$ is a homomorphism from A onto B , and B is a field, then the kernel of f is a maximal ideal.

The kernel K is a subset of the ideal for A . As shown above $f(J)$ is an ideal of B , which by D6 can only be $\{0\}$ or B itself. Since the homomorphism is onto, then $f(A)$ maps to B , but A is a trivial ideal of A . Thus K , the kernel of f is the proper ideal for A which maps to $\{0\}$ in B .

9.3 Q3

There are no nontrivial homomorphisms from \mathbb{Z} to \mathbb{Z} .

$$\begin{aligned}
f(1) &= f(1 \cdot 1) = f(1) \cdot f(1) \\
f(1) &= 1 \text{ or } f(1) = 0 \\
f(n) &= f(1 + \cdots + 1) = f(1) + \cdots + f(1) = nf(1)
\end{aligned}$$

So $f(n) = n$ or $f(n) = 0$

See also [here](#) and [here](#)

9.4 Q4

If n is a multiple of m , then \mathbb{Z}_m is a homomorphic image of \mathbb{Z}_n .

$f : \mathbb{Z}_n \rightarrow \mathbb{Z}_m$ by $f(a) = a(\text{mod } m)$ obeys the homomorphic properties.

See also [here](#)

9.5 Q5

If n is odd, there is an injective homomorphism from \mathbb{Z}_2 into \mathbb{Z}_{2n} .

$$f(x) = nx$$

Above homomorphism is injective since $f(0) = 0$ and $f(1) = n$.

10 J. A Ring of Endomorphisms

10.1 Q1

$$\begin{aligned}
\pi_a(x) &= ax \\
\pi_a(x + y) &= a(x + y) = ax + ay = \pi_a(x) + \pi_a(y)
\end{aligned}$$

10.2 Q2

$$\pi_a(x) = \pi_a(y) \implies x = y$$

a is not a divisor of zero $\implies \forall x \in A, ax \neq 0$, thus ring A has cancellation property

$$\pi_a(x) = \pi_a(y) = ax = ay \implies x = y$$

10.3 Q3

If a is invertible then $\forall y \in A, y = a(a^{-1}y)$ so $x = a^{-1}y, f(x) = y$, thus π_a is surjective.

10.4 Q4

$$\begin{aligned}
\mathcal{A} &= \{\pi_a : a \in A\} \\
[\pi_a + \pi_b](x) &= \pi_a(x) + \pi_b(x) \\
\pi_a \pi_b &= \pi_a \cdot \pi_b
\end{aligned}$$

1. Addition is abelian
2. Multiplication is associative: $(\pi_a \cdot \pi_b \cdot \pi_c)(x) = (abc)x = a(bcx) = \pi_a((\pi_b \cdot \pi_c)(x))$
3. Distributive over addition

10.5 Q5

$\phi : A \rightarrow \mathcal{A}$ given by $\phi(a) = \pi_a$

As shown above this is homomorphic.

10.6 Q6

$$\phi(a) = \phi(b) \implies \pi_a = \pi_b$$

$$\pi_a(1) = \pi_b(1) \implies a = b$$

$\forall \pi_a \in \mathcal{A}, \exists a \in A : \pi_a = \phi(a)$ by definition.

If a has no divisors of zero, then to show injective property, note that

$$ax = bx \implies a = b$$

$$\pi_a = \pi_b \implies \pi_a(x) = ax = \pi_b(x) = bx \implies a = b$$

From the cancellation property since it has no divisors of zero.