



Residues and duality

June 22, 2023

Contents



1	one inch + \hoffset	2	one inch + \voffset
3	\oddsidemargin = 35pt	4	\topmargin = 22pt
5	\headheight = 12pt	6	\headsep = 18pt
7	\textheight = 550pt	8	\textwidth = 345pt
9	\marginparsep = 7pt	10	\marginparwidth = 125pt
11	\footskip = 25pt		\marginparpush = 5pt (not shown)
	\hoffset = 0pt		\voffset = 0pt
	\paperwidth = 614pt		\paperheight = 794pt



1	one inch + \hoffset	2	one inch + \voffset
3	\oddsidemargin = 35pt	4	\topmargin = 22pt
5	\headheight = 12pt	6	\headsep = 18pt
7	\textheight = 550pt	8	\textwidth = 345pt
9	\marginparsep = 7pt	10	\marginparwidth = 125pt
11	\footskip = 25pt		\marginparpush = 5pt (not shown)
	\hoffset = 0pt		\voffset = 0pt
	\paperwidth = 614pt		\paperheight = 794pt

Chapter 1

Introduction

The main purpose of these notes is to prove a duality theorem for cohomology of quasi-coherent sheaves, with respect to a proper morphism of locally noetherian preschemes. Various such theorems are already known. Typical is the duality theorem for a non-singular complete curve X over an algebraically closed field k , which says that

$$h^0(D) = h^1(K - D),$$

where D is a divisor, K is the canonical divisor, and

$$h^i(D) = \dim_K H^i(X, L(D))$$

for any i , and any divisor D . (See e.g. [16, Ch. II] for a proof.)

Various attempts were made to generalize this theorem to varieties of higher dimension, and as Zariski points out in his report [20], his generalization of a lemma of Enriques-Severi [19] is equivalent to the statement that

Bibliography

- [Hartshorne1] R. Hartshorne, *Residues and duality*, Lecture Notes in Mathematics, Vol. 20, Springer-Verlag, Berlin, 1966.
- [Sato1] Mikio Sato, *Theory of hyperfunctions II*, J. Fac. Sci. Univ. Tokyo, **8** (1960), 387–437.
- [16] Jean-Pierre Serre, *Groupes algébriques et corps de classes*, Paris, Herman (1959).