

Truncated Microlocal Morse Theory

Toshihiro Oshiba

August 5, 2026*

1 Introduction

2 Background Material

In this section, we recall some definitions and results of homological algebra and sheaf theory including the theory of derived categories from [KS90], following its notations with slight modifications.

Let X be a topological space, and Z a locally closed set of X . Let us denote by $i: Z \hookrightarrow X$ the inclusion. We use the following functors:

$$\mathrm{R}\Gamma_Z(\cdot) := \mathrm{R}i_*i^!, \quad (\cdot)_Z := \mathrm{R}i_!i^{-1}.$$

If there is no risk of confusion, we will write $\mathbf{k}_Z$ rather than $\mathbf{k}_{X,Z}$.

2.1 Geometrical Notions

Let M be a real manifold of class C^∞ . We denote by $\tau_M: TM \rightarrow M$ the tangent bundle of M , and by $\pi_M: T^*M \rightarrow M$ the cotangent bundle of M . If there is no risk of confusion, we will write π rather than π_M . If $f: M \rightarrow N$ is a morphism of manifolds (i.e., a C^∞ -map), then we have the natural diagrams.

$$\begin{array}{ccccc} TM & \xrightarrow{f'} & M \times_N TN & \xrightarrow{f_\tau} & TN \\ & \searrow \tau_M & \downarrow & \square & \downarrow \tau_N \\ & & M & \xrightarrow{f} & N, \end{array} \quad \begin{array}{ccccc} T^*M & \xleftarrow{f_d} & M \times_N T^*N & \xrightarrow{f_\pi} & T^*N \\ & \searrow \pi_M & \downarrow & \square & \downarrow \pi_N \\ & & M & \xrightarrow{f} & N. \end{array}$$

In these diagrams, f_τ and f_π denote the natural projections, f' is defined by $(x; v) \mapsto (x; df_x v)$, and f_d is induced by the transposition of f' .

*First draft: 2026/03/29

2.2 Approximation by Inductive Limits

Proposition 2.1 ([KS90, Proposition 2.5.1]). Let X be a topological space, Z a subspace, and F a sheaf on X . Consider the canonical morphism:

$$\psi: \varinjlim_U \Gamma(U; F) \rightarrow \Gamma(Z; F),$$

where U ranges through the family of open neighborhoods of Z in X . Then the following hold.

- (i) The morphism ψ is injective.
- (ii) Assume that X is Hausdorff and Z is compact. Then ψ is an isomorphism.
- (iii) Assume that X is paracompact and Z is closed. Then ψ is an isomorphism.

Note that a paracompact space is, by definition, a Hausdorff space.

2.3 Complement of Homological Algebra

Consider a projective system of abelian groups $(M_n, \rho_{n,p})_n$ indexed by $\mathbf{N}$. One says that this system satisfies the Mittag-Leffler condition (ML condition for short) if for any $n \in \mathbf{N}$, the decreasing sequence $(\rho_{n,p}(M_p))_p$ of subgroups of M_n is stationary. For a projective system of abelian groups $(M_n)_n$, we set

$$M_\infty := \varprojlim_n M_n.$$

Consider a projective system of complexes

$$(2.1) \quad E_n^\bullet: \cdots \rightarrow M_n^{j-1} \rightarrow M_n^j \rightarrow M_n^{j+1} \rightarrow \cdots,$$

and its projective limit

$$(2.2) \quad E_\infty^\bullet: \cdots \rightarrow M_\infty^{j-1} \rightarrow M_\infty^j \rightarrow M_\infty^{j+1} \rightarrow \cdots.$$

Denote by

$$(2.3) \quad \Phi_k: H^k(E_\infty^\bullet) \rightarrow \varprojlim_n H^k(E_n^\bullet)$$

the natural morphism.

Proposition 2.2 ([KS90, Proposition 1.12.4]). Assume that for each $k \in \mathbf{Z}$, the system $(M_n^k)_n$ satisfies the ML condition. Then

- (a) for each $k \in \mathbf{Z}$, the map Φ_k is surjective,
- (b) if moreover, for a given i the system $(H^{i-1}(E_n^\bullet))_n$ satisfies the ML condition, then Φ_i is bijective.

The following is called Kashiwara's lemma.

Proposition 2.3 ([KS90, Proposition 1.12.6]). Let $(X_s, \rho_{s,t})$ be a projective system of sets indexed by $\mathbf{R}$. Assume that for each $s \in \mathbf{R}$, the canonical maps

$$\lambda_s: X_s \rightarrow \varprojlim_{r < s} X_r$$

and

$$\mu_s: \varinjlim_{t > s} X_t \rightarrow X_s$$

are both injective (resp. surjective). Then all the maps ρ_{s_0, s_1} ($s_0 \leq s_1$) are injective (resp. surjective).

2.4 ML Procedure for Sheaves

We use the following in the proof of Theorem 4.2.

Proposition 2.4 ([KS90, Proposition 2.7.1]). Let X be a topological space and let $F \in \mathbf{D}^+(\mathbf{k}_X)$. Let $(U_n)_{n \in \mathbf{N}}$ be an increasing sequence of open subsets of X , $(Z_n)_{n \in \mathbf{N}}$ a decreasing sequence of closed subsets of X . Set $U := \bigcup_n U_n$, $Z := \bigcap_n Z_n$. Then the following hold.

- (i) For any j , the natural map $\varphi_j: H_Z^j(U; F) \rightarrow \varprojlim_n H_{Z_n}^j(U_n; F)$ is surjective.
- (ii) Assume that for a given j , the projective system $(H_{Z_n}^{j-1}(U_n; F))_n$ satisfies the ML condition. Then φ_j is bijective.
- (iii) Let now $(X_n)_{n \in \mathbf{N}}$ be an increasing family of subsets of X satisfying $X = \bigcup_n X_n$ and $X_n \subset \text{Int}(X_{n+1})$ for all n . Assume that for a given j , the projective system $(H^{j-1}(X_n; F))_n$ satisfies the ML condition. Then the natural map $H^j(X; F) \rightarrow \varprojlim_n H^j(X_n; F)$ is bijective.

3 Truncated Microsupport

In this section, we review the definition of a truncated microsupport and its functorial property under direct images of proper morphisms. First, we review the original definition of a microsupport.

Definition 3.1 ([KS90, Definition 5.1.2]). Let $F \in \mathbf{D}^b(\mathbf{k}_X)$. The closed conic subset $\mathrm{SS}(F)$ of T^*X is defined by following condition: $p \notin \mathrm{SS}(F)$ if and only if there exists an open neighborhood U of p such that for any $x \in \pi(U)$ and for any real-valued C^1 -function φ defined on a neighborhood of x such that $\varphi(x) = 0$, $d\varphi(x) \in U$, one has

$$(3.1) \quad \mathrm{R}\Gamma_{\{\varphi \geq 0\}}(F)_x = 0.$$

Kashiwara, Monteiro-Fernandes, and Schapira modified this definition and gave the notion of truncated microsupport.

Definition 3.2 ([KFS03a, Definition 3.4]). Let $F \in \mathbf{D}^b(\mathbf{k}_X)$, and fix an integer $k \in \mathbf{Z}$. The *truncated microsupport* $\mathrm{SS}_k(F)$ is the closed conic subset of T^*X defined by the following condition: $p \notin \mathrm{SS}_k(F)$ if and only if there exists an open conic neighborhood U of p such that for any $x \in \pi(U)$ and for any real-valued C^1 -function φ defined on a neighborhood of x such that $\varphi(x) = 0$, $d\varphi(x) \in U$, one has

$$(3.2) \quad H_{\{\varphi \geq 0\}}^j(F)_x = 0 \quad \text{for any } j \leq k.$$

For other equivalent conditions of the definition, see [KFS03a]. Truncated microsupports satisfy the following functorial property under proper direct images.

Proposition 3.3. Let $f: X \rightarrow Y$ be a morphism of manifolds and let $F \in \mathbf{D}^b(\mathbf{k}_X)$ such that f is proper on $\mathrm{supp}(F)$. Then for any $k \in \mathbf{Z}$,

$$(3.3) \quad \mathrm{SS}_k(\mathrm{R}f_*F) \subset f_{\pi}f_d^{-1}(\mathrm{SS}_k(F)).$$

Moreover if f is a closed embedding, the inclusion above is an equality.

For other functorial properties, see [KFS03a, FM07].

4 Truncated Noncharacteristic Deformation Lemma

In this section, we prove a truncated version of the noncharacteristic deformation lemma. First we state the result. After giving several examples,

we prove Theorem 4.2. In the course of the proof, we use the key lemma (Lemma 4.4). Since its proof is rather long, we postpone the proof to Section 6.

Before we state the truncated version, let us review the original version of the noncharacteristic deformation lemma.

Proposition 4.1 ([KS90, Proposition 2.7.2]). Let X be a Hausdorff space. Let $F \in \mathbf{D}^+(\mathbf{k}_X)$, and let $(U_t)_{t \in \mathbf{R}}$ be a family of open subsets of X . We assume the following conditions.

- (i) $U_t = \bigcup_{s < t} U_s$, for all $t \in \mathbf{R}$.
- (ii) For all pairs (s, t) with $s \leq t$, the set $\overline{U_t \setminus U_s} \cap \text{supp } F$ is compact.
- (iii) Setting $Z_s := \bigcap_{t > s} \overline{U_t \setminus U_s}$, we have for all pairs (s, t) with $s \leq t$ and all $x \in Z_s \setminus U_t$,
$$(\mathbf{R}\Gamma_{X \setminus U_t}(F))_x = 0.$$

Then, for all $t \in \mathbf{R}$, we have the isomorphisms

$$\mathbf{R}\Gamma \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \xrightarrow{\sim} \mathbf{R}\Gamma(U_t; F).$$

Our result is as follows.

Theorem 4.2 (Truncated Noncharacteristic Deformation Lemma). Let X be a Hausdorff space. Let $F \in \mathbf{D}^+(\mathbf{k}_X)$, and let $(U_t)_{t \in \mathbf{R}}$ be a family of open subsets of X . Let $k \in \mathbf{Z}$. For $s \in \mathbf{R}$, set $Z_s := \bigcap_{t > s} \overline{U_t \setminus U_s}$. We assume the following conditions.

- (i) $U_t = \bigcup_{s < t} U_s$, for all $t \in \mathbf{R}$.
- (ii) For all pairs (s, t) with $s \leq t$, the set $\overline{U_t \setminus U_s} \cap \text{supp } F$ is compact.
- (iii) For all $s \in \mathbf{R}$ and all $x \in Z_s \setminus U_s$, we have

$$(H_{X \setminus U_s}^j(F))_x = 0 \quad \text{for all } j \leq k.$$

- (iv) For all pairs (s, t) with $s < t$, we have $Z_s \subset U_t$.

Then, for all $t \in \mathbf{R}$, we have the isomorphisms

$$H^j \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \xrightarrow{\sim} H^j(U_t; F) \quad \text{for all } j < k,$$

and the monomorphisms

$$H^k \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \hookrightarrow H^k(U_t; F).$$

Before we prove Theorem 4.2, let us give several examples of families of deformation.

Example 4.3. The condition (iv) of Theorem 4.2 is not contained in the original version (Proposition 4.1). Therefore let us give an example of $(U_t)_{t \in \mathbf{R}}$ that does not satisfy the condition (iv).

(i) Let $X = \mathbf{R}^2$, and for a real number t define the function

$$\varphi_t(x) := \begin{cases} \max(0, t(2 - |x|)) & (|x| < 2), \\ 0 & (|x| \geq 2). \end{cases}$$

Then let us define the open subset U_t of X by

$$U_t := \{(x, y) \in X; y < \varphi_t(x)\}.$$

Then the family $(U_t)_t$ does not satisfy the condition (iv). Indeed, if $0 < s < t$, we have

$$Z_s = \{(x, y) \in X; y = \varphi_s(x), |x| \leq 2\}$$

and the point $(2, 0) \in Z_s$ is not contained in U_t . Moreover, let us define a locally closed subset S of X by

$$S := \{(x, y) \in X; (x + 2)^2 + (y - 1)^2 = 1\} \setminus \{(-1, 0)\}.$$

We denote by $\mathbf{k}_S$ the sheaf on X as in Section 2. If $j \leq 0$, we have $H_{X \setminus U_t}^j(\mathbf{k}_S)_x = 0$ for all $t \geq s$ and $x \in Z_s \setminus U_t$, but the morphism $H^0(U_1; \mathbf{k}_S) = \mathbf{k} \rightarrow 0 = H^0(U_0; \mathbf{k}_S)$ is not injective. This gives an example that satisfies the conditions (i)–(iii) but not (iv), which results in that the consequence of the proposition does not hold. Hence the condition (iv) is necessary.

Let us, in turn, give two examples of $(U_t)_{t \in \mathbf{R}}$ satisfying the condition (iv). In all the examples below, X is $\mathbf{R}^2$.

(ii) For a real number t , define the open set U_t of X by

$$U_t := \begin{cases} \{(x, y) \in X; x^2 + y^2 < t\} & (t > 0), \\ \emptyset & (t \leq 0). \end{cases}$$




Figure 1: example (i)




Figure 2: example (ii)

- (iii) For a real number t , define the function $\varphi_t: \mathbf{R} \rightarrow \mathbf{R}$ as follows. If $t \leq 0$, set $\varphi_t(x) \equiv 0$. If $t > 0$, set

$$\varphi_t(x) := \begin{cases} \sqrt{t^2 - x^2} & (|x| < t), \\ 0 & (|x| \geq t). \end{cases}$$

Then define the open set U_t of X by

$$U_t := \{(x, y) \in X; y < \varphi_t(x)\}.$$




Figure 3: example (iii)

Proof of Theorem 4.2 Consider the following conditions for $j \in \mathbf{Z}$ and

$s \in \mathbf{R}$:

$$\begin{aligned}
(a)_s^j & \quad \mu_s^j: \varinjlim_{t>s} H^j(U_t; F) \xrightarrow{\sim} H^j(U_s; F), \\
(a')_s^k & \quad \mu_s^k: \varinjlim_{t>s} H^k(U_t; F) \hookrightarrow H^k(U_s; F), \\
(b)_s^j & \quad \lambda_s^j: H^j(U_s; F) \xrightarrow{\sim} \varprojlim_{r<s} H^j(U_r; F).
\end{aligned}$$

Let us prove, for all $s \in \mathbf{R}$, $(a)_s^j$ for $j < k$ and $(a')_s^k$. We may assume that $\overline{U_t \setminus U_s}$ is compact for any $s \leq t$ by replacing X by $\text{supp } F$. Consider the distinguished triangle

$$\mathrm{R}\Gamma_{X \setminus U_s}(F) \rightarrow F \rightarrow \mathrm{R}\Gamma_{U_s}(F) \xrightarrow{+1}.$$

Let us endow with $Z_s \cup U_s$ the induced topology, and denote by $i_s: Z_s \cup U_s \hookrightarrow X$ the natural continuous map. Applying the functor

$$(\cdot)|_{Z_s \cup U_s} = i_s^{-1}$$

to the distinguished triangle above, we have

$$\mathrm{R}\Gamma_{X \setminus U_s}(F)|_{Z_s \cup U_s} \rightarrow F|_{Z_s \cup U_s} \rightarrow \mathrm{R}\Gamma_{U_s}(F)|_{Z_s \cup U_s} \xrightarrow{+1}.$$

Again applying to this triangle the functor $\mathrm{R}\Gamma(Z_s \cup U_s; \cdot)$, we have

$$(4.1) \quad \mathrm{R}\Gamma(Z_s \cup U_s; \mathrm{R}\Gamma_{X \setminus U_s}(F)|_{Z_s \cup U_s}) \rightarrow \mathrm{R}\Gamma(Z_s \cup U_s; F|_{Z_s \cup U_s}) \rightarrow \mathrm{R}\Gamma(U_s; F) \xrightarrow{+1}.$$

By the hypothesis (iii) and the definition of a local cohomology, we see that

$$H_{X \setminus U_s}^j(F)|_{Z_s \cup U_s} = 0 \quad \text{for any } j \leq k.$$

Therefore we have

$$H^j(Z_s \cup U_s; \mathrm{R}\Gamma_{X \setminus U_s}(F)|_{Z_s \cup U_s}) = 0 \quad \text{for any } j \leq k.$$

Thus we have from (4.1)

$$(4.2) \quad H^j(Z_s \cup U_s; F|_{Z_s \cup U_s}) \xrightarrow{\sim} H^j(U_s; F) \quad \text{for any } j < k,$$

$$(4.3) \quad H^k(Z_s \cup U_s; F|_{Z_s \cup U_s}) \hookrightarrow H^k(U_s; F).$$

Therefore, if we show Lemma 4.4 below, whose proof will be given in Section 6, we have from Lemma 4.4 and (4.2)

$$\begin{aligned} \varinjlim_{t>s} H^j(U_t; F) &\cong H^j(Z_s \cup U_s; F|_{Z_s \cup U_s}) \\ &\xrightarrow{\sim} H^j(U_s; F) \end{aligned}$$

for any $j < k$, and $(a)_s^j$ follows. Similarly, we have from Lemma 4.4 and (4.3)

$$\begin{aligned} \varinjlim_{t>s} H^k(U_t; F) &\cong H^k(Z_s \cup U_s; F|_{Z_s \cup U_s}) \\ &\hookrightarrow H^k(U_s; F) \end{aligned}$$

and $(a')_s^k$ follows.

Lemma 4.4. For any $j \in \mathbf{Z}$ and $s \in \mathbf{R}$, there exists a natural isomorphism

$$\varinjlim_{t>s} H^j(U_t; F) \cong H^j(Z_s \cup U_s; F|_{Z_s \cup U_s}).$$

We prove $(b)_s^j$ for all $j \in \mathbf{Z}$ and $s \in \mathbf{R}$ by induction on j . Since $F \in \mathbf{D}^+(\mathbf{k}_X)$, we have $(b)_s^j$ for $j \ll 0$ and $s \in \mathbf{R}$. Hence there exists an integer $j_0 < k$ such that $(b)_s^j$ holds for any $j < j_0$ and $s \in \mathbf{R}$. Then assume $(b)_s^j$ is proved for all $j < j_0$ and $s \in \mathbf{R}$. Since we have $(a)_s^j$ and $(b)_s^j$ for each $j < j_0$ and each $s \in \mathbf{R}$, we can apply Kashiwara's lemma (Proposition 2.3) and have

$$H^j(U_t; F) \xrightarrow{\sim} H^j(U_s; F) \quad \text{for } j < j_0 \text{ and } s \leq t.$$

Fix $s \in \mathbf{R}$ and consider the projective system $(H^{j_0-1}(U_{s-1/n}))_{n \in \mathbf{N}}$. This system satisfies ML condition. Therefore, by applying Proposition 2.4 (iii), we have

$$H^j(U_s; F) \xrightarrow{\sim} \varprojlim_n H^j(U_{s-1/n}; F).$$

Since $(s-1/n)_n$ is a cofinal family of $(t)_{s>t}$, we have $(b)_s^{j_0}$. Then by induction on j , we have $(b)_s^j$ for $j < k$ and $s \in \mathbf{R}$. Again by applying Proposition 2.4 to the projective system $(H^j(U_n; F))_n$, we obtain

$$\begin{aligned} H^j\left(\bigcup_{s \in \mathbf{R}} U_s; F\right) &\cong H^j\left(\bigcup_{n \in \mathbf{N}} U_n; F\right) \\ &\xrightarrow{\sim} \varprojlim_n H^j(U_n; F) \end{aligned}$$

and for each n , we have by Kashiwara's lemma (Proposition 2.3) that

$$H^j(U_n; F) \cong H^j(U_s; F)$$

for $n \geq s$.

Finally we replace $(a)_s^j$ by $(a')_s^k$ for $j = k$, and obtain

$$H^k\left(\bigcup_{s \in \mathbf{R}} U_s; F\right) \hookrightarrow H^k(U_s; F).$$

This completes the proof. $\square$

5 Truncated microlocal Morse theory

Let $f: Y \rightarrow X$ be a morphism of manifolds, and φ a real valued function of class C^1 . For a real number $t \in \mathbf{R}$, we set

$$\begin{aligned} Y_t &:= \{y \in Y ; \varphi(y) < t\} = \varphi^{-1}(]-\infty, t[), \\ \tilde{Y}_t &:= \{y \in Y ; \varphi(y) \leq t\} = \varphi^{-1}(]-\infty, t]). \end{aligned}$$

Let us define the following morphisms:

$$\begin{array}{ccc} Y_t & \xrightarrow{j_t} & Y \\ & \searrow f_t & \downarrow f \\ & & X, \end{array} \quad \begin{array}{ccc} \tilde{Y}_t & \xrightarrow{\tilde{j}_t} & Y \\ & \searrow f_t & \downarrow f \\ & & X. \end{array}$$

Proposition 5.1. Let $G \in D^b(Y)$, and assume $f_t: \text{supp}(G) \cap \overline{Y}_t \rightarrow X$ is proper for all $t \in \mathbf{R}$. Let k be an integer and t_o a real number.

(i) Assume that, for any $y \in Y \setminus \tilde{Y}_{t_o}$, we have

$$d\varphi(y) \notin \left(\text{SS}_k(G) + f_d \left(Y \times_X T^*X \right) \right).$$

Then the following statements hold.

(a) For $t > t_o$, the natural morphism

$$R^j f_* G \rightarrow R^j f_{t*} j_t^{-1} G$$

is isomorphic for $j < k$ and injective for $j = k$.

(a') For $t \geq t_o$, the natural morphism

$$R^j \tilde{f}_* G \rightarrow R^j \tilde{f}_{t*} \tilde{j}_t^{-1} G$$

is isomorphic for $j < k$ and injective for $j = k$.

(b) We have

$$\begin{aligned} \mathrm{SS}_k(Rf_* G) &\subset f_\pi \left(f_d^{-1} \left(\mathrm{SS}_k(G) \cap \pi^{-1}(\tilde{Y}_{t_o}) \right) \right) \\ &= f_\pi f_d^{-1} (\mathrm{SS}_k(G)). \end{aligned}$$

(ii) Assume that, for any $y \in Y \setminus \tilde{Y}_{t_o}$, we have

$$-d\varphi(y) \notin \left(\mathrm{SS}_k(G) + f_d \left(Y \times_X T^* X \right) \right).$$

Then

$$\begin{aligned} (c) \quad & R^j f_! G \cong R^j f_{t!} j_t^{-1} G \quad \text{for } j < k \text{ and } t > t_o, \\ (c_k) \quad & R^k f_! G \hookrightarrow R^k f_{t!} j_t^{-1} G \quad \text{for } t > t_o, \\ (c') \quad & R^j \tilde{f}_! G \cong R^j \tilde{f}_{t*} \tilde{j}_t^! G \quad \text{for } j < k \text{ and } t \geq t_o, \\ (c'_k) \quad & R^k \tilde{f}_! G \hookrightarrow R^k \tilde{f}_{t*} \tilde{j}_t^! G \quad \text{for } t \geq t_o, \\ (d) \quad & \mathrm{SS}_k(Rf_! G) \subset f_\pi \left(f_d^{-1} \left(\mathrm{SS}_k(G) \cap \pi^{-1}(\tilde{Y}_{t_o}) \right) \right) \\ &= f_\pi f_d^{-1} (\mathrm{SS}_k(G)). \end{aligned}$$

Proof. Let $\tilde{f}$ be the morphism $(f, \varphi): Y \rightarrow X \times \mathbf{R}$, and let q and $\tilde{\varphi}$ be the first and second projections defined on $X \times \mathbf{R}$ respectively. Then $\tilde{f}$ is proper on $\mathrm{supp}(G)$ and $f = q \circ \tilde{f}$. Let i_t be the open embedding $X \times]-\infty, t[\hookrightarrow X \times \mathbf{R}$ and $\tilde{i}_t$ the closed embedding $X \times]-\infty, t] \hookrightarrow X \times \mathbf{R}$. Set $q_t = q \circ i_t$ and $\tilde{q}_t = q \circ \tilde{i}_t$. These morphisms can be visualized by the following diagrams, in both of which the left squares are cartesian.

$$\begin{array}{ccc} Y & \xrightarrow{\tilde{f}} & X \times \mathbf{R} \xrightarrow{q} X, \\ \uparrow j_t & \square & \uparrow i_t \nearrow q_t \\ Y_t & \xrightarrow{\tilde{f}_t} & X \times]-\infty, t[\end{array} \quad , \quad \begin{array}{ccc} Y & \xrightarrow{\tilde{f}} & X \times \mathbf{R} \xrightarrow{q} X. \\ \uparrow \tilde{j}_t & \square & \uparrow \tilde{i}_t \nearrow \tilde{q}_t \\ Y_t & \xrightarrow{\tilde{f}_t} & X \times]-\infty, t] \end{array}$$

Then, setting $\tilde{G} := Rf_*G$, we have

$$\begin{aligned} Rf_*G &\cong Rq_*\tilde{G}, & R\tilde{f}_*G &\cong R\tilde{q}_*\tilde{G}, \\ Rf_{t*}j_t^{-1}G &\cong Rq_{t*}i_t^{-1}\tilde{G}, & R\tilde{f}_{t*}\tilde{j}_t^{-1}G &\cong R\tilde{q}_{t*}\tilde{i}_t^{-1}\tilde{G}, \\ Rf_{t!}j_t^{-1}G &\cong Rq_{t!}i_t^{-1}\tilde{G}, & R\tilde{f}_{t*}\tilde{j}_t^!G &\cong R\tilde{q}_{t*}\tilde{i}_t^!\tilde{G}. \end{aligned}$$

Thus taking j -th cohomology, we have

$$\begin{aligned} R^j f_*G &\cong R^j q_*\tilde{G}, & R^j \tilde{f}_*G &\cong R^j \tilde{q}_*\tilde{G}, \\ R^j f_{t*}j_t^{-1}G &\cong R^j q_{t*}i_t^{-1}\tilde{G}, & R^j \tilde{f}_{t*}\tilde{j}_t^{-1}G &\cong R^j \tilde{q}_{t*}\tilde{i}_t^{-1}\tilde{G}, \\ R^j f_{t!}j_t^{-1}G &\cong R^j q_{t!}i_t^{-1}\tilde{G}, & R^j \tilde{f}_{t*}\tilde{j}_t^!G &\cong R^j \tilde{q}_{t*}\tilde{i}_t^!\tilde{G}. \end{aligned}$$

Therefore, it is enough to prove the result for $\tilde{G}$, q , i_t , $\tilde{\varphi}$ instead of G , f , j_t , φ .

(i) It follows from the hypothesis that

$$(5.1) \quad \text{SS}_k(\tilde{G}) \cap (T^*X \times \{(t; \tau); t > t_o\}) \subset T^*X \times \{(t; \tau); \tau \leq 0\}.$$

Indeed, by hypothesis, for any $(y; \eta) \in \text{SS}_k(G)$ and $(f(x); \xi) \in Y \times_X T^*X$, we have

$$(5.2) \quad d\varphi(y) \neq \eta + f^*\xi.$$

If $(x, t; \xi, \tau)$ is a point of $\text{SS}_k(\tilde{G}) \cap (T^*X \times \{(t; \tau); t > t_o\})$, then, by the formula (3.3), there exists $(y; \eta) \in \text{SS}_k(G)$ such that

$$\tilde{f}(y) = (x, t) \text{ and } \eta = \tilde{f}^*(\xi, \tau).$$

For this η , we have

$$\eta = \tilde{f}^*(\xi, \tau) = f^*\xi + \varphi^*\tau = f^*\xi + \tau d\varphi(y).$$

If $\tau > 0$, we get

$$d\varphi(y) = \frac{1}{\tau}\eta - \frac{1}{\tau}f^*\xi$$

and this contradicts (5.2) because $\text{SS}_k(G)$ is conic.

By (5.1), Let U be an open subset of X contained in a local chart, and convex in such a chart.

□

6 Proof of Lemma 4.4

In this section, we prove Lemma 4.4, which is postponed in the proof of Theorem 4.2.

Proof of Lemma 4.4 First we prove that for $F \in \text{Mod}(\mathbf{k}_X)$, the natural morphism

$$(6.1) \quad \varinjlim_{t>s} \Gamma(U_t; F) \xrightarrow{\sim} \Gamma(Z_s \cup U_s; F).$$

is isomorphic. Consider the following diagram.

$$\begin{array}{ccc} \varinjlim_{t>s} \Gamma(U_t; F) & & \\ \downarrow \tilde{\phi} & \searrow & \\ \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F) & \xrightarrow{\phi} & \Gamma(Z_s \cup U_s; F|_{Z_s \cup U_s}) \\ \downarrow & & \downarrow \\ \varinjlim_{U \supset Z_s} \Gamma(U; F) & \xrightarrow[\psi]{\sim} & \Gamma(Z_s; F|_{Z_s}). \end{array}$$

Let us begin by showing that the morphism

$$\tilde{\phi}: \varinjlim_{t>s} \Gamma(U_t; F) \rightarrow \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F)$$

is an isomorphism. It is enough to show that $(U_t)_{t>s}$ is a cofinal family of $(U \cup U_s)_{U \supset Z_s}$. Let U be an open neighborhood of Z_s . Then we shall show by contradiction that there exists a real number $t > s$ such that $U_t \subset U \cup U_s$. Let us assume that $U_t - (U \cup U_s) \neq \emptyset$ for any $t > s$. Then define the family of subsets $(A_r)_{s < r < t}$ of $\overline{U_t - U_s}$ by $A_r := U_r - (U \cup U_s)$. This $(A_r)_{s < r < t}$ satisfies the finite intersection property. Indeed, for finite real numbers $s < r_0 < \dots < r_m < t$, it follows from the assumption for contradiction that

$$\begin{aligned} A_{r_0} \cap \dots \cap A_{r_m} &= A_{r_0} \\ &= U_{r_0} - (U \cup U_s) \\ &\neq \emptyset. \end{aligned}$$

Therefore, since $\overline{U_t - U_s}$ is compact, we have $\bigcap_{s < r < t} \overline{A_r} \neq \emptyset$. However, we have from the hypothesis $Z_s \subset U$ that

$$\begin{aligned}
\emptyset &\neq \bigcap_{s < r < t} \overline{A_r} = \bigcap_{s < r < t} \overline{U_r - (U \cup U_s)} \\
&= \bigcap_{s < r < t} (\overline{U_r - U_s} \cap \overline{U_r - U}) \\
&= \bigcap_{s < r < t} \overline{U_r - U_s} \cap \bigcap_{s < r < t} \overline{U_r - U} \\
&= Z_s \cap \bigcap_{s < r < t} \overline{U_r - U} \\
&= Z_s \cap \bigcap_{s < r < t} \overline{U_r} \cap \overline{X - U} \\
&= Z_s \cap \bigcap_{s < r < t} \overline{U_r} \cap (X - U) \\
&= Z_s \cap (X - U) \cap \bigcap_{s < r < t} \overline{U_r} \\
&= \emptyset.
\end{aligned}$$

This is a contradiction, and it follows that $\tilde{\phi}$ is an isomorphism.

Since $\tilde{\phi}$ is an isomorphism, it is enough to show that

$$\phi: \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F) \rightarrow \Gamma(Z_s \cup U_s; F)$$

is isomorphic. Since X is Hausdorff and Z_s is compact, the natural morphism

$$\psi: \varinjlim_{U \supset Z_s} \Gamma(U; F) \rightarrow \Gamma(Z_s; F)$$

is isomorphic by Proposition 2.1. Since U_s is open, we have another isomorphism

$$\psi': \varinjlim_{V \supset U_s} \Gamma(V; F) \rightarrow \Gamma(U_s; F).$$

The injectivity of ϕ follows from the definition of a sheaf and the injectivities of ψ and ψ' .

Let us prove that ϕ is surjective. Let $\sigma \in \Gamma(Z_s \cup U_s; F)$ be a section of $F|_{Z_s \cup U_s}$. By the isomorphism ψ above, we can extend $\sigma|_{Z_s}$ to an open neighborhood U of Z_s in X . More precisely, there exists a section τ on U such that $\tau|_{Z_s} = \sigma|_{Z_s}$. On the other hand, it follows from the compatibility

of restrictions of a section that $\tau|_{Z_s} = (\tau|_{(Z_s \cup U_s) \cap U})|_{Z_s}$. By the definition of a section on the closed subset Z_s of $(Z_s \cup U_s) \cap U$, we can find an open subset W of $(Z_s \cup U_s) \cap U$ such that

$$\tau|_W = \sigma|_W.$$

By the definition of the induced topology of $(Z_s \cup U_s) \cap U$, there exists an open set $\widetilde{W}$ of X such that

$$W = ((Z_s \cup U_s) \cap U) \cap \widetilde{W} \quad \text{and} \quad \widetilde{W} \subset U.$$

Thus we have

$$(6.2) \quad U_s \cap W = U_s \cap \widetilde{W}, \quad \sigma|_{U_s \cap \widetilde{W}} = \tau|_{U_s \cap \widetilde{W}}.$$

For this $\widetilde{W}$, the pair $(U_s, \widetilde{W})$ is an open covering of $\widetilde{W} \cup U_s$. Thus by (6.2), we can glue $\sigma|_{U_s}$ and $\tau|_{\widetilde{W}}$ together and get a section $\tilde{\sigma}$ such that

$$\tilde{\sigma}|_{Z_s \cup U_s} = \sigma, \quad \tilde{\sigma}|_{\widetilde{W}} = \tau|_{\widetilde{W}}.$$

This proves the surjectivity of ϕ .

We show the the following statement.

Next let $F \in D^+(\mathbf{k}_X)$. Let $I^\bullet$ be a complex of flabby sheaves quasi-isomorphic to F . the restrictions of a flabby sheaf F , denoting by $F|_S$, is also flabby. Then it follows from (6.1) that

$$\begin{aligned} \varinjlim_{t>s} H^j(U_t; F) &\cong \varinjlim_{t>s} H^j \Gamma(U_t; I^\bullet) \\ &\cong H^j \left(\varinjlim_{t>s} \Gamma(U_t; I^\bullet) \right) \\ &\cong H^j \Gamma(Z_s \cup U_s; I^\bullet) \\ &\cong H^j(Z_s \cup U_s; F). \end{aligned}$$

This completes the proof. □

References

- [FM07] Teresa Monteiro Fernandes, Ana Rita Martins. *Truncated Microsupport and Hyperbolic Inequalities*, Nagoya Mathematical Journal, 2007, Vol. 185, pp.63–91.

- [GKS12] Stéphane Guillermou, Masaki Kashiwara, Pierre Schapira, *Sheaf Quantization of Hamiltonian Isotopies and Applications to Nondisplaceability Problems*, Duke. Math. J. 161, No. 2, pp.201–245, 2012.
- [KFS03a] Masaki Kashiwara, Teresa Monteiro Fernandes, Pierre Schapira, *Truncated Microsupport and Holomorphic Solutions of D -modules*, Ann. Scient. Éc. Norm. Sup., série 4, tome 36, pp.583–399, 2003.
- [KFS03b] Masaki Kashiwara, Teresa Monteiro Fernandes, Pierre Schapira, *Involutivity of Truncated microsupports*, Bull. Soc. math. France, 131 (2), 2003, pp.259–266.
- [KS90] Masaki Kashiwara, Pierre Schapira, *Sheaves on Manifolds*, Grundlehren der Mathematischen Wissenschaften, 292, Springer, 1990.