

修士論文

Microlocal Morse Theory and Truncated
Microsupport
(超局所モース理論と打切り超局所台)

大柴寿浩

2026.3

北海道大学大学院理学院
数学専攻博士前期課程

Microlocal Morse Theory and Truncated Microsupport (超局所モース理論と打切り超局所台)

大柴寿浩

目次

1	Introduction	1
1.1	Organization	1
2	Background Material	1
2.1	The Mittag-Leffler Condition	2
3	Truncated Microsupport	3
4	Truncated Noncharacteristic Deformation Lemma	4
5	An Application to Microlocal Morse Theory	9

1 Introduction

In this paper, We study microlocal Morse theory, which is presented, for instance, in [GKS12]. However, we replace the notion of microsupports with that of truncated microsupprts, introduced by Kashiwara, Monteiro-Fernandes, and Schapira in [KFS03]. In [KFS03], they do not study the functorial property of truncated microsupport of direct images of non proper morphisms. We prove this functorial property under non proper direct images.

1.1 Organization

This paper is organized as follows. In Section 4, we prove a truncated version of the noncharacteristic deformation lemma.

2 Background Material

In this section, we recall some definitions and results from [KS90], following its notations with slight modifications.

We consider a commutative unital ring $\mathbf{k}$ of finite global dimension (e.g. $\mathbf{Z}$ or a field). Let X be a

topological space. We denote by $D^b(\mathbf{k}_X)$ the bounded derived category of sheaves of $\mathbf{k}$ -modules on X .

Let us denote Z a locally closed set of X and $i: Z \hookrightarrow X$ an inclusion. We use the following functors:

$$R\Gamma_Z(\cdot) := Ri_*i^!, \quad (\cdot)_Z := Ri_!i^{-1}.$$

If there is no risk, we will write $\mathbf{k}_Z$ rather than $\mathbf{k}_{X,Z}$.

Geometrical Notations

Let M be a real manifold of class C^∞ . We denote by $\pi: T^*M \rightarrow M$ the cotangent bundle of M . If $f: M \rightarrow N$ is a morphism of manifolds, then we have the natural diagram.

$$\begin{array}{ccccc} TM & \xrightarrow{f'} & M \times_N TN & \xrightarrow{f_\tau} & TN \\ & \searrow \tau_M & \downarrow & \square & \downarrow \tau_N \\ & & M & \xrightarrow{f} & N, \end{array} \quad \begin{array}{ccccc} T^*M & \xleftarrow{f_d} & M \times_N T^*N & \xrightarrow{f_\pi} & T^*N \\ & \searrow \pi_M & \downarrow & \square & \downarrow \pi_N \\ & & M & \xrightarrow{f} & N. \end{array}$$

In this diagram, f_τ and f_π denote the natural projections. f' is defined by $(x; v) \mapsto (x; df_x v)$, and f_d is induced by the transposition of f' .

Approximation by Inductive Limits

Proposition 2.1 ([KS90, Proposition 2.5.1]). Let X be a topological space, Z a subspace, F a sheaf on X . Consider the canonical morphism:

$$\psi: \varinjlim_U \Gamma(U; F) \rightarrow \Gamma(Z; F),$$

where U ranges through the family of open neighborhoods of Z in X .

- (i) The morphism ψ is injective.
- (ii) Assume that X is Hausdorff and Z is compact. Then ψ is an isomorphism.
- (iii) Assume that X is paracompact and Z is closed. Then ψ is an isomorphism.

A paracompact space is by definition a Hausdorff space.

2.1 The Mittag-Leffler Condition

Consider a projective system of abelian groups $(M_n, \rho_{n,p})_n$ indexed by $\mathbf{N}$. One says that this system satisfies the Mittag-Leffler condition (ML for short) if for any $n \in \mathbf{N}$, the decreasing sequence $(\rho_{n,p}(M_p))_p$ of subgroups of M_n is stationary.

For a projective system of abelian groups $(M_n)_n$, we set

$$M_\infty := \varprojlim_n M_n.$$

Consider a projective system of complexes

$$(2.1) \quad E_n^\bullet: \cdots \rightarrow M_n^{j-1} \rightarrow M_n^j \rightarrow M_n^{j+1} \rightarrow \cdots,$$

and projective limit

$$(2.2) \quad E_\infty^\bullet: \cdots \rightarrow M_\infty^{j-1} \rightarrow M_\infty^j \rightarrow M_\infty^{j+1} \rightarrow \cdots.$$

Denote by

$$(2.3) \quad \Phi_k: H^k(E_\infty^\bullet) \rightarrow \varprojlim_n H^k(E_n^\bullet)$$

the natural morphism.

Proposition 2.2 ([KS90, Proposition 1.12.4]). Assume that for each $j \in \mathbf{Z}$, the system $(M_n^{j-1})_n$ satisfies the ML condition. Then

- (a) for each $k \in \mathbf{Z}$, the map Φ_k is surjective,
- (b) if moreover, for a given i the system $(H^{i-1}(E_n^\bullet))_n$ satisfies the ML condition, then Φ_i is bijective.

The following is called Kashiwara's lemma.

Proposition 2.3 ([KS90, Proposition 1.12.6]). Let $(X_s, \rho_{s,t}: X_t \rightarrow X_s)_{(s, (s \leq t))}$ be a projective system of sets indexed by $\mathbf{R}$. Assume that for each $s \in \mathbf{R}$, the canonical maps

$$\lambda_s: X_s \rightarrow \varprojlim_{r < s} X_r$$

and

$$\mu_s: \varinjlim_{t > s} X_t \rightarrow X_s$$

are both injective (resp. surjective). Then all the maps ρ_{s_0, s_1} ($s_0 \leq s_1$) are injective (resp. surjective).

ML procedure for sheaves

Proposition 2.4 ([KS90, Proposition 2.7.1]). Let X be a topological space and let $F \in \mathbf{D}^+(\mathbf{Z}_X)$. Let $(U_n)_{n \in \mathbf{N}}$ be an increasing sequence of open subsets of X , $(Z_n)_{n \in \mathbf{N}}$ a decreasing sequence of closed subsets of X . Set $U := \bigcup_n U_n$, $Z := \bigcap_n Z_n$.

- (i) For any j , the natural map $\varphi_j: H_Z^j(U; F) \rightarrow \varprojlim_n H_{Z_n}^j(U_n; F)$ is surjective.
- (ii) Assume that for a given j , the projective system $(H_{Z_n}^{j-1}(U_n; F))_n$ satisfies the ML condition. Then φ_j is bijective.
- (iii) Let now $(X_n)_{n \in \mathbf{N}}$ be an increasing family of subsets of X satisfying $X = \bigcup_n X_n$ and $X_n \subset \text{Int}(X_{n+1})$ for all n . Assume that for a given j , the projective system $(H^{j-1}(X_n; F))_n$ satisfies the ML condition. Then the natural map $H^j(X; F) \rightarrow \varprojlim_n H^j(X_n; F)$ is bijective.

3 Truncated Microsupport

Definition 3.1. Let $F \in \mathbf{D}^b(\mathbf{k}_X)$. The closed conic subset $\text{SS}_k(F)$ of T^*X is defined by: $p \notin \text{SS}_k(F)$ if and only if there exists an open conic neighborhood U of p such that for any $x \in \pi(U)$ and for any

real-valued C^1 -function φ defined on a neighborhood of x such that $\varphi(x) = 0$, $d\varphi(x) \in U$, one has

$$(3.1) \quad H_{\{\varphi \geq 0\}}^j(F)_x = 0 \quad \text{for any } j \leq k.$$

Functorial Properties

Proposition 3.2. Let $f: X \rightarrow Y$ be a morphism of real analytic manifolds and let $F \in D^b(\mathbf{k}_X)$ such that f is proper on $\text{supp}(F)$. Then for any $k \in \mathbf{Z}$,

$$(3.2) \quad \text{SS}_k(Rf_*F) \subset f_\pi f_d^{-1}(\text{SS}_k(F)).$$

The equality holds in case f is a closed embedding.

4 Truncated Noncharacteristic Deformation Lemma

Proposition 4.1. Let X be a Hausdorff space. Let $F \in D^+(\mathbf{k}_X)$, and $(U_t)_{t \in \mathbf{R}}$ be a family of open subsets of X . Let $k \in \mathbf{Z}$. If s is a real number, set $Z_s := \bigcap_{t > s} \overline{U_t - U_s}$. We assume the following conditions.

- (i) $U_t = \bigcup_{s < t} U_s$, for all $t \in \mathbf{R}$.
- (ii) For all pairs (s, t) with $s \leq t$, the set $\overline{U_t - U_s} \cap \text{supp } F$ is compact.
- (iii) For all $s \in \mathbf{R}$ and all $x \in Z_s - U_s$, we have

$$(H_{X-U_s}^j(F))_x = 0 \quad \text{for all } j \leq k.$$

- (iv) For all pairs (s, t) with $s < t$, $Z_s \subset U_t$.

Then, for all $t \in \mathbf{R}$, we have the isomorphisms

$$H^j \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \xrightarrow{\sim} H^j(U_t; F) \quad \text{for all } j < k,$$

and the injections

$$H^k \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \hookrightarrow H^k(U_t; F).$$

Example 4.2. The condition (iv) of Proposition 4.1 is not contained in the original version ([KS90, Proposition 2.7.2]). Therefore let us give an example of $(U_t)_t$ that does not satisfy the condition (iv).

- (i) Let $X = \mathbf{R}^2$, and for a real number t define the function

$$\varphi_t(x) := \begin{cases} \max(0, t(2 - |x|)) & (|x| < 2), \\ 0 & (|x| \geq 2). \end{cases}$$

Then let us denote by U_t the open subset of X

$$U_t := \{(x, y) \in X; y < \varphi_t(x)\}.$$




Fig.1 example (i)

Let us, in turn, give three examples of $(U_t)_t$ satisfying the condition (iv). In all the examples below, X is $\mathbf{R}^2$.

(ii) For a real number t , define the open set U_t by

$$U_t := \begin{cases} \{(x, y) \in X; \|(x, y)\| < t\} & (t > 0), \\ \emptyset & (t \leq 0). \end{cases}$$




Fig.2 example (ii)

(iii) For a real number t , define the function $\varphi_t: \mathbf{R} \rightarrow \mathbf{R}$ as follows. If $t \leq 0$, set $\varphi_t(x) \equiv 0$. If $t > 0$, set

$$\varphi_t(x) := \begin{cases} \sqrt{t^2 - x^2} & (|x| < t), \\ 0 & (|x| \geq t). \end{cases}$$

Then define the open set U_t by

$$U_t := \{(x, y) \in X; y < \varphi_t(x)\}.$$

(iv) For a real number t , define the open set U_t as follows. If $t \leq 0$, set

$$U_t := \{(x, y) \in X; y < 0\}.$$

If $t > 0$, set

$$U_t := \left\{ (x, y) \in X; \begin{array}{ll} \text{If } |x| < \frac{1}{t}, & y < t. \\ \text{Otherwise,} & y < \frac{1}{|x|}. \end{array} \right\}.$$




Fig.3 example (iii)




Fig.4 example (iv)

Proof. Consider the following conditions for $j \in \mathbf{Z}$ and $s \in \mathbf{R}$:

$$\begin{aligned}
 (a)_s^j & \quad \mu_s^j: \varinjlim_{t>s} H^j(U_t; F) \xrightarrow{\sim} H^j(U_s; F), \\
 (a')_s^k & \quad \mu_s^k: \varinjlim_{t>s} H^k(U_t; F) \hookrightarrow H^k(U_s; F), \\
 (b)_s^j & \quad \lambda_s^j: H^j(U_s; F) \xrightarrow{\sim} \varprojlim_{r<s} H^j(U_r; F).
 \end{aligned}$$

Let us prove, for all $s \in \mathbf{R}$, $(a)_s^j$ for $j < k$ and $(a')_s^k$. We may assume that $\overline{U_t - U_s}$ is compact for any $s \leq t$ by replacing X by $\text{supp } F$. Consider the distinguished triangle

$$\text{R}\Gamma_{X-U_s}(F) \rightarrow F \rightarrow \text{R}\Gamma_{U_s}(F) \xrightarrow{+1}.$$

Let us endow with $Z_s \cup U_s$ the induced topology, and denote by $i_s: Z_s \cup U_s \hookrightarrow X$ the natural continuous map. Applying the functor

$$(\cdot)|_{Z_s \cup U_s} = i_s^{-1}$$

to the distinguished triangle above, we have

$$\text{R}\Gamma_{X-U_s}(F)|_{Z_s \cup U_s} \rightarrow F|_{Z_s \cup U_s} \rightarrow \text{R}\Gamma_{U_s}(F)|_{Z_s \cup U_s} \xrightarrow{+1}.$$

Again applying to this triangle the functor $\text{R}\Gamma(Z_s \cup U_s; \cdot)$, we have

$$(4.1) \quad \text{R}\Gamma(Z_s \cup U_s; \text{R}\Gamma_{X-U_s}(F)) \rightarrow \text{R}\Gamma(Z_s \cup U_s; Z_s \cup U_s) \rightarrow \text{R}\Gamma(U_s; F) \xrightarrow{+1}.$$

By the hypothesis (iii) and the definition of a local cohomology, we see that

$$H_{X-U_s}^j(F)|_{Z_s \cup U_s} = 0 \quad \text{for any } j \leq k,$$

and then

$$H^j(Z_s \cup U_s; \text{R}\Gamma_{X-U_s}(F)) = 0 \quad \text{for any } j \leq k.$$

Thus we have

$$(4.2) \quad H^j(Z_s \cup U_s; F) \xrightarrow{\sim} H^j(U_s; F) \quad \text{for any } j < k,$$

$$(4.3) \quad H^k(Z_s \cup U_s; F) \hookrightarrow H^k(U_s; F).$$

Therefore, if we show Lemma 4.3 below, we have

$$\begin{aligned} \varinjlim_{t>s} H^j(U_t; F) &\cong H^j(Z_s \cup U_s; F) \\ &\xrightarrow{\sim} H^j(U_s; F) \end{aligned}$$

for any $j < k$, and $(a)_s^j$ follows. Similarly, we have

$$\begin{aligned} \varinjlim_{t>s} H^k(U_t; F) &\cong H^k(Z_s \cup U_s; F) \\ &\hookrightarrow H^k(U_s; F) \end{aligned}$$

and $(a')_s^k$ follows.

Lemma 4.3. For any $j \in \mathbf{Z}$ and $s \in \mathbf{R}$, there exists a natural isomorphism

$$\varinjlim_{t>s} H^j(U_t; F) \cong H^j(Z_s \cup U_s; F).$$

Proof of Lemma 4.3 First we prove that for $F \in \text{Mod}(\mathbf{k}_X)$, the natural morphism

$$(4.4) \quad \varinjlim_{t>s} \Gamma(U_t; F) \xrightarrow{\sim} \Gamma(Z_s \cup U_s; F).$$

is isomorphic. Consider the following diagram.

$$\begin{array}{ccc} \varinjlim_{t>s} \Gamma(U_t; F) & & \\ \downarrow & \searrow & \\ \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F) & \longrightarrow & \Gamma(Z_s \cup U_s; F|_{Z_s \cup U_s}) \\ \downarrow & & \downarrow \\ \varinjlim_{U \supset Z_s} \Gamma(U; F) & \longrightarrow & \Gamma(Z_s; F|_{Z_s}) \end{array}$$

Note that $(U_t)_{t>s}$ is a cofinal family of $(U \cup U_s)_{U \supset Z_s}$. Indeed, かく .

Therefore, we have

$$\varinjlim_{t>s} \Gamma(U_t; F) \xrightarrow{\sim} \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F).$$

By this isomorphism, it is enough to show that

$$\phi: \varinjlim_{U \supset Z_s} \Gamma(U \cup U_s; F) \rightarrow \Gamma(Z_s \cup U_s; F)$$

is isomorphic. Since X is Hausdorff and Z_s is compact, the natural morphism

$$\psi: \varinjlim_{U \supset Z_s} \Gamma(U; F) \rightarrow \Gamma(Z_s; F)$$

is isomorphic by Proposition 2.1. Since U_s is open, we have another isomorphism

$$\psi': \varinjlim_{V \supset U_s} \Gamma(V; F) \rightarrow \Gamma(U_s; F).$$

The injectivity of ϕ follows from the definition of a sheaf and the injectivities of ψ and ψ' .

Let us prove that ϕ is surjective. Let $\sigma \in \Gamma(Z_s; F)$ be a section of $F|_{Z_s \cup U_s}$. By the isomorphism ψ above, we can extend $\sigma|_{Z_s}$ to an open neighborhood U of Z_s in X . More precisely, there exists a section τ on U such that $\tau|_{Z_s} = \sigma|_{Z_s}$. Now we see that $\tau|_{Z_s} = (\tau|_{(Z_s \cup U_s) \cap U})|_{Z_s}$, and by the definition of a section on the closed subset Z_s of $(Z_s \cup U_s) \cap U$, we can find an open subset W of $(Z_s \cup U_s) \cap U$ such that

$$(4.5) \quad \tau|_W = \sigma|_W.$$

By the definition of the induced topology of $(Z_s \cup U_s) \cap U$, there exists an open set $\widetilde{W}$ of X such that

$$W = ((Z_s \cup U_s) \cap U) \cap \widetilde{W} \quad \text{and} \quad \widetilde{W} \subset U.$$

Thus we have $W = ((Z_s \cup U_s) \cap \widetilde{W})$. For this $\widetilde{W}$, the pair $(Z_s \cup U_s, \widetilde{W})$ is an open covering of $\widetilde{W} \cup U_s$. Thus by (4.5), we can glue σ and $\tau|_{\widetilde{W}}$ together and get a section $\tilde{\sigma}$ such that

$$\tilde{\sigma}|_{Z_s \cup U_s} = \sigma, \quad \tilde{\sigma}|_{\widetilde{W}} = \tau|_{\widetilde{W}}.$$

This proves the surjectivity of ϕ .

Next Let $F \in D^+(\mathbf{k}_X)$. Let $I^\bullet$ be a complex of flabby sheaves quasi-isomorphic to F . Then it follows from (4.4) that

$$\begin{aligned} \varinjlim_{t > s} H^j(U_t; F) &\cong \varinjlim_{t > s} H^j \Gamma(U_t; I^\bullet) \\ &\cong H^j \left(\varinjlim_{t > s} \Gamma(U_t; I^\bullet) \right) \\ &\cong H^j \Gamma(Z_s \cup U_s; I^\bullet) \\ &\cong H^j(Z_s \cup U_s; F). \end{aligned}$$

This completes the proof. $\square$

Let us come back to the proof of Proposition 4.1. We prove $(b)_s^j$ for all $j \in \mathbf{Z}$ and $s \in \mathbf{R}$ by induction on j . Since $F \in D^+(\mathbf{k}_X)$, we have $(b)_s^j$ for $j \ll 0$ and $s \in \mathbf{R}$. Hence there exists an integer $j_0 < k$ such that $(b)_s^j$ holds for any $j < j_0$ and $s \in \mathbf{R}$. Then assume $(b)_s^j$ is proved for all $j < j_0$ and $s \in \mathbf{R}$. Since we have $(a)_s^j$ and $(b)_s^j$ for each $j < j_0$ and each $s \in \mathbf{R}$, we can apply Kashiwara's lemma (Proposition 2.3) and have

$$H^j(U_t; F) \xrightarrow{\sim} H^j(U_s; F) \quad \text{for } j < j_0 \text{ and } s \leq t.$$

Fix $s \in \mathbf{R}$ and consider the projective system $(H^{j_0-1}(U_{s-1/n}))_{n \in \mathbf{N}}$. This system satisfies ML condition. Therefore, by applying Proposition 2.4 (iii), we have

$$H^j(U_s; F) \xrightarrow{\sim} \varprojlim_n H^j \left(U_{s-\frac{1}{n}}; F \right).$$

Since $(s - 1/n)_n$ is a cofinal family of $(t)_{s>t}$, we have $(b)_s^{j_0}$. Then by induction on j , we have $(b)_s^j$ for $j < k$ and $s \in \mathbf{R}$. Again by applying Proposition 2.4 to the projective system $(H^j(U_n; F))_n$, we obtain

$$H^j \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \cong H^j \left(\bigcup_{n \in \mathbf{N}} U_n; F \right) \xrightarrow{\sim} \varprojlim_n H^j(U_n; F)$$

and for each n , we have by Kashiwara's lemma that

$$H^j(U_n; F) \cong H^j(U_s; F)$$

for $n \geq s$.

Finally we replace $(a)_s^j$ by $(a')_s^k$ for $j = k$, and obtain

$$H^k \left(\bigcup_{s \in \mathbf{R}} U_s; F \right) \hookrightarrow H^k(U_s; F).$$

This completes the proof. □

5 An Application to Microlocal Morse Theory

In this section, let M be a manifold of class C^∞ .

Lemma 5.1. Let $-\infty < a < b < +\infty$ be real numbers and $G \in \mathbf{D}^b(\mathbf{R})$. Let k be an integer. If

$$\left(H_{[t, +\infty)}^j(G) \right)_t \cong 0 \quad \text{for all } t \in [a, b), \text{ and all } j \leq k.$$

Then one has

$$H^j((-\infty, b); G) \xrightarrow{\sim} H^j((-\infty, a); G)$$

for all $j < k$, and this morphism is injective for $j = k$.

Proof. We will use Proposition 4.1. Let us put $X = (-\infty, b)$, $F = G|_{I_b}$, and $U_t = I_t$, and verify the hypotheses of the proposition.

- (i) $\bigcup_{s < t} I_s = \bigcup_{s < t} (-\infty, s) = (-\infty, t) = I_t$ for all $t \in \mathbf{R}$.
- (ii) If $s \leq t$, then $\overline{I_t - I_s} \cap \text{supp } G = \overline{[s, t)} \cap \text{supp } G = [s, t] \cap \text{supp } G$ is compact.
- (iii) For $s \in \mathbf{R}$, we have

$$Z_s = \bigcap_{t > s} \overline{I_t - I_s} = \bigcap_{t > s} \overline{[s, t)} = \bigcap_{t > s} [s, t] = \{s\},$$

and

$$Z_s - I_s = \{s\}.$$

By the hypothesis, we have

$$H_{X - I_s}^j(F)_s = H_{[s, \infty)}^j(F)_s = 0$$

for $s \in \{s\}$.

(iv) It follows that $Z_s = \{s\} \subset (-\infty, t) = I_t$ if $s < t$.

Then we can apply Proposition 4.1, and we have the isomorphisms for all $t \in [a, b)$ and $j < k$,

$$H^j(I_b; F) = H^j\left(\bigcup_s I_s; F\right) \xrightarrow{\sim} H^j(I_t; F)$$

and the injections

$$H^k(I_b; F) = H^k\left(\bigcup_s I_s; F\right) \hookrightarrow H^k(I_t; F).$$

□

Theorem 5.2. Let F in $D^b(\mathbf{k}_M)$, and $\psi: M \rightarrow \mathbf{R}$ be a real-valued function of class C^1 , proper on $\text{supp}(F)$. Let $a < b$ be real numbers. We assume $d\psi(x) \notin \text{SS}_k(F)$ for $a \leq \psi(x) < b$. For $t \in \mathbf{R}$, we set $M_t := \psi^{-1}((-\infty, t))$. Then, the restriction morphisms

$$H^j(M_b; F) \rightarrow H^j(M_a; F)$$

are isomorphic for each $j < k$, and injective for $j = k$.

Proof. First, we have

$$H^j(M_t; F) \cong H^j \text{R}\Gamma((-\infty, t); \text{R}\psi_* F)$$

for $t \in I$. Since we assume that $d\psi(x) \notin \text{SS}_k(F)$ for $a \leq \psi(x) < b$, and by (3.2), the formula $\text{SS}_k(\text{R}\psi_* F) \subset \psi_\pi \psi_d^{-1} \text{SS}_k(F)$ holds. Hence we have

$$\left(H^j_{[\psi(x), +\infty)}(\text{R}\psi_* F) \right)_{\psi(x)} = 0$$

for each $j \leq k$. Then we can apply Lemma 5.1 and have

$$H^j((-\infty, b); \text{R}\psi_* F) \xrightarrow{\sim} H^j((-\infty, a); \text{R}\psi_* F)$$

for all $j < k$ and

$$H^k((-\infty, b); \text{R}\psi_* F) \hookrightarrow H^k((-\infty, a); \text{R}\psi_* F).$$

Then we can conclude that

$$H^j(M_b; F) \xrightarrow{\sim} H^j(M_a; F)$$

for all $j < k$ and

$$H^k(M_b; F) \hookrightarrow H^k(M_a; F).$$

□

参考文献

- [GKS12] Stéphane Guillermou, Masaki Kashiwara, Pierre Schapira, *Sheaf Quantization of Hamiltonian Isotopies and Applications to Nondisplaceability Problems*, Duke. Math. J. 161, No. 2, pp.201–245, 2012.

- [KFS03] Masaki Kashiwara, Teresa Monteiro Fernandes, Pierre Schapira, *Truncated Microsupport and Holomorphic Solutions of D -modules*, Ann. Scient. Éc. Norm. Sup., série 4, tome 36, pp.583–399, 2003.
- [KS90] Masaki Kashiwara, Pierre Schapira, *Sheaves on Manifolds*, Grundlehren der Mathematischen Wissenschaften, 292, Springer, 1990.
- [ST92] Pierre Schapira, Nobuyuki Tose, *Morse Inequalities for $\mathbf{R}$ -constructible Sheaves*, Adv. Math. 93, pp.1–8, 1992.