

$$\frac{16}{20}$$

Untitled

November 4, 2022

[]:

[17]: `# -*- coding: utf-8 -*-
 """
 Created on Sat Oct 29 12:36:03 2022

 @author: bene-
 """

 import numpy as np
 from math import *
 import matplotlib.pyplot as plt
 import scipy.optimize as so
 import scipy.special as sp
 import mpmath as mp

 N_x=10
 N_y=10
 h=0.0
 T=1
 J=0.5/T`

Que-1)
 10/10

[18]: `def init_2D():
 global N_x
 global N_y
 lattice = np.zeros((N_x,N_y))
 return lattice`

Creates a $N_x \times N_y$ Array which will be the lattice.

[19]: `def step(lattice): #Does one step in the Markov-Chain and return the next
 ↳constallation
 global N_x
 global N_y
 global J
 global T
 beta=1/T
 for i in range (N_x):`

```

M=i
for j in range (N_y):
    N=j #selects a random spin in the chain
    if np.random.rand()>0.5:#generates a new random orientation of
    → the spin
        spin_new=1
    else:
        spin_new=-1
    delta_H=calc_delta_H(lattice,M,N,spin_new) #calculates the
    → energy difference from the old and new constellation
    if delta_H<0:
        lattice[M,N]=spin_new
    else:
        if np.exp(-beta * delta_H)>np.random.rand():
            lattice[M,N]=spin_new
return(lattice)

```

Input: $N_x \times N_y$ Array (lattice); Output: $N_x \times N_y$ Array (lattice after one step)

A step in the underlying Markov chain is defined by a complete loop of the lattice. Every spin on the lattice is randomly changed. This change is accepted if the energy of the new configuration is lower than before and if the energy for the new configuration is higher than before, the new configuration is only accepted if the Boltzmann probability is larger than a randomly drawn number between 0 and 1.

[20]: `def calc_delta_H(lattice,M,N,spin_new):`

```

global J
global N_x
global N_y
global h

spin_old=lattice[M][N]
delta_H=0
delta_H+=J* (lattice[(M-1)%N_x][N] * spin_new) \
            -J* (lattice[(M-1)%N_x][N] * spin_old)
delta_H+=J* (lattice[(M+1)%N_x][N] * spin_new) \
            -J* (lattice[(M+1)%N_x][N] * spin_old)
delta_H+=J* (lattice[M][(N-1)%N_y] * spin_new) \
            -J* (lattice[M][(N-1)%N_y] * spin_old)
delta_H+=J* (lattice[M][(N+1)%N_y] * spin_new) \
            -J* (lattice[M][(N+1)%N_y] * spin_old)
delta_H+=h*(spin_new-spin_old)

return -delta_H

```

Tip: you wouldn't have to write such big expression for ΔE if you use the fact $\delta \text{spin}_{\text{old}} = -\delta \text{spin}_{\text{new}}$ then it just becomes $2J \times \text{neighbour spin} \times \delta \text{spin}_{\text{new}}$

Input: $N_x \times N_y$ Array (lattice), Position of the changed spin (M,N), value of the changed spin (spin_new); Output: Energy difference from old and new configuration.

The cost of the energy-calculation for the entire configuration scales -in our case- with $5 \cdot (N_x \cdot N_y)$ which is $\mathcal{O}(N_x \cdot N_y)$.

Que-3

$\sqrt{1}$

The calculation itself is straight forward and the periodic boundary conditions are implemented through the Modulo calculations.

Input: $N_x N_y$ Array (lattice); Output: Magnetization per Spin

Input: $N_x N_y$ Array (lattice); Output: Energy per Spin

We create four new versions of the lattice which are individually rolled in each possible direction (periodic boundary conditions are taken into account correctly by this method). By taking the direct multiplication (no matrix-multiplication) we can calculate the energy for the entire configuration in each direction pretty quickly.

```
[23]: def eps_theo():
    global J

    A=mp.sech(2*J)**2
    B=mp.tanh(2*J)**2

    return -2*J*mp.coth(2*J)*(1+(2/np.pi)*(2*(mp.tanh(2*J)**2)-1)\
        *mp.ellipk(4*A*B))
```

Input: None; Output: Theoretical value of energy per Spin in the thermodynamic limit.


```
[24]: def abs_m_theo():
    global J
    if J>0.4406867935:
        return (1-(1/mp.sinh(2*J)**4))**(1/8)
    else:
        return 0
```

Input: None; Output: Theoretical value of magnetization per Spin in the thermodynamic limit.

```
[25]: def m_h_5():
    global N_x
    global N_y
    global h
    E=20 #amount of points for the graph
    I=10 #initialization
    A=2 #autocorrelationcorrection
    C=100 #Number of components for an ensemble
    D=5
    final_M=np.zeros(E+1)
    M=0
    lattice=init_2D()
    colours=['b','g','r','c','m','y','k']
    for m in range(D):
        N_x=N_y=(m+1)*4
        final_M=np.zeros(E+1)
        for j in range(E+1):
            h=-1+(j/10)
            lattice=init_2D()
            for i in range(I):
                (lattice)=step(lattice)
            for k in range(C):
                for i in range(A):
                    (lattice)=step(lattice)
                M+=calc_magn(lattice)
            final_M[j]=M/C
            M=0
        plt.plot(np.arange(-1.0,1.1,0.1),final_M,color=colours[m+1],\
            use J = 0.2 / T to see smoother transition
```

Also with increasing lattice size these values typically go up so its not a good idea to use fixed values for all lattice sizes!

Thermalize for $I=100$ for better results
Typical Autocorrelation size $4w$ 10-20 is used.
Use atleast 1000 for smoother curve.



Result your code gives after all suggested changes.

```

label='$N_x=N_y=%s'%str((m+1)*4))
plt.legend()
plt.xlabel(r"$h$")
plt.ylabel(r"$m(h)$")
plt.show()


```

Input: None; Output: None

This function calculates the thermodynamically expected value of the magnetization per spin. This is done for $N_x = N_y \in [4, 20]$ and $h \in [-1, 1]$ and $J = 0.5$. We plot our results as shown below.

We can see that the magnetization highly depends on h . This is because the spins align according to the external magnetic field as long as the coupling J is strong enough and the temperature T is low enough. For the sake of dimensionlessness we use the $[J] = \frac{1}{T}$ and therefor $J = 0.5$ fulfills these conditions.

[26]: `m_h_5()`



Point
deduced
as the
sampled values
are not
normalized
properly!

```

[27]: def eps_J_6():
    global N_x
    global N_y
    global h
    global J

```

E=16 #amount of points for the graph

```
I=10 #initialization
A=2 #autocorrelationcorrection
C=100 #Number of components for an ensemble
D=5
```


```
lattice=init_2D()
colours=['b','g','r','c','m','y','k']
h=0.0
```

```
final_eps=np.zeros(E)
eps_theo_array=np.zeros(E)
for i in range(E):
    J=0.25+(i*0.75/(E-1))
    eps_theo_array[i]=eps_theo()
```

```
for m in range(D):
    N_x=N_y=(m+1)*4
    final_eps=np.zeros(E)
    for j in range(E):
        J=0.25+(j*0.75/(E-1))
        lattice=init_2D()
        En=0
        for i in range(I):
            (lattice)=step(lattice)
        for k in range(C):
            for i in range(A):
                (lattice)=step(lattice)
            En+=calc_eps(lattice)
        final_eps[j]=En/C
    plt.plot(np.arange(0.25,1.05,0.
```

```
→05),final_eps,color=colours[m+1],label='$N_x=N_y=%s'%str((m+1)*4))
    plt.plot(np.arange(0.25,1.05,0.05),eps_theo_array,label='$N_x=N_y=\infty$,
→(theo)')
    plt.legend()
    plt.xlabel(r"$J$")
    plt.ylabel(r"$\epsilon(J)$")
    plt.show()
```

Implementing same suggestion
here gives you a better
plot.



Input: None; Output: None


This function calculates the thermodynamically expected value of the energy per spin. This is done for $N_x = N_y \in [4, 20]$ and $J \in [0.25, 1]$ (We don't use $J \in [0.25, 2]$ because for $J \in [1, 2]$ there is nothing interesting to see and this costs valuable computation time). We set $h = 0$. We plot our results as well as the theoretically expected solution in the thermodynamic limit.

As one can see our result matches the theoretical expected solution very well. A close look at the curve shows a little kink in the area $J \in [0.4, 0.5]$. The curve shows a almost linear behaviour which is not surprising because with rising J the spins align pretty quickly and for four nearest neighbors the energy per spin should be close to $4 \cdot J$ when $h = 0$.

but there
is interesting
stuff below
0.25

nice!

[28]: `eps_J_6()`



```
[29]: def abs_m_J_7():
    global N_x
    global N_y
    global h
    global J

    E=16 #amount of points for the graph
    I=10 #initialization
    A=2 #autocorrelationcorrection
    C=100 #Number of components for an ensemble
    D=5
    lattice=init_2D()
    colours=['b','g','r','c','m','y','k']
    h=0.0
    final=np.zeros(E)
    abs_m_theo_array=np.zeros(E)
    for i in range(E):
        J=0.25+(i*0.75/(E-1))
        abs_m_theo_array[i]=abs_m_theo()
```



```

for m in range(D):
    N_x=N_y=(m+1)*4
    final=np.zeros(E)
    for j in range(E):
        J=(0.25+(j*0.75/(E-1)))/T
        lattice=init_2D()
        abs_m=0
        for i in range(I):
            (lattice)=step(lattice)
        for k in range(C):
            for i in range(A):
                (lattice)=step(lattice)
            abs_m+=np.abs(calc_magn(lattice))
        final[j]=abs_m/C
    plt.plot((np.arange(0.25,1.05,0.05)/T)**(-1),final,color=colours[m+1],\
            label='$N_x=N_y=%s'%str((m+1)*4))
plt.plot((np.arange(0.25,1.05,0.05)/T)**(-1),abs_m_theo_array,\
        label='$N_x=N_y=\infty$ (theo)')
plt.legend()
plt.xlabel(r"$J^{-1}$")
plt.ylabel(r"$\langle |m| \rangle$ (J)")
plt.show()

```

Input: None; Output: None

This function calculates the thermodynamically expected value of the absolute magnetization per spin. This is done for $N_x = N_y \in [4, 20]$ and $J \in [0.25, 1]$. We set $h = 0$. We plot our results as a function of J^{-1} as well as the theoretically expected solution in the thermodynamic limit.


Apparently the calculated expected value of the absolute magnetization per spin converges to the theoretically expected solution for increasing size of the system. Here the significance of the critical coupling-value J_c gets apparent. The critical coupling corresponds to a phasetransition, which is visible in the drastic drop of the absolute magnetization around the critical coupling value. We defined our J as $\frac{J}{T}$ for the dimensionlessness and so $\frac{1}{J_c}$ is therefor also a critical temperature.

Its important to plot the absolute value of the magnetization here because as we have seen before the magnetization for $h = 0$ is always 0. This is because the system is isotropic in the beginning and would produce a magnetization in a random direction which would cancel out over multiple constellations.

[30]: `abs_m_J_7()`

Que-4 not answered

2/3




As usual you can get a better plot!

[]:

[]:

drastic drop of the absolute magnetization around the critical coupling
critical temperature.
It's important to plot the absolute value of the magnetization here bec
the system is isotropic in the beginning and would produce a magnet

abs_m_J_7()



phase transition
more
visible in
 $N = 8$!