

Ontologies Classes Object Properties Data Properties Annotation Properties Individuals Datatypes Clouds

## Class: Warnings\_and\_Precautions\_AFINI

### Annotations (1)

- `rdfs:comment` "Non-Infectious Pneumonitis: Monitor for clinical symptoms or radiological changes. Withhold or permanently discontinue based on severity. (2.9, 5.1) Infections: Monitor for signs and symptoms of infection. Withhold or permanently discontinue based on severity. (2.9, 5.2) Severe Hypersensitivity Reactions: Permanently discontinue for clinically significant hypersensitivity. (5.3) Angioedema: Patients taking concomitant angiotensin-converting-enzyme (ACE) inhibitors may be at increased risk for angioedema. Permanently discontinue for angioedema. (5.4, 7.2) Stomatitis: Initiate dexamethasone alcohol-free mouthwash when starting treatment. (5.5, 6.1) Renal Failure: Monitor renal function prior to treatment and periodically thereafter. (5.6) Risk of Impaired Wound Healing: Withhold for at least 1 week prior to elective surgery. Do not administer for at least 2 weeks following major surgery and until adequate wound healing. The safety of resumption of treatment after resolution of wound healing complications has not been established. (5.7) Geriatric Patients: Monitor and adjust dose for adverse reactions. (5.8) Metabolic Disorders: Monitor serum glucose and lipids prior to treatment and periodically thereafter. Withhold or permanently discontinue based on severity. (2.9, 5.9) Myelosuppression: Monitor hematologic parameters prior to treatment and periodically thereafter. Withhold or permanently discontinue based on severity. (2.9, 5.10) Risk of Infection or Reduced Immune Response with Vaccination: Avoid live vaccines and close contact with those who have received live vaccines. Complete recommended childhood vaccinations prior to starting treatment. (5.11) Radiation Sensitization and Radiation Recall: Severe radiation reactions may occur. (5.12, 6.2) Embryo-Fetal Toxicity: Can cause fetal harm. Advise patients of reproductive potential of the potential risk to a fetus and to use effective contraception. (5.13, 8.1, 8.3)"(`xsd:string`)

### Superclasses (2)

- `'\Afinitor_(Everolimus)\'`
- `'\Afinitor_(Everolimus)_\'`

### Disjoints (8)

Adverse\_Reactions\_AFINI, Contraindications\_AFINI, Dosage\_and\_Administration\_AFINI, Dosage\_Forms\_and\_Strengths\_AFINI, Drug\_Interactions\_AFINI, Indications\_and\_Usage\_AFINI, Use\_in\_Specific\_Populations\_AFINI, **Warnings\_and\_Precautions\_AFINI**

OWL HTML inside