

Contents

5	Sheaves of Modules	1
5.1	\mathcal{O}_X -modules	1
5.2	locally free \mathcal{O}_X -modules	2
5.3	\mathcal{O}_X -modules on affine schemes	2
5.4	Quasicoherent \mathcal{O}_X -modules	3
5.5	Coherent sheaves	4
5.6	Functoriality	4
5.7	\mathcal{O}_X -modules on a projective scheme	5
5.8	Morphisms to projective spaces	6
5.9	Divisors	8
5.10	Differentials	8
6	Cohomology	10
6.1	Some (quick) homological algebra	10
6.2	Cech cohomology	11
6.3	Cohomology of projective space	12
6.4	Higher direct images	12
6.5	Duality	13
6.6	Flat base change	13
7	Riemann-Roch and Serre duality	14
7.1	Divisors on curves	14
7.2	Riemann-Roch	14
7.3	Residues	15
7.4	Serre duality	15
7.5	Some consequences of Riemann-Roch	15
7.6	Embeddings in projective space	16
7.7	Hyperelliptic curves	16
7.8	Coverings	17

Propositions and Theorems

5 Sheaves of Modules

5.1 \mathcal{O}_X -modules

Fact. For any \mathcal{O}_X -module \mathcal{F} and any $x \in X$ the stalk $\mathcal{F}_x = \varinjlim \mathcal{F}(U)$ is a module over $\mathcal{O}_{X,x}$

Lemma. For any presheaf of \mathcal{O}_X -modules \mathcal{F} , the sheafification $\tilde{\mathcal{F}}$ is naturally a sheaf of \mathcal{O}_X -modules and it satisfies the analogous universal property:

For all sheaves of \mathcal{O}_X -modules \mathcal{G} : $\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \cong \text{Hom}_{\mathcal{O}_X}(\tilde{\mathcal{F}}, \mathcal{G})$

Fact. for \mathcal{F}_i \mathcal{O}_X -modules, $\prod \mathcal{F}_i$ is a sheaf (since \prod is a \varprojlim)

Fact. $\bigoplus \mathcal{F}_i$ is the sheafification of $U \mapsto \bigoplus \mathcal{F}_i(U)$

It is a subsheaf of $\prod \mathcal{F}_i$ with equality if $|I| < \infty$

Universal Property. $\text{Hom}_{\mathcal{O}_X}(\bigoplus \mathcal{F}_i, \mathcal{G}) \cong \prod \text{Hom}_{\mathcal{O}_X}(\mathcal{F}_i, \mathcal{G})$ for all \mathcal{O}_X -modules \mathcal{G}

Fact. There is a natural homomorphism $\mathcal{F}(U) \otimes \mathcal{G}(U) \rightarrow \mathcal{F} \otimes \mathcal{G}(U)$

Prop. Basic Properties:

1. $\mathcal{F} \otimes \mathcal{G} \cong \mathcal{G} \otimes \mathcal{F}$
2. $(\mathcal{F} \otimes \mathcal{G}) \otimes \mathcal{H} \cong \mathcal{F} \otimes (\mathcal{G} \otimes \mathcal{H})$
3. $\mathcal{O}_X \otimes \mathcal{F} \cong \mathcal{F}$
4. $(\bigoplus \mathcal{F}_i) \otimes \mathcal{G} \cong \bigoplus (\mathcal{F}_i \otimes \mathcal{G})$

Prop. for any homomorphism of \mathcal{O}_X -modules $\varphi : \mathcal{F} \rightarrow \mathcal{G}$, $\ker(\varphi)$, $\text{im}(\varphi)$ and $\text{coker}(\varphi)$ are \mathcal{O}_X -modules

5.2 locally free \mathcal{O}_X -modules

Prop. \mathcal{F}, \mathcal{G} locally free of rank $r, s \Rightarrow \mathcal{F} \oplus \mathcal{G}$ free of rank $r + s$ and $\mathcal{F} \otimes \mathcal{G}$ free of rank $r \cdot s$

Prop. 1. \mathcal{L} invertible $\Rightarrow \mathcal{L}^\vee \otimes \mathcal{L} \cong \mathcal{O}_X$

2. The set of isomorphism classes of invertible sheaves on X form an abelian group $\text{Pic}(X)$ called the picard group of X

5.3 \mathcal{O}_X -modules on affine schemes

Take $X = \text{Spec } R$ and M an R -module and $\mathcal{B} = \{D_f | f \in R\}$ the basis of the Zariski topology.

For all $f \in R$ set $\tilde{M}(D_f) := M_f = M \otimes_R R_f$
and all $f, g \in R$ with $D_g \subset D_f \rightsquigarrow R_f \rightarrow R_g$
 $\rightsquigarrow \tilde{M}(D_f) = R_f \otimes_R M \rightarrow R_g \otimes_R M = \tilde{M}(D_g)$
 $\Rightarrow \tilde{M}$ is a presheaf on \mathcal{B}

Prop. \tilde{M} is a sheaf and extends uniquely to a sheaf of abelian groups on X .
 $R \times M \rightarrow M$ induces a scalar $\mathcal{O}_X \times \tilde{M} \rightarrow \tilde{M} \Rightarrow \tilde{M}$ is an \mathcal{O}_X -module

Fact. $\tilde{M}|_{D_f} \cong \tilde{M}_f$

$\forall x \in X : (\tilde{M})_x = \varinjlim \tilde{M}(U) = \varinjlim_{f \in R \setminus \mathfrak{p}} M_f = M_f$

Functoriality:

Any R -homomorphism $M \xrightarrow{\varphi} N$ induces $\text{id} \otimes \varphi : R_f \otimes M \rightarrow R_f \otimes N$
 $\rightsquigarrow \mathcal{O}_X$ -module homomorphism $\tilde{\varphi} : \tilde{M} \rightarrow \tilde{N}$
 $\Rightarrow \text{Functor } ((R\text{-mod})) \rightarrow ((\mathcal{O}_X\text{-mod}))$

Lemma. For any \mathcal{O}_X -module \mathcal{G} there is a natural isomorphism:

$\text{Hom}_{\mathcal{O}_X}(\tilde{M}, \mathcal{G}) \xrightarrow{\sim} \text{Hom}_R(M, \mathcal{G}(X))$

$\psi \mapsto (\psi(X) : \tilde{M}(X) = M \rightarrow \mathcal{G}(X))$ is an $\mathcal{O}_X(X)$ -module homomorphism i.e. an R -mod hom

Prop. *The Functor $M \rightarrow \tilde{M}$ is fully faithful*

Prop. 1. *the functor is exact*

2. *commutes with direct sum*

3. *commutes with tensor product, i.e. $\tilde{M} \otimes_{\mathcal{O}_X} \tilde{N} \cong (M \otimes_R N)^\sim$*

Important: $\tilde{M}(X) = M$ so the global sections functor is exact on sheaves of the form \tilde{M}

5.4 Quasicoherent \mathcal{O}_X -modules

Let X be an arbitrary scheme.

Theorem. *TFAE:*

1. *For all open affine $U = \text{Spec } R \subset X$ there exists an R -module M such that $\mathcal{F}|_U \cong \tilde{M}$*
2. *There exists an open affine covering of such U*
3. *\mathcal{F} is quasicoherent*
4. *for all open affine $U = \text{Spec } R \subset X$ and all $f \in R$ the homomorphism $R_f \otimes_R \mathcal{F}(U) \xrightarrow{\text{res, mult}} \mathcal{F}(D_f^U)$ is an isomorphism*

- Theorem.** 1. *For all homomorphisms of quasicoherent \mathcal{O}_X -modules, \ker , im , coker are quasicoherent*
2. *Direct sum of quasicoherent \mathcal{O}_X -modules are quasicoherent.*
3. *Let \mathcal{F}_i be quasicoherent submodules of a quasicoherent \mathcal{O}_X -module \mathcal{F} then $\sum \mathcal{F}_i, \bigcap \mathcal{F}_i$ are quasicoherent.*
4. *if \mathcal{F}, \mathcal{G} are quasicoherent then $\mathcal{F} \otimes \mathcal{G}$ is quasicoherent and for all open affine $U : \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U) \xrightarrow{\sim} (\mathcal{F} \otimes \mathcal{G})(U)$ is an iso*

Corollary. *The closed subschemes of X correspond to the quasicoherent ideal sheaves of \mathcal{O}_X*

Theorem. *For $X = \text{Spec } R$ we have an equivalence of categories*

$$\begin{aligned} ((R - \text{mod})) &\Leftrightarrow ((\text{quasicoherent } \mathcal{O}_X - \text{modules})) \\ M &\longmapsto \tilde{M} \\ \mathcal{F}(x) &\longleftarrow \mathcal{F} \end{aligned}$$

and both functors are exact

5.5 Coherent sheaves

Prop. For any quasicoherent \mathcal{O}_X -module we have the following implications:

1. \mathcal{F} is coherent

\Downarrow

2. \mathcal{F} is finitely generated

\Updownarrow

3. $\forall U \subset X$ open affine $\mathcal{F}(U)$ is a finitely generated $\mathcal{O}_X(U)$ -module

If X is locally noetherian all are equivalent.

Prop. Assume X is locally noetherian. Then for all coherent sheaves \mathcal{F}, \mathcal{G} :
 $\mathcal{F} \oplus \mathcal{G}$, $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$, $\mathcal{H}om(\mathcal{F}, \mathcal{G})$ as well as \ker , im and coker of any homomorphism are coherent.

5.6 Functoriality

Consider a morphism $f : X \rightarrow Y$ and sheaves \mathcal{F}, \mathcal{G} on X and Y respectively.

Prop. If f is affine¹ and \mathcal{F} is quasicoherent then $f_*\mathcal{F}$ is quasicoherent.

Prop. If f is finite, X, Y are locally noetherian and \mathcal{F} is coherent then $f_*\mathcal{F}$ is coherent.

Prop. f_* naturally extends to a functor which commutes with products and is left exact.

Prop. There exists a natural isomorphism $(f^*\mathcal{G})_x \cong \mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,f(x)}} \mathcal{G}_{f(x)}$

Prop. f^* naturally extends to a functor which commutes with \oplus and is right exact and satisfies:

$$f^*(\mathcal{G} \otimes_{\mathcal{O}_Y} \mathcal{H}) \cong (f^*\mathcal{G}) \otimes_{\mathcal{O}_X} (f^*\mathcal{H})$$

Prop. If $X = \text{Spec } R$ and $Y = \text{Spec } S$ and $\mathcal{G} = \tilde{N}$ on Y then $f^*\mathcal{G} \cong (R \otimes_S N)^\sim$

Prop. If \mathcal{G} is

- quasicoherent
- finitely generated
- free
- locally free
- coherent (if X, Y are locally noetherian)

then so is $f^*\mathcal{G}$

Prop. Adjunction:

$$\text{Hom}_{\mathcal{O}_X}(f^*\mathcal{G}, \mathcal{F}) \cong \text{Hom}_{\mathcal{O}_Y}(\mathcal{G}, f_*\mathcal{F})$$

this is functorial in \mathcal{F} and \mathcal{G} .

¹ f is affine if the inverse image of every open affine is open affine

5.7 \mathcal{O}_X -modules on a projective scheme

Take $X = \text{Proj } R$ for a graded ring $R = \bigoplus_{d \geq 0} R_d$

Prop. *There exists a functor*

$$((\text{graded } R - \text{mod})) \longrightarrow ((\text{quasicoherent } \mathcal{O}_X - \text{module}))$$

$$M \longrightarrow \tilde{M}$$

this functor is exact and commutes with \bigoplus and \bigotimes

Prop. 1. $\mathcal{O}_X(n)$ is an invertible sheaf

$$2. \tilde{M}(n) \cong \tilde{M} \otimes_{\mathcal{O}_X} \mathcal{O}_X(n)$$

Corollary. $\forall m, n \in \mathbb{Z} : \mathcal{O}_X(m+n) \cong \mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{O}_X(n)$

$$\Rightarrow \mathcal{O}_X(n) \cong \mathcal{O}_X(1)^{\otimes n} \text{ for all } n \geq 1$$

$$\mathcal{O}_X(-1) \cong \mathcal{O}_X(1)^\vee$$

Lemma. *Let \mathcal{F} be a quasicoherent \mathcal{O}_X -module and $f \in \mathcal{F}(X)$*

1. *If X is quasicompact and $f|_{D_f} = 0$ there exists an $n \geq 0$ such that $f \otimes s^n = 0$ in $(\mathcal{F} \otimes \mathcal{L}^{\otimes n})(X)$*
2. *If X is quasicompact and separated for any $g \in \mathcal{F}(D_s)$ there exists n_0 such that $\forall n \geq n_0 \quad g \otimes s^n|_{D_s} \in (\mathcal{F} \otimes \mathcal{L}^{\otimes n})(D_s)$ extends to an element of $(\mathcal{F} \otimes \mathcal{L}^{\otimes n})(X)$*

(if $X = \text{Spec } R, \mathcal{L} = \mathcal{O}_X, \mathcal{F} = \tilde{M}$)

1. *$f = m \in M$ if $\frac{m}{1} = 0$ in M_s then $\exists n \quad m \cdot s^n = 0 \in M$*
2. *Take $n \in M_s$ write $n = \frac{m}{s^{n_0}}$*

Theorem. *For any finitely generated quasicoherent sheaf \mathcal{F} on X :*

1. *$\exists n_0 \forall n \geq n_0 \quad \mathcal{F}(n)$ is generated by global sections.*
2. *$\exists m \exists r \geq 0 \exists$ surjective homomorphism $\mathcal{O}_X(m)^{\oplus r} \rightarrow \mathcal{F}$*

Construction:

1. for a graded R -module M and $n \in \mathbb{Z}$ there exists a natural map

$$M_n \longrightarrow \tilde{M}(n)(X) = \widetilde{M(n)}(X)$$

this map is R_0 -linear

2. this is multiplicative

3. Set $R' = \bigoplus_{d \geq 0} \mathcal{O}_X(d)(X)$. it is a graded ring with a graded ring homomorphism $R \rightarrow R'$ and a graded R -module homomorphism $M \rightarrow M' = \bigoplus_{n \in \mathbb{Z}} \tilde{M}(n)(X)$

4. for any \mathcal{O}_X -module \mathcal{F} and any $n_0 \in \mathbb{Z} \cup \{-\infty\}$

set: $M' := \bigoplus_{n \geq n_0} \mathcal{F}(n)(X)$

This is naturally a graded R' -module; hence a graded R -module

Theorem. For any quasicoherent \mathcal{O}_X -module \mathcal{F} there is a natural isomorphism

$\widetilde{M'} \longrightarrow \mathcal{F}$ for M' as in (4.)

If $\mathcal{F} = \widetilde{M}$ this is the inverse to the homomorphism
 $M \longrightarrow M'$ induced by (3.)

Corollary. Every quasicoherent \mathcal{O}_X -module on $\text{Proj } R$ is isomorphic to \widetilde{M} for some graded R -module M

5.8 Morphisms to projective spaces

Prop. For all schemes and all $n \geq 0$ there is a natural bijection:

$\text{Mor}(X, \mathbb{P}^n) \cong \{(\mathcal{L}, l_0, \dots, l_n) | \mathcal{L} \text{ invertible sheaf on } X, l_0, \dots, l_n \in \mathcal{L}(X) \text{ generate } \mathcal{L}\}$

\cong isos of $\mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ mapping $l_i \mapsto l'_i$

Prop. Let $(\mathcal{L}, l_0, \dots, l_n)$ be as above and let $l_{n+1}, \dots, l_m \in \mathcal{L}(X)$ then we have the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{(\mathcal{L}l_0 \dots l_n)} & \mathbb{P}^n \\ & \searrow (\mathcal{L}l_0 l_m) & \uparrow \text{proj} \\ & & \bigcup_{i=0}^n \overline{D}_{X_i}^{p_m} \subset \mathbb{P}^m \end{array}$$

Now let X be a scheme over $\text{Spec } R$

Prop. For X noetherian we have for all very ample sheaves \mathcal{L} and all invertible sheaves \mathcal{L}' on X that is generated by global sections: $\mathcal{L} \otimes \mathcal{L}'$ is very ample.

Note. very ample \Rightarrow generated by global sections

Corollary. For $\mathcal{L}, \mathcal{L}'$ very ample $\Rightarrow \mathcal{L} \otimes \mathcal{L}'$ and $\mathcal{L}^{\otimes n} \quad \forall n \geq 1$ are very ample

Prop. TFAE:

1. \mathcal{L} ample
2. $\forall m \geq 1 : \mathcal{L}^{\otimes m}$ ample
3. $\exists m \geq 1 : \mathcal{L}^{\otimes m}$ ample

Prop. very ample \Rightarrow ample

Theorem. *If X is separated of finite type over $\text{Spec } R$ then:*

\mathcal{L} ample $\Rightarrow \exists m \geq 1 : \mathcal{L}^{\otimes m}$ very ample

Corollary. *same assumptions \Rightarrow TFAE:*

1. \mathcal{L} ample
2. $\exists m \geq 1 : \mathcal{L}^{\otimes m}$ very ample
3. $\exists m_0 \forall m \geq m_0 : \mathcal{L}^{\otimes m}$ very ample

Note. *very ample is a relative notion for X over $\text{Spec } R$. Ample is an absolute notion*

want: true relative notions for X over arbitrary scheme S

Prop. *If $X \rightarrow Y$ is separated of finite type with Y noetherian, TFAE:*

1. \mathcal{L} is relatively ample over Y
2. $\exists m \geq 1 : \mathcal{L}^{\otimes m}$ is relatively ample over Y
3. $\exists m_0 : \forall m \geq m_0 : \mathcal{L}^{\otimes m}$ is relatively very ample over Y

Prop. *let the following diagram be cartesian:*

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & \square & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

with all schemes noetherian and f separated of finite type \mathcal{L} invertible on X which is relatively {ample/ very ample} over Y then $g'^\mathcal{L}$ is as well over Y'*

Lemma. *Take morphisms $X \xrightarrow{f} Y \xrightarrow{g} Z = \text{Spec } R$ which are separated of finite type with R noetherian. Let \mathcal{L} be an invertible sheaf on X which is relatively very ample over Y and let \mathcal{M} be an invertible sheaf on Y which is very ample.*

Then: $\exists l_0 \forall l \geq l_0 :$

1. $\mathcal{L} \otimes f^*\mathcal{M}^{\otimes l} \cong \varepsilon^*\text{pr}_1^*\mathcal{O}(1)$ for some locally closed embedding $\varepsilon : X \hookrightarrow \mathbb{P}^N \times Y$
2. $\mathcal{L} \otimes f^*\mathcal{M}^{\otimes(l+1)}$ is very ample for X over S .

Prop. *Let $f : X \rightarrow Y$ be a quasicompact morphism and \mathcal{L} an invertible sheaf on X . Let f be separated of finite type with Y noetherian then relatively {ample / very ample} is equivalent to saying for all open coverings fulfill the respective ampleness property.*

Prop. *Take $X \xrightarrow{f} Y \xrightarrow{g} Z$ separated of finite type with Z noetherian, \mathcal{L} an invertible sheaf on X relatively {ample / very ample} over Y and \mathcal{M} ditto over Z*

Then $\exists l_0 \forall l > l_0 : \mathcal{L} \otimes f^\mathcal{M}^{\otimes l}$ is ditto over Z*

Prop. *$X \xrightarrow{f} Y \xrightarrow{g} Z$ both {projective/ quasiprojective} with Z noetherian. Then $g \circ f$ ditto.*

5.9 Divisors

Prop. $\mathcal{O}_X(D)$ is an invertible sheaf and every invertible sheaf, which is a subsheaf of \mathcal{K}_X , arises like this from a unique Cartier divisor.

Prop. 1. $\forall D, D' : \mathcal{O}_X(D + D') \cong \mathcal{O}_X(D) \otimes \mathcal{O}_X(D')$

2. $\forall D : \mathcal{O}_X(-D) \cong \mathcal{O}_X(D)^\vee$

Theorem. The map $D \mapsto \mathcal{O}_X(D)$ induces an isomorphism

$$\text{Div Cl}(X) \longrightarrow \text{Pic}(X)$$

Prop. X regular \implies locally factorial

Now assume X is noetherian integral and locally factorial

Prop. For any Cartier divisor $D = ([f_x])_{x \in X}$ on X the sum $\text{cyc}(D) := \sum_{\substack{y \in X \\ \dim \mathcal{O}_{X,y} = 1}} \text{ord}_y(f_y) \cdot \overline{\{y\}}^{\text{red}}$ is finite, so is $Z^1(X)$.

Theorem. there is a group isomorphism $\text{Div}(X) \rightarrow Z^1(X) \quad D \mapsto \text{cyc}(D)$

Prop. For any field k , $\text{Cl}(\mathbb{A}_k^n) = 0 \implies \text{Pic}(\mathbb{A}_k^n) = 0 \quad \forall n \geq 0$

Theorem. For any field k and any $n \geq 1$

$$\text{Cl}(\mathbb{P}_k^n) \cong \text{Pic}(\mathbb{P}_k^n) \cong \mathbb{Z}$$

generated by $[H] \longleftrightarrow [\mathcal{O}_X(1)]$

Prop. D is equivalent to an effective divisor iff $\Gamma(X, \mathcal{O}_X(D)) \neq 0$

Prop. If X possesses a very ample divisor, every divisor is a difference of very ample divisors.

Prop. $\varphi^* : \text{Div}(Y) \rightarrow \text{Div}(X) \quad ([f_y])_y \mapsto ([\varphi^* f_{\varphi(x)}])_{x \in X}$
is a well defined homomorphism and $\varphi^* \mathcal{O}_Y(D) \cong \mathcal{O}_X(\varphi^* D)$

Prop. If D is the effective divisor corresponding to the locally principal closed subscheme $T \subset Y$ then $\varphi^* D$ corresponds to $\varphi^{-1}(T) := X \times_Y T$

5.10 Differentials

Fact. $\text{Der}_A(B, M)$ is a B -module by the action of B on M .

Fact. Any B -module homomorphism $f : M \rightarrow N$ induces a B -module homomorphism $\text{Der}_A(B, M) \rightarrow \text{Der}_A(B, N) \quad d \mapsto f \circ d$.

Fact. There is a natural bijection

$$\text{Der}_A(B, M) \cong \{\varphi : B \rightarrow B \ltimes M \text{ } A\text{-algebra homos with } \pi \circ \varphi = \text{id}_B\} \quad d \leftrightarrow (\text{id}, d)$$

Prop. $(\Omega_{B/A}, d)$ is unique up to unique isomorphism.

Prop. I/I^2 with d is a module of differentials of B over A .

Prop. $\Omega_{A[X_1, \dots, X_n]/A}$ is a free $A[X_1, \dots, X_n]$ module with basis dX_1, \dots, dX_n

Fact. $\Omega_{A/A} = 0$

Prop. If $B' = B \otimes_A A'$ then $B' \otimes_B \Omega_{B/A} \xrightarrow{\sim} \Omega_{B'/A'}$

Prop. For any multiplicative system $S \subset B : S^{-1}\Omega_{B/A} \cong \Omega_{S^{-1}B/A}$.

Prop. For any ring homomorphisms $A \rightarrow B \rightarrow C$ there exists a natural exact sequence $C \otimes_B \Omega_{B/A} \rightarrow \Omega_{C/A} \rightarrow \Omega_{C/B} \rightarrow 0$

Prop. For all ideals $J \subset B$ set $C := B/J \Rightarrow$ there is a natural exact sequence $J/J^2 \rightarrow C \otimes_B \Omega_{B/A} \rightarrow \Omega_{C/A} \rightarrow 0$

Corollary. If B is a finitely generated A -algebra, then $\Omega_{B/A}$ is a finitely generated B -module.

Prop. L/K finite field extension $\Omega_{L/K} = 0 \Leftrightarrow$ separable.

Prop. For any local k -algebra B with $B/\mathfrak{m} \cong k$ there exists a natural isomorphism $\mathfrak{m}/\mathfrak{m}^2 \rightarrow (B/\mathfrak{m}) \otimes_B \Omega_{B/k}$

Theorem. Let $B = k \oplus \mathfrak{m}$ local ring, k perfect, B is a localization of a finitely generated k -algebra.

Then $\Omega_{B/k}$ is a free B -module of finite rank $\dim(B)$ iff B is a regular local ring.

Prop. $Y \rightarrow X$ locally of finite type $\Rightarrow \Omega_{Y/X}$ finitely generated.

Prop.
$$\begin{array}{ccc} Y & \longrightarrow & X \\ \cup & & \cup \\ V & \longrightarrow & U \end{array} \text{ open} \Rightarrow \Omega_{V/U} = \Omega_{Y/X}|_V$$

Prop.
$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \uparrow g' & \square & \uparrow \\ Y' & \longrightarrow & X' \end{array} \Rightarrow \Omega_{Y'/X'} \cong g'^*\Omega_{Y/X}$$

Prop. $Z \xrightarrow{g} Y \xrightarrow{f} X \Rightarrow$ exact sequence: $g^*\Omega_{Y/X} \rightarrow \Omega_{Z/X} \rightarrow \Omega_{Z/Y} \rightarrow 0$

Prop. If $Z \xrightarrow{i} Y$ is a closed embedding with sheaf of ideals $J \Rightarrow$ exact sequence $J/J^2 \rightarrow i^*\Omega_{Y/X} \rightarrow \Omega_{Z/X} \rightarrow 0$

Prop. $Y = Y_1 \times_X Y_2 \Rightarrow \Omega_{Y/X} \cong \text{pr}_1^*\Omega_{Y_1/X} \oplus \text{pr}_2^*\Omega_{Y_2/X}$

Theorem. There is a natural short exact sequence for $X := \mathbb{P}_Y^n$:

$$0 \rightarrow \Omega_{X/Y} \rightarrow \mathcal{O}_X(-1)^{n+1} \rightarrow \mathcal{O}_X \rightarrow 0$$

Theorem. Any localization of a regular local ring at a prime ideal is regular.

Theorem. Let X be connected of finite type over an algebraically closed field k . Then X is regular iff $\Omega_{X/k}$ is locally free of rank $\dim(X)$

Corollary. X reduced of finite type over k algebraically closed \Rightarrow there is an open dense subscheme $U \subset X$ which is regular

Theorem. Let X be regular of finite type over k algebraically closed. Let $Y \subset X$ be the closed subscheme associated to $J \subset \mathcal{O}_X$

Then Y is regular iff $\Omega_{Y/k}$ is locally free and the second sequence

$$0 \rightarrow J/J^2 \rightarrow i^* \mathcal{O}_{X/k} \rightarrow \Omega_{Y/k} \rightarrow 0$$

is also left exact.

Then J is locally generated by $\text{codim}(Y) =: r$ elements and J/J^2 is locally free of rank r

Prop. $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ short exact sequence of locally free sheaves of rank n', n, n'' .

There is a natural isomorphism $\bigwedge^n \mathcal{F} \cong \bigwedge^{n'} \mathcal{F}' \otimes \bigwedge^{n''} \mathcal{F}''$

Lemma. $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ short exact sequence of locally free sheaves \Rightarrow

There is a natural isomorphism $\bigwedge^{\text{top}} \mathcal{F} \cong \bigwedge^{\text{top}} \mathcal{F}' \otimes \bigwedge^{\text{top}} \mathcal{F}''$

Corollary. $i^* \omega_{X/Y} \cong i^* \bigwedge^{\text{top}} (J/J^2) \otimes \omega_{Y/k}$

Prop. For any nonsingular closed hypersurface $Y \subset \mathbb{P}_k^n$ for k algebraically closed of degree d i.e. $Y = V(f)$ for $f \neq 0$ homogeneous of degree d

Then $\omega_{Y/k} \cong i^* \mathcal{O}(-n-1+d)$

Corollary. $\omega_{Y/k} \cong \mathcal{O}_Y$ iff $d = n+1$

Prop. Let $Y \subset \mathbb{P}_k^2$ be a nonsingular closed curve of degree d with k algebraically closed. Then:

1. $\deg \omega_{Y/k} = d(d-3)$
2. $\omega_{Y/k}$ is ample iff $d \geq 4$
3. $Y \cong \mathbb{P}_k^1$ iff $d \leq 2$

6 Cohomology

6.1 Some (quick) homological algebra

Fact. Any homomorphism $f : X \rightarrow Y$ induces a homomorphism in homology:

$$H^n(f) : H^n(X) \rightarrow H^n(Y)$$

Prop. $0 \rightarrow X^\bullet \xrightarrow{f} Y^\bullet \xrightarrow{g} Z^\bullet \rightarrow 0$ short exact sequence of complexes \Rightarrow exists natural homomorphisms δ_n yielding a long exact sequence in cohomology:

$$\dots \xrightarrow{\delta_{n-1}} H^n X \xrightarrow{H^n f} H^n Y \xrightarrow{H^n g} H^n Z \xrightarrow{\delta_n} H^{n+1} X \rightarrow \dots$$

Prop. f, g homotopic $\Rightarrow \forall n : H^n f = H^n g$

6.2 Čech cohomology

Prop. The projection $C_{\text{alt}}^\bullet(\mathcal{U}, \mathcal{F}) \rightarrow C_{\text{ord}}^\bullet(\mathcal{U}, \mathcal{F})$ is an isomorphism of complexes.

Prop. The inclusion $C_{\text{alt}}^\bullet(\mathcal{U}, \mathcal{F}) \xrightarrow{\varepsilon} C^\bullet(\mathcal{U}, \mathcal{F})$ is a quasiisomorphism.

Corollary.

$$H^n(\mathcal{U}, \mathcal{F}) \xleftarrow[H^n \varepsilon]{\sim} H^n(C_{\text{alt}}^\bullet(\mathcal{U}, \mathcal{F})) \xrightarrow[H^n \pi]{\sim} H^n(C_{\text{ord}}^\bullet(\mathcal{U}, \mathcal{F}))$$

Prop. $H^n(\sigma^*)$ is independent of σ^*

Prop. If \mathcal{U} and \mathcal{V} are refinements of each other then $H^n(\sigma^*)$ is an isomorphism.

Prop. Any homomorphism of sheaves of abelian groups $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ induces a natural homomorphism in cohomology $H^n \varphi : H^n(X, \mathcal{F}) \rightarrow H^n(X, \mathcal{G})$

Now let \mathcal{F} be a quasicoherent sheaf on a scheme X

Theorem. For X affine $\Rightarrow H^n(X, \mathcal{F}) = 0$ for all $n > 0$

Prop. For any closed embedding $i : Y \hookrightarrow X$ and any \mathcal{O}_X -module \mathcal{F} and any $p : H^p(Y, \mathcal{F}) \cong H^p(X, i_* \mathcal{F})$

Theorem. X projective over $\text{Spec } A$ with A noetherian, \mathcal{L} very ample invertible sheaf on X , \mathcal{F} coherent sheaf on X

1. $\forall : H^p(X, \mathcal{F})$ is a finitely generated A -module

2. $\exists m_0 \forall m \geq m_0 \forall p > 0 : H^p(X, \mathcal{F} \otimes \mathcal{L}^{\otimes m}) = 0$

Theorem. Let \mathcal{F} be a quasicoherent sheaf on a separated scheme X . Then for any affine open covering \mathcal{U} of X :

$$H^p(\mathcal{U}, \mathcal{F}) \xrightarrow{\sim} H^p(X, \mathcal{F})$$

Theorem. Let X be separated which possesses an open covering by $d+1$ open affines. Then $\forall p > d \forall \mathcal{F}$ quasicoherent on $X : H^p(X, \mathcal{F}) = 0$

Prop. X quasiprojective of dimension d over $k \Rightarrow X$ can be covered by $d+1$ open affines.

Corollary. X quasiprojective over k , \mathcal{F} quasicoherent $\Rightarrow \forall p > \dim(X) : H^p(X, \mathcal{F}) = 0$

Prop. X separated $0 \rightarrow \mathcal{F}' \xrightarrow{\varepsilon} \mathcal{F} \xrightarrow{\pi} \mathcal{F}'' \rightarrow 0$ short exact sequence of quasicoherent sheaves $\Rightarrow \exists$ natural long exact sequence in cohomology

$$\begin{aligned} 0 \rightarrow \mathcal{F}'(X) \rightarrow \mathcal{F}(X) \rightarrow \mathcal{F}''(X) \rightarrow H^1(X, \mathcal{F}') \rightarrow \dots \\ \dots \rightarrow H^p(X, \mathcal{F}') \xrightarrow{H^p \varepsilon} H^p(X, \mathcal{F}) \xrightarrow{H^p \pi} H^p(X, \mathcal{F}'') \rightarrow H^{p+1}(X, \mathcal{F}') \end{aligned}$$

Theorem. For X separated and quasicoherent: TFAE:

1. X is affine

2. $\forall \mathcal{F}$ quasicoherent on $X : \forall p > 0 : H^p(X, \mathcal{F}) = 0$

3. dito for $p = 1$

6.3 Cohomology of projective space

Theorem.

$$H^p(\mathbb{P}_A^n, \mathcal{O}_{\mathbb{P}_A^n}(m)) = \begin{cases} \bigoplus_{\substack{d \in \mathbb{Z}^{n+1} \\ \text{all } d_i \geq 0}} A \cdot \underline{X}^d = A[\underline{X}]_m & \text{if } p = 0 \\ \bigoplus_{\substack{d \in \mathbb{Z}^{d+1} \\ \sum d_i = m \\ \text{all } d_i < 0}} A \cdot \underline{X}^d & \text{if } p = n \\ 0 & \text{else} \end{cases}$$

Theorem. Let X be proper over $\text{Spec } A$ for A noetherian and \mathcal{L} an invertible sheaf on X . Then

TFAE:

1. \mathcal{L} is ample
2. \forall coherent $\mathcal{F} \exists m_0 \forall m \geq m_0 \forall p \geq 1 : H^p(X, \mathcal{F}\mathcal{L}^m) = 0$
3. \forall coherent sheaves of ideals $J \subset \mathcal{O}_X \exists m_0 \forall m \geq m_0 : H^1(X, J(m)) = 0$

6.4 Higher direct images

Take $f : X \rightarrow Y$ and sheaves of modules \mathcal{F} on X and \mathcal{G} on Y

Fact. 1. There is a natural homo : $H^p(Y, f_*\mathcal{F}) \rightarrow H^p(X, \mathcal{F})$

2. There is a natural homo

$$\begin{array}{ccc} H^p(Y, \mathcal{G}) & \xrightarrow{\quad} & H^p(X, f^*\mathcal{G}) \\ & \searrow^{H^p(\text{adj})} & \nearrow_{(1.)} \\ & H^p(Y, f_*f^*\mathcal{G}) & \end{array}$$

3. Both are compatible with composition and functorial in \mathcal{F}, \mathcal{G}

Prop. f separated, quasicompact, \mathcal{F} quasicoherent \Rightarrow each $R^p f_*\mathcal{F}$ is quasicoherent and for all open affine $V \subset Y$:

$$(R^p f_*\mathcal{F})(V) = H^p(f^{-1}(V), \mathcal{F}|_{f^{-1}(V)})$$

Theorem (Direct Image Theorem). Let f separated, Y locally noetherian, \mathcal{F} coherent $\Rightarrow \forall p : R^p f_*\mathcal{F}$ is coherent.

Fact. f affine, \mathcal{F} quasicoherent $\Rightarrow \forall p > 0 : R^p f_*\mathcal{F} = 0$

Prop. For $X \xrightarrow{f} Y \xrightarrow{g} Z$ there is a relation between $R^p g_* R^p f_*\mathcal{F}$ and $R^r(g \circ f)_*\mathcal{F}$ (spectral sequences)

6.5 Duality

Prop. For $X = \mathbb{P}_A^n$ and all $m \in \mathbb{Z}$ the map

$$H^0(X; \mathcal{O}_X(m)) \otimes H^n(X, \mathcal{O}_X(-n-1-m)) \longrightarrow H^n(X, \mathcal{O}_X(-n-1)) \cong A$$

is a perfect pairing.

Fact. The isomorphism $H^n(\mathbb{P}_A^n, \mathcal{O}(-n-1)) \cong A$ is not canonical but after $\mathcal{O}(-n-1) \cong \omega_{\mathbb{P}_A^n/A}$ the isomorphism

$$H^n(\mathbb{P}_A^n, \omega_{\mathbb{P}_A^n/A}) \cong A$$

is

Prop. If an r -dualizing sheaf (ω_f, tr_f) exists it is unique up to unique isomorphism.

Theorem. Y locally noetherian $\implies (\omega_{\mathbb{P}_Y^n/Y, \text{tr}_f})$ is an n -dualizing sheaf for $f : \mathbb{P}_Y^n \longrightarrow Y$

Theorem. For any projective morphism $f : X \rightarrow Y$ with Y locally noetherian and all fibers of $\dim \leq r$ the r -dualizing sheaf for f exists.

Theorem. $f : X \rightarrow Y$ projective smooth of relative dimension r , Y locally noetherian $\implies \omega_f := \omega_{X/Y} := \Lambda^r \Omega_{X/Y}$ together with some tr_f is an r -dualizing sheaf.

6.6 Flat base change

Prop. \widetilde{M} on $\text{Spec } A$ is flat iff M is flat over A .

Prop. 1. \mathcal{F} locally free $\implies \mathcal{F}$ flat

2. \mathcal{F} finitely generated, X locally noetherian

Prop. $f : \text{Spec } B \rightarrow \text{Spec } A$ is flat iff B is a flat A -module.

Prop.
$$\begin{array}{ccc} X' & \xrightarrow{g'} & X \\ \downarrow f' & \square & \downarrow f \\ Y' & \xrightarrow{g} & Y \end{array}$$
 For a cartesian diagram with f separated and quasicompact

and \mathcal{F} quasicoherent and g flat then base change is an isomorphism.

Prop. $f : X \rightarrow Y$ projective all fibers of dimension $\leq d$; \mathcal{F} quasicoherent \implies

1. $\forall p > d : R^p f_* \mathcal{F} = 0$

2. $R^d f_*$ is rightexact and commutes with base change

7 Riemann-Roch and Serre duality

7.1 Divisors on curves

Let X be regular integral projective scheme of $\dim 1$ over a field k

Prop. *If D is the effective divisor associated to a finite closed subscheme*

1. $T \subset X$ then $D = \sum_{P \in T} \text{length}(\mathcal{O}_{T,P}) \cdot P$
2. $\deg_k(D) = \sum_{P \in T} \text{length}(\mathcal{O}_{T,P}) \cdot [k(P)/k] = \dim_k \Gamma(T, \mathcal{O}_T) =: \deg T$

Prop. *If k is a finite extension of $k' \Rightarrow \deg_{k'}(D) = \deg_k(D) \cdot [k/k']$*

Prop. *For any field extension k'/k . Let $X' := X_{k'} = X \times_{\text{Spec } k} \text{Spec } k' \xrightarrow{\pi} X$. Assume X' integral then*

$$\deg_{k'}(\pi^* D) = \deg_k(D)$$

Lemma. *For all $P, Q \in |X| \exists R \in |X| : P + Q + R \sim 3P_0$*

Theorem. *The map $|X| \rightarrow Cl^0(X) \quad P \mapsto [P - P_0]$ is bijective.*

7.2 Riemann-Roch

Prop. *For any short exact sequence $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ of coherent sheaves*

$$\chi(X, \mathcal{F}) = \chi(X, \mathcal{F}') + \chi(X, \mathcal{F}'')$$

Lemma. *For any exact sequence of finite dimensional k -vector spaces V^i for $i \in \mathbb{Z}$ and almost all zero:*

$$\sum_i (-1)^i \dim_k V^i = 0$$

Fact. $\forall L/k$ field extension: $X_L \xrightarrow{\pi} X \rightsquigarrow \chi(X, \mathcal{F}) = \chi(X_L, \pi^* \mathcal{F})$

Theorem (Riemann-Roch Version 1). *For any invertible sheaf \mathcal{L} on X :*

$$\chi(X, \mathcal{L}) = 1 - g + \deg(\mathcal{L})$$

Theorem (Serre Duality). *The sheaf $\Omega_{X/k}$ is a dualizing sheaf for X over k i.e. there exists a natural $\text{tr} : H^1(X, \Omega_{X/k}) \rightarrow k$ such that for all quasicoherent \mathcal{F} we have perfect duality*

$$\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \Omega_{X/k}) \times H^1(X, \mathcal{F}) \longrightarrow H^1(X, \Omega_{X/k}) \xrightarrow{\text{tr}} k$$

Theorem (Riemann-Roch Version 2). \mathcal{L} invertible sheaf on $X \implies$

$$h^0(X, \mathcal{L}) - h^0(X, \mathcal{L}^\vee \otimes \Omega_{X/k}) = 1 - g + \deg(\mathcal{L})$$

7.3 Residues

Prop. $\forall \omega \in \widehat{\Omega}_{\mathcal{F}/k} : \text{res}_t(\omega) = \text{res}_t(\omega \circ \varphi)$

Prop. $\forall \omega \in \widehat{\Omega}_{\mathcal{E}/k} : \text{res}_u(\omega) = \text{res}_t(\text{tr}_{\mathcal{E}/\mathcal{F}} \omega)$

Prop. The composite map $\Omega_{K/k} \hookrightarrow \widehat{\Omega}_{\mathcal{F}/k} \xrightarrow{\text{res}_t} k$ is independent of t

Prop. Let $\varphi : X \rightarrow Y$ be a finite separated morphism of such curves corresponding to the finite separated extension of function fields K/L .

$$\begin{array}{ccc} \Omega_{K/k} & \xrightarrow{\quad} & \Omega_{L/K} \\ \cong \nearrow & & \uparrow \text{tr}_{K/L} \otimes \text{id}_{\Omega_{L/k}} \\ K \otimes_L \Omega_{K/L} & & \end{array}$$

Define $\text{tr}_{K/L} :$

Then $\forall \omega \in \Omega_{K/k} \forall Q \in |Y|$

$$\sum_{P \in \varphi^{-1}(Q)} \text{res}_P(\omega) = \text{res}_Q(\text{tr}_{K/L} \omega)$$

Theorem (Residue Theorem).

$$\forall \omega \in \Omega_{K/k} : \sum_{P \in |X|} \text{res}_P \omega = 0$$

7.4 Serre duality

Theorem (Serre Duality). The sheaf $\Omega_{X/k}$ is a dualizing sheaf for X over k i.e. there exists a natural $\text{tr} : H^1(X, \Omega_{X/k}) \rightarrow k$ such that for all quasicoherent \mathcal{F} we have perfect duality

$$\text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \Omega_{X/k}) \times H^1(X, \mathcal{F}) \longrightarrow H^1(X, \Omega_{X/k}) \xrightarrow{\text{tr}} k$$

7.5 Some consequences of Riemann-Roch

Now always: X smooth connected projective curve over k algebraically closed field.

Abbreviate: $h^i(\mathcal{L}) := h^i(X, \mathcal{L})$ $\Omega_X := \Omega_{X/k}$

Prop. $\deg(\Omega_X) = 2g - 2$

Prop. Any smooth closed curve of degree d in \mathbb{P}_k^2 has $g = \frac{(d-1)(d-2)}{2}$

Prop. TFAE:

1. $X \cong \mathbb{P}_k^1$
2. $g = 0$
3. $\deg \Omega_X < 0$

4. $\forall \mathcal{L}$ of degree 1: $h^0(\mathcal{L}) > 1$

5. $\exists \mathcal{L}$ of degree 1: $h^0(\mathcal{L}) > 1$

Prop. 1. $\forall P \in |X| \exists f \in K(X)$ f has a pole at P and no other pole

2. $\forall P \in |X| : X \setminus \{P\}$ is affine

3. \exists finite morphism $X \rightarrow \mathbb{P}_k^1$ of degree $\leq g + 1$

Prop. Any locally free coherent sheaf on $X = \mathbb{P}_k^1$ is isomorphic to $\bigoplus_{i=1}^r \mathcal{O}_X(n_i)$ for unique $n_1 \geq \dots \geq n_r$

7.6 Embeddings in projective space

Let \mathcal{L} be an invertible sheaf on X .

Prop. 1. \mathcal{L} is generated by global sections iff $\forall P \in |X| : h^0(\mathcal{L}(-P)) = h^0(\mathcal{L}) - 1$

2. \mathcal{L} is very ample iff $\forall P, Q \in |X| : h^0(\mathcal{L}(-P - Q)) = h^0(\mathcal{L}) - 2$

Fact. $\forall \mathcal{L} :$

$$\deg(\mathcal{L}) < 0 \Rightarrow h^0(\mathcal{L}) = 0$$

$$\deg(\mathcal{L}) > 2g - 2 \Rightarrow h^1(\mathcal{L}) = h^0(\mathcal{L}^\vee \otimes \Omega_X) = 0$$

Prop. $\forall \mathcal{L} :$

1. $\deg(\mathcal{L}) \geq 2g \Rightarrow \mathcal{L}$ is generated by global sections.

2. $\deg(\mathcal{L}) \geq 2g + 1 \Rightarrow \mathcal{L}$ is very ample.

Corollary. \mathcal{L} is ample iff $\deg(\mathcal{L}) > 0$

Prop. $\forall g \geq 1 : \Omega_X$ is generated by global sections.

Corollary. $g \geq 1 \Rightarrow$ get the canonical morphism $X \rightarrow \mathbb{P}_k^{g-1}$ (unique up to $PGL_g(k)$)
If $g \geq 2$ this is non constant and hence finite

Prop. X is hyperelliptic iff $\exists \mathcal{L}$ with $h^0(\mathcal{L}) = \deg(\mathcal{L}) = 2$

Prop. For $g \geq 2$ the canonical morphism $X \rightarrow \mathbb{P}_k^{g-1}$ is a closed embedding iff X is not hyperelliptic

7.7 Hyperelliptic curves

Prop. $\Omega_{X/k} = f^* \mathcal{O}_{\mathbb{P}_k^1}((g-1)\infty) \cdot \omega$ and

$$H^0(X, \Omega_{X/k}) = H^0(\mathbb{P}_k^1, \mathcal{O}_{\mathbb{P}_k^1}((g-1)\infty)) \cdot \omega \text{ for } \omega := \frac{dx}{2y-a(x)} \in \Omega_{K/k} \text{ for } K = k(X)$$

Corollary. The canonical morphism factors as $(g-1)$ -uple embedding if $g \geq 2$

$$\begin{array}{ccc} X & \longrightarrow & \mathbb{P}_k^{g-1} \\ \downarrow f & \nearrow & \\ \mathbb{P}_k^1 & & \end{array}$$

Prop. $\text{char } k \neq 2$:

1. for any separated $c \in k[X]$ of degree $\{2g + 1, 2g + 2\}$ the equation $y^2 = c(x)$ defines a hyperelliptic curve of genus g .
2. in particular there exists a hyperelliptic curve of any genus ≥ 1

7.8 Coverings

Let $f : X \rightarrow Y$ be a finite separated morphism of smooth irreducible projective curves over k algebraically closed

Prop. $\forall Q \in |Y| : \sum_{P \in f^{-1}(Q)} e_P = \deg(f)$

Prop. exact sequence

$$0 \rightarrow f^* \Omega_{Y/k} \rightarrow \Omega_{X/k} \rightarrow \Omega_{X/Y} \rightarrow 0$$

Prop. $\forall P \in |X| :$

$$\text{length}(\Omega_{X/Y,P}) = \begin{cases} e_P - 1 & \text{if } \text{char}(k) \nmid e_P \\ \geq e_P & \text{if } \text{char}(k) | e_P \end{cases}$$

Theorem (Hurwitz).

$$2g(X) - 2 = \deg(f) \cdot (2g(Y) - 2) + \deg(\text{Ram}_f)$$

Prop. $\text{Aut}_k(X)$ is finite.