
TrackDX — Physician Summary

Condition: Epilepsy · Generated Nov 7, 2026 · Data since Oct 10, 2022

LAST 90 DAYS VS. PREVIOUS 90 DAYS

	Episodes	Avg severity	Avg duration	Time in episode
Last 90 days	2	5.5	2.5 h	5 h
Previous 90 days	2	4.0	1.5 h	3 h
Change	**0%**	**+38%**	**+67%**	**+67%**

EPISODES PER 90-DAY PERIOD



Preventatives during each period: 1 = Melatonin + Vimpat · 2 = Vimpat

PREVENTATIVE MEDICATIONS — ODDS OF AN EPISODE WHILE ON (ADJUSTED FOR TRIGGERS AND TIME)

Vimpat: ×0.5 odds of an episode while on it (662 days on)

Melatonin: ×0.7 odds of an episode while on it (796 days on)

Lamotrigine: ×1.3 odds of an episode while on it (432 days on)

Vitamin B6: ×1.6 odds of an episode while on it (284 days on)

Ridge-penalized logistic regression across 1378 days, adjusting for 5 triggers and for changes in episode frequency over time; odds < 1 = fewer episodes while on the medication. First 28 days of each medication period excluded (ramp-up) once it has run longer than that. Observational data — association, not causation.

RELIEF MEDICATION EFFICACY (ALL DATA)

Rescue midazolam: 79% (81 uses; average of the patient's 0–10 relief ratings)

Clobazam: 46% (28 uses; average of the patient's 0–10 relief ratings)

** Limited data — low confidence. Estimates marked with an asterisk are based on few recorded days or uses and should be interpreted cautiously.*

TRIGGERS MOST ASSOCIATED WITH EPISODES

Missed medication: ×3.6 odds of an episode on days it occurred (294 days)

Sleep deprivation: ×2.9 odds of an episode on days it occurred (279 days)

Hormonal changes: ×1.2 odds of an episode on days it occurred (272 days)

Illness: ×0.9 odds of an episode on days it occurred (311 days)

Flashing lights: ×0.7 odds of an episode on days it occurred (268 days)

Ridge-penalized logistic regression over 1490 days (109 episode days); each trigger adjusted for the others and for changes over time.

PREVENTATIVE MEDICATIONS – SEVERITY OF AN EPISODE WHILE ON (ADJUSTED FOR TRIGGERS AND TIME)

Vimpat: -0.1 points on the 1–10 scale while on it [2% decrease] (21 episodes)

Lamotrigine: -0.1 points on the 1–10 scale while on it [2% decrease] (40 episodes)

Melatonin: +0.2 points on the 1–10 scale while on it [3% increase] (77 episodes)

Vitamin B6: +0.3 points on the 1–10 scale while on it [4% increase] (48 episodes)

Linear regression on 109 episodes, adjusted for triggers and changes over time; negative = milder episodes while on the medication. Association, not causation.

TRIGGERS – SEVERITY OF AN EPISODE (ADJUSTED FOR PREVENTATIVES)

Missed medication: +1.1 points on the 1–10 scale when present [17% increase] (49 episodes)

Sleep deprivation: +0.8 points on the 1–10 scale when present [12% increase] (43 episodes)

Illness: +0.5 points on the 1–10 scale when present [7% increase] (21 episodes)

Flashing lights: +0.2 points on the 1–10 scale when present [2% increase] (15 episodes)

Hormonal changes: -0.1 points on the 1–10 scale when present [2% decrease] (22 episodes)

Linear regression on 109 episodes, adjusted for preventatives and changes over time; positive = worse episodes when the trigger is present. Association, not causation.

WHEN EPISODES START

12am–5:59am: 7% · 6am–11:59am: 36% · 12pm–5:59pm: 34% · 6pm–11:59pm: 23%

Episodes without a start time are left out.

TrackDX is a personal tracking tool, not a medical device. The insights are statistics based on what you log — they are not medical advice, a diagnosis, or a treatment recommendation. Always consult a qualified healthcare professional about your condition.