

Supplementary Material

Decodable but Not Accessible: Auditing Distance-Based Reliability Estimation on Disentangled Skin-Lesion Representations

Supplementary Material

S1. Full geometry metrics across the ladder

Extends main-text Figure 2, which shows only condition number and the decoupled Fisher-ratio scalar. Figure S1 shows all five pre-specified geometry metrics (condition number, Fisher ratio—Hotelling–Lawley and decoupled-scalar forms—and Mardia’s multivariate kurtosis, b and z statistics), one gray point per checkpoint ($n = 13$), mean $\pm$ SD in blue, across the three rungs of the disentanglement ladder. Table S1 reports the corresponding exact-permutation Kendall’s τ and Jonckheere–Terpstra statistics for all five metrics.




Figure S1: All five geometry metrics across the disentanglement ladder.

Table S1: Geometry vs. rung order: full statistics for all five metrics (exact-permutation tests, Section 2.7).

Metric	τ (full, $n = 13$)	p (exact)	τ (common-seed, $n = 9$)	p (exact)	J	p_{JT} (exact)
Condition number	0.840	2.8×10^{-5}	0.866	0.0012	55.0	1.4×10^{-5}
Fisher ratio (HL)	0.168	0.521	-0.096	0.831	33.0	0.261
Fisher ratio (scalar)	0.076	0.798	-0.289	0.388	30.0	0.399
Mardia’s kurtosis (b)	-0.137	0.608	-0.160	0.668	23.0	0.739
Mardia’s kurtosis (z)	-0.168	0.521	-0.160	0.668	22.0	0.780

S2. Full geometry-vs-AUROC statistics

Extends main-text Figure 3, which shows only the condition-number panel. Figure S2 shows all five geometry metrics plotted against Mahalanobis AUROC, one point per checkpoint, colored by rung. Table S2 reports the corresponding statistics, including a 10^5 -draw Monte Carlo cross-check of the exact Kendall’s τ p -values.

S3. Baseline reference: implementation detail

The **baseline_soft** checkpoint (main text, Section 3.6) is reported as a descriptive reference, not a fourth ladder point. Table S3 gives its full distance-scorer and probe AUROC (single checkpoint, seed 42). Its geom-

Figure 2 -- Geometry vs. Mahalanobis AUROC, primary disentanglement ladder
(baseline_soft categorical reference not shown -- see experiment_contract.md E2b)




Figure S2: All five geometry metrics vs. Mahalanobis AUROC.

Table S2: Geometry vs. Mahalanobis AUROC: full statistics for all five metrics ($n = 13$, exact-permutation Kendall’s τ ; Monte Carlo cross-check at 10^5 draws in parentheses).

Metric	τ	p (exact / MC)
Condition number	−0.128	0.590 / 0.589
Fisher ratio (HL)	−0.282	0.204 / 0.202
Fisher ratio (scalar)	−0.154	0.510 / 0.509
Mardia’s kurtosis (b)	−0.128	0.590 / 0.590
Mardia’s kurtosis (z)	−0.128	0.590 / 0.592

AUROC itself vs. rung order (Jonckheere–Terpstra): $J = 25.0$ (null mean 27.5), $p_{\text{exact}} = 0.650$.

etry metrics (condition number, Fisher ratio, Mardia’s kurtosis) were not computed: because **baseline_soft** was never load-bearing for any statistical claim in this study, computing its geometry was out of scope.

Table S3: **baseline_soft**: distance-scorer and domain-probe AUROC ($n = 1$ checkpoint).

Scorer / probe	AUROC
Mahalanobis	0.863
Cosine-to-centroid	0.715
k -NN ($k = 1$)	0.865
k -NN ($k = 10$)	0.853
k -NN ($k = 50$)	0.844
Distance-scorer mean	0.828
Probe: logistic regression	0.998
Probe: linear SVM	0.997
Probe: random forest	0.977

S4. ID/OOD Mahalanobis distance distributions

Figure S4 shows the full histogram, kernel density estimate, and empirical CDF of the ID (ISIC-test) and OOD (PAD-UFES) Mahalanobis squared-distance distributions, pooled across seeds within each rung, supporting the median-gap numbers reported in main-text Section 3.3. The distributions are visibly overlapping, not cleanly separated.

S5. Full nearest-centroid confusion matrix

Figure S5 shows the complete 8×8 nearest-centroid confusion matrix for ISIC-test samples (true label vs. predicted centroid), supporting the majority-class-only summary in main-text Figure 4B.

ID vs OOD Mahalanobis squared distance distributions, pooled across seeds per rung
(baseline_soft categorical reference not shown, same scope as E1a/E2a)




Figure S4: ID vs. OOD Mahalanobis squared-distance distributions, all three rungs.




Figure S5: Full ISIC-test nearest-centroid confusion matrix, all 8 classes.

S6. Full predicted-class distribution for PAD-UFES

Figure S6 shows the complete nearest-centroid predicted-class distribution for PAD-UFES samples across all 8 ISIC classes, extending main-text Figure 4B’s majority-class (Nevus)-only summary.




Figure S6: Full PAD-UFES nearest-centroid predicted-class distribution, all 8 classes.

S7. Full statistical test results: scorers and probes vs. rung order

Table S7 reports exact-permutation Kendall’s τ (and, for the five distance-based scorers, Jonckheere–Terpstra) for AUROC vs. rung order, for every scorer variant (main-text Section 3.4) and every probe (main-text Section 3.5) individually. No metric or scorer reaches significance at $n = 13$; none was omitted from this table.

Table S7: AUROC vs. rung order, per scorer and per probe ($n = 13$, exact-permutation tests).

	Scorer / probe	τ	p (Kendall, exact)	p_{JT} (exact)
Distance-based	Mahalanobis	−0.076	0.798	0.650
	Cosine-to-centroid	0.321	0.194	0.097
	k -NN ($k = 1$)	−0.046	0.898	0.601
	k -NN ($k = 10$)	0.015	1.000	0.500
	k -NN ($k = 50$)	0.076	0.798	0.399
Non-distance-based	Energy	0.443	0.065	0.032
	ViM	0.076	0.798	0.399
	Density (KDE)	0.015	1.000	0.500
Domain probes	Logistic regression	0.260	0.302	—
	Linear SVM	0.260	0.302	—
	Random forest	−0.107	0.701	—

S8. Detectability and power of the permutation designs

Supports main-text Section 2.8 and the Limitations discussion in Section 4.4. Figure S8A shows the power of each permutation design as a function of the expected Kendall’s τ under the alternative, with the design’s critical value τ_{crit} and the pre-specified $|\tau| \geq 0.3$ threshold marked; Figure S8B shows every observed τ with its 95% bootstrap interval against the band of values no design at this sample size can resolve. Table S8 gives the design-level numbers; the per-test intervals plotted in panel B are tabulated in Section S9.

Effect-size and power bounds on this study's null associations (exact permutation tests, $\alpha = 0.05$)




Figure S8: (A) Power of each exact permutation design against expected τ (2×10^4 simulated datasets per point, common random numbers; $\alpha = 0.05$, two-sided). Dotted vertical lines mark each design's τ_{crit} ; the dashed red line marks the pre-specified $|\tau| \geq 0.3$ criterion, which falls below every τ_{crit} . (B) Observed τ and 95% BCa bootstrap intervals for all 21 reported associations. The shaded band spans $\pm \tau_{crit}$ for the 5/5/3 ladder: no $|\tau|$ inside it can reach $p \leq 0.05$ at this design. Condition number vs. λ_{orth} sits at the design ceiling and is shown as a point without an interval (see Table S9).

Table S8: Detectability of the two permutation designs used in this study. τ_{crit} is the smallest $|\tau|$ attainable at that design whose two-sided exact permutation p -value reaches 0.05: no dataset, however clean, can produce a significant association below it. Power is computed by simulation (2×10^4 datasets per point, common random numbers) under an alternative whose expected value of the reported statistic equals the stated τ — a bivariate-normal alternative for Design A, a linear dose-shift alternative $y = \theta d + \varepsilon$ for Design B. $MDE_{80\%}$ is the expected τ at which power first reaches 0.80.

Design	n	Permutation null space	Max. attainable $ \tau $	τ_{crit}	Power		$MDE_{80\%}$
					$\tau = 0.3$	$\tau = 0.5$	
A: continuous–continuous (E2: geometry vs. AUROC)	13	13! orderings	1.000	0.436	0.27	0.71	0.54
B: 5/5/3 ordered ladder (E1, E2.6, E2.7 vs. λ_{orth})	13	72,072	0.840	0.473	0.23	0.63	0.57
B: 3/3/3 common-seed subset	9	1,680	0.866	0.609	0.14	0.39	0.69

The success criterion this study pre-specified (Section 2.7, $|\tau| \geq 0.3$ and significant at $\alpha = 0.05$) lies below τ_{crit} for all three designs, and is therefore unsatisfiable at these sample sizes: an association of exactly the pre-specified magnitude could not have been declared significant regardless of the data. Only the condition-number effect ($\tau = 0.840$, the maximum attainable value at this design) clears τ_{crit} .

S9. Interval estimates for every reported association

Table S9 tabulates the point estimates and intervals plotted in Figure S8B: every Kendall’s τ reported anywhere in this study, with a 95% stratified-bootstrap BCa interval and the largest effect size that interval leaves unexcluded. No association tested is omitted.

Table S9: Every reported Kendall’s τ with a 95% bootstrap confidence interval (2×10^4 stratified resamples, BCa; resampling is within-rung so the 5/5/3 design is held fixed). “Largest $|\tau|$ not excluded” is the interval endpoint furthest from zero — the effect size each null result still permits.

	Test	τ	95% CI	Largest $ \tau $ not excluded
E1 vs. λ_{orth}	Condition number	0.840	at design ceiling — see note	
	Fisher ratio (HL)	0.168	[−0.461, 0.620]	0.62
	Fisher ratio (scalar)	0.076	[−0.532, 0.532]	0.53
	Mardia’s kurtosis (b)	−0.137	[−0.400, 0.206]	0.40
	Mardia’s kurtosis (z)	−0.168	[−0.412, 0.174]	0.41
E2 vs. AUROC	Condition number	−0.128	[−0.507, 0.200]	0.51
	Fisher ratio (HL)	−0.282	[−0.750, 0.183]	0.75
	Fisher ratio (scalar)	−0.154	[−0.611, 0.324]	0.61
	Mardia’s kurtosis (b)	−0.128	[−0.500, 0.233]	0.50
	Mardia’s kurtosis (z)	−0.128	[−0.534, 0.216]	0.53
E2.6 scorers	Mahalanobis	−0.076	[−0.524, 0.432]	0.52
	Cosine-to-centroid	0.321	[−0.210, 0.688]	0.69
	k -NN ($k = 1$)	−0.046	[−0.500, 0.556]	0.56
	k -NN ($k = 10$)	0.015	[−0.564, 0.489]	0.56
	k -NN ($k = 50$)	0.076	[−0.524, 0.545]	0.55
	Energy	0.443	[0.048, 0.747]	0.75
	ViM	0.076	[−0.268, 0.270]	0.27
	Density (KDE)	0.015	[−0.560, 0.471]	0.56
E2.7 probes	Logistic regression	0.260	[−0.210, 0.656]	0.66
	Linear SVM	0.260	[−0.210, 0.656]	0.66
	Random forest	−0.107	[−0.560, 0.368]	0.56

Note. Condition number vs. λ_{orth} attains $\tau = 0.840$, the maximum value the 5/5/3 design permits (a perfectly monotone ladder). Every bootstrap resample is therefore at or below the observed value and the resampling distribution is one-sided by construction, so no bootstrap interval is valid there; the exact permutation p -value (2.8×10^{-5}) is the meaningful inference for that effect. ViM (bold) is the only null result in this study whose interval excludes the pre-specified $|\tau| = 0.3$; the other nineteen do not.