



# Review

- 有理函数  $\frac{p(x)}{q(x)}$  ( $p, q$  为多项式) 的积分
- 三角有理式  $R(\sin x, \cos x)$     万能变换  $t = \tan \frac{x}{2}$
- 可化为有理式的简单无理式

$$1) \int R(x, \sqrt[n]{\frac{ax+b}{cx+d}}) dx \quad (ad-bc \neq 0) \quad \text{令 } t = \sqrt[n]{\frac{ax+b}{cx+d}}$$

$$2) \int R(x, \sqrt{ax^2+bx+c}) dx, (a \neq 0)$$

三角变换开根号、Euler变换



## § 6. 定积分的计算

Newton-Leibniz公式  $\int_a^b F'(x)dx = F(x)\Big|_{x=a}^b$

Thm.(定积分的换元法)  $f \in C[a, b]$ ,  $\varphi \in C^1[\alpha, \beta]$ ,  $\varphi(\alpha) = a$ ,  $\varphi(\beta) = b$ ,  $a \leq \varphi(t) \leq b$ , 则  $\int_a^b f(x)dx = \int_\alpha^\beta f(\varphi(t))\varphi'(t)dt$ .

Proof.  $f \in C[a, b]$ , 则  $f$  在  $[a, b]$  上有原函数  $F(x)$ ,

$$\frac{d}{dt} F(\varphi(t)) = f(\varphi(t))\varphi'(t), \quad \forall t \in [\alpha, \beta].$$

$$\int_\alpha^\beta f(\varphi(t))\varphi'(t)dt = F(\varphi(t))\Big|_\alpha^\beta = F(b) - F(a) = \int_a^b f(x)dx. \square$$



## Ex. 判断正误

$$\int_{-1}^1 \frac{1}{1+x^2} dx = \arctan x \Big|_{-1}^1 = \frac{\pi}{2} \quad (\checkmark)$$

$$\int_{-1}^1 \frac{1}{1+x^2} dx = -\int_{-1}^1 \frac{1}{1+\frac{1}{x^2}} d\frac{1}{x} = -\arctan \frac{1}{x} \Big|_{-1}^1 = -\frac{\pi}{2}. \quad (\times)$$

$$\left( -\arctan \frac{1}{x} \Big| \text{在 } x=0 \text{ 处不连续; } x=\frac{1}{t} \text{ 在 } t=0 \text{ 处不连续} \right)$$

**Question.** 不定积分的第一、二换元法以及定积分换元法中, 对  $x = \varphi(t)$  分别有什么要求?




Ex.  $\int_0^a \sqrt{a^2 - x^2} dx$  ( $a > 0$ )      几何意义?

解: 令  $x = a \sin t, t \in [0, \pi/2]$ ,

$$\int_0^a \sqrt{a^2 - x^2} dx = \int_0^{\pi/2} a^2 \cos^2 t dt$$

$$= \frac{1}{2} a^2 \int_0^{\pi/2} (1 + \cos 2t) dt$$

$$= \frac{1}{2} a^2 \left( t + \frac{1}{2} \sin 2t \right) \Big|_0^{\pi/2} = \frac{\pi}{4} a^2. \square$$





**Ex.**  $f \in C[-a, a]$  为偶函数, 则  $\int_{-a}^a \frac{f(x)}{1+e^x} dx = \int_0^a f(x) dx$ .

**Proof.**

$$\begin{aligned} \int_{-a}^a \frac{f(x)}{1+e^x} dx &= \int_0^a \frac{f(x)}{1+e^x} dx + \int_{-a}^0 \frac{f(x)}{1+e^x} dx \\ &= \int_0^a \frac{f(x)}{1+e^x} dx - \int_a^0 \frac{1}{1+e^{-t}} f(-t) d(t) \quad (x = -t) \\ &= \int_0^a \frac{f(x)}{1+e^x} dx + \int_0^a \frac{e^t}{1+e^t} f(t) dt \quad (f \text{ 偶}) \\ &= \int_0^a \frac{f(x)}{1+e^x} dx + \int_0^a \frac{e^x}{1+e^x} f(x) dx = \int_0^a f(x) dx. \square \end{aligned}$$



**Ex.**  $f \in C[0, a]$ ,  $f(x) + f(a-x) \neq 0$ , 则  $\int_0^a \frac{f(x)}{f(x) + f(a-x)} dx = \frac{a}{2}$ .

**Proof.**  $I = \int_0^a \frac{f(x)}{f(x) + f(a-x)} dx$

(令  $x = a - t$ )  
$$= - \int_a^0 \frac{f(a-t)}{f(a-t) + f(t)} dt = \int_0^a \frac{f(a-t)}{f(a-t) + f(t)} dt$$

(令  $t = x$ )  
$$= \int_0^a \frac{f(a-x)}{f(a-x) + f(x)} dx$$

$$2I = \int_0^a \left( \frac{f(x)}{f(x) + f(a-x)} + \frac{f(a-x)}{f(a-x) + f(x)} \right) dx = a. \square$$



**Ex.**  $f \in C[1, +\infty)$ ,  $a > 1$ , 则  $\int_1^a f(x^2 + \frac{a^2}{x^2}) \frac{dx}{x} = \int_1^a f(x + \frac{a^2}{x}) \frac{dx}{x}$ .

**Proof.**  $\int_1^a f(x^2 + \frac{a^2}{x^2}) \frac{dx}{x} = \frac{1}{2} \int_1^{a^2} f(t + \frac{a^2}{t}) \frac{dt}{t} \quad (t = x^2)$

$$= \frac{1}{2} \int_1^a f(t + \frac{a^2}{t}) \frac{dt}{t} + \frac{1}{2} \int_a^{a^2} f(t + \frac{a^2}{t}) \frac{dt}{t}$$

$$\triangleq \frac{1}{2} (I_1 + I_2).$$

$$I_2 = -\int_a^1 f(s + \frac{a^2}{s}) \frac{ds}{s} = \int_1^a f(s + \frac{a^2}{s}) \frac{ds}{s} \quad (s = \frac{a^2}{t}) \quad \square$$



**Ex.**(1)  $f \in C[a, b]$ , 则  $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$ ;

$$(2) I = \int_{\pi/6}^{\pi/3} \frac{\cos^2 x}{x(\pi-2x)} dx = \frac{1}{\pi} \ln 2.$$

**Proof.**(1)  $\int_a^b f(a+b-x)dx$

$$\underline{\underline{t = a+b-x}} - \int_b^a f(t)dt = \int_a^b f(t)dt = \int_a^b f(x)dx.$$

$$\begin{aligned} (2) \text{利用(1), } I &= \int_{\pi/6}^{\pi/3} \frac{\sin^2 x}{x(\pi-2x)} dx = \frac{1}{2} \int_{\pi/6}^{\pi/3} \frac{1}{x(\pi-2x)} dx \\ &= \frac{1}{\pi} \int_{\pi/6}^{\pi/3} \left( \frac{1}{2x} + \frac{1}{\pi-2x} \right) dx = \frac{1}{2\pi} \ln \frac{2x}{\pi-2x} \Big|_{\pi/6}^{\pi/3} = \frac{1}{\pi} \ln 2. \square \end{aligned}$$





Ex.  $I = \int_0^1 \frac{\ln(1+x)}{1+x^2} dx.$

解:  $I = \int_0^{\pi/4} \ln(1+\tan t) dt \quad (t = \arctan x)$

$$= \int_0^{\pi/4} \ln(\sin t + \cos t) dt - \int_0^{\pi/4} \ln(\cos t) dt = I_1 - I_2.$$

$$I_1 = \int_0^{\pi/4} \left( \ln \sqrt{2} + \ln \sin\left(t + \frac{\pi}{4}\right) \right) dt$$

$$= \frac{\pi}{8} \ln 2 + \int_0^{\pi/4} \ln \cos\left(\frac{\pi}{4} - t\right) dt = \frac{\pi}{8} \ln 2 + I_2.$$

$$I = \frac{\pi}{8} \ln 2. \square$$



Thm.(定积分的分部积分法)  $u, v \in C^1[a, b]$ , 则

$$\int_a^b u(x)v'(x)dx = u(x)v(x)\Big|_a^b - \int_a^b v(x)u'(x)dx.$$

Proof.  $(u(x)v(x))' = u'(x)v(x) + u(x)v'(x)$

$$\Rightarrow \int_a^b u'(x)v(x)dx + \int_a^b u(x)v'(x)dx = u(x)v(x)\Big|_a^b$$

$$\Rightarrow \int_a^b u(x)v'(x)dx = u(x)v(x)\Big|_a^b - \int_a^b v(x)u'(x)dx. \square$$



Ex. 证明  $I_n \triangleq \int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx$ , 并求  $I_n$ .

Proof. 令  $t = \frac{\pi}{2} - x$ , 则

$$\int_0^{\pi/2} \sin^n x dx = -\int_{\pi/2}^0 \sin^n \left(\frac{\pi}{2} - t\right) dt = \int_0^{\pi/2} \cos^n t dt.$$

$$\begin{aligned} I_n &= -\int_0^{\pi/2} \sin^{n-1} x d \cos x \\ &= -\sin^{n-1} x \cos x \Big|_0^{\pi/2} + (n-1) \int_0^{\pi/2} \cos^2 x \sin^{n-2} x dx \\ &= (n-1) \int_0^{\pi/2} (1 - \sin^2 x) \sin^{n-2} x dx \\ &= (n-1) I_{n-2} - (n-1) I_n. \end{aligned}$$



$$I_n = \frac{n-1}{n} I_{n-2}.$$

$$I_0 = \int_0^{\pi/2} dx = \frac{\pi}{2},$$

$$I_1 = \int_0^{\pi/2} \sin x dx = -\cos x \Big|_0^{\pi/2} = 1,$$

$$I_{2n} = \frac{2n-1}{2n} \cdot \frac{2n-3}{2n-2} \cdots \frac{1}{2} \cdot \frac{\pi}{2} = \frac{(2n-1)!!}{(2n)!!} \cdot \frac{\pi}{2},$$

$$I_{2n-1} = \frac{2n-2}{2n-1} \cdot \frac{2n-4}{2n-3} \cdots \frac{2}{3} \cdot 1 = \frac{(2n-2)!!}{(2n-1)!!}. \square$$



Ex. 计算  $\int_0^1 x^n \ln^n x dx$ ,  $n$  为正整数.

解:  $\int_0^1 x^n \ln^n x dx = \frac{1}{n+1} \int_0^1 \ln^n x dx^{n+1}$

$$= \frac{1}{n+1} x^{n+1} \ln^n x \Big|_0^1 - \frac{n}{n+1} \int_0^1 x^n \ln^{n-1} x dx = -\frac{n}{n+1} \int_0^1 x^n \ln^{n-1} x dx$$

$$= -\frac{n}{(n+1)^2} \int_0^1 \ln^{n-1} x dx^{n+1} = \frac{n(n-1)}{(n+1)^2} \int_0^1 x^n \ln^{n-2} x dx$$

$$= \cdots = (-1)^n \frac{n!}{(n+1)^n} \int_0^1 x^n dx = (-1)^n \frac{n!}{(n+1)^{n+1}}. \square$$



**Ex.**  $f, g \in C[a, b]$ ,  $\int_a^x f(x)dx \geq \int_a^x g(x)dx$  ( $a \leq x \leq b$ ),

$\int_a^b f(x)dx = \int_a^b g(x)dx$ , 则  $\int_a^b xf(x)dx \leq \int_a^b xg(x)dx$ .

**Proof.** 令  $F(x) = \int_a^x f(x)dx$ ,  $G(x) = \int_a^x g(x)dx$ , 则

$F(a) = G(a) = 0$ ,  $F(b) = G(b)$ ,  $F(x) \geq G(x)$  ( $a \leq x \leq b$ ).

$$\begin{aligned}\int_a^b x(f(x) - g(x))dx &= \int_a^b x d(F(x) - G(x)) \\ &= x(F(x) - G(x)) \Big|_a^b - \int_a^b (F(x) - G(x))dx \\ &= -\int_a^b (F(x) - G(x))dx \leq 0. \square\end{aligned}$$



Ex.  $f \in C^1[a, b]$ ,  $f(a) = 0$ , 则

$$\int_a^b f^2(x) dx \leq \frac{(b-a)^2}{2} \int_a^b (f'(x))^2 dx - \frac{1}{2} \int_a^b (x-a)^2 (f'(x))^2 dx.$$

Proof.  $f^2(x) = \left( \int_a^x 1 \cdot f'(t) dt \right)^2 \leq (x-a) \int_a^x (f'(t))^2 dt$

$$\begin{aligned} \int_a^b f^2(x) dx &\leq \int_a^b \left( \int_a^x (f'(t))^2 dt \right) d \frac{(x-a)^2}{2} \\ &= \frac{(x-a)^2}{2} \int_a^x (f'(t))^2 dt \Big|_{x=a}^b - \int_a^b \frac{(x-a)^2}{2} (f'(x))^2 dx \\ &= \frac{(b-a)^2}{2} \int_a^b (f'(x))^2 dx - \frac{1}{2} \int_a^b (x-a)^2 (f'(x))^2 dx. \square \end{aligned}$$



**Ex.**  $f \in C^1[a, b]$ ,  $f(a) = f(b) = 0$ ,  $\int_a^b f^2(x) dx = 1$ , 则

$$\int_a^b (f'(x))^2 dx \cdot \int_a^b x^2 f^2(x) dx > \frac{1}{4}.$$

**Proof.** 若  $f'(x) + \lambda x f(x) \equiv 0$ , 则

$$\left( f(x) e^{\frac{1}{2} \lambda x^2} \right)' = (f'(x) + \lambda x f(x)) e^{\frac{1}{2} \lambda x^2} \equiv 0,$$

$$f(x) e^{\frac{1}{2} \lambda x^2} \equiv C, \quad f(x) = C e^{-\frac{1}{2} \lambda x^2},$$

$f(a) = f(b) = 0$ , 则  $C = 0$ ,  $f(x) \equiv 0$ , 与  $\int_a^b f^2(x) dx = 1$  矛盾.

故  $\forall \lambda \in \mathbb{R}$ ,  $\int_a^b (f'(x) + \lambda x f(x))^2 dx > 0$ , 即






$$\int_a^b (f'(x))^2 dx + 2\lambda \int_a^b xf(x)f'(x)dx + \lambda^2 \int_a^b x^2 f^2(x)dx > 0.$$

$$\text{于是} \int_a^b (f'(x))^2 dx \cdot \int_a^b x^2 f^2(x)dx > \left( \int_a^b xf(x)f'(x)dx \right)^2.$$

$$\text{而} \int_a^b xf(x)f'(x)dx = \frac{1}{2} \int_a^b x df^2(x)$$

$$= \frac{1}{2} xf^2(x) \Big|_a^b - \frac{1}{2} \int_a^b f^2(x) dx = -\frac{1}{2}$$

$$\text{故} \int_a^b (f'(x))^2 dx \cdot \int_a^b x^2 f^2(x)dx > \frac{1}{4}. \square$$



Ex.  $\left(\frac{2n-1}{e}\right)^{\frac{2n-1}{2}} < 1 \cdot 3 \cdot 5 \cdots (2n-1) < \left(\frac{2n+1}{e}\right)^{\frac{2n+1}{2}}.$

Proof.  $S_n \triangleq 1 \cdot 3 \cdot 5 \cdots (2n-1),$

$$2 \ln S_n = 2 \sum_{k=2}^n \ln(2k-1) < \sum_{k=2}^n \int_{2k-1}^{2k+1} \ln x dx$$

$$= \int_3^{2n+1} \ln x dx = x \ln x \Big|_3^{2n+1} - \int_3^{2n+1} x \cdot \frac{1}{x} dx$$

$$= (x \ln x - x) \Big|_3^{2n+1} = x \ln \frac{x}{e} \Big|_3^{2n+1} < (2n+1) \ln \frac{2n+1}{e}.$$

同理,  $2 \ln S_n > \sum_{k=2}^n \int_{2k-3}^{2k-1} \ln x dx = \int_1^{2n-1} \ln x dx = (2n-1) \ln \frac{2n-1}{e} + 1. \square$



**Thm.**(带积分余项的Taylor公式)  $f \in C^{n+1}[a, b], x_0 \in [a, b],$

则  $\forall x \in [a, b],$  有

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{1}{n!} \int_{x_0}^x (x - t)^n f^{(n+1)}(t) dt.$$

**Proof.**  $n = 0$  时, 即Newton-Leibniz公式.

假设  $n = m - 1$  时, 定理成立, 即

$$f(x) = \sum_{k=0}^{m-1} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{1}{(m-1)!} \int_{x_0}^x (x - t)^{m-1} f^{(m)}(t) dt.$$

对余项分部积分, 得



$$\begin{aligned} \frac{1}{(m-1)!} \int_{x_0}^x (x-t)^{m-1} f^{(m)}(t) dt &= \frac{-1}{m!} \int_{x_0}^x f^{(m)}(t) d(x-t)^m \\ &= -\frac{1}{m!} f^{(m)}(t)(x-t)^m \Big|_{t=x_0}^x + \frac{1}{m!} \int_{x_0}^x (x-t)^m f^{(m+1)}(t) dt \\ &= \frac{1}{m!} f^{(m)}(x_0)(x-x_0)^m + \frac{1}{m!} \int_{x_0}^x (x-t)^m f^{(m+1)}(t) dt. \end{aligned}$$

$$\text{故 } f(x) = \sum_{k=0}^m \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k + \frac{1}{m!} \int_{x_0}^x (x-t)^m f^{(m+1)}(t) dt,$$

即  $n = m$  时, 定理成立.  $\square$



**作业：习题5.6**  
**No.1-3(单),5,8,10,11,13.**