



Review article

The human microbiome

Hubert E. Blum

Department of Medicine II, University Hospital Freiburg, Hugstetter Strasse 55, D-79106 Freiburg, Germany



ARTICLE INFO

Article history:

Received 24 March 2017

Accepted 26 April 2017

Available online 13 July 2017

Keywords:

Microbiota

Fecal transplantation

Inflammatory bowel disease

Obesity

Atherosclerosis

Neurodegenerative diseases

ABSTRACT

Until recently, human microbiology was based on the identification of single microbes, such as bacteria, fungi and viruses, frequently isolated from patients with acute or chronic infections. Novel culture-independent molecular biochemical analyses (genomics, transcriptomics, proteomics, metabolomics) allow today to detect and classify the diverse microorganisms in a given ecosystem (microbiota), such as the gastrointestinal tract, the skin, the airway system, the urogenital tract and others, and to assess all genomes in these ecosystems (microbiome) as well as their gene products. These analyses revealed that each individual has its own microbiota that plays a role in health and disease. In addition, they greatly contributed to the recent advances in the understanding of the pathogenesis of a wide range of human diseases. It is to be expected that these new insights will translate into diagnostic, therapeutic and preventive measures in the context of personalized/precision medicine.

© 2017 Medical University of Białystok. Published by Elsevier B.V. All rights reserved.

Contents

1. Introduction	414
2. Review	415
2.1. The human microbiome project	415
2.2. The intestinal microbial community	415
2.2.1. Inflammatory bowel diseases and colon cancer	416
2.2.2. Obesity	417
2.2.3. Atherosclerosis and thrombosis risk	418
2.2.4. Neurodevelopmental, psychiatric and neurodegenerative diseases	418
3. Conclusions and perspectives	419
Conflict of interests	419
Financial disclosure	419
Acknowledgement	419
References	419

1. Introduction

The basic aspects of molecular and cell biology are not only integral part of biomedical research but are also translated into patient care. Several global consortia have been launched and in part completed during the last decades. All of them continuously transform basic biomedical research and translate into medical applications and, after evaluation in randomized clinical trials,

enter clinical practice with a tremendous potential to advance the diagnosis, treatment and prevention of human diseases.

More than 15 years ago, the international human genome organization (HUGO) project established the complete sequence of the ca. 3 billion base pairs that make up the human genome [1,2]. In order to utilize these data from the HUGO project for research as well as for clinical applications and to define the functions of newly identified genes, collectively termed 'functional genomics', strategies were developed to globally analyze genomic DNA sequences as well as their cell-, tissue- or organ-specific expression profile. Using chips, so-called 'microarrays',

E-mail address: hubert.blum@uniklinik-freiburg.de (H.E. Blum).

thousands or ten thousands of single-stranded DNA species, reverse transcribed RNA (cDNA) or oligonucleotides of known sequence can provide a global gene (genomics), gene expression (transcriptomics, proteomics) or metabolite (metabolomics) profile ('signature') that is characteristic for the disease of individual patients, including its natural course/prognosis and response to therapy.

In 2005 the international haplotype map (HapMap) project was initiated to identify, based on genome-wide association studies (GWAS) in ethnically different populations, single nucleotide polymorphisms (SNPs) and their association with specific human diseases and individual phenotypic characteristics, respectively [3,4]. Through GWAS an increasing number of gene loci have been identified that are associated with individual (future) phenotypic traits, such as hair or eye color, height, body mass index and others as well as with the disposition to develop a specific disease [3,4]. Further, genetic variants are associated with the individuals' response to drug treatment, e.g., to lithium [5]. Overall, GWAS allow an increasingly better understanding of disease pathogenesis and a more accurate assessment of the individual risk to develop a specific disease. Clinically, this may eventually translate into clinical advances in disease prevention, early diagnosis and therapy. It should be cautioned, however, that the contribution of a defined SNP to the risk assessment for a given disease must be weighed against established clinical parameters and needs to be carefully evaluated before entering clinical practice.

2. Review

2.1. The human microbiome project

The human microbiome project (HMP) was established in 2007 as another global consortium [6–10]. The HMP and the 'Meta-genomics of the Human Intestinal Tract (Meta-HIT) Consortium Europe' aim at the sequencing of all microbes (eukaryotes, archaea, bacteria, viruses) that inhabit specific body sites, such as the mouth, throat and airways, stomach and intestine, the urogenital system and the skin, respectively (Fig. 1A). Recent data demonstrate that specific compositions of the microbial community are associated with health and disease [6–10] and suggest that the detailed characterization, function and variation of the microbial community will reveal important commensal host-microbe as well as microbe-microbe interactions with diagnostic, therapeutic and preventive implications [11,12].

While the HMP has meanwhile developed into a major field of biomedical research, the intestinal microbial community in particular has turned out to play a major role in human health and disease pathogenesis as will be discussed in more detail below [13].

2.2. The intestinal microbial community

In recent years the intestinal microbial community has been studied in great detail. It represents a microbial ecosystem




Fig. 1. (A) Different microbiomes in humans; (B) The intestinal microbiome in healthy individuals and patients [7].

consisting of trillion microbial cells with an aggregate 9.9 million microbial genes across the fecal microbiome [14]. While until recently, the environment *in utero* has been considered sterile, DNA-based analyses identified bacterial species in maternal placenta, amniotic fluid and meconium. The colonization of the human gut begins at birth with a rapid expansion of bacterial diversity and is characterized by a successively changing composition that eventually becomes relatively stable in adulthood [15]. While the specific microbial species and subspecies and their proportions vary greatly from person to person the individual microbiome is unique and becomes more diverse in the elderly.

Important factors for the composition of the intestinal microbial community are endogenous and exogenous factors [16,17]. Examples are mode of delivery of the neonates, diet (dietary supplements, breast-feeding, formula-feeding), xenobiotics, including antibiotics and other drugs [18–21]. Further, infections and exposure to environmental microbial agents are established risk factors for childhood diseases, such as obesity and allergy [22,23]. Recent evidence further suggests that human genetic variation also influences the abundance of specific members of the intestinal microbial community [24].

Taken together, the emerging data suggest that the detailed characterization of the human intestinal microbiome composition, function and variation across different body sites will reveal important commensal host-microbe as well as microbe-, microbe interactions that may play a role in human health and disease.

In view of the numerous and diverse physiological functions of the intestinal microbiota in human health (Table 1) it is not surprising that it is also involved in gastrointestinal as well as non-gastrointestinal diseases, such as obesity/metabolic syndrome, and atherosclerosis/cardiovascular as well as neurologic/psychiatric diseases, making it one of the most dynamic current topics in biomedical research (Table 2). In the following, a few examples will be discussed in more detail.

2.2.1. Inflammatory bowel diseases and colon cancer

The inflammatory bowel diseases (IBD) in humans include ulcerative colitis (UC) and Crohn disease (CD). These are characterized by inflammation limited to the mucosal layer of the colon in UC and the transmural involvement of the gastrointestinal tract, including extraintestinal sites in CD. While the pathogenesis of IBD is not fully understood [25], it is clear that its pathology depends among others on the intestinal microbial community [26,27]. Further, a case-control study identified ‘IBD-

Table 2

Disease associations with the intestinal microbial community (examples).

	References
Allergies/Allergy protection	[23,59,64–66]
Atherosclerosis/Thrombosis/Cardiovascular Diseases	[46–48,67–70]
Cancer	[40,71,72]
Diabetes mellitus	[42,73]
Immune-Mediated Inflammatory Diseases	
–Inflammatory bowel diseases	[26,27,30,31,74,75]
–Multiple sclerosis	[76,77]
–Rheumatoid arthritis	[78]
–Psoriasis	[79]
Kwashiorkor	[43,44]
Liver Diseases	[80,81]
Metabolic Syndrome/Obesity	[41,82–85]
Neurodevelopmental, Psychiatric and Neuro-degenerative diseases	
–Autism	[49,86]
–Depression	[49,87]
–Alzheimer’s Disease, Parkinson’s Disease	[49,53,54,88,89]

specific’ alterations of the intestinal microbiota that may serve as biomarkers for the prediction of disease predisposition, activity/severity and responsiveness to therapy [28,29].

Host genes with effects on the composition of the intestinal microbiota are the IgA locus and the HLA genes as well as the defensin genes, the NOD2 gene, the resistin-like molecule beta gene, the apolipoprotein I gene, the MEFV gene and the myeloid differentiation primary response protein 88 gene. The three components – environment, host genetics and the microbial community – interact to maintain homeostasis in the intestine [7]. The disruption of the stability of this interaction may be a trigger for disease development (Fig. 1B). Two recent publications shed a new light on the pathogenesis of IBD through the change of the intestinal microbial composition involving two different pathways: helminth infection [30] and lipocalin-2 expression [31], respectively.

2.2.1.1. Helminth infection, microbial community and IBD. Epidemiologic studies demonstrated a major increase of the incidence of IBD in the developed world, suggesting a change in the environment, including an alteration of the intestinal microbiome [32] and a decreased exposure to intestinal parasites, such as helminths [33]. In mice deficient for the CD susceptibility gene Nod2 (Nod2^{−/−} knockout) [34] it could be demonstrated that small intestinal abnormalities develop in the face of a sustained colonization with the inflammatory bacterium *Bacteroides vulgatus*, an ubiquitous member of the intestina microbial community [35]. Chronic infection of Nod2^{−/−} mice with the parasitic worm *Trichuris muris*, however, inhibited colonization with inflammatory *Bacteroides* species and promoted the establishment of a protective microbial environment enriched in *Clostridiales* [30]. Further, the authors demonstrated that individuals from helminth-endemic regions harbour a similar protective microbial community and deworming treatment reduced *Clostridiales* and increased *Bacteroidales*, resulting in an increased IBD incidence. These data support the hygiene hypothesis whereby certain individuals are genetically susceptible to the consequences of a changing intestinal microbial community that favors IBD development IBD.

2.2.1.2. Lipocalin-2 protection from IBD and colon cancer. Lipocalin-2 (Lcn2) is an antimicrobial peptide with high mucosal and fecal concentrations in patients with IBD. It is produced by various cell types, including epithelial cells, and acts as an antimicrobial defense mediator by binding to a subset of bacterial siderophores, thereby preventing bacterial iron acquisition and growth of

Table 1

Functions of the intestinal microbial community in human health (examples).

	References
Host Physiology	
Adaptive immunity	[58]
Autoimmunity	[59]
Innate immunity	[60]
Cell proliferation	[61]
Bone density	[20]
Vascularization	[62]
Neurological signalling	[63]
Biosynthesis	
Neurotransmitters	
Steroid hormones	
Vitamins	
Metabolism	
Dietary components	
Bile salts	
Drugs	
Xenobiotics	

siderophore-dependent strains. While it has been implicated in several biologic processes, such as acute phase response, erythropoiesis and iron metabolism, its functional role in contributing to IBD development remained unclear.

To decipher the role of Lcn2 in colon inflammation, mice double deficient in Lcn2 and IL-10 (Lcn2^{-/-}/IL10^{-/-} double knockout) were generated and compared to single knockouts and wild-type animals. The experimental data indicate that Lcn2 expression protects from early onset colitis and the spontaneous emergence of right-sided colonic tumors that result from IL-10 deficiency. The inflammation is driven by IL-6 which also controls tumorigenesis. The Lcn2^{-/-}/IL10^{-/-} double knockout mice showed major alterations of their intestinal microbial community, especially with respect to the facultative pathogenic *Alistipes* spp. These contribute to inflammation and tumorigenesis as shown by the transmissibility of the phenotype and the protection by antibiotic therapy. Taken together, the authors demonstrate that Lcn2 protects against intestinal inflammation and tumorigenesis in the face of an altered intestinal microbial composition [31].

While donor feces have previously been shown to be effective for the treatment of *Clostridium difficile* infection [36,37] recent randomized controlled trials demonstrated that fecal microbiota transplantation can also induce clinical remission and endoscopic improvement in patients with active UC not responding to conventional medical therapy and is associated with distinct microbial changes that relate to outcome [12,38,39].

Recently, it was discovered that the probiotic *Lactobacillus casei* strain ATCC 334 produces ferrichrome that inhibits colon cancer progression by apoptosis mediated through the c-Jun N-terminal kinase pathway [40], possibly representing a novel tumor-suppressing strategy.

2.2.2. Obesity

Obesity [41], insulin resistance [42] and khwashiorakor [43,44] are examples for which a correlation between microbial dysbiosis and the clinical state has been demonstrated. Further, in genetically susceptible hosts, the transplantation of fecal microbiota from healthy donors to patients resulted in clinical improvement [45]. The underlying concept is a 'common ground




Fig. 2. The intestinal microbial metabolite trimethylamine (TMA) and the atherosclerosis/thrombosis risk [47].

hypothesis' that involves a leaky mucosa caused by various endogenous or exogenous factors, the expansion of opportunistic microbes (dysbiotic pathobionts) with the generation of pathogenic microbial gene products which can be transferred to genetically susceptible individuals [13].

2.2.3. Atherosclerosis and thrombosis risk

Recent studies suggest that intestinal microbes are involved in atherosclerosis. In this context, foods rich in choline, phosphatidylcholine and carnitine such as meat, egg yolk and high-fat dairy products, serve as precursors of trimethylamine (TMA) and TMA N-oxide (TMAO) that accelerates atherosclerosis [46] and elevated TMAO blood levels are associated with an increased risk for atherosclerotic heart disease and incident major adverse cardiac events. Further, TMAO enhances platelet hyperreactivity and thrombotic events in studies of animal model, employing dietary choline or TMAO, germ-free mice and microbial transplantation, collectively confirming the key role of intestinal microbiota [47]. Taken together, these results reveal a previously unknown link between specific dietary nutrients, intestinal microbes and thrombosis risk (Fig. 2).

The intestinal TMAO formation is a two-step process involving the generation of TMA by intestinal microbes after food ingestion and the hepatic conversion of TMA to TMAO by host flavin monooxygenases. In a recent, study Wang et al. [48] demonstrated that 3,3-dimethyl-1-butanol (DMB), a structural analog of choline, blocks the intestinal TMA formation through the inhibition of the microbial TMA lyase, and results in reduced TMAO levels. Thus, 'drugging the microbiome' with DMB may be a novel approach for the prevention/treatment of atherosclerosis (Fig. 3).

2.2.4. Neurodevelopmental, psychiatric and neurodegenerative diseases

Studies investigating the intestinal microbial-brain communication (gut-brain axis) demonstrate a critical role of the intestinal microbial community in modulating the maturation and function of tissue-resident immune cells in the central nervous system (CNS) as well as the activation of peripheral immune cells involved in neuroinflammation, brain injury, autoimmunity and neurogenesis [49]. Germ-free mice raised under sterile conditions or mice depleted of their intestinal microbiota by antibiotics show major alterations in behaviours or neuropathologies that characterize neurodevelopmental, psychiatric and neurodegenerative disorders [50]. These include among others autism spectrum disorders, depression and Alzheimer's or Parkinson's disease (Table 2).

An impressive example for a pathogenic role of the intestinal microbial community is Parkinson's disease (PD). In patients with PD, plaques in brain cells as well as in the intestine containing the neurotoxic protein alpha-synuclein (AS) are a hallmark of the disease. For example, in PD patients gastric motility is frequently impaired [51] and AS levels in the intestine are elevated [52].

In an animal model, mice overexpressing AS indeed develop neurologic deficits resembling those of patients with PD. Recently, three lines of evidence demonstrated a central role of the intestinal microbial community in the pathogenesis of PD: (1) in the PD mouse model germ-free animals developed fewer plaques and almost no neurological deficits as compared to conventionally colonized controls, (2) treatment of PD mice with antibiotics resulted in an improvement of the neurological deficits and (3) fecal transplantation from patients with PD to germ-free mice resulted in neurological deficits resembling PD [53].




Fig. 3. Inhibition of intestinal microbial trimethylamine (TMA) synthesis by 3,3-dimethyl-1-butanol (DMB) and attenuation of atherosclerosis [48].

The underlying concept is the central contribution of the microbial community to a defect of the microglia *via* short-chain fatty acids (SCFAs) that represent bacterial fermentation products [54]. While germ-free mice showed a reduced microglia mice, SCFAs modulate microglia and enhance PD pathophysiology [53]. The identification of the disease-causing bacteria and the mechanism leading to the deposition of the neurotoxic AS plaques await further clarification.

3. Conclusions and perspectives

Basic biomedical research has made major advances in recent years and holds the promise to increasingly provide individual diagnostic, preventive as well as therapeutic options for patients with inherited or acquired, malignant or non-malignant diseases. Apart from an increasing number of host genetic susceptibility loci and environmental factors, the individual microbial community is central for the barrier between microbes and hosts. In particular, the intestinal microbial community is involved in a large number of normal biological functions in health (Table 1) as well as in numerous common, gastrointestinal and non-gastrointestinal diseases, such as obesity/metabolic syndrome, atherosclerosis/thrombosis or neurodevelopmental, psychiatric and neurodegenerative diseases (Table 2). Dietary interventions targeting intestinal microbiota, such as calory restricted diets rich in fiber and vegetables as well as fecal microbial transplantation are examples for health benefits in humans. In recent years, the intestinal microbiome thus has become one of the most dynamic areas of biomedical research that holds an enormous potential for interventions regarding human health and disease.

In addition to the intestinal microbiome, other sites of the human microbiome are of increasing interest. In particular, the skin microbiome has been shown to be highly personalized with a high degree of interindividual variability. Thus, the HMP yielded not only important contributions to the fields of microbiology and medicine, but also unexpected and novel applications to other fields and disciplines, such as forensic medicine [55–57].

Conflict of interests

The author declares no conflict of interest.

Financial disclosure

The author has no financing to disclose.

Acknowledgement

The excellent secretarial assistance of Mrs. K. Bigot and Mrs. M. Gutsell is gratefully acknowledged.

References

- [1] Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. *Nature* 2001;409:860–921.
- [2] Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, et al. The sequence of the human genome. *Science* 2001;291:1304–51.
- [3] Manolio TA, Brooks LD, Collins FS. A HapMap harvest of insights into the genetics of common disease. *J Clin Invest* 2008;118:1590–605.
- [4] Manolio TA, Collins FS. The HapMap and genome-wide association studies in diagnosis and therapy. *Annu Rev Med* 2009;60:443–56.
- [5] Hou L, Heilbronner U, Degenhardt F, Adli M, Akiyama K, Akula N, et al. Genetic variants associated with response to lithium treatment in bipolar disorder: a genome-wide association study. *Lancet* 2016;387(10023):1085–93.
- [6] Proctor LM. The human microbiome project in 2011 and beyond. *Cell Host Microbe* 2011;10:287–91.
- [7] Spor A, Koren O, Ley R. Unravelling the effects of the environment and host genotype on the gut microbiome. *Nat Rev Microbiol* 2011;9:279–90.
- [8] Human Microbiome Project C. Structure, function and diversity of the healthy human microbiome. *Nature* 2012;486:207–14.
- [9] Human Microbiome Project C. A framework for human microbiome research. *Nature* 2012;486:215–21.
- [10] Gevers D, Knight R, Petrosino JF, Huang K, McGuire AL, Birren BW, et al. The Human Microbiome Project: a community resource for the healthy human microbiome. *PLoS Biol* 2012;10:e1001377.
- [11] Morgan XC, Segata N, Huttenhower C. Biodiversity and functional genomics in the human microbiome. *Trends Genet* 2013;29:51–8.
- [12] Rossen NG, Fuentes S, van der Spek MJ, Tijssen JG, Hartman JH, et al. Findings from a randomized controlled trial of fecal transplantation for patients with ulcerative colitis. *Gastroenterology* 2015;149:110–8.
- [13] Lynch SV, Pedersen O. The human intestinal microbiome in health and disease. *N Engl J Med* 2016;375:2369–79.
- [14] Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S, et al. An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* 2014;32:834–41.
- [15] Yatsunenoto T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al. Human gut microbiome viewed across age and geography. *Nature* 2012;486:222–7.
- [16] Zhernakova A, Kurilshikov A, Bonder MJ, Tigchelaar EF, Schirmer M, Vatanen T, et al. Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. *Science* 2016;352:565–9.
- [17] Falony G, Joossens M, Vieira-Silva S, Wang J, Darzi Y, Faust K, et al. Population-level analysis of gut microbiome variation. *Science* 2016;352:560–4.
- [18] De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, et al. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci U S A* 2010;107:14691–6.
- [19] Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med* 2014;20:159–66.
- [20] Cho I, Yamanishi S, Cox L, Methe BA, Zavadil J, Li K, et al. Antibiotics in early life alter the murine colonic microbiome and adiposity. *Nature* 2012;488:621–6.
- [21] Maurice CF, Haiser HJ, Turnbaugh PJ. Xenobiotics shape the physiology and gene expression of the active human gut microbiome. *Cell* 2013;152:39–50.
- [22] Fujimura KE, Demoor T, Rauch M, Faruqi AA, Jang S, Johnson CC, et al. House dust exposure mediates gut microbiome Lactobacillus enrichment and airway immune defense against allergens and virus infection. *Proc Natl Acad Sci U S A* 2014;111:805–10.
- [23] Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, et al. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med* 2016;22:1187–91.
- [24] Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhan R, et al. Human genetics shape the gut microbiome. *Cell* 2014;159:789–99.
- [25] Xavier RJ, Podolsky DK. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448:427–34.
- [26] Hall LJ, Walshaw J, Watson AJ. Gut microbiome in new-onset Crohn's disease. *Gastroenterology* 2014;147:932–4.
- [27] Sha S, Xu B, Wang X, Zhang Y, Wang H, Kong X, et al. The biodiversity and composition of the dominant fecal microbiota in patients with inflammatory bowel disease. *Diagn Microbiol Infect Dis* 2013;75:245–51.
- [28] Frank DN, St Amand AL, Feldman RA, Boedeker EC, Harpaz N, Pace NR. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci U S A* 2007;104:13780–5.
- [29] Peterson DA, Frank DN, Pace NR, Gordon JI. Metagenomic approaches for defining the pathogenesis of inflammatory bowel diseases. *Cell Host Microbe* 2008;3:417–27.
- [30] Ramanan D, Bowcutt R, Lee SC, Tang MS, Kurtz ZD, Ding Y, et al. Helminth infection promotes colonization resistance via type 2 immunity. *Science* 2016;352:608–12.
- [31] Moschen AR, Gerner RR, Wang J, Klepsch V, Adolph TE, Reider SJ, et al. Lipocalin 2 protects from inflammation and tumorigenesis associated with gut microbiota alterations. *Cell Host Microbe* 2016;19:455–69.
- [32] Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell* 2014;157:121–41.
- [33] Weinstock JV, Elliott DE. Helminths and the IBD hygiene hypothesis. *Inflamm Bowel Dis* 2009;15:128–33.
- [34] Cleyne I, Boucher G, Jostins L, Schumm LP, Zeissig S, Ahmad T, et al. Inherited determinants of Crohn's disease and ulcerative colitis phenotypes: a genetic association study. *Lancet* 2016;387:156–67.
- [35] Ramanan D, Tang MS, Bowcutt R, Loke P, Cadwell K. Bacterial sensor Nod2 prevents inflammation of the small intestine by restricting the expansion of the commensal *Bacteroides vulgatus*. *Immunity* 2014;41:311–24.
- [36] van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, et al. Duodenal infusion of donor feces for recurrent *Clostridium difficile*. *N Engl J Med* 2013;368:407–15.
- [37] Youngster I, Russell GH, Pindar C, Ziv-Baran T, Sauk J, et al. Oral, capsulized, frozen fecal microbiota transplantation for relapsing *Clostridium difficile* infection. *JAMA* 2014;312:1772–8.
- [38] Moayyedi P, Suree MG, Kim PT, Libertucci J, Wolfe M, et al. Fecal microbiota transplantation induces remission in patients with active ulcerative colitis in a randomized controlled trial. *Gastroenterology* 2015;149:102–9.

- [39] Paramsothy S, Kamm MA, Kaakoush NO, Walsh AJ, van den Bogaerde J, et al. Multi-donor intensive faecal microbiota transplantation for active ulcerative colitis: a randomised placebo-controlled trial. *Lancet* 2017;389:1218–28.
- [40] Konishi H, Fujiya M, Tanaka H, Ueno N, Moriichi K, Sasajima J, et al. Probiotic-derived ferrichrome inhibits colon cancer progression via JNK-mediated apoptosis. *Nat Commun* 2016;7:12365.
- [41] Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, et al. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science* 2013;341:1241214.
- [42] Pedersen HK, Gudmundsdottir V, Nielsen HB, Hyötyläinen T, Nielsen T, Jensen BA, et al. Human gut microbes impact host serum metabolome and insulin sensitivity. *Nature* 2016;535:376–81.
- [43] Smith MI, Yatsunenko T, Manary MJ, Trehan I, Mkakosya R, Cheng J, et al. Gut microbiomes of Malawian twin pairs discordant for kwashiorkor. *Science* 2013;339:548–54.
- [44] Subramanian S, Huq S, Yatsunenko T, Haque R, Mahfuz M, Alam MA, et al. Persistent gut microbiota immaturity in malnourished Bangladeshi children. *Nature* 2014;510:417–21.
- [45] Vrieze A, Van Nood E, Holleman F, Salojarvi J, Kootte RS, Bartelsman JF, et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 2012;143: e913–6 e7.
- [46] Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med* 2013;19:576–85.
- [47] Zhu W, Gregory JC, Org E, Buffa JA, Gupta N, Wang Z, et al. Gut microbial metabolite TMAO enhances platelet hyperactivity and thrombosis risk. *Cell* 2016;165:111–24.
- [48] Wang Z, Roberts AB, Buffa JA, Levison BS, Zhu W, Org E, et al. Non-lethal inhibition of gut microbial trimethylamine production for the treatment of atherosclerosis. *Cell* 2015;163:1585–95.
- [49] Fung TC, Olson CA, Hsiao EY. Interactions between the microbiota, immune and nervous systems in health and disease. *Nat Neurosci* 2017;20:145–55.
- [50] Sampson TR, Mazmanian SK. Control of brain development, function, and behavior by the microbiome. *Cell Host Microbe*. 2015;17:565–76.
- [51] Fasano A, Visanji NP, Liu LW, Lang AE, Pfeiffer RF. Gastrointestinal dysfunction in Parkinson's disease. *Lancet Neurol* 2015;14:625–39.
- [52] Shannon KM, Keshavarzian A, Mutlu E, Dodiya HB, Daian D, Jaglin JA, et al. Alpha-synuclein in colonic submucosa in early untreated Parkinson's disease. *Mov Disord* 2012;27:709–15.
- [53] Sampson TR, Debelius JW, Thron T, Janssen S, Shastri GG, Ilhan ZE, et al. Gut microbiota regulate motor deficits and neuroinflammation in a model of Parkinson's disease. *Cell* 2016;167: 1469–80 e12.
- [54] Erny D, Hrabé de Angelis AL, Jaitin D, Wieghofer P, Staszewski O, David E, et al. Host microbiota constantly control maturation and function of microglia in the CNS. *Nat Neurosci* 2015;18:965–77.
- [55] Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, Knight R. Bacterial community variation in human body habitats across space and time. *Science* 2010;326:1694–7.
- [56] Edmonds-Wilson SL, Nurinova NI, Zapka CA, Fierer N, Wilson M. Review of human hand microbiome research. *J Dermatol Sci* 2015;80:3–12.
- [57] Fierer N, Lauber CL, Zhou N, McDonald D, Costello EK, Knight R. Forensic identification using skin bacterial communities. *Proc Natl Acad Sci USA* 2010;107:6477–81.
- [58] Honda K, Littman DR. The microbiota in adaptive immune homeostasis and disease. *Nature* 2016;535:75–84.
- [59] Vatanen T, Kostic AD, d'Hennezel E, Siljander H, Franzosa EA, Yassour M, et al. Variation in microbiome LPS immunogenicity contributes to autoimmunity in humans. *Cell* 2016;165:842–53.
- [60] Thaïs CA, Zmora N, Levy M, Elinav E. The microbiome and innate immunity. *Nature* 2016;535:65–74.
- [61] Ijssennagger N, Belzer C, Hoiveld GJ, Dekker J, van Mil SW, Muller M, et al. Gut microbiota facilitates dietary heme-induced epithelial hyperproliferation by opening the mucus barrier in colon. *Proc Natl Acad Sci U S A* 2015;112:10038–43.
- [62] Reinhardt C, Bergentall M, Greiner TU, Schaffner F, Ostergren-Lunden G, Petersen LC, et al. Tissue factor and PAR1 promote microbiota-induced intestinal vascular remodelling. *Nature* 2012;483:627–31.
- [63] Yano JM, Yu K, Donaldson GP, Shastri GG, Ann P, Ma L, et al. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell* 2015;161:264–76.
- [64] Ege MJ, Mayer M, Normand AC, Genuneit J, Cookson WO, Braun-Fahrlander C, et al. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med*. 2011;364:701–9.
- [65] Lynch SV, Wood RA, Boushey H, Bacharier LB, Bloomberg GR, Kattan M, et al. Effects of early-life exposure to allergens and bacteria on recurrent wheeze and atopy in urban children. *J Allergy Clin Immunol* 2014;134: 593–601 e12.
- [66] Stein MM, Hrusch CL, Gozdz J, Igartua C, Pivniouk V, Murray SE, et al. Innate immunity and asthma risk in amish and hutterite farm children. *N Engl J Med*. 2016;375:411–21.
- [67] Jonsson AL, Backhed F. Drug the Bug!. *Cell* 2015;163:1565–6.
- [68] Tilg H. A gut feeling about thrombosis. *N Engl J Med* 2016;374:2494–6.
- [69] Wang Z, Klipfelf E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* 2011;472:57–63.
- [70] Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med* 2013;368:1575–84.
- [71] Feng Q, Liang S, Jia H, Stadlmayr A, Tang L, Lan Z, et al. Gut microbiome development along the colorectal adenoma-carcinoma sequence. *Nat Commun* 2015;6:6528.
- [72] Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, et al. Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 2015;350:1084–9.
- [73] Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 2012;490:55–60.
- [74] Forbes JD, Van Domselaar G, Bernstein CN. Microbiome survey of the inflamed and noninflamed gut at different compartments within the gastrointestinal tract of inflammatory bowel disease patients. *Inflamm Bowel Dis* 2016;22:817–25.
- [75] Forbes JD, Van Domselaar G, Bernstein CN. The gut microbiota in immune-mediated inflammatory diseases. *Front Microbiol* 2016;7:1081.
- [76] Cantarel BL, Waubant E, Chehoud C, Kuczynski J, DeSantis TZ, Warrington J, et al. Gut microbiota in multiple sclerosis: possible influence of immunomodulators. *J Invest Med* 2015;63:729–34.
- [77] Jangi S, Gandhi R, Cox LM, Li N, von Glehn F, Yan R, et al. Alterations of the human gut microbiome in multiple sclerosis. *Nat Commun* 2016;7:12015.
- [78] Chen J, Wright K, Davis JM, Jeraldo P, Marietta EV, Murray J, et al. An expansion of rare lineage intestinal microbes characterizes rheumatoid arthritis. *Genome Med*. 2016;8:43.
- [79] Scher JU, Ubeda C, Artacho A, Attur M, Isaac S, Reddy SM, et al. Decreased bacterial diversity characterizes the altered gut microbiota in patients with psoriatic arthritis, resembling dysbiosis in inflammatory bowel disease. *Arthritis Rheumatol*. 2015;67:128–39.
- [80] Haque TR, Barritt AS. Intestinal microbiota in liver disease. *Best Pract Res Clin Gastroenterol* 2016;30:133–42.
- [81] Qin N, Yang F, Li A, Prifti E, Chen Y, Shao L, et al. Alterations of the human gut microbiome in liver cirrhosis. *Nature* 2014;513:59–64.
- [82] Turnbaugh PJ, Hamady M, Yatsunenkov T, Cantarel BL, Duncan A, Ley RE, et al. A core gut microbiome in obese and lean twins. *Nature* 2009;457:480–4.
- [83] Shen J, Obin MS, Zhao L. The gut microbiota, obesity and insulin resistance. *Mol Aspects Med* 2013;34:39–58.
- [84] Mazidi M, Rezaie P, Kengne AP, Mobarhan MG, Ferns GA. Gut microbiome and metabolic syndrome. *Diabetes Metab Syndr* 2016;10:S150–7.
- [85] Sonnenburg JL, Backhed F. Diet-microbiota interactions as moderators of human metabolism. *Nature* 2016;535:56–64.
- [86] Choi GB, Yim YS, Wong H, Kim S, Kim H, Kim SV, et al. The maternal interleukin-17a pathway in mice promotes autism-like phenotypes in offspring. *Science* 2016;351:933–9.
- [87] Aizawa E, Tsuji H, Asahara T, Takahashi T, Teraishi T, Yoshida S, et al. Possible association of Bifidobacterium and Lactobacillus in the gut microbiota of patients with major depressive disorder. *J Affect Disord* 2016;202:254–7.
- [88] Bu XL, Yao XQ, Jiao SS, Zeng F, Liu YH, Xiang Y, et al. A study on the association between infectious burden and Alzheimer's disease. *Eur J Neurol* 2015;22:1519–25.
- [89] Scheperjans F, Aho V, Pereira PA, Koskinen K, Paulin L, Pekkonen E, et al. Gut microbiota are related to Parkinson's disease and clinical phenotype. *Mov Disord* 2015;30:350–8.