

CSC412 / CSC2506 Solutions to Sample Problems for Midterm

1. ~~Let $p(k)$ be a one-dimensional discrete distribution that we wish to approximate, with support on non-negative integers. One way to fit an approximating distribution $q(k)$ is to minimize the Kullback-Leibler divergence:~~

$$KL(p||q) = \sum_{k=0}^{\infty} p(k) \log \frac{p(k)}{q(k)}$$

Show that when $q(k)$ is a Poisson distribution,

$$q(k) = \frac{\lambda^k \exp(-\lambda)}{k!}$$

this KL divergence is minimized by setting λ to the mean of $p(k)$.

Solution:

$$\frac{\partial KL}{\partial \lambda} = 0 \implies \lambda = E[p(k)]$$

2. ~~Recall that the definition of an exponential family model is:~~

$$f(x|\eta) = h(x)g(\eta) \exp(\eta^\top T(x))$$

where:

η are the parameters
 $T(x)$ are the sufficient statistics
 $h(x)$ is the base measure
 $g(\eta)$ is the normalizing constant

Consider the univariate Gaussian, with mean μ and precision $\lambda = \frac{1}{\sigma^2}$:

$$p(D|\mu, \lambda) = \prod_{i=1}^N \left(\frac{\lambda}{2\pi}\right)^{\frac{1}{2}} \exp\left(-\frac{1}{2}(x_i - \mu)^2\right)$$

What are η and $T(x)$ for this distribution when represented in exponential family form?


Solution:

$$p(D|\mu, \lambda) = (2\pi)^{-N/2} [\lambda^{1/2} \exp(-\frac{\lambda}{2}\mu^2)]^N \exp[\mu\lambda \sum_i x_i - \lambda/2 \sum_i x_i^2]$$

$$\eta = \begin{bmatrix} \mu\lambda \\ -\lambda/2 \end{bmatrix}$$


$$T(x) = \begin{bmatrix} \sum_i x_i \\ \sum_i x_i^2 \end{bmatrix}$$

3. Consider the following directed graphical model:



- (a) List all variables that are independent of A given evidence on B .
 (b) Write down the factorized normalized joint distribution that this graphical model represents.


Solution: a) $\{\}$



b) $P(G)P(H)P(I|G, H)P(D|G)P(E|G)P(F|H)P(B|D)P(C|E, F)P(A|B, C)$

4. ~~Murphy 20.1~~


Correction! Older copies of the Murphy book have a typo pointing you to an incorrect figure.
Do this question but with this MRF:



Solution:


a) Largest intermediate term has size 3 (1,2,3) and (2,3,4)

b) Largest maximal clique has size 3.




c) The largest intermediate term has size 4 (2,3,4,5)

d) Largest maximal clique has size 4.



5. Consider the Factor Graph:



(a) Write down the normalized joint distribution $P(X_1, X_2, X_3, X_4, X_5)$ in terms of the potentials.

(b) Write down any conditional independence relationships given by the graph.

Solution:

a) $\frac{1}{Z} \phi_A(X_1) \phi_B(X_2) \phi_C(X_1, X_2, X_3) \phi_D(X_3, X_4) \phi_E(X_5)$

where $Z = \sum_{\mathbf{x}} \phi_A(X_1) \phi_B(X_2) \phi_C(X_1, X_2, X_3) \phi_D(X_3, X_4) \phi_E(X_5)$

b) $X_1, X_2 \perp X_4, X_5 | X_3$

$X_4 \perp X_5 | X_3$