

Lecture Note: Week 4

MATH 461: Probability Theory, Spring 2021
Daesung Kim

Lecture 10. Random Variables (Sec 4.1-2, 10)

Suppose that our experiment consists of tossing 3 fair coins. Let X denote the number of heads that appear. For instance, if the outcome is (H, H, T) , then the corresponding X is 2. That means, X is a function of outcomes in the sample space.

Definition: Random Variables

A random variable is a real-valued function defined on the sample space.

If X is 2, then possible outcomes are $(H, H, T), (H, T, H), (T, H, H)$. We use the notation $\{X = 2\} = \{(H, H, T), (H, T, H), (T, H, H)\}$. The probability that $X = 2$ is then defined by

$$\mathbb{P}(X = 2) = \mathbb{P}(\{(H, H, T), (H, T, H), (T, H, H)\}) = \frac{3}{8}.$$

Example 1. Three balls are to be randomly selected without replacement from an urn containing 20 balls numbered 1 through 20. Let X be the largest ball selected. Indicate what values it takes and with what probabilities.

Definition: Discrete Random Variables

A discrete random variable is a random variable that takes at most a countable number of possible values. For a discrete random variable, we define the probability mass function $p(a)$ by $p(a) = \mathbb{P}(X = a)$. If X takes the values x_1, x_2, \dots , then

$$\begin{aligned} p(x_i) &\geq 0, \text{ for } i = 1, 2, \dots, \\ p(x) &= 0, \text{ otherwise.} \end{aligned}$$

Example 2. Independent trials consisting of the flipping of a coin having probability p of coming up heads are continually performed until either a head occurs or a total of n flips is made. Let X be the number of times the coin is flipped. Indicate what values it takes and with what probabilities.

Example 3. Let $\lambda > 0$. The probability mass function of a random variable X is given by

$$p(k) = \begin{cases} c \frac{\lambda^k}{k!} & \text{if } k = 0, 1, 2, \dots, \\ 0, & \text{otherwise} \end{cases}$$

for some c .

- (i) Find c in terms of λ .
- (ii) Find $\mathbb{P}(X = 0)$.
- (iii) Find $\mathbb{P}(X > 2)$.

Distribution function

The (cumulative) distribution function (d.f.) is

$$F(x) = \mathbb{P}(X \leq x)$$

for $-\infty < x < \infty$. We have the following properties:

- (i) F is nondecreasing.
- (ii) $\lim_{b \rightarrow \infty} F(b) = 1$.
- (iii) $\lim_{b \rightarrow -\infty} F(b) = 0$.
- (iv) F is right continuous. (That is, for any $x \in \mathbb{R}$ and a sequence $\{x_n\}$ with $x_n \geq x$ and $\lim_n x_n \rightarrow \infty = x$, we have $F(x) = \lim_{n \rightarrow \infty} F(x_n)$.)

Remark 4. If we define the cdf by $F(x) = \mathbb{P}(X < x)$, then $F(x)$ is left continuous. The right continuity of the cdf is useful when we determine if a given function is the cdf of some random variable.

Example 5. If the distribution function of X is given by

$$F(x) = \begin{cases} 0, & x < 0, \\ \frac{1}{2}, & 0 \leq x < 1, \\ \frac{3}{5}, & 1 \leq x < 2, \\ \frac{4}{5}, & 2 \leq x < 3, \\ \frac{9}{10}, & 3 \leq x < 3.5, \\ 1, & x \geq 3.5, \end{cases}$$

calculate the probability mass function of X .

Lecture 11. Expectation (Sec 4.3, 4, 6)

Definition

If X is a discrete random variable taking values x_i with probability $p(x_i)$, its expected value (or mean or expectation) is defined as

$$\mathbb{E}[X] = \sum_i x_i p(x_i).$$

One can think that the expected value of X is a weighted average of the possible values that X takes on.

Example 6. Find $\mathbb{E}[X]$, where X is the outcome when we roll a fair die.

The expectation can be understood as a long run average of values of X in n repeated experiments. That is,

$$\begin{aligned} \mathbb{E}[X] &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n X^{(i)} \\ &= \lim_{n \rightarrow \infty} x_k \frac{\text{Number of } x_k \text{ in } n \text{ repeated experiments}}{n} \\ &= \sum_k x_k p(x_k). \end{aligned}$$

Example 7. We say that I_A is an indicator variable for the event A if

$$I_A = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{if } A^c \text{ occurs.} \end{cases}$$

Find $\mathbb{E}[I_A]$.

Example 8. If X is a discrete random variable and g is a function, then $g(X)$ is also a discrete random variable. Suppose X takes values $-1, 0$ and 1 with probabilities $0.2, 0.5$ and 0.3 . Let $Y = X^2$. Then, Y takes values either 0 or 1 with probabilities $0.5, 0.5$. Thus, the expected value is $\mathbb{E}[Y] = 0.5$. Indeed, we have

$$\mathbb{E}[Y] = 1 \cdot 0.5 + 0 \cdot 0.5 = 1^2 \cdot 0.3 + 0^2 \cdot 0.5 + (-1)^2 \cdot 0.2.$$

Expectation of a function of RV

If X is a discrete random variable taking values x_i with probability $p(x_i)$, and g is a function, then

$$\mathbb{E}[g(X)] = \sum_i g(x_i)p(x_i).$$

In particular, for $g(x) = x^n$ and a positive integer n , we call $\mathbb{E}[g(X)] = \mathbb{E}[X^n]$ the n -th moment of X .

Proof. Let $Y = g(X)$ and $p_X(a), p_Y(a)$ be the probability mass functions of X and Y respectively. Then,

$$p_Y(y_j) = \sum_{i: g(x_i)=y_j} p_X(x_i).$$

$$\begin{aligned} \mathbb{E}[g(X)] &= \mathbb{E}[Y] = \sum_j y_j p_Y(y_j) = \sum_j y_j \sum_{i: g(x_i)=y_j} p_X(x_i) \\ &= \sum_j \sum_{i: g(x_i)=y_j} g(x_i) p_X(x_i) \\ &= \sum_i g(x_i) p_X(x_i). \end{aligned}$$

□

Linearity of Expectation

Let X be a discrete random variable with probability mass function $p(a)$. If a and b are real numbers, then

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b.$$

Proof. Using $g(x) = ax + b$ and the fact that $\sum_i p(x_i) = 1$, we have

$$\mathbb{E}[aX + b] = \sum_i (ax_i + b)p(x_i) = a \sum_i x_i p(x_i) + b \sum_i p(x_i) = a\mathbb{E}[X] + b.$$

□

References

[SR] Sheldon Ross, *A First Course in Probability*, 9th Edition, Pearson

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN
E-mail address: daesungk@illinois.edu