

A solutions manual for Set Theory by Thomas Jech

<https://github.com/9beach>

In December 2017, for no special reason I started studying mathematics and writing a solutions manual for [Set Theory by Thomas Jech](#).

GitHub repository [here](#), HTML versions [here](#), and PDF version [here](#).

Contents

Part I: Basic Set Theory

1. Axioms of Set Theory
2. Ordinal Numbers
3. Cardinal Numbers
4. Real Numbers *wip*
5. The Axiom of Choice and Cardinal Arithmetic
6. The Axiom of Regularity
7. Filters, Ultrafilters and Boolean Algebras
8. Stationary Sets
9. Combinatorial Set Theory
10. Measurable Cardinals
11. Borel and Analytic Sets
12. Models of Set Theory

Part II: Advanced Set Theory

13. Constructible Sets
14. Forcing
15. Applications of Forcing
16. Iterated Forcing and Martin's Axiom
17. Large Cardinals
18. Large Cardinals and L
19. Iterated Ultrapowers and $L[U]$
20. Very Large Cardinals
21. Large Cardinals and Forcing
22. Saturated Ideals
23. The Nonstationary Ideal
24. The Singular Cardinal Problem
25. Descriptive Set Theory
26. The Real Line

Part III: Selected Topics

- 27. Combinatorial Principles in L
- 28. More Applications of Forcing
- 29. More Combinatorial Set Theory
- 30. Complete Boolean Algebras
- 31. Proper Forcing
- 32. More Descriptive Set Theory
- 33. Determinacy
- 34. Supercompact Cardinals and the Real Line
- 35. Inner Models for Large Cardinals
- 36. Forcing and Large Cardinals
- 37. Martin's Maximum
- 38. More on Stationary Sets

Part I: Basic Set Theory

1. Axioms of Set Theory

1.1. Verify (1.1) $(a, b) = (c, d)$ if and only if $a = c$ and $b = d$.

Proof. If $a = c$ and $b = d$, then $(a, b) = \{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\} = (c, d)$. Assume that $(a, b) = (c, d)$ and $a = b$. Then $\{\{c\}, \{c, d\}\} = \{\{a\}\}$; thus $\{c, d\} \in \{\{a\}\}$ and $\{c\} \in \{\{a\}\}$, so $\{c, d\} = \{a\} = \{c\}$. Hence $c = d = a$. Since it was assumed that $a = b$, $a = c$ and $b = d$. Assume that $(a, b) = (c, d)$ and $a \neq b$. Since $\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$ and $\{a\} \neq \{a, b\}$, $\{c\} = \{a\}$ and $\{a, b\} = \{c, d\}$; thus $c = a$ and $d = b$. \square

1.2. There is no set X such that $P(X) \subset X$.

Proof. Suppose $P(X) \subset X$, then we have a function f from X onto $P(X)$. But the set $Y = \{x \in X : x \notin f(x)\}$ is not in the range of f : If $z \in X$ were such that $f(z) = Y$, then $z \in Y$ if and only if $z \notin Y$, a contradiction. Thus f is not a function of X onto $P(X)$; also a contradiction. \square

Let

$$\mathbb{N} = \bigcap \{X : X \text{ is inductive}\}.$$

\mathbb{N} is the smallest inductive set. Let us use the following notation:

$$0 = \emptyset, \quad 1 = \{0\}, \quad 2 = \{0, 1\}, \quad 3 = \{0, 1, 2\}, \quad \dots$$

If $n \in \mathbb{N}$, let $n + 1 = n \cup \{n\}$. Let us define $<$ (on \mathbb{N}) by $n < m$ if and only if $n \in m$.

A set T is transitive if $x \in T$ implies $x \subset T$.

1.3. If X is inductive, then the set $\{x \in X : x \subset X\}$ is inductive. Hence \mathbb{N} is transitive, and for each $n, n = \{m \in \mathbb{N} : m < n\}$.

Proof. Let $Y = \{x \in X : x \subset X\}$. Since $\emptyset \subset X$, and $\emptyset \in X$, $\emptyset \in Y$. Now let $y \in Y$. Since $Y \subset X$, and X is inductive, $y \in X$, i.e., $\{y\} \subset X$, and $y \cup \{y\} \in X$, and since $y \subset X$, $y \cup \{y\} \subset X$; thus $y \cup \{y\} \in Y$. Therefore, Y is inductive.

Let $Y_{\mathbb{N}} = \{x \in \mathbb{N} : x \subset \mathbb{N}\}$, then $Y_{\mathbb{N}} \subset \mathbb{N}$, and since $Y_{\mathbb{N}}$ is inductive, $\mathbb{N} \subset Y_{\mathbb{N}}$; thus $\mathbb{N} = Y_{\mathbb{N}}$, and so we have that $x \in \mathbb{N}$ implies $x \subset \mathbb{N}$. Therefore, \mathbb{N} is transitive.

It's obvious that $k \in n \cup \{n\}$ if and only if $k \in n$ or $k = n$. So it follows that for all $k, n \in \mathbb{N}$, $k < n + 1$ if and only if $k < n$ or $k = n$. Now we show that for each $n, n = \{m \in \mathbb{N} : m < n\}$ by induction. Let $P(x)$ be the property " $x = \{m \in \mathbb{N} : m < x\}$ ". $P(0)$ holds, and assume that $P(n)$ holds. $n + 1 = n \cup \{n\} = \{m \in \mathbb{N} : m < n\} \cup \{n\} = \{m \in \mathbb{N} : m < n \text{ or } m = n\} = \{m \in \mathbb{N} : m < n + 1\} = P(n + 1)$ holds. Therefore, for each $n, n = \{m \in \mathbb{N} : m < n\}$. \square

1.4. If X is inductive, then the set $\{x \in X : x \text{ is transitive}\}$ is inductive. Hence every $n \in \mathbb{N}$ is transitive.

Proof. Let $Y = \{x \in X : x \text{ is transitive}\}$. Since $\emptyset \in X$, and \emptyset is transitive, $\emptyset \in Y$. Now let $y \in Y$. Since $Y \subset X$, and X is inductive, $y \in X$, and $y \cup \{y\} \in X$. Let $z \in y \cup \{y\}$, then either $z \in y$ or $z = y$; since y is transitive, in any case, $z \subset y \cup \{y\}$. Thus $y \cup \{y\}$ is transitive, and so $y \cup \{y\} \in Y$. Therefore, Y is inductive.

Let $Y_{\mathbb{N}} = \{x \in \mathbb{N} : x \text{ is transitive}\}$, then $Y_{\mathbb{N}} \subset \mathbb{N}$, and since $Y_{\mathbb{N}}$ is inductive, $\mathbb{N} \subset Y_{\mathbb{N}}$; thus $\mathbb{N} = Y_{\mathbb{N}}$. Therefore, every $n \in \mathbb{N}$ is transitive. \square

1.5. If X is inductive, then the set $\{x \in X : x \text{ is transitive and } x \notin x\}$ is inductive. Hence $n \notin n$ and $n \neq n + 1$ for each $n \in \mathbb{N}$.

Proof. Let $Y = \{x \in X : x \text{ is transitive and } x \notin x\}$. Since $\emptyset \in X$, and \emptyset is transitive and $\emptyset \notin \emptyset$, $\emptyset \in Y$. Now let $y \in Y$. Since $Y \subset X$, and X is inductive, $y \in X$, and $y \cup \{y\} \in X$. We already have that $y \cup \{y\}$ is transitive. Suppose $y \cup \{y\} \in y \cup \{y\}$, then $y \cup \{y\} = y$ or $y \cup \{y\} \in y$, i.e., $y \cup \{y\} \subset y$; in any case, $\{y\} \subset y$, i.e., $y \in y$; a contradiction. Thus $y \cup \{y\} \notin y \cup \{y\}$, and so $y \cup \{y\} \in Y$. Therefore, Y is inductive.

Let $Y_{\mathbb{N}} = \{x \in \mathbb{N} : x \text{ is transitive and } x \notin x\}$, then $Y_{\mathbb{N}} \subset \mathbb{N}$, and since $Y_{\mathbb{N}}$ is inductive, $\mathbb{N} \subset Y_{\mathbb{N}}$; thus $\mathbb{N} = Y_{\mathbb{N}}$, and so $n \notin n$. Suppose $n + 1 = n$, i.e., $n \cup \{n\} = n$, then $\{n\} \subset n$, i.e., $n \in n$; a contradiction. Therefore, $n \notin n$ and $n \neq n + 1$ for each $n \in \mathbb{N}$. \square

1.6. If X is inductive, then $\{x \in X : x \text{ is transitive and every nonempty } z \subset x \text{ has an } \in\text{-minimal element}\}$ is inductive (t is \in -minimal in z if there is no $s \in z$ such that $s \in t$).

Proof. Let $P(x)$ be the property “every nonempty $z \subset x$ has an \in -minimal element”; let $Y = \{x \in X : x \text{ is transitive and } P(x)\}$. $\emptyset \in X$, and is transitive, and has no nonempty subset, thus $\emptyset \in Y$. Let $y \in Y$. Since $Y \subset X$, and X is inductive, $y \in X$, and $y \cup \{y\} \in X$. We already have that $y \cup \{y\}$ is transitive. Now we show that $P(y \cup \{y\})$ holds. $y \notin y$; otherwise $\{y\} \subset y$ does not have an \in -minimal element ($\cdots y \in y \in y \cdots$), a contradiction. There is no $a \in y$ such that $y \in a$; otherwise $y \in a \in y \Rightarrow y \in a \subset y \Rightarrow y \in y$. Hence for every nonempty $z \subset y$ if m is an \in -minimal element in z then so is in $z \cup \{y\}$; otherwise $y \in m$, a contradiction. Similarly, $P(\{y\})$ holds; otherwise $\cdots y \in y \in y \cdots$. Therefore, $P(y \cup \{y\})$ holds, and so Y is inductive. \square

1.7. Every nonempty $X \subset \mathbb{N}$ has an \in -minimal element.

[Pick $n \in X$ and look at $X \cap n$.]

Proof. Since \mathbb{N} is the smallest inductive set, from 1.6, we have that every $n \in \mathbb{N}$ has an \in -minimal element. Let $n \in X$. If $n \cap X = \emptyset$, then n is an \in -minimal element. Suppose not. There exists $m \in X \setminus n$ such that $m \in n$, but then since $n = \{m \in \mathbb{N} : m < n\}$, $n \cap X \neq \emptyset$; a contradiction. If $n \cap X \neq \emptyset$, then $n \cap X \subset n$ has an \in -minimal element, and it's an \in -minimal element of X ; otherwise similarly to the previous, a contradiction. \square

1.8. If X is inductive then so is $\{x \in X : x = \emptyset \text{ or } x = y \cup \{y\} \text{ for some } y\}$. Hence each $n \neq 0$ is $m + 1$ for some m .

1. Axioms of Set Theory

Proof. Let $A = \{x \in X : x = \emptyset \text{ or } x = y \cup \{y\} \text{ for some } y\}$; let $a \neq \emptyset \in A$. Since $a = y \cup \{y\}$ for some y , so is $a \cup \{a\}$ for a ; thus $a \cup \{a\} \in A$. Therefore, A is inductive, and each $n \neq 0$ is $m + 1$ for some m . \square

1.9 (Induction). Let A be a subset of \mathbb{N} such that $0 \in A$, and if $n \in A$ then $n + 1 \in A$. Then $A = \mathbb{N}$.

Proof. By definition, A is an inductive subset of \mathbb{N} . Therefore, $A = \mathbb{N}$. \square

A set X has n elements (where $n \in \mathbb{N}$) if there is a one-to-one mapping of n onto X . A set is *finite* if it has n elements for some $n \in \mathbb{N}$, and *infinite* if it is not finite.

A set S is *T-finite* if every nonempty $X \subset P(S)$ has a \subset -maximal element, i.e., $u \in X$ such that there is no $v \in X$ with $u \subset v$ and $u \neq v$. S is *T-infinite* if it is not T-finite. (T is for Tarski.)

1.10. Each $n \in \mathbb{N}$ is T-finite.

Proof. Let $A = \{n \in \mathbb{N} : n \text{ is T-finite}\}$. We show that $A = \mathbb{N}$ by induction.

$P(\emptyset) = \{\emptyset\}$ has the only nonempty subset $\{\emptyset\}$ which has a \subset -maximal element \emptyset .

Let $n \in A$; let $X \subset P(n+1)$. For some $Y \subset P(n)$, X is either Y or $Z = \{x \cup \{n\} : x \in Y\}$. For the latter case, let a be a \subset -maximal element of Y . Then it's obvious that $a \cup \{n\}$ is a \subset -maximal element of Z ; thus X is T-finite. \square

1.11. \mathbb{N} is T-infinite; the set $\mathbb{N} \subset P(\mathbb{N})$ has no \subset -maximal element.

Proof. For any $n \in \mathbb{N}$, there exists $n + 1$ such that $n \subsetneq n + 1$; thus $\mathbb{N} \subset P(\mathbb{N})$ has no \subset -maximal element. \square

Note that $\mathbb{N} \in P(\mathbb{N})$, $\mathbb{N} \subset P(\mathbb{N})$, and $\bigcup \mathbb{N} = \mathbb{N}$.

1.12. Every finite set is T-finite.

Proof. Let F be a finite set, then there is a one-to-one mapping f of F onto $n \in \mathbb{N}$. Let $A \subset P(F)$ be a nonempty set. Then $B = \{f(X) \subset P(n) : X \in A\}$ is nonempty, and has a \subset -maximal element. It's obvious that $\forall X, Y \in A (X \subset Y \iff f(X) \subset f(Y))$; A has a \subset -maximal element. \square

1.13. Every infinite set is T-infinite.

[If S is infinite, consider $X = \{u \subset S : u \text{ is finite}\}$.]

Proof. Since $\emptyset \in X$, X is nonempty. Suppose X has a \subset -maximal element m . Then $S \setminus m \neq \emptyset$; otherwise S is a subset of a finite set; a contradiction, and so there exists $x \in S \setminus m$. Then $m \subsetneq m \cup \{x\} \in X$; a contradiction. \square

1.14. The Separation Axioms follow from the Replacement Schema.

[Given ϕ , let $F = \{(x, x) : \phi(x)\}$. Then $\{x \in X : \phi(x)\} = F(X)$, for every X .]

Proof. Let $\varphi(x, y)$ be $x = y \wedge \phi(x)$. Then $F = \{(x, x) : \phi(x)\} = \{(x, y) : \varphi(x, y)\}$. Since $\forall x \forall y \forall z (\varphi(x, y) \wedge \varphi(x, z) \rightarrow y = z)$ holds, $\varphi(x, y)$ is a functional formula. Therefore, we have that The Separation Axioms follow from the Replacement Schema.

$$F(X) = \{y : (\exists x \in X)\varphi(x, y)\} = \{y : (\exists x \in X)x = y \wedge \phi(x)\} = \{x : (\exists x \in X)\phi(x)\} = \{x \in X : \phi(x)\}. \quad \square$$

1.15. Instead of Union, Power Set, and Replacement Axioms consider the following weaker versions:

$$(1.8) \quad \forall X \exists Y \bigcup X \subset Y, \text{ i.e., } \forall X \exists Y (\forall x \in X)(\forall u \in x)u \in Y,$$

$$(1.9) \quad \forall X \exists Y P(X) \subset Y, \text{ i.e., } \forall X \exists Y \forall u (u \subset X \rightarrow u \in Y),$$

$$(1.10) \quad \text{If a class } F \text{ is a function, then } \forall X \exists Y F(X) \subset Y.$$

Then axioms 1.4, 1.5, and 1.7 can be proved from (1.8), (1.9), and (1.10), using the Separation Schema (1.3).

Proof. Using the Separation Schema,

$$(1.8) \implies \{x \in Y : (\exists a \in X)x \in a\} = \bigcup X,$$

$$(1.9) \implies \{x \in Y : x \subset X\} = P(X),$$

$$(1.10) \implies \{y \in Y : (\exists x \in X)\varphi(x, y, p)\} = F(X). \quad \square$$

2. Ordinal Numbers

2.1. The relation “ $(P, <)$ is isomorphic to $(Q, <)$ ” is an equivalence relation (on the class of all partially ordered sets).

Proof. Let $(P, <)$, $(Q, <)$, and $(G, <)$ be partially ordered sets. P is isomorphic to P by $f : x \mapsto x$. If P is isomorphic to Q by f , then Q is isomorphic to P by f^{-1} . If P is isomorphic to Q by f , and Q is isomorphic to G by g , then P is isomorphic to G by $g \circ f$. \square

2.2. α is a limit ordinal if and only if $\beta < \alpha$ implies $\beta + 1 < \alpha$, for every β .

Proof. Suppose $\beta + 1 \geq \alpha$. Since $\beta < \alpha$, $\alpha = \beta + 1$; thus α is a successor ordinal, a contradiction. Conversely, suppose that α is a successor ordinal. Then $\alpha = \beta + 1$ for some $\beta < \alpha$, a contradiction. \square

2.3. If a set X is inductive, then $X \cap \text{Ord}$ is inductive. The set $\mathbb{N} = \bigcap \{X : X \text{ is inductive}\}$ is the least limit ordinal $\neq 0$.

Proof. $\emptyset \in X \cap \text{Ord}$. If $x \in X \cap \text{Ord}$, then $x + 1 = x \cup \{x\} \in X \cap \text{Ord}$. Therefore, $X \cap \text{Ord}$ is inductive.

By exercise 1.3, \mathbb{N} is transitive. Since $\mathbb{N} \cap \text{Ord} = \mathbb{N}$, by exercise 1.7 and lemma 2.11, \mathbb{N} is well-ordered by \in . Thus \mathbb{N} is an ordinal number. And, if $n \in \mathbb{N}$, then $n + 1 \in \mathbb{N}$. Therefore, \mathbb{N} is the least limit ordinal $\neq \emptyset$. \square

2.4. (Without the Axiom of Infinity). Let $\omega = \text{least limit } \alpha \neq 0$ if it exists, $\omega = \text{Ord}$ otherwise. Prove that the following statements are equivalent:

- (i) There exists an inductive set.
- (ii) There exists an infinite set.
- (iii) ω is a set.

[For (ii) \rightarrow (iii), apply Replacement to the set of all finite subsets of X .]

Proof. *Case (i) \Rightarrow (ii).* By exercise 1.11, there exists an inductive T-infinite set \mathbb{N} , and by exercise 1.12 and 1.13, \mathbb{N} is infinite.

Case (ii) \Rightarrow (iii). Let $Y = \{x \subset X : x \text{ is a finite set}\}$. Then $Y \subset P(X)$ is a set, and since $x \in Y$ is finite, there is a mapping f from Y into ω given by $x \mapsto n$, n such that x has n elements. Since X is infinite, if $x \in Y$ has n elements, then there exists $y \in Y$ which has $n + 1$ elements; thus $\text{ran}(f) = \omega$. Therefore, ω is a set.

Case (iii) \Rightarrow (i). By definition, $\emptyset \in \omega$, and if $\alpha \in \omega$, then $\alpha + 1 \in \omega$. Therefore, ω is inductive. \square

2.5. If W is a well-ordered set, then there exists no sequence $\langle a_n : n \in \mathbb{N} \rangle$ in W such that $a_0 > a_1 > a_2 > \dots$.

Proof. Otherwise there is no least element. \square

2.6. There are arbitrarily large limit ordinals; i.e., $\forall \alpha \exists \beta > \alpha$ (β is a limit).

[Consider $\lim_{n \rightarrow \omega} \alpha_n$, where $\alpha_{n+1} = \alpha_n + 1$.]

Proof. Given $\alpha_0 \in \text{Ord}$, we define $\alpha_{n+1} = \alpha_n + 1$, and $\beta = \lim_{n \rightarrow \omega} \alpha_n$, i.e., $\beta = \bigcup \{\alpha_n : n < \omega\} = \sup \{\alpha_n : n < \omega\}$. Then since the union of ordinals is an ordinal, β is an ordinal. And for every $\gamma < \beta$, there exists α_n such that $\alpha_n > \gamma$; otherwise $\gamma \geq \sup \{\alpha_n : n < \omega\}$, a contradiction. Thus $\gamma + 1 < \alpha_n + 1 = \alpha_{n+1} < \beta$, and so β is a limit ordinal. Therefore, there are arbitrarily large limit ordinals. \square

2.7. Every normal sequence $\langle \gamma_\alpha : \alpha \in \text{Ord} \rangle$ has arbitrarily large *fixed points*, i.e., α such that $\gamma_\alpha = \alpha$.

[Let $\alpha_{n+1} = \gamma_{\alpha_n}$, and $\alpha = \lim_{n \rightarrow \omega} \alpha_n$.]

Proof. Since $\langle \gamma_\alpha : \alpha \in \text{Ord} \rangle$ is increasing, for $\beta \in \text{Ord}$, there exists $m \in \text{Ord}$ such that $\gamma_m > \beta$. Let $\alpha_0 = \gamma_m$, $\alpha_{n+1} = \gamma_{\alpha_n}$. Then $\langle \alpha_n : n \in \omega \rangle$ is increasing, so we let $\alpha = \lim_{n \rightarrow \omega} \alpha_n$; similarly to 2.6, α is a limit ordinal. Hence we have that $\alpha = \lim_{n \rightarrow \omega} \alpha_{n+1} = \lim_{n \rightarrow \omega} \gamma_{\alpha_n} = \lim_{\xi \rightarrow \alpha} \gamma_\xi = \gamma_{\lim_{\xi \rightarrow \alpha} \xi} = \gamma_\alpha$. Therefore, $\gamma_\alpha = \alpha$ \square

2.8. For all α, β and γ ,

- (i) $\alpha \cdot (\beta + \gamma) = \alpha \cdot \beta + \alpha \cdot \gamma$,
- (ii) $\alpha^{\beta+\gamma} = \alpha^\beta \cdot \alpha^\gamma$,
- (iii) $(\alpha^\beta)^\gamma = \alpha^{\beta \cdot \gamma}$.

Proof. *Case (i).* We show by induction on γ . $\alpha \cdot (\beta + 0) = \alpha \cdot \beta = \alpha \cdot \beta + \alpha \cdot 0$. $\alpha \cdot (\beta + (\gamma + 1)) = \alpha \cdot ((\beta + \gamma) + 1) = \alpha \cdot (\beta + \gamma) + \alpha = \alpha \cdot \beta + \alpha \cdot \gamma + \alpha = \alpha \cdot \beta + \alpha \cdot (\gamma + 1)$. For all limit $\gamma > 0$, $\alpha \cdot (\beta + \gamma) = \alpha \cdot \lim_{\xi \rightarrow \gamma} (\beta + \xi) = \lim_{\xi \rightarrow \gamma} \alpha \cdot (\beta + \xi) = \lim_{\xi \rightarrow \gamma} (\alpha \cdot \beta + \alpha \cdot \xi) = \alpha \cdot \beta + \lim_{\xi \rightarrow \gamma} (\alpha \cdot \xi) = \alpha \cdot \beta + \alpha \cdot \lim_{\xi \rightarrow \gamma} \xi = \alpha \cdot \beta + \alpha \cdot \gamma$

Case (ii) and (iii). Similarly to the previous. \square

2.9. (i) Show that $(\omega + 1) \cdot 2 \neq \omega \cdot 2 + 1 \cdot 2$.

(ii) Show that $(\omega \cdot 2)^2 \neq \omega^2 \cdot 2^2$.

Proof. *Case (i).* $(\omega + 1) \cdot 2 = \omega + 1 + \omega + 1 = \omega + \omega + 1 = \omega \cdot 2 + 1 < \omega \cdot 2 + 2 = \omega \cdot 2 + 1 \cdot 2$

Case (ii). $(\omega \cdot 2)^2 = \omega \cdot 2 \cdot \omega \cdot 2 = \omega \cdot (2 \cdot \omega) \cdot 2 = \omega \cdot \omega \cdot 2 < \omega \cdot \omega \cdot 4 = \omega^2 \cdot 2^2$ \square

2.10. If $\alpha < \beta$ then $\alpha + \gamma \leq \beta + \gamma$, $\alpha \cdot \gamma \leq \beta \cdot \gamma$, and $\alpha^\gamma \leq \beta^\gamma$.

Proof. We show that if $\alpha < \beta$ then $\alpha + \gamma \leq \beta + \gamma$ by induction on γ .

$\alpha + 0 \leq \beta + 0$. $\alpha + \gamma + 1 \leq \alpha + 1 + \gamma + 1 \leq \beta + \gamma + 1$. Let a limit ordinal > 0 be γ . For all $\xi < \gamma$, if $\alpha + \xi < \beta + \xi$ then $\sup \{\alpha + \xi : \xi < \gamma\} \leq \sup \{\beta + \xi : \xi < \gamma\}$. Therefore, $\alpha + \gamma \leq \beta + \gamma$.

Similarly to the previous, $\alpha \cdot \gamma \leq \beta \cdot \gamma$, and $\alpha^\gamma \leq \beta^\gamma$. \square

2.11. Find α, β, γ such that

- (i) $\alpha < \beta$ and $\alpha + \gamma = \beta + \gamma$,
- (ii) $\alpha < \beta$ and $\alpha \cdot \gamma = \beta \cdot \gamma$,
- (iii) $\alpha < \beta$ and $\alpha^\gamma = \beta^\gamma$.

Proof. *Case (i).* $0 + \omega = 1 + \omega$

Case (ii). $1 \cdot \omega = 2 \cdot \omega$

Case (iii). $2^\omega = 3^\omega$ \square

2. Ordinal Numbers

2.12. Let $\varepsilon_0 = \lim_{n \rightarrow \omega} \alpha_n$ where $\alpha_0 = \omega$ and $\alpha_{n+1} = \omega^{\alpha_n}$ for all n . Show that ε_0 is the least ordinal ε such that $\omega^\varepsilon = \varepsilon$.

Proof. $\varepsilon_0 = \sup \{\omega, \omega^\omega, \omega^{\omega^\omega}, \dots\} = \sup \{\omega^\omega, \omega^{\omega^\omega}, \omega^{\omega^{\omega^\omega}}, \dots\} = \omega^{\sup \{\omega, \omega^\omega, \omega^{\omega^\omega}, \dots\}} = \omega^{\varepsilon_0}$.

Suppose that there exists $\varepsilon < \varepsilon_0$ such that $\omega^\varepsilon = \varepsilon$. Then since for every finite number a , $a \neq \omega^a$, $\varepsilon \geq \omega$, and so there exists the least n such that $n > 0, n \in \omega$, and $\alpha_n > \varepsilon$. Then $\alpha_n = \omega^{\alpha_{n-1}} > \varepsilon = \omega^\varepsilon$. But since $\alpha_{n-1} < \varepsilon$, a contradiction. \square

A limit ordinal $\gamma > 0$ is called *indecomposable* if there exist no $\alpha < \gamma$ and $\beta < \gamma$ such that $\alpha + \beta = \gamma$.

2.13. A limit ordinal $\gamma > 0$ is indecomposable if and only if $\alpha + \gamma = \gamma$ for all $\alpha < \gamma$ if and only if $\gamma = \omega^\alpha$ for some α .

Proof. $\gamma > 0$ is indecomposable if and only if $\alpha + \beta < \gamma$ for all $\alpha < \gamma$ and $\beta < \gamma$; otherwise $\alpha < \gamma < \alpha + \beta$, and so there is δ such that $\alpha + \delta = \gamma$ and $\delta < \gamma$; a contradiction. Hence for all $\alpha < \gamma$, $\alpha + \gamma = \sup \{\alpha + \xi : \xi < \gamma\} \leq \gamma$, but we know that $\sup \{\alpha + \xi : \xi < \gamma\} \geq \sup \{\xi : \xi < \gamma\} = \gamma$. Thus $\alpha + \gamma = \gamma$. Conversely, if $\alpha + \gamma = \gamma$ for all $\alpha < \gamma$ then $\sup \{\alpha + \xi : \xi < \gamma\} = \gamma$. It follows that for all $\alpha < \gamma$ and $\beta < \gamma$, $\alpha + \beta < \gamma$.

Let $\gamma = \omega^{\beta_1} \cdot k_1 + \dots + \omega^{\beta_n} \cdot k_n$ be Cantor's normal form. If a limit ordinal $\gamma \neq \omega^\alpha$ for all α , i.e., $n > 1$ or $k_n > 1$ for some n then clearly not indecomposable, i.e., *decomposable*. Conversely, let $\beta_1, \beta_2 < \gamma = \omega^\alpha$ for all $\alpha > 0$. There exist $\alpha' < \alpha$ and $k < \omega$ such that $\beta_1, \beta_2 < \omega^{\alpha'} \cdot k$ (Consider Cantor's normal forms of β_1 and β_2). Hence $\beta_1 + \beta_2 < \omega^{\alpha'} \cdot (k + k) < \omega^\alpha = \gamma$. \square

2.14. If E is a well-founded relation on P , then there is no sequence $\langle a_n : n \in \mathbb{N} \rangle$ in P such that $a_1 E a_0, a_2 E a_1, a_3 E a_2, \dots$

Proof. Otherwise, $\dots E a_3 E a_2 E a_1 E a_0$; there is no E -minimal elements. \square

2.15. (Well-Founded Recursion). Let E be a well-founded relation on a set P , and let G be a function. Then there exists a function F such that for all $x \in P, F(x) = G(x, F \upharpoonright \{y \in P : y E x\})$.

Proof. A set $B \subset A$ is called E -transitive in A if $\{y \in A : y E x\} \subset B$ holds for all $x \in B$. Let $T = \{g : g \text{ is a function. } \text{dom}(g) \text{ is a } E\text{-transitive in } P, \text{ and } (\forall x \in \text{dom}(g)) g(x) = G(x, g \upharpoonright x)\}$. T is nonempty, since for every E -minimal element $m \in P, \{m\}$ is E -transitive, and a function g of $\{m\}$ is given by $m \mapsto G(m, \emptyset)$.

We claim that $\bigcup T$ is a function. Suppose not. There is a E -minimal element m of the set $\{x \in \text{dom}(g_1) \cap \text{dom}(g_2) : g_1(x) \neq g_2(x) \text{ for some } g_1, g_2 \in T\}$. Then $g_1(m) = G(m, g_1 \upharpoonright \{y \in \text{dom}(g_1) : y E m\}) = G(m, g_1 \upharpoonright \{y \in \text{dom}(g_2) : y E m\}) = G(m, g_2 \upharpoonright \{y \in \text{dom}(g_2) : y E m\}) = g_2(m)$, a contradiction. Similarly, $\text{dom}(\bigcup T) = P$. Therefore, $\bigcup T = F$. \square

3. Cardinal Numbers

- 3.1.** (i) A subset of a finite set is finite.
 (ii) The union of a finite set of finite sets is finite.
 (iii) The power set of a finite set is finite.
 (iv) The image of a finite set (under a mapping) is finite.

Proof. (i) Let X be a finite set, and $Y \subset X$. Suppose that Y is infinite. Then Y is T-infinite, so there is $S \subset P(Y)$ such that S has no \subset -maximal element. But by definition, $P(Y) \subset P(X)$, and so $S \subset P(X)$, a contradiction.

(ii) For p such that $0 < p < \omega$, let S_i be a finite set, and f_i be a function of S_i onto a finite ordinal n_i for each $i < p$. Let $S = \bigcup_{i < p} S_i$; let $f : S \rightarrow \sum_{i < p} n_i$ given by $x \mapsto \sum_{i < k} n_i + f_k(x)$ where k is the least number such that $x \in S_k$. Then f is one-to-one function of S into $\sum_{i < p} n_i$ which is bounded. Thus S is finite.

(iii) Let X be a finite set. $|P(X)| = 2^{|X|} < \aleph_0$.

(iv) Let f be a function of a finite set X onto Y . Then there is a one-to-one function g of X onto $n < \omega$. Clearly, a function h of $f(X)$ into n given by $y \mapsto \bigcap g_{-1} \circ f_{-1}(y)$ exists. \square

- 3.2.** (i) A subset of a countable set is at most countable.
 (ii) The union of a finite set of countable sets is countable.
 (iii) The image of a countable set (under a mapping) is at most countable.

Proof. (i) Let X be a countable set, and $Y \subset X$. Then there is a one-to-one function f of X onto ω . Let id_Y be a function of Y into X given by $x \mapsto x$. Clearly the function $f \cdot id_Y$ is a function of Y into ω , and so $|Y| \leq \aleph_0$. Therefore, by definition of \aleph_0 , Y is at most countable.

(ii) For some n such that $0 < n < \omega$, Let $S = \bigcup_{i < n} S_i$ where S_i is a countable set; for each $I < n$, let f_i be a function of S_i onto ω . Let $S = \bigcup_{i < n} S_i$; let $f : S \rightarrow \omega$ given by $x \mapsto 2^i 3^{f_i(x)}$ where i is the least number $x \in S_i$. Then f is one-to-one function of S into ω . Thus S is countable.

(iii) Let f be a function of a countable set X onto Y . Then there is a one-to-one function g of X onto ω . Clearly, a function h of $f(X)$ into ω given by $y \mapsto \bigcap g_{-1} \circ f_{-1}(y)$ exists. \square

3.3. $\mathbb{N} \times \mathbb{N}$ is countable.

$$[f(m, n) = 2^m(2n + 1) - 1.]$$

Proof. (i) Let f be a function of $\mathbb{N} \times \mathbb{N}$ into \mathbb{N} given by $(m, n) \mapsto 2^m(2n + 1) - 1$. Let $x \in \omega$, and $m = \sup \{a \in \omega : 2^a \text{ divides } x + 1\}$. Then $(x + 1)/2^m$ is odd, so there is $n \in \omega$ such that $2n + 1 = (x + 1)/2^m$. Thus f is a function onto \mathbb{N} . Suppose that $2^{m_1}(2n_1 + 1) = 2^{m_2}(2n_2 + 1)$. Since $2x + 1 \neq 2y$ for all $x, y \in \mathbb{N}$, the prime factorization of $2x + 1$ does not have 2 as a factor. Thus $m_1 = m_2$ and $n_1 = n_2$, and so f is a one-to-one function onto \mathbb{N} . Therefore, $\mathbb{N} \times \mathbb{N}$ is countable. \square

- 3.4.** (i) The set of all finite sequences in \mathbb{N} is countable.
 (ii) The set of all finite subsets of a countable set is countable.

3. Cardinal Numbers

Proof. (i) Let f be a function of all finite sequences in \mathbb{N} into \mathbb{N} given by, for some $k \in \mathbb{N}$, $\langle s_i \in \mathbb{N} : i < k \rangle \mapsto \prod_{i < k} p_{i+1}^{s_i+1} - 1$ where p_i is the i -th prime number. Clearly, f is a one-to-one function onto \mathbb{N} .

(ii) Let X be a countable set; let Y be a set of all finite subsets of X . Then there is a one-to-one function f of X onto \mathbb{N} given by $x \mapsto n$ for some $n < \omega$, and so for $S \in Y$, there is a unique increasing finite sequence $\langle f(x) : x \in S \rangle$. Thus there is a one-to-one function of Y into all finite sequences in \mathbb{N} ; $Y \leq \aleph_0$, and $\aleph_0 = |X| = \{S \in Y : S \text{ is singleton}\} \subset Y$; thus $\aleph_0 \leq Y$. Therefore, $Y = \aleph_0$. \square

3.5. Show that $\Gamma(\alpha \times \alpha) \leq \omega^\alpha$.

Proof. We show this by induction of α . $\Gamma(0 \times 0) \leq \omega^0$. $\Gamma(\alpha \times \alpha) \leq \omega^\alpha \Leftrightarrow \Gamma((\alpha + 1) \times (\alpha + 1)) = \Gamma(\alpha \times \alpha) + \alpha \cdot 2 + 1 \leq \omega^\alpha + \omega^\alpha \cdot 2 + \omega^\alpha = \omega^\alpha \cdot 4 \leq \omega^{\alpha+1}$. For a limit ordinal $\gamma > 0$, by definition $\Gamma(\gamma \times \gamma) = \sup \{\Gamma(\alpha \times \alpha) : \alpha < \gamma\} \leq \omega^\gamma$. \square

3.6. There is a well-ordering of the class of all finite sequences of ordinals such that for each α , the set of all finite sequences in ω_α is an initial segment and its order-type is ω_α .

Proof. We define:

$$\begin{aligned} \langle \alpha_0, \dots \rangle \prec \langle \beta_0, \dots \rangle &\leftrightarrow \\ \text{there is } k \text{ such that} & \\ \alpha_k < \beta_k \text{ and } \alpha_i = \beta_i \text{ for all } i < k, & \\ \langle \alpha_i : i < m \rangle < \langle \beta_i : i < n \rangle &\leftrightarrow \\ \text{either } \sum_{i < m} \alpha_i + m < \sum_{i < n} \beta_i + n & \\ \text{or } \sum_{i < m} \alpha_i + m = \sum_{i < n} \beta_i + n & \\ \text{and } \langle \alpha_i : i < m \rangle \prec \langle \beta_i : i < n \rangle. & \end{aligned}$$

Let X be the class of all finite sequences of ordinals. The relation $<$ defined above is a linear ordering of X . Moreover, if $S \subset X$ is nonempty, then S has a least element. If we let $\Gamma(\alpha) =$ the order-type of the set $\{\beta \in X : \beta < \alpha\}$ for $\alpha \in X$, then Γ is a one-to-one mapping of X onto Ord . Note that for a finite sequence α in ω , $\Gamma(\alpha) \in \omega$, and so $\langle \omega \rangle$ is the least element α of X such that α is not a finite sequence in ω ; thus $\Gamma(\langle \omega \rangle) = \omega$.

Let $\gamma(\alpha) = \Gamma(\langle \alpha \rangle)$. Note that $\gamma(\alpha)$ is an increasing function of α , and also that since each infinite cardinal is indecomposable, by definition of $(X, <)$, $\gamma(\omega_\alpha)$ is the set of all finite sequences in ω_α . Let $\eta(\alpha) =$ the order-type of the set of all finite sequences in α . Then $\gamma(\alpha) \leq \eta(\alpha)$ and $\gamma(\omega_\alpha) = \eta(\omega_\alpha)$ for each α . We show that $\gamma(\omega_\alpha) = \omega_\alpha$ by induction of α . This is true for $\alpha = 0$. Thus let α be the least ordinal such that $\gamma(\omega_\alpha) \neq \omega_\alpha$. Since γ is increasing, $\gamma(\omega_\alpha) \geq \omega_\alpha$; thus $\gamma(\omega_\alpha) > \omega_\alpha$, and so there is a sequence β such that $\Gamma(\beta) = \omega_\alpha$ and $\beta < \langle \omega_\alpha \rangle$. Then there is an ordinal δ such that $\beta < \langle \delta \rangle < \langle \omega_\alpha \rangle$; thus $\Gamma(\beta) = \omega_\alpha < \gamma(\delta) \leq \eta(\delta)$

$\Leftrightarrow \aleph_\alpha \leq |\eta(\delta)| = |\eta(|\delta|)| \leq \eta(|\delta|)$. But since $\delta < \omega_\alpha$, by the minimality of α , $\eta(|\delta|) = |\delta| < \aleph_\alpha$. A contradiction. Finally, by definition of γ , for each nonzero limit ordinal α , $\gamma(\omega_\alpha) = \sup \{\gamma(\omega_\xi) : \xi < \alpha\} = \omega_\alpha$. \square

We say that a set B is a *projection* of a set A if there is a mapping of A onto B . Note that B is a projection of A if and only if there is a partition P of A such that $|P| = |B|$. If $|A| \geq |B| > 0$, then B is a projection of A . Conversely, using the Axiom of Choice, one shows that if B is a projection of A , then $|A| \geq |B|$. This, however, cannot be proved without the Axiom of Choice.

3.7. If B is a projection of ω_α , then $|B| \leq \aleph_\alpha$.

Proof. Let f be a function of ω_α onto B . Then a one-to-one function g of B into ω_α is given by $x \mapsto \min f^{-1}(x)$. \square

3.8. The set of all finite subsets of ω_α has cardinality \aleph_α .

[The set is a projection of the set of finite sequences.]

Proof. Let X be the set of all finite sequences in ω_α ; let Y be the set of all finite subsets of ω_α . Then there is a function of X onto Y given by $\langle \alpha_0, \dots, \alpha_n \rangle \mapsto \{\alpha_0, \dots, \alpha_n\}$. Thus $\aleph_\alpha = |X| \geq |Y|$. But there is a one-to-one mapping of $S \subset Y$ such that each $x \in S$ is singleton onto a set of cardinality \aleph_α . Thus $Y \geq \aleph_\alpha$. Therefore, $|Y| = \aleph_\alpha$. \square

3.9. If B is a projection of A , then $|P(B)| \leq |P(A)|$.

[Consider $g(X) = f^{-1}(X)$, where f maps A onto B .]

Proof. Since for each $S \subset B$, there is unique $f^{-1}(S) \subset A$, there is a one-to-one function of $P(B)$ into $P(A)$ given by $S \mapsto f^{-1}(S)$. \square

3.10. $\omega_{\alpha+1}$ is a projection of $P(\omega_\alpha)$.

[Use $|\omega_\alpha \times \omega_\alpha| = \omega_\alpha$ and project $P(\omega_\alpha \times \omega_\alpha)$: If $R \subset \omega_\alpha \times \omega_\alpha$ is a well-ordering, let $f(R)$ be its order-type.]

Proof. Since $\omega_{\alpha+1}$ is a set of possible well-orderings of subsets of X such that $|X| = \aleph_\alpha$, there is $R \in P(\omega_\alpha \times \omega_\alpha)$ such that $f(R) = \beta$ for each $\beta \in \omega_{\alpha+1}$. Let $g(R) = f(R)$ if R is a well-ordering; otherwise $g(R) = 0$. Then we have a mapping of $P(\omega_\alpha \times \omega_\alpha)$ onto $\omega_{\alpha+1}$ given by $R \mapsto g(R)$. \square

3.11. $\aleph_{\alpha+1} < 2^{\aleph_\alpha}$.

[Use Exercises 3.10 and 3.9.]

Proof. By exercises 3.10 and 3.9, $\aleph_{\alpha+1} \leq 2^{\aleph_\alpha}$, and by Cantor's theorem, $\aleph_{\alpha+1} < 2^{2^{\aleph_\alpha}}$. \square

3.12. If \aleph_α is an uncountable limit cardinal, then $\text{cf } \omega_\alpha = \text{cf } \alpha$; ω_α is the limit of a cofinal sequence $\langle \omega_\xi : \xi < \text{cf } \alpha \rangle$ of cardinals.

Proof. $\text{cf } \omega_\alpha = \text{cf } \text{cf } \alpha = \text{cf } \alpha$. \square

3.13 (ZF). Show that ω_2 is not a countable union of countable sets.

[Assume that $\omega_2 = \bigcup_{n < \omega} S_n$ with S_n countable and let α_n be the order-type of S_n . Then $\alpha = \sup_n \alpha_n \leq \omega_1$ and there is a mapping of $\omega \times \alpha$ onto ω_2 .]

Proof. We can assume that S_n is disjoint for each $n \leq \omega$. Then we have a one-to-one function of $\omega \times \alpha$ onto ω_2 given by $(n, \beta) \mapsto$ the β -th element of S_n if $\beta \in \alpha_n$ otherwise 0. Thus $\aleph_2 = |\omega_2| \leq |\omega \times \alpha| \leq \aleph_0 \cdot \aleph_1 = \aleph_1$. A contradiction. \square

A set S is *Dedekind-finite* (D-finite) if there is no one-to-one mapping of S onto a proper subset of S . Every finite set is D-finite. Using the Axiom of Choice, one proves that every infinite set is D-infinite, and so D-finiteness is the same as finiteness. Without the Axiom of Choice, however, one cannot prove that every D-finite set is finite.

The set \mathbb{N} of all natural numbers is D-infinite and hence every S such that $|S| \geq \aleph_0$, is D-infinite.

3.14. S is D-infinite if and only if S has a countable subset.

[If S is D-infinite, let $f : S \rightarrow X \subset S$ be one-to-one. Let $x_0 \in S - X$ and $x_{n+1} = f(x_n)$. Then $S \supset \{x_n : n < \omega\}$.]

Proof. If S is D-infinite, since f is one-to-one, for each m and n such that $0 \leq m < n < \omega$, $x_m \neq x_n$. Thus we have a countable set $X = \{x_n : n < \omega\} \subsetneq S$.

Conversely, if S has a countable subset $X = \{x_n : n < \omega\}$. We have a one-to-one mapping of S onto $S \setminus \{x_0\}$ given by $x \mapsto x$ if $x \notin X$; otherwise $x_n \mapsto x_{n+1}$. \square

3.15. (i) If A and B are D-finite, then $A \cup B$ and $A \times B$ are D-finite.

(ii) The set of all finite one-to-one sequences in a D-finite set is D-finite.

(iii) The union of a disjoint D-finite family of D-finite sets is D-finite.

Proof. (i) Suppose that $X \subset A \cup B$ is countable. Then since a subset of a countable set is at most countable, $X \cap A$ and $X \cap B$ are at most countable. Since $X = (X \cap A) \cup (X \cap B)$, and the union of a finite set of finite sets is finite, $X \cap A$ or $X \cap B$ are countable. Thus A or B are D-infinite. A contradiction. Suppose that $X = \{(x_i, y_i) : i < \omega\} \subset A \times B$ is countable. Consider $C = \{x \in A : (x, y) \in X \text{ for some } y\}$ and $D = \{y \in B : (x, y) \in X \text{ for some } x\}$. Since $\aleph_0 = |X| \leq |C| \times |D|$, $|C| \leq |X| = \aleph_0$, and, $|D| \leq |X| = \aleph_0$, C or D are countable. But $C \subset A$ and $D \subset B$, a contradiction.

(ii) Let A be a D-finite set; let $X = \{X_i : i < \omega\}$ be a subset of all finite one-to-one sequences in A . Suppose that X is countable. Consider the cardinality of $S = \bigcup_{i < \omega} \text{ran}(X_i)$. Since X_i is a finite one-to-one sequence for all $i < \omega$, $|S| \leq |\omega| \cdot |\omega| = \aleph_0$. So suppose that $|S| = n < \omega$. then $X_i \in \bigcup \{S^1, S^2, \dots, S^n\}$ for all $i < \omega \Leftrightarrow X \subset \bigcup \{S^1, S^2, \dots, S^n\}$. By (i), and induction of n , the union of a finite family of D-finite sets is D-finite, and a finite product of D-finite sets is D-finite, thus $\bigcup \{S, S^2, \dots, S^n\}$ is D-finite. But $X \subset \bigcup \{S, S^2, \dots, S^n\}$, a contradiction. Thus S is countable. But $S \subset A$, also a contradiction.

(iii) Let $X = \bigcup_{i \in I} X_i$ for some D-finite I be a union of a disjoint D-finite family of D-finite sets. Suppose that $S = \{\alpha_n : n < \omega\} \subset X$ is countable. Consider the

cardinality of $T = \{i \in I : i \text{ such that } \alpha \in X_i \text{ for some } \alpha \in S\}$. Since X_i is disjoint for each $i \in I$, $|T| \leq |S| = \aleph_0$. Now suppose that $|T| = n < \omega$. Then S is a union of a finite family of D-finite set, thus finite; a contradiction. So T is countable. But $T \subset I$, also a contradiction. \square

On the other hand, one cannot prove without the Axiom of Choice that a projection, power set, or the set of all finite subsets of a D-finite set is D-finite, or that the union of a D-finite family of D-finite sets is D-finite.

3.16. If A is an infinite set, then $PP(A)$ is D-infinite.

[Consider the set $\{\{X \subset A : |X| = n\} : n < \omega\}$.]

Proof. The set $\{\{X \subset A : |X| = n\} : n < \omega\} \subset PP(A)$ is countable. \square

4. Real Numbers

4.1. The set of all continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ has cardinality \mathfrak{c} (while the set of all functions has cardinality $2^{\mathfrak{c}}$).

[A continuous function on \mathbb{R} is determined by its values at rational points.]

4.2. There are at least \mathfrak{c} countable order-types of linearly ordered sets.

[For every sequence $a = \langle a_n : n \in \mathbb{N} \rangle$ of natural numbers consider the order-type

$$\tau_a = a_0 + \xi + a_1 + \xi + a_2 + \dots$$

where ξ is the order-type of the integers. Show that if $a \neq b$, then $\tau_a \neq \tau_b$.]

A real number is algebraic if it is a root of a polynomial whose coefficients are integers. Otherwise, it is transcendental.

4.3. The set of all algebraic reals is countable.

4.4. If S is a countable set of reals, then $|\mathbb{R} - S| = \mathfrak{c}$.

[Use $\mathbb{R} \times \mathbb{R}$ rather than \mathbb{R} (because $|\mathbb{R} \times \mathbb{R}| = 2^{\aleph_0}$).]

4.5. (i) The set of all irrational numbers has cardinality \mathfrak{c} .

(ii) The set of all transcendental numbers has cardinality \mathfrak{c} .

4.6. The set of all open sets of reals has cardinality \mathfrak{c} .

4.7. The Cantor set is perfect.

4.8. If P is a perfect set and (a, b) is an open interval such that $P \cap (a, b) \neq \emptyset$, then $|P \cap (a, b)| = \mathfrak{c}$.

4.9. If $P_2 \not\subset P_1$ are perfect sets, then $|P_2 - P_1| = \mathfrak{c}$.

[Use Exercise 4.8.]

If A is a set of reals, a real number a is called a *condensation point* of A if every neighborhood of a contains uncountably many elements of A . Let A^* denote the set of all condensation points of A .

4.10. If P is perfect then $P^* = P$.

[Use Exercise 4.8.]

4.11. If F is closed and $P \subset F$ is perfect, then $P \subset F^*$.

[$P = P^* \subset F^*$.]

4.12. If F is an uncountable closed set and P is the perfect set constructed in Theorem 4.6, then $F^* \subset P$; thus $F^* = P$.

[Every $a \in F^*$ is a limit point of P since $|F - P| \leq \aleph_0$.]

4.13. If F is an uncountable closed set, then $F = F^* \cup (F - F^*)$ is the unique partition of F into a perfect set and an at most countable set.

[Use Exercise 4.9.]

4.14. \mathbb{Q} is not the intersection of a countable collection of open sets.

[Use the Baire Category Theorem.]

4.15. If B is Borel and f is a continuous function then $f_{-1}(B)$ is Borel.

4.16. Let $f : \mathbb{R} \rightarrow \mathbb{R}$. Show that the set of all x at which f is continuous is a G_δ set.

4.17. (i) $\mathbb{N} \times \mathbb{N}$ is homeomorphic to \mathbb{N} .

(ii) \mathbb{N}^ω is homeomorphic to \mathbb{N} .

4.18. The tree T_F in (4.6) has no maximal node, i.e., $s \in T$ such that there is no $t \in T$ with $s \subset t$. The map $F \mapsto T_F$ is a one-to-one correspondence between closed sets in \mathbb{N} and sequential trees without maximal nodes.

4.19. Every perfect Polish space has a closed subset homeomorphic to the Cantor space.

4.20. Every Polish space is homeomorphic to a G_δ subspace of the Hilbert cube.

[Let $\{x_n : n \in \mathbb{N}\}$ be a dense set, and define $f(x) = \langle d(x, x_n) : n \in \mathbb{N} \rangle$.]