

Contents

Chapter 1

Real Analysis

(1.5)

$$\text{OrderTrichotomy}[\prec, S] := \forall_{x,y \in S} (x \prec y \vee x = y \vee y \prec x)$$

$$\text{OrderTransitivity}[\prec, S] := \forall_{x,y,z \in S} ((x \prec y \wedge y \prec z) \implies x \prec z)$$

$$\text{Order}[\prec, S] := (\text{OrderTrichotomy}[\prec, S]) \wedge (\text{OrderTransitivity}[\prec, S])$$

(1.7)

$$\text{BoundedAbove}[E, S, \prec] := (\text{Order}[\prec, S]) \wedge (E \subset S) \wedge \left(\exists_{\beta \in S} \forall_{x \in E} (x \leq \beta) \right)$$

$$\text{BoundedBelow}[E, S, \prec] := (\text{Order}[\prec, S]) \wedge (E \subset S) \wedge \left(\exists_{\beta \in S} \forall_{x \in E} (\beta \leq x) \right)$$

$$\text{UpperBound}[\beta, E, S, \prec] := (\text{Order}[\prec, S]) \wedge (E \subset S) \wedge (\beta \in S \wedge \forall_{x \in E} (x \leq \beta))$$

$$\text{LowerBound}[\beta, E, S, \prec] := (\text{Order}[\prec, S]) \wedge (E \subset S) \wedge (\beta \in S \wedge \forall_{x \in E} (\beta \leq x))$$

(1.8)

$$\text{LUB}[\alpha, E, S, \prec] := (\text{UpperBound}[\alpha, E, S, \prec]) \wedge \left(\forall_{\gamma} (\gamma \prec \alpha \implies \neg \text{UpperBound}[\gamma, E, S, \prec]) \right)$$

$$\text{GLB}[\alpha, E, S, \prec] := (\text{LowerBound}[\alpha, E, S, \prec]) \wedge \left(\forall_{\beta} (\alpha \prec \beta \implies \neg \text{LowerBound}[\beta, E, S, \prec]) \right)$$

(1.10)

$$\text{LUBProperty}[S, \prec] := \forall_E \left(((\emptyset \neq E \subset S) \wedge (\text{BoundedAbove}[E, S, \prec]) \implies \exists_{\alpha \in S} (\text{LUB}[\alpha, E, S, \prec])) \right)$$

$$\text{GLBProperty}[S, \prec] := \forall_E \left(((\emptyset \neq E \subset S) \wedge (\text{BoundedBelow}[E, S, \prec]) \implies \exists_{\alpha \in S} (\text{GLB}[\alpha, E, S, \prec])) \right)$$

(1.11)

$$\text{LUBPropertyImpliesGLBProperty} := \text{LUBProperty}[S, \prec] \implies \text{GLBProperty}[S, \prec]$$

$$(1) \quad \text{LUBProperty}[S, \prec] \implies \dots$$

wts: 2

$$(1.1) \quad (\emptyset \neq B \subset S \wedge \text{BoundedBelow}[B, S, \prec]) \implies \dots$$

wts: 1.2

$$(1.1.1) \quad \text{Order}[\prec, S] \wedge \exists_{\delta' \in S} (\text{LowerBound}[\delta', B, S, \prec])$$

from: [BoundedBelow](#), 1.1

$$(1.1.2) \quad |B| = 1 \implies \dots$$

wts: 1.1.3

$$(1.1.2.1) \quad \exists_{u'} (u' \in B) \quad \blacksquare \quad u := \text{choice}(\{u' : u' \in B\}) \quad \blacksquare \quad B = \{u\}$$

from: 1.1.2

$$(1.1.2.2) \quad \text{GLB}[u, B, S, \prec] \quad \blacksquare \quad \exists_{\epsilon_0 \in S} (\text{GLB}[\epsilon_0, B, S, \prec])$$

$$(1.1.3) \quad |B| = 1 \implies \exists_{\epsilon_0 \in S} (\text{GLB}[\epsilon_0, B, S, \prec])$$

$$(1.1.4) \quad |B| \neq 1 \implies \dots$$

wts: 1.1.5

$$(1.1.4.1) \quad \forall_E ((\emptyset \neq E \subset S \wedge \text{BoundedAbove}[E, S, \prec]) \implies \exists_{\alpha \in S} (\text{LUB}[\alpha, E, S, \prec]))$$

from: [LUBProperty](#), 1

$$(1.1.4.2) \quad L := \{s \in S : \text{LowerBound}[s, B, S, \prec]\}$$

$$(1.1.4.3) \quad |B| > 1 \wedge \text{OrderTrichotomy}[\prec, S] \quad \blacksquare \quad \exists_{b_1' \in B} \exists_{b_0' \in B} (b_0' < b_1')$$

from: [Order](#), 1.1.1
wts: 1.1.4.7

$$(1.1.4.4) \quad b_1 := \text{choice}(\{b_1' \in B : \exists_{b_0' \in B} (b_0' < b_1')\}) \quad \blacksquare \quad \neg \text{LowerBound}[b_1, B, S, \prec]$$

from: 1.1.4.2

$$(1.1.4.5) \quad b_1 \notin L \quad \blacksquare \quad L \subset S$$

$$(1.1.4.6) \quad \delta := \text{choice}(\{\delta' \in S : \text{LowerBound}[\delta', B, S, \prec]\}) \quad \blacksquare \quad \delta \in L \quad \blacksquare \quad \emptyset \neq L$$

from: 1.1.1

$$(1.1.4.7) \quad \emptyset \neq L \subset S$$

from: 1.1.4.5, 1.1.4.6

$$(1.1.4.8) \quad \forall_{y \in L} (\text{LowerBound}[y_0, B, S, \prec]) \quad \blacksquare \quad \forall_{y \in L} \forall_{x \in B} (y_0 \leq x)$$

from: [LowerBound](#), 1.1.4.2
wts: 1.1.4.10

$$(1.1.4.9) \quad \forall_{x \in B} \left(x \in S \wedge \forall_{y \in L} (y_0 \leq x) \right) \quad \blacksquare \quad \forall_{x \in B} (\text{UpperBound}[x, L, S, <])$$

from: *UpperBound*

$$(1.1.4.10) \quad \exists_{x \in S} (\text{UpperBound}[x, L, S, <]) \quad \blacksquare \quad \text{BoundedAbove}[L, S, <]$$

$$(1.1.4.11) \quad \emptyset \neq L \subset S \wedge \text{BoundedAbove}[L, S, <]$$

from: 1.1.4.7, 1.1.4.10

$$(1.1.4.12) \quad \exists_{\alpha' \in S} (\text{LUB}[\alpha', L, S, <]) \quad \blacksquare \quad \alpha := \text{choice}(\{\alpha' \in S : (\text{LUB}[\alpha', L, S, <])\})$$

from: 1.1.4.1
wts: 1.1.4.21

$$(1.1.4.13) \quad \forall_x (x \in B \implies \text{UpperBound}[x, L, S, <])$$

from: 1.1.4.9
wts: 1.1.4.17

$$(1.1.4.14) \quad \forall_x (\neg \text{UpperBound}[x, L, S, <] \implies x \notin B)$$

$$(1.1.4.15) \quad \gamma < \alpha \implies \dots$$

wts: 1.1.4.16

$$(1.1.4.15.1) \quad \neg \text{UpperBound}[\gamma, L, S, <] \quad \blacksquare \quad \gamma \notin B$$

from: *LUB*, 1.1.4.12, 1.1.4.14

$$(1.1.4.16) \quad \gamma < \alpha \implies \gamma \notin B \quad \blacksquare \quad \gamma \in B \implies \gamma \geq \alpha$$

$$(1.1.4.17) \quad \forall_{\gamma \in B} (\alpha \leq \gamma) \quad \blacksquare \quad \text{LowerBound}[\alpha, B, S, <]$$

from: *LowerBound*

$$(1.1.4.18) \quad \alpha < \beta \implies \dots$$

wts: 1.1.4.19

$$(1.1.4.18.1) \quad \forall_{y \in L} (y_0 \leq \alpha < \beta) \quad \blacksquare \quad \forall_{y \in L} (y_0 \neq \beta)$$

from: *LUB*, 1.1.4.12, 1.1.4.18

$$(1.1.4.18.2) \quad \beta \notin L \quad \blacksquare \quad \neg \text{LowerBound}[\beta, B, S, <]$$

from: 1.1.4.2

$$(1.1.4.19) \quad \alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <] \quad \blacksquare \quad \forall_{\beta \in S} (\alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <])$$

$$(1.1.4.20) \quad \text{LowerBound}[\alpha, B, S, <] \wedge \forall_{\beta \in S} (\alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <])$$

from: 1.1.4.17, 1.1.4.19

$$(1.1.4.21) \quad \text{GLB}[\alpha, B, S, <] \quad \blacksquare \quad \exists_{\epsilon_1 \in S} (\text{GLB}[\epsilon_1, B, S, <])$$

$$(1.1.5) \quad |B| \neq 1 \implies \exists_{\epsilon_1 \in S} (\text{GLB}[\epsilon_1, B, S, <])$$

$$(1.1.6) \quad \left(|B| = 1 \implies \exists_{\epsilon_0 \in S} (\text{GLB}[\epsilon_0, B, S, <]) \right) \wedge \left(|B| \neq 1 \implies \exists_{\epsilon_1 \in S} (\text{GLB}[\epsilon_1, B, S, <]) \right)$$

from: 1.1.3, 1.1.5

$$(1.1.7) \quad (|B| = 1 \vee |B| \neq 1) \implies \exists_{\epsilon \in S} (\text{GLB}[\epsilon, B, S, <]) \quad \blacksquare \quad \exists_{\epsilon \in S} (\text{GLB}[\epsilon, B, S, <])$$

$$(1.2) \quad (\emptyset \neq B \subset S \wedge \text{BoundedBelow}[B, S, <]) \implies \exists_{\epsilon \in S} (\text{GLB}[\epsilon, B, S, <])$$

$$(1.3) \quad \forall_B ((\emptyset \neq B \subset S \wedge \text{BoundedBelow}[B, S, <]) \implies \exists_{\epsilon \in S} (\text{GLB}[\epsilon, B, S, <]))$$

$$(1.4) \quad \text{GLBProperty}[S, <]$$

$$(2) \quad \text{LUBProperty}[S, <] \implies \text{GLBProperty}[S, <]$$

$$(1.12)$$

$$\text{Field}[F, +, *] := \exists_{0, 1 \in F} \forall_{x, y, z \in F} \left(\begin{array}{l} x + y \in F \quad \wedge \quad x * y \in F \quad \wedge \\ x + y = y + x \quad \wedge \quad x * y = y * x \quad \wedge \\ (x + y) + z = x + (y + z) \quad \wedge \quad (x * y) * z = x * (y * z) \quad \wedge \\ 1 \neq 0 \quad \wedge \quad x * (y + z) = (x * y) + (x * z) \quad \wedge \\ 0 + x = x \quad \wedge \quad 1 * x = x \quad \wedge \\ \exists_{-x \in F} (x + (-x) = 0) \wedge \left(x \neq 0 \implies \exists_{1/x \in F} (x * (1/x) = 1) \right) \end{array} \right)$$

$$\text{***** (Field}[F, +, *] \wedge x, y, z \in F) \implies \dots \text{*****}$$

$$(1.14)$$

$$\text{AdditiveCancellation} := (x + y = x + z) \implies y = z$$

$$(1) \quad y = 0 + y = (x + (-x)) + y = ((-x) + x) + y = (-x) + (x + y) = \dots$$

from: *Field*

$$(2) \quad (-x) + (x + z) = ((-x) + x) + z = (x + (-x)) + z = 0 + z = z$$

from: *Field*

$$\text{AdditiveIdentityUniqueness} := (x + y = x) \implies y = 0$$

$$(1) \quad x + y = x = 0 + x = x + 0$$

from: *Field*

$$(2) \quad y = 0$$

from: *AdditiveCancellation*

$$\text{AdditiveInverseUniqueness} := (x + y = 0) \implies y = -x$$

$$(1) \quad x + y = 0 = x + (-x)$$

from: *Field*

$$(2) \quad y = -x$$

from: *AdditiveCancellation*

$$\text{DoubleNegative} := x = -(-x)$$

$$(1) \quad 0 = x + (-x) = (-x) + x \quad \blacksquare \quad 0 = (-x) + x$$

from: *Field*

$$(2) \quad x = -(-x)$$

from: [AdditiveInverseUniqueness](#)

(1.15)

$$\textcolor{red}{\textit{MultiplicativeCancellation}} := (x \neq 0 \wedge x * y = x * z) \implies y = z \quad \text{---}$$

$$\textcolor{red}{\textit{MultiplicativeIdentityUniqueness}} := (x \neq 0 \wedge x * y = x) \implies y = 1 \quad \text{---}$$

$$\textcolor{red}{\textit{MultiplicativeInverseUniqueness}} := (x \neq 0 \wedge x * y = 1) \implies y = 1/x \quad \text{---}$$

$$\textcolor{red}{\textit{DoubleReciprocal}} := (x \neq 0) \implies x = 1/(1/x) \quad \text{---}$$

(1.16)

$$\textcolor{red}{\textit{Domination}} := 0 * x = 0$$

from: [Field](#)

$$(1) \quad 0 * x = (0 + 0) * x = 0 * x + 0 * x \quad \blacksquare \quad 0 * x = 0 * x + 0 * x$$

$$(2) \quad 0 * x = 0$$

from: [AdditiveIdentityUniqueness](#)

$$\textcolor{red}{\textit{NonDomination}} := (x \neq 0 \wedge y \neq 0) \implies x * y \neq 0$$

$$(1) \quad (x \neq 0 \wedge y \neq 0) \implies \dots$$

$$(1.1) \quad (x * y = 0) \implies \dots$$

$$(1.1.1) \quad 1 = 1 * 1 = (x * (1/x)) * (y * (1/y)) = (x * y) * ((1/x) * (1/y)) = 0 * ((1/x) * (1/y)) = 0$$

from: [Field](#), [Domination](#), 1, 1.1

$$(1.1.2) \quad 1 = 0 \wedge 1 \neq 0 \quad \blacksquare \quad \perp$$

from: [Field](#)

$$(1.2) \quad (x * y = 0) \implies \perp \quad \blacksquare \quad x * y \neq 0$$

$$(2) \quad (x \neq 0 \wedge y \neq 0) \implies x * y \neq 0$$

$$\textcolor{red}{\textit{NegationCommutativity}} := (-x) * y = -(x * y) = x * (-y)$$

$$(1) \quad x * y + (-x) * y = (x + -x) * y = 0 * y = 0 \quad \blacksquare \quad x * y + (-x) * y = 0$$

from: [Field](#), [Domination](#)
wts: 2

$$(2) \quad (-x) * y = -(x * y)$$

from: [AdditiveInverseUniqueness](#)

$$(3) \quad x * y + x * (-y) = x * (y_0 + -y) = x * 0 = 0 \quad \blacksquare \quad x * y + x * (-y) = 0$$

from: [Field](#), [Domination](#)
wts: 4

$$(4) \quad x * (-y) = -(x * y)$$

from: [AdditiveInverseUniqueness](#)

$$(5) \quad (-x) * y = -(x * y) = x * (-y)$$

from: 2, 4

$$\textcolor{red}{\textit{NegativeMultiplication}} := (-x) * (-y) = x * y$$

$$(1) \quad (-x) * (-y) = -(x * (-y)) = -(-(x * y)) = x * y$$

from: [NegationCommutativity](#), [DoubleNegative](#)

(1.17)

$$\textcolor{red}{\textit{OrderedField}}[F, +, *, <] := \left(\begin{array}{l} \textcolor{blue}{\textit{Field}}[F, +, *] \quad \wedge \quad \textcolor{blue}{\textit{Order}}[<, F] \quad \wedge \\ \forall_{x,y,z \in F} (y_0 < z \implies x + y < x + z) \quad \wedge \\ \forall_{x,y \in F} ((x > 0 \wedge y > 0) \implies x * y > 0) \end{array} \right)$$

$$\textcolor{blue}{\textit{OrderedField}}[F, +, *, <] \wedge x, y, z \in F \implies \dots$$

(1.18)

$$\textcolor{red}{\textit{NegationOnOrder}} := x > 0 \iff -x < 0$$

$$(1) \quad x > 0 \implies \dots$$

$$(1.1) \quad 0 = (-x) + x > (-x) + 0 = -x \quad \blacksquare \quad 0 > -x \quad \blacksquare \quad -x < 0$$

from: [OrderedField](#)

$$(2) \quad x > 0 \implies -x < 0$$

$$(3) \quad -x < 0 \implies \dots$$

$$(3.1) \quad 0 = x + (-x) < x + 0 = x \quad \blacksquare \quad 0 < x \quad \blacksquare \quad x > 0$$

from: [OrderedField](#)

$$(4) \quad -x < 0 \implies x > 0$$

$$(5) \quad x > 0 \implies -x < 0 \wedge -x < 0 \implies x > 0 \quad \blacksquare \quad x > 0 \iff -x < 0$$

from: 2, 4

$$\textcolor{red}{\textit{PositiveFactorPreservesOrder}} := (x > 0 \wedge y < z) \implies x * y < x * z$$

$$(1) \quad (x > 0 \wedge y < z) \implies \dots$$

$$(1.1) \quad (-y) + z > (-y) + y = 0 \quad \blacksquare \quad z + (-y) = 0$$

from: [OrderedField](#)

$$(1.2) \quad x * (z + (-y)) > 0 \quad \blacksquare \quad x * z + x * (-y) > 0$$

from: [OrderedField](#)

$$(1.3) \quad x * z = 0 + x * z = (x * y + -(x * y)) + x * z = (x * y + x * (-y)) + x * z = \dots$$

from: *Field, NegationCommutativity*

$$(1.4) \quad x * y + (x * z + x * (-y)) > x * y + 0 = x * y$$

from: *Field, 1.2*

$$(1.5) \quad x * z > x * y$$

from: 1.3, 1.4

$$(2) \quad (x > 0 \wedge y < z) \implies x * z > x * y$$

$$\textcolor{red}{NegativeFactorFlipsOrder} := (x < 0 \wedge y < z) \implies x * y > x * z$$

$$(1) \quad (x < 0 \wedge y < z) \implies \dots$$

$$(1.1) \quad -x > 0$$

from: *NegationOnOrder*

$$(1.2) \quad (-x) * y < (-x) * z \quad \blacksquare \quad 0 = x * y + (-x) * y < x * y + (-x) * z \quad \blacksquare \quad 0 < x * y + (-x) * z$$

from: *PositiveFactorPreservesOrder*

$$(1.3) \quad 0 < (-x) * (-y + z) \quad \blacksquare \quad 0 > x * (-y + z) \quad \blacksquare \quad 0 > -(x * y) + x * z$$

from: *NegationOnOrder*

$$(1.4) \quad x * y > x * z$$

$$(2) \quad (x < 0 \wedge y < z) \implies x * y > x * z$$

$$\textcolor{red}{SquareIsPositive} := (x \neq 0) \implies x * x > 0$$

$$(1) \quad (x > 0) \implies x * x > 0$$

from: *OrderedField*

$$(2) \quad (x < 0) \implies \dots$$

$$(2.1) \quad -x > 0 \quad \blacksquare \quad x * x = (-x) * (-x) > 0 \quad \blacksquare \quad x * x > 0$$

from: *NegationOnOrder, OrderedField, NegativeMultiplication*

$$(3) \quad (x < 0) \implies x * x > 0$$

$$(4) \quad x \neq 0 \implies (x > 0 \vee x < 0) \implies x * x > 0 \quad \blacksquare \quad x \neq 0 \implies x * x > 0$$

from: *OrderTrichotomy, 1.3*

$$\textcolor{red}{OneIsPositive} := 1 > 0$$

$$(1) \quad 1 \neq 0 \quad \blacksquare \quad 1 = 1 * 1 > 0$$

from: *Field, SquareIsPositive*

$$\textcolor{red}{ReciprocationOnOrder} := (0 < x < y) \implies 0 < 1/y < 1/x$$

$$(1) \quad (0 < x < y) \implies \dots$$

$$(1.1) \quad x * (1/x) = 1 > 0 \quad \blacksquare \quad x * (1/x) > 0$$

from: *Field, OneIsPositive*

$$(1.2) \quad 1/x < 0 \implies x * (1/x) < 0 \wedge x * (1/x) > 0 \implies \perp \quad \blacksquare \quad 1/x > 0$$

from: *NegativeFactorFlipsOrder, 1*

$$(1.3) \quad y * (1/y) = 1 > 0 \quad \blacksquare \quad y * (1/y) > 0$$

from: *Field, OneIsPositive*

$$(1.4) \quad 1/y < 0 \implies y * (1/y) < 0 \wedge y * (1/y) > 0 \implies \perp \quad \blacksquare \quad 1/y > 0$$

from: *NegativeFactorFlipsOrder, 1*

$$(1.5) \quad (1/x) * (1/y) > 0$$

from: *OrderedField*

$$(1.6) \quad 0 < 1/y = ((1/x) * (1/y)) * x < ((1/x) * (1/y)) * y = 1/x$$

from: *OrderedField, 1, 1.4, 1.5*

$$(1.19)$$

$$\textcolor{red}{OrderedFieldQ} := \textcolor{teal}{OrderedField}[\mathbb{Q}, +, *, <] \quad \text{---}$$

$$\textcolor{red}{Subfield}[K, F, +, *] := \textcolor{teal}{Field}[F, +, *] \wedge K \subset F \wedge \textcolor{teal}{Field}[K, +, *]$$

$$\textcolor{red}{OrderedSubfield}[K, F, +, *, <] := \textcolor{teal}{OrderedField}[F, +, *, <] \wedge K \subset F \wedge \textcolor{teal}{OrderedField}[K, +, *, <]$$

$$\textcolor{red}{CutI}[\alpha] := \emptyset \neq \alpha \subset \mathbb{Q}$$

$$\textcolor{red}{CutII}[\alpha] := \forall_{p \in \alpha} \forall_{q \in \mathbb{Q}} (q < p \implies q \in \alpha)$$

$$\textcolor{red}{CutIII}[\alpha] := \forall_{p \in \alpha} \exists_{r \in \alpha} (p < r)$$

$$\mathbb{R} := \{\alpha \in \mathbb{Q} : \textcolor{teal}{CutI}[\alpha] \wedge \textcolor{teal}{CutII}[\alpha] \wedge \textcolor{teal}{CutIII}[\alpha]\}$$

$$\textcolor{red}{CutCorollaryI} := (\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies p < q$$

$$(1) \quad (\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies \dots$$

$$(1.1) \quad \forall_{p' \in \alpha} \forall_{q' \in \mathbb{Q}} (q' < p' \implies q' \in \alpha)$$

from: *CutII, 1*

$$(1.2) \quad q < p \implies q \in \alpha \quad \blacksquare \quad q \notin \alpha \implies q \geq p$$

from: 1

$$(1.3) \quad (q \notin \alpha) \implies \dots$$

$$(1.3.1) \quad q \geq p$$

from: 1.2

$$(1.3.2) \quad (q = p) \implies (p \in \alpha \wedge p \notin \alpha) \implies \perp \quad \blacksquare \quad q \neq p$$

from: 1, 1.3

(1.3.3)	$q \geq p \wedge q \neq p \blacksquare p < q$	
(1.4)	$q \notin \alpha \implies p < q \blacksquare p < q$	from: 1
(2)	$(\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies p < q$	

CutCorollaryI $:= (\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies s \notin \alpha$

(1)	$(\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies \dots$	
(1.1)	$\forall_{s' \in \alpha} \forall_{r' \in \mathbb{Q}} (r' < s' \implies r' \in \alpha)$	from: CutII, 1
(1.2)	$s \in \alpha \implies (r \in \mathbb{Q} \implies (r < s \implies r \in \alpha)) \blacksquare s \in \alpha \implies r \in \alpha$	from: 1, 1.1
(1.3)	$r \notin \alpha \implies s \notin \alpha \blacksquare s \notin \alpha$	from: 1, 1.2
(2)	$(\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies s \notin \alpha$	

$<_{\mathbb{R}}[\alpha, \beta] := \alpha, \beta \in \mathbb{R} \wedge \alpha \subset \beta$

OrderTrichotomyOfR $:= \text{OrderTrichotomy}[\mathbb{R}, <_{\mathbb{R}}]$

(1)	$(\alpha, \beta \in \mathbb{R}) \implies \dots$	
(1.1)	$\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \implies \dots$	
(1.1.1)	$\alpha \not\subset \beta \wedge \alpha \neq \beta$	from: $<_{\mathbb{R}}$, 1.1
(1.1.2)	$\exists_{p'} (p' \in \alpha \wedge p' \notin \beta) \blacksquare p := \text{choice}(\{p' : p' \in \alpha \wedge p' \notin \beta\})$	
(1.1.3)	$q \in \beta \implies \dots$	
(1.1.3.1)	$p, q \in \mathbb{Q}$	
(1.1.3.2)	$q < p$	from: CutCorollaryI
(1.1.3.3)	$q \in \alpha$	from: CutII
(1.1.4)	$q \in \beta \implies q \in \alpha$	
(1.1.5)	$\forall_{q \in \beta} (q \in \alpha) \blacksquare \beta \subseteq \alpha$	
(1.1.6)	$\beta \subset \alpha \blacksquare \beta <_{\mathbb{R}} \alpha$	
(1.2)	$\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \implies \beta <_{\mathbb{R}} \alpha$	
(1.3)	$\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \vee (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \blacksquare (\beta <_{\mathbb{R}} \alpha) \vee (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta)$	
(1.4)	$\alpha = \beta \implies \neg(\alpha <_{\mathbb{R}} \beta \vee \beta <_{\mathbb{R}} \alpha)$	
(1.5)	$\alpha <_{\mathbb{R}} \beta \implies \neg(\alpha = \beta \vee \beta <_{\mathbb{R}} \alpha)$	
(1.6)	$\beta <_{\mathbb{R}} \alpha \implies \neg(\alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$	
(1.7)	$\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta$	
(2)	$(\alpha, \beta \in \mathbb{R}) \implies (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$	
(3)	$\forall_{\alpha, \beta \in \mathbb{R}} (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$	
(4)	OrderTrichotomy $[\mathbb{R}, <_{\mathbb{R}}]$	

OrderTransitivityOfR $:= \text{OrderTransitivity}[\mathbb{R}, <_{\mathbb{R}}]$

(1)	$(\alpha, \beta, \gamma \in \mathbb{R}) \implies \dots$	
(1.1)	$(\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \dots$	
(1.1.1)	$\alpha \subset \beta \wedge \beta \subset \gamma$	
(1.1.2)	$\forall_{a \in \alpha} (a \in \beta) \wedge \forall_{b \in \beta} (b \in \gamma)$	
(1.1.3)	$\forall_{a \in \alpha} (\alpha \in \gamma) \blacksquare \alpha \subset \gamma \blacksquare \alpha <_{\mathbb{R}} \gamma$	
(1.2)	$(\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma$	
(2)	$(\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma)$	
(3)	$\forall_{\alpha, \beta, \gamma \in \mathbb{R}} ((\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma)$	
(4)	OrderTransitivity $[\mathbb{R}, <_{\mathbb{R}}]$	

OrderOfR $:= \text{Order}[<_{\mathbb{R}}, \mathbb{R}]$

LUBPropertyOfR $:= \text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}]$

(1)	$(\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \dots$	from: OrderTrichotomyR , OrderTransitivityR wts:
(1.1)	$\gamma := \{p \in \mathbb{Q} : \exists_{\alpha \in A} (p \in \alpha)\}$	

(1.2)	$A \neq \emptyset \quad \blacksquare \quad \exists_{\alpha}(\alpha \in A) \quad \blacksquare \quad \alpha_0 := \text{choice}(\{\alpha : \alpha \in A\})$
(1.3)	$\alpha_0 \neq \emptyset \quad \blacksquare \quad \exists_a(a \in \alpha_0) \quad \blacksquare \quad a_0 := \text{choice}(\{a : a \in \alpha_0\}) \quad \blacksquare \quad a_0 \in \gamma \quad \blacksquare \quad \gamma \neq \emptyset$
(1.4)	$\text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}] \quad \blacksquare \quad \exists_{\beta}(\text{UpperBound}[\beta, A, \mathbb{R}, <_{\mathbb{R}}])$
(1.5)	$\beta_0 := \text{choice}(\{\beta : \text{UpperBound}[\beta, A, \mathbb{R}, <_{\mathbb{R}}]\})$
(1.6)	$\text{UpperBound}[\beta_0, A, \mathbb{R}, <_{\mathbb{R}}] \quad \blacksquare \quad \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \beta_0) \quad \blacksquare \quad \forall_{\alpha \in A}(\alpha \subseteq \beta_0) \quad \blacksquare \quad \forall_{\alpha \in A} \forall_{a \in \alpha}(a \in \beta_0)$
(1.7)	$(\alpha \in A \wedge a \in \alpha) \iff a \in \gamma \quad \blacksquare \quad \forall_{a \in \gamma}(a \in \beta_0) \quad \blacksquare \quad \gamma \subseteq \beta_0$
(1.8)	$\beta_0 \subset \mathbb{Q} \quad \blacksquare \quad \gamma \subseteq \beta_0 \subset \mathbb{Q} \quad \blacksquare \quad \gamma \subset \mathbb{Q}$
(1.9)	$\emptyset \neq \gamma \subset \mathbb{Q} \quad \blacksquare \quad \text{CutI}[\gamma]$
(1.10)	$(p \in \gamma \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$
(1.10.1)	$p \in \gamma \quad \blacksquare \quad \exists_{\alpha \in A}(p \in \alpha) \quad \blacksquare \quad \alpha_1 := \text{choice}(\{\alpha \in A : p \in \alpha\})$
(1.10.2)	$p \in \alpha_1 \wedge q \in \mathbb{Q} \wedge q < p \quad \blacksquare \quad q \in \alpha_1 \quad \blacksquare \quad q \in \gamma$
(1.11)	$(p \in \gamma \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \gamma \quad \blacksquare \quad \forall_{p \in \gamma} \forall_{q \in \mathbb{Q}}(q < p \implies q \in \gamma) \quad \blacksquare \quad \text{CutII}[\gamma]$
(1.12)	$p \in \gamma \implies \dots$
(1.12.1)	$\exists_{\alpha \in A}(p \in \alpha) \quad \blacksquare \quad \alpha_2 := \text{choice}(\{\alpha \in A : p \in \alpha\})$
(1.12.2)	$\alpha_2 \in \mathbb{R} \quad \blacksquare \quad \text{CutII}[\alpha_2] \quad \blacksquare \quad \exists_{r \in \alpha_2}(p < r) \quad \blacksquare \quad r_0 := \text{choice}(\{r \in \alpha_2 : p < r\})$
(1.12.3)	$r_0 \in \alpha_2 \quad \blacksquare \quad r_0 \in \gamma$
(1.12.4)	$p < r_0 \quad \blacksquare \quad p < r_0 \wedge r_0 \in \gamma \quad \blacksquare \quad \exists_{r \in \gamma}(p < r)$
(1.13)	$p \in \gamma \implies \exists_{r \in \gamma}(p < r) \quad \blacksquare \quad \forall_{p \in \gamma} \exists_{r \in \gamma}(p < r) \quad \blacksquare \quad \text{CutIII}[\gamma]$
(1.14)	$\text{CutI}[\gamma] \wedge \text{CutII}[\gamma] \wedge \text{CutIII}[\gamma] \quad \blacksquare \quad \gamma \in \mathbb{R}$
(1.15)	$\forall_{\alpha \in A}(\alpha \subseteq \gamma) \quad \blacksquare \quad \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \gamma)$
(1.16)	$\forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \gamma) \wedge \gamma \in \mathbb{R} \quad \blacksquare \quad \text{UpperBound}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}]$
(1.17)	$\delta <_{\mathbb{R}} \gamma \implies \dots$
(1.17.1)	$\delta \subset \gamma \quad \blacksquare \quad \exists_s(s \in \gamma \wedge s \notin \delta) \quad \blacksquare \quad s_0 := \text{choice}(\{s \in \mathbb{Q} : s \in \gamma \wedge s \notin \delta\})$
(1.17.2)	$s_0 \in \gamma \quad \blacksquare \quad \exists_{\alpha \in A}(s_0 \in \alpha) \quad \blacksquare \quad \alpha_3 := \text{choice}(\{\alpha \in A : s_0 \in \alpha\})$
(1.17.3)	$s_0 \in \alpha_3 \wedge s_0 \notin \delta \quad \blacksquare \quad \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta)$
(1.17.4)	$\delta \geq_{\mathbb{R}} \alpha_3 \implies \dots$
(1.17.4.1)	$\alpha_3 \subseteq \delta \quad \blacksquare \quad \forall_{s \in \mathbb{Q}}(s \in \alpha_3 \implies s \in \delta) \quad \blacksquare \quad \neg \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta)$
(1.17.4.2)	$\neg \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta) \wedge \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta) \quad \blacksquare \quad \perp$
(1.17.5)	$\delta \geq_{\mathbb{R}} \alpha_3 \implies \perp \quad \blacksquare \quad \delta <_{\mathbb{R}} \alpha_3 \quad \blacksquare \quad \exists_{\alpha \in A}(\delta <_{\mathbb{R}} \alpha) \quad \blacksquare \quad \exists_{\alpha \in A}(\neg(\alpha \leq_{\mathbb{R}} \delta))$
(1.17.6)	$\neg \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \delta) \quad \blacksquare \quad \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}]$
(1.18)	$\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}] \quad \blacksquare \quad \forall_{\delta}(\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}])$
(1.19)	$\text{UpperBound}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}] \wedge \forall_{\delta}(\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}])$
(1.20)	$\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}] \quad \blacksquare \quad \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}])$
(2)	$(\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}])$
(3)	$\forall_A \left((\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}]) \right) \quad \blacksquare \quad \text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}]$

$$+_{\mathbb{R}}[\alpha, \beta] := \alpha, \beta \in \mathbb{R} \wedge (\alpha +_{\mathbb{R}} \beta) = \{r + s : r \in \alpha \wedge s \in \beta\}$$

$$0_{\mathbb{R}} := \{x \in \mathbb{Q} : x < 0\}$$

$$\text{ZeroInR} := 0_{\mathbb{R}} \in \mathbb{R}$$

(1)	$-1 \in 0_{\mathbb{R}} \wedge 1 \notin 0_{\mathbb{R}} \quad \blacksquare \quad \emptyset \neq 0_{\mathbb{R}} \subseteq \mathbb{Q} \quad \blacksquare \quad \text{CutI}[0_{\mathbb{R}}]$
(2)	$(x \in 0_{\mathbb{R}} \wedge y \in \mathbb{Q} \wedge y < x) \implies y < x < 0 \implies y < 0 \implies y \in 0_{\mathbb{R}} \quad \blacksquare \quad \forall_{x \in 0_{\mathbb{R}}} \forall_{y \in \mathbb{Q}}(y_0 < x \implies y \in 0_{\mathbb{R}}) \quad \blacksquare \quad \text{CutII}[0_{\mathbb{R}}]$
(3)	$y := x/2 \quad \blacksquare \quad (x \in 0_{\mathbb{R}}) \implies (x < y < 0) \implies \exists_{y \in 0_{\mathbb{R}}}(x < y) \quad \blacksquare \quad \forall_{x \in 0_{\mathbb{R}}} \exists_{y \in 0_{\mathbb{R}}}(x < y) \quad \blacksquare \quad \text{CutIII}[0_{\mathbb{R}}]$
(4)	$\text{CutI}[0_{\mathbb{R}}] \wedge \text{CutII}[0_{\mathbb{R}}] \wedge \text{CutIII}[0_{\mathbb{R}}] \quad \blacksquare \quad 0_{\mathbb{R}} \in \mathbb{R}$

$$\text{FieldAdditionClosureOfR} := (\alpha, \beta \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) \in \mathbb{R})$$

(1)	$(\alpha, \beta \in \mathbb{R}) \implies \dots$
(1.1)	$(\alpha +_{\mathbb{R}} \beta) = \{r + s : r \in \alpha \wedge s \in \beta\}$
(1.2)	$\emptyset \neq \alpha \subset \mathbb{Q} \wedge \emptyset \neq \beta \subset \mathbb{Q}$

- (1.3) $\exists_a(a \in \alpha) ; \exists_b(b \in \beta) \blacksquare a_0 := choice(\{a : a \in \alpha\}) ; b_0 := choice(\{b : b \in \beta\}) \blacksquare a_0 + b_0 \in \alpha +_{\mathbb{R}} \beta$
- (1.4) $\exists_x(x \notin \alpha) ; \exists_y(y_0 \notin \beta) \blacksquare x_0 := choice(\{x : x \notin \alpha\}) ; y_0 := choice(\{y : y \notin \beta\})$
- (1.5) $\forall_{r \in \alpha}(r < x_0) ; \forall_{s \in \beta}(s < y_0) \blacksquare \forall_{r \in \alpha} \forall_{s \in \beta}(r + s < x_0 + y_0) \blacksquare x_0 + y_0 \notin \alpha +_{\mathbb{R}} \beta$
- (1.6) $\emptyset \neq \alpha +_{\mathbb{R}} \beta \subset \mathbb{Q} \blacksquare \textcolor{blue}{CutI}[\alpha +_{\mathbb{R}} \beta]$
- (1.7) $(p \in \alpha +_{\mathbb{R}} \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$
- (1.7.1) $\exists_{r \in \alpha} \exists_{s \in \beta}(p = r + s) \blacksquare (r_0, s_0) := choice((r, s) \in \alpha \times \beta : p = r + s)$
- (1.7.2) $q < p = r_0 + s_0 \blacksquare (q - s_0) < r_0 \blacksquare (q - s_0) \in \alpha$
- (1.7.3) $s_0 \in \beta \blacksquare q = (q - s_0) + s_0 \in \alpha +_{\mathbb{R}} \beta \blacksquare q \in \alpha +_{\mathbb{R}} \beta$
- (1.8) $(p \in \alpha +_{\mathbb{R}} \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \alpha +_{\mathbb{R}} \beta \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} \beta} \forall_{q \in \mathbb{Q}}(q < p \implies q \in \alpha +_{\mathbb{R}} \beta) \blacksquare \textcolor{blue}{CutII}[\alpha +_{\mathbb{R}} \beta]$
- (1.9) $p \in \alpha \implies \dots$
- (1.9.1) $\exists_{r \in \alpha} \exists_{s \in \beta}(p = r + s) \blacksquare (r_1, s_1) := choice(\{(r, s) \in \alpha \times \beta : p = r + s\})$
- (1.9.2) $r_1 \in \alpha \blacksquare \exists_{t \in \alpha}(r_1 < t) \blacksquare t_0 := choice(\{t \in \alpha : r_1 < t\})$
- (1.9.3) $s_1 \in \beta \blacksquare t + s_1 \in \alpha +_{\mathbb{R}} \beta \wedge p = r_1 + s_1 < t + s_1 \blacksquare \exists_{r \in \alpha +_{\mathbb{R}} \beta}(p < r)$
- (1.10) $p \in \alpha \implies \exists_{r \in \alpha +_{\mathbb{R}} \beta}(p < r) \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} \beta} \exists_{r \in \alpha +_{\mathbb{R}} \beta}(p < r) \blacksquare \textcolor{blue}{CutIII}[\alpha +_{\mathbb{R}} \beta]$
- (1.11) $\textcolor{blue}{CutI}[\alpha +_{\mathbb{R}} \beta] \wedge \textcolor{blue}{CutII}[\alpha +_{\mathbb{R}} \beta] \wedge \textcolor{blue}{CutIII}[\alpha +_{\mathbb{R}} \beta] \blacksquare \alpha +_{\mathbb{R}} \beta \in \mathbb{R}$
- (2) $(\alpha, \beta \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) \in \mathbb{R})$

FieldAdditionCommutativityOf \mathbb{R} $:= (\alpha, \beta \in \mathbb{R}) \implies (\alpha +_{\mathbb{R}} \beta = \beta +_{\mathbb{R}} \alpha)$

- (1) $\alpha +_{\mathbb{R}} \beta = \{r + s : r \in \alpha \wedge s \in \beta\} = \{s + r : s \in \beta \wedge r \in \alpha\} = \beta +_{\mathbb{R}} \alpha$

FieldAdditionAssociativityOf \mathbb{R} $:= (\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma))$

- (1) $(\alpha, \beta, \gamma \in \mathbb{R}) \implies \dots$
- (1.1) $(\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \{(a + b) + c : a \in \alpha \wedge b \in \beta \wedge c \in \gamma\} = \dots$
- (1.2) $\{a + (b + c) : a \in \alpha \wedge b \in \beta \wedge c \in \gamma\} = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma)$
- (2) $(\alpha, \beta, \gamma \in \mathbb{R}) \implies (\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma)$

FieldAdditionIdentityOf \mathbb{R} $:= (\alpha \in \mathbb{R}) \implies 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$

- (1) $\alpha \in \mathbb{R} \implies \dots$
- (1.1) $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \implies \dots$
- (1.1.1) $s < 0 \blacksquare r + s < r + 0 = r \blacksquare r + s < r \blacksquare r + s \in \alpha$
- (1.2) $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \implies r + s \in \alpha \blacksquare \forall_{r \in \alpha} \forall_{s \in 0_{\mathbb{R}}}(r + s \in \alpha)$
- (1.3) $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \iff (r + s \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}) \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}}(p \in \alpha) \blacksquare \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \subseteq \alpha$
- (1.4) $p \in \alpha \implies \dots$
- (1.4.1) $\exists_{r \in \alpha}(p < r) \blacksquare r_2 := choice(\{r \in \alpha : p < r\})$
- (1.4.2) $p < r_2 \blacksquare p - r_2 < r_2 - r_2 = 0 \blacksquare (p - r_2) < 0 \blacksquare (p - r_2) \in 0_{\mathbb{R}}$
- (1.4.3) $r_2 \in \alpha \blacksquare p = r_2 + (p - r_2) \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}$
- (1.5) $p \in \alpha \implies p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare \forall_{p \in \alpha}(p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}) \blacksquare \alpha \subseteq \alpha +_{\mathbb{R}} 0_{\mathbb{R}}$
- (1.6) $\alpha +_{\mathbb{R}} 0_{\mathbb{R}} \subseteq \alpha \wedge \alpha \subseteq \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$
- (2) $\alpha \in \mathbb{R} \implies 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$

FieldAdditionInverseOf \mathbb{R} $:= (\alpha \in \mathbb{R}) \implies \exists_{-\alpha \in \mathbb{R}}(\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$

- (1) $\alpha \in \mathbb{R} \implies \dots$
- (1.1) $\beta := \{p \in \mathbb{Q} : \exists_{r > 0}(-p - r \notin \alpha)\}$
- (1.2) $\alpha \subset \mathbb{Q} \blacksquare \exists_{s \in \mathbb{Q}}(s \notin \alpha) \blacksquare s_0 := choice(\{s : s \notin \alpha\}) \blacksquare p_0 := -s_0 - 1$
- (1.3) $-p_0 - 1 = -(-s_0 - 1) - 1 = s_0 \notin \alpha \blacksquare -p_0 - 1 \notin \alpha \blacksquare \exists_{r > 0}(-p_0 - r \notin \alpha) \blacksquare p_0 \in \beta$
- (1.4) $\emptyset \neq \alpha \blacksquare \exists_{q \in \alpha} \blacksquare q_0 := choice(\{q \in \mathbb{Q} : q \in \alpha\})$
- (1.5) $r > 0 \implies \dots$
- (1.5.1) $q_0 \in \alpha \blacksquare -(-q_0) - r = q_0 - r < q_0 \blacksquare -(-q_0) - r < q_0 \blacksquare -(-q_0) - r \in \alpha$
- (1.6) $\forall_{r > 0}(-(-q_0) - r \in \alpha) \blacksquare \neg \exists_{r > 0}(-(-q_0) - r \notin \alpha) \blacksquare -q_0 \notin \beta$
- (1.7) $\emptyset \neq \beta \subset \mathbb{Q} \blacksquare \textcolor{blue}{CutI}[\beta]$

(1.8)	$(p \in \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$	
(1.8.1)	$p \in \beta \blacksquare \exists_{r>0}(-p-r \notin \alpha) \blacksquare r_0 := \text{choice}(\{r > 0 : -p-r \notin \alpha\})$	
(1.8.2)	$q < p \blacksquare -p-r < -q-r$	
(1.8.3)	$-q-r \notin \alpha \blacksquare q \in \beta$	
(1.9)	$(p \in \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \beta \blacksquare \forall_{p \in \beta} \forall_{q \in \mathbb{Q}}(q < p \implies q \in \beta) \blacksquare \text{CutII}[\beta]$	
(1.10)	$p \in \beta \implies \dots$	
(1.10.1)	$p \in \beta \blacksquare \exists_{r>0}(-p-r \notin \alpha) \blacksquare r_1 := \text{choice}(\{r > 0 : -p-r \notin \alpha\})$	
(1.10.2)	$t_0 := p + (r_1/2)$	
(1.10.3)	$r_1 > 0 \blacksquare r_1/2 > 0$	
(1.10.4)	$t_0 > t_0 - (r_1/2) = p \blacksquare t_0 > p$	
(1.10.5)	$-t_0 - (r_1/2) = -(p + (r_1/2)) - (r_1/2) = -p - r_1$	
(1.10.6)	$-p - r_1 \notin \alpha \blacksquare -t_0 - (r_1/2) \notin \alpha \blacksquare \exists_{r>0}(-t_0 - r \notin \alpha) \blacksquare t_0 \in \beta$	
(1.10.7)	$t_0 > p \wedge t_0 \in \beta \blacksquare \exists_{t \in \beta}(p < t)$	
(1.11)	$p \in \beta \implies \exists_{t \in \beta}(p < t) \blacksquare \forall_{p \in \beta} \exists_{t \in \beta}(p < t) \blacksquare \text{CutIII}[\beta]$	
(1.12)	$\text{CutI}[\beta] \wedge \text{CutII}[\beta] \wedge \text{CutIII}[\beta] \blacksquare \beta \in \mathbb{R}$	
(1.13)	$(r \in \alpha \wedge s \in \beta) \implies \dots$	
(1.13.1)	$s \in \beta \blacksquare \exists_{t>0}(-s-t \notin \alpha) \blacksquare t_1 := \text{choice}(\{t > 0 : -s-t \notin \alpha\}) \blacksquare -s - t_1 < -s$	
(1.13.2)	$\alpha \in \mathbb{R} \wedge s, t_1 \in \mathbb{Q} \wedge -s - t_1 < -s \wedge -s - t_1 \notin \alpha \blacksquare -s \notin \alpha$	
(1.13.3)	$\alpha \in \mathbb{R} \wedge r \in \alpha \wedge -s \notin \alpha \blacksquare r < -s \blacksquare r + s < 0 \blacksquare r + s \in 0_{\mathbb{R}}$	
(1.14)	$(r \in \alpha \wedge s \in \beta) \implies r + s \in 0_{\mathbb{R}} \blacksquare \forall_{(r,s) \in \alpha \times \beta}(r + s \in 0_{\mathbb{R}}) \blacksquare \alpha +_{\mathbb{R}} \beta \subseteq 0_{\mathbb{R}}$	
(1.15)	$v \in 0_{\mathbb{R}} \implies \dots$	
(1.15.1)	$v < 0 \blacksquare w_0 := -v/2 \blacksquare w > 0$	
(1.15.2)	$\exists_{n \in \mathbb{Z}}(nw_0 \in \alpha \wedge (n+1)w_0 \notin \alpha) \blacksquare n_0 := \text{choice}(\{n \in \mathbb{Z} : nw_0 \in \alpha \wedge (n+1)w_0 \notin \alpha\})$	from: ARCHIMEDEANPROPERTYOFQ + LUB???
(1.15.3)	$p_0 := -(n_0 + 2)w_0 \blacksquare -p_0 - w_0 = (n_0 + 2)w_0 - w_0 = (n_0 + 1)w_0 \notin \alpha \blacksquare -p_0 - w_0 \notin \alpha \blacksquare p_0 \in \beta$	
(1.15.4)	$n_0 w_0 \in \alpha \wedge p_0 \in \beta \blacksquare n_0 w_0 + p_0 = n_0(-v/2) + -(n_0 + 2) - v/2 = v \in \alpha +_{\mathbb{R}} \beta$	
(1.16)	$v \in 0_{\mathbb{R}} \implies v \in \alpha +_{\mathbb{R}} \beta \blacksquare \forall_{v \in 0_{\mathbb{R}}}(v \in \alpha +_{\mathbb{R}} \beta) \blacksquare 0_{\mathbb{R}} \subseteq \alpha +_{\mathbb{R}} \beta$	
(1.17)	$\alpha +_{\mathbb{R}} \beta \subseteq 0_{\mathbb{R}} \wedge 0_{\mathbb{R}} \subseteq \alpha +_{\mathbb{R}} \beta \blacksquare \alpha +_{\mathbb{R}} \beta = 0_{\mathbb{R}}$	
(1.18)	$\beta \in \mathbb{R} \wedge \alpha +_{\mathbb{R}} \beta = 0_{\mathbb{R}} \blacksquare \exists_{-\alpha \in \mathbb{R}}(\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$	
(2)	$\alpha \in \mathbb{R} \implies \exists_{-\alpha \in \mathbb{R}}(\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$	

$*_{\mathbb{R}}[\alpha, \beta] := \quad \text{---}$
 $1_{\mathbb{R}} := \{x \in \mathbb{Q} : x < 1\}$

$11s\text{Not}0 := 0_{\mathbb{R}} \neq 1_{\mathbb{R}} \quad \text{---}$
 $11nR := 1_{\mathbb{R}} \in \mathbb{R} \quad \text{---}$

$\text{FieldMultiplicationClosureOf } \mathbb{R} := (\alpha, \beta \in \mathbb{R}) \implies (\alpha *_{\mathbb{R}} \beta) \in \mathbb{R} \quad \text{---}$
 $\text{FieldMultiplicationCommutativityOf } \mathbb{R} := (\alpha, \beta \in \mathbb{R}) \implies (\alpha *_{\mathbb{R}} \beta = \beta *_{\mathbb{R}} \alpha) \quad \text{---}$
 $\text{FieldMultiplicationAssociativityOf } \mathbb{R} := (\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha *_{\mathbb{R}} \beta) *_{\mathbb{R}} \gamma = \alpha *_{\mathbb{R}} (\beta *_{\mathbb{R}} \gamma)) \quad \text{---}$
 $\text{FieldMultiplicationIdentityOf } \mathbb{R} := (\alpha \in \mathbb{R}) \implies 1_{\mathbb{R}} *_{\mathbb{R}} \alpha = \alpha \quad \text{---}$
 $\text{FieldMultiplicationInverseOf } \mathbb{R} := (\alpha \in \mathbb{R}) \implies \exists_{1/\alpha \in \mathbb{R}}(\alpha *_{\mathbb{R}} (1/\alpha) = 1_{\mathbb{R}}) \quad \text{---}$

$\text{FieldDistributivityOf } \mathbb{R} := (\alpha, \beta, \gamma \in \mathbb{R}) \implies \gamma *_{\mathbb{R}} (\alpha +_{\mathbb{R}} \beta) = \gamma *_{\mathbb{R}} \alpha + \gamma *_{\mathbb{R}} \beta \quad \text{---}$

$\text{FieldWith } \mathbb{R} := \text{Field}[\mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}] \quad \text{---}$
 $\text{OrderedFieldWith } \mathbb{R} := \text{OrderedField}[\mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}] \quad \text{---}$

$\mathbb{Q}_{\mathbb{R}} := \{\{r \in \mathbb{Q} : r < q\} : q \in \mathbb{Q}\}$
 $\text{QROrderedSubfieldOf } \mathbb{R} := \text{OrderedSubfield}[\mathbb{Q}_{\mathbb{R}}, \mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}] \quad \text{---}$
 $\text{QIsomorphicToQR} := \mathbb{Q}_{\mathbb{R}} \simeq \mathbb{Q} \quad \text{---}$
 $\text{CompletenessOf } \mathbb{R} := \exists_{\mathbb{R}}(\text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}] \wedge \text{OrderedSubfield}[\mathbb{Q}, \mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}]) \quad \text{---}$

(1.20)
 $\text{ArchimedeanPropertyOf } \mathbb{R} := \forall_{x,y \in \mathbb{R}}(x > 0 \implies \exists_{n \in \mathbb{N}^+}(nx > y))$

(1) $(x, y \in \mathbb{R} \wedge x > 0) \implies \dots$

(1.1) $A := \{nx : n \in \mathbb{N}^+\} \blacksquare (\emptyset \neq A \subset \mathbb{R}) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a))$

(1.2) $\neg \exists_{n \in \mathbb{N}^+}(nx > y) \implies \dots$

(1.2.1) $\neg \exists_{n \in \mathbb{N}^+}(nx > y) \blacksquare \forall_{n \in \mathbb{N}^+}(nx \leq y) \blacksquare \text{UpperBound}[y_0, A, \mathbb{R}, <] \blacksquare \text{BoundedAbove}[A, \mathbb{R}, <]$

(1.2.2) $\text{CompletenessOf } \mathbb{R} \blacksquare \text{LUBProperty}[\mathbb{R}, <]$

(1.2.3) $(\text{LUBProperty}[\mathbb{R}, <]) \wedge (\emptyset \neq A \subset \mathbb{R}) \wedge (\text{BoundedAbove}[A, \mathbb{R}, <]) \blacksquare \exists_{\alpha \in \mathbb{R}}(\text{LUB}[\alpha, A, \mathbb{R}, <]) \dots$

(1.2.4) $\dots \alpha_0 := \text{choice}(\{\alpha \in \mathbb{R} : \text{LUB}[\alpha, A, \mathbb{R}, <]\}) \blacksquare \text{LUB}[\alpha_0, A, \mathbb{R}, <]$

(1.2.5) $x > 0 \blacksquare \alpha_0 - x < \alpha_0$

(1.2.6) $(\alpha_0 - x < \alpha_0) \wedge (\text{LUB}[\alpha_0, A, \mathbb{R}, <]) \blacksquare \neg \text{UpperBound}[\alpha_0 - x, A, \mathbb{R}, <]$

(1.2.7) $\neg \text{UpperBound}[\alpha_0 - x, A, \mathbb{R}, <] \blacksquare \exists_{c \in A}(\alpha_0 - x < c) \dots$

(1.2.8) $\dots c_0 := \text{choice}(\{c \in A : \alpha_0 - x < c\}) \blacksquare (c_0 \in A) \wedge (\alpha_0 - x < c_0)$

(1.2.9) $(c_0 \in A) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a)) \blacksquare \exists_{m \in \mathbb{N}^+}(mx = c_0) \dots$

(1.2.10) $\dots m_0 := \text{choice}(\{m \in \mathbb{N}^+ : mx = c_0\}) \blacksquare (m_0 \in \mathbb{N}^+) \wedge (m_0 x = c_0)$

(1.2.11) $(\alpha_0 - x < c_0) \wedge (m_0 x = c_0) \blacksquare \alpha_0 - x < c_0 = m_0 x \blacksquare \alpha_0 < m_0 x + x \blacksquare \alpha_0 < (m_0 + 1)x$

(1.2.12) $m_0 \in \mathbb{N}^+ \blacksquare m_0 + 1 \in \mathbb{N}^+$

(1.2.13) $(m_0 + 1 \in \mathbb{N}^+) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a)) \blacksquare (m_0 + 1)x \in A$

(1.2.14) $(\alpha_0 < (m_0 + 1)x) \wedge ((m_0 + 1)x \in A) \blacksquare \exists_{c \in A}(\alpha_0 < c)$

(1.2.15) $\text{LUB}[\alpha_0, A, \mathbb{R}, <] \blacksquare \text{UpperBound}[\alpha_0, A, \mathbb{R}, <] \blacksquare \forall_{c \in A}(c \leq \alpha_0) \blacksquare \neg \exists_{c \in A}(c > \alpha_0) \blacksquare \neg \exists_{c \in A}(\alpha_0 < c)$

(1.2.16) $(\exists_{c \in A}(\alpha_0 < c)) \wedge (\neg \exists_{c \in A}(\alpha_0 < c)) \blacksquare \perp$

(1.3) $\neg \exists_{n \in \mathbb{N}^+}(nx > y) \implies \perp \blacksquare \exists_{n \in \mathbb{N}^+}(nx > y)$

(2) $(x, y \in \mathbb{R} \wedge x > 0) \implies \exists_{n \in \mathbb{N}^+}(nx > y) \blacksquare \forall_{x, y \in \mathbb{R}}(x > 0 \implies \exists_{n \in \mathbb{N}^+}(nx > y))$

QDenseInR := $\forall_{x, y \in \mathbb{R}}(x < y \implies \exists_{p \in \mathbb{Q}}(x < p < y))$

(1) $(x, y \in \mathbb{R} \wedge x < y) \implies \dots$

(1.1) $x < y \blacksquare (0 < y - x) \wedge (y - x \in \mathbb{R})$

(1.2) $\text{ArchimedeanPropertyOf } \mathbb{R} \wedge (0 < y - x) \wedge (y - x, 1 \in \mathbb{R}) \blacksquare \exists_{n \in \mathbb{N}^+}(n(y - x) > 1) \dots$

(1.3) $\dots n_0 := \text{choice}(\{n \in \mathbb{N}^+ : n(y - x) > 1\}) \blacksquare (n_0 \in \mathbb{N}^+) \wedge (n_0(y - x) > 1)$

(1.4) $(n_0 \in \mathbb{N}^+) \wedge (x \in \mathbb{R}) \blacksquare n_0 x, -n_0 x \in \mathbb{R}$

(1.5) $\text{ArchimedeanPropertyOf } \mathbb{R} \wedge (1 > 0) \wedge (n_0 x, 1 \in \mathbb{R}) \blacksquare \exists_{m \in \mathbb{N}^+}(m(1) > n_0 x) \dots$

(1.6) $\dots m_1 := \text{choice}(\{m \in \mathbb{N}^+ : m(1) > n_0 x\}) \blacksquare (m_1 \in \mathbb{N}^+) \wedge (m_1 > n_0 x)$

(1.7) $\text{ArchimedeanPropertyOf } \mathbb{R} \wedge (1 > 0) \wedge (-n_0 x, 1 \in \mathbb{R}) \blacksquare \exists_{m \in \mathbb{N}^+}(m(1) > -n_0 x) \dots$

(1.8) $\dots m_2 := \text{choice}(\{m \in \mathbb{N}^+ : m(1) > -n_0 x\}) \blacksquare (m_2 \in \mathbb{N}^+) \wedge (m_2 > -n_0 x)$

(1.9) $(m_1 > n_0 x) \wedge (m_2 > -n_0 x) \blacksquare -m_2 < n_0 x < m_1$

(1.10) $m_1, m_2 \in \mathbb{N}^+ \blacksquare |m_1 - (-m_2)| \geq 2$

(1.11) $(-m_2 < n_0 x < m_1) \wedge (|m_1 - (-m_2)| \geq 2) \blacksquare \exists_{m \in \mathbb{Z}}((-m_2 < m < m_1) \wedge (m - 1 \leq n_0 x < m)) \dots$

(1.12) $\dots m_0 := \text{choice}(\{m \in \mathbb{Z} : (-m_2 < m < m_1) \wedge (m - 1 \leq n_0 x < m)\}) \blacksquare (-m_2 < m_0 < m_1) \wedge (m_0 - 1 \leq n_0 x < m_0)$

(1.13) $(n_0(y - x) > 1) \wedge (m_0 - 1 \leq n_0 x < m_0) \blacksquare n_0 x < m_0 \leq 1 + n_0 x < n_0 y \blacksquare n_0 x < m_0 < n_0 y$

(1.14) $(n_0 \in \mathbb{N}^+) \wedge (n_0 x < m_0 < n_0 y) \blacksquare x < m_0/n_0 < y$

(1.15) $m_0, n_0 \in \mathbb{Z} \blacksquare m_0/n_0 \in \mathbb{Q}$

(1.16) $(m_0/n_0 \in \mathbb{Q}) \wedge (x < m_0/n_0 < y) \blacksquare \exists_{p \in \mathbb{Q}}(x < p < y)$

(2) $(x, y \in \mathbb{R} \wedge x < y) \implies \exists_{p \in \mathbb{Q}}(x < p < y) \blacksquare \forall_{x, y \in \mathbb{R}}(x < y \implies \exists_{p \in \mathbb{Q}}(x < p < y))$

(1.21)

Root Lemma := $(0 < a < b) \implies (b^n - a^n \leq (b - a)nb^{n-1})$

(1) $(0 < a < b) \implies \dots$

(1.1) $b^n - a^n = (b - a) \sum_{i=1}^n (b^{n-i} a^{i-1})$

(1.2) $0 < a < b \blacksquare b/a > 1$

$$(1.3) \quad b/a > 1 \quad \blacksquare \quad \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq \sum_{i=1}^n \left(b^{n-i} a^{i-1} (b/a)^{i-1} \right) = \sum_{i=1}^n (b^{n-1}) = nb^{n-1} \quad \blacksquare \quad \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq \sum_{i=1}^n (b^{n-1}) = nb^{n-1}$$

$$(1.4) \quad b^n - a^n = (b - a) \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq (b - a) nb^{n-1} \quad \blacksquare \quad b^n - a^n \leq (b - a) nb^{n-1}$$

$$(2) \quad (0 < a < b) \implies \left(b^n - a^n \leq (b - a) nb^{n-1} \right)$$

$$\text{Root Existence In } \mathbb{R} := \forall_{0 < x \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} \exists!_{0 < y \in \mathbb{R}} (y_0^n = x)$$

$$(1) \quad (0 < x \in \mathbb{R} \wedge 0 < n \in \mathbb{Z}) \implies \dots$$

$$(1.1) \quad E := \{t \in \mathbb{R} : t > 0 \wedge t^n < x\} \quad \blacksquare \quad t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)$$

$$(1.2) \quad t_0 := x/(1+x) \quad \blacksquare \quad (t_0 = x/(1+x)) \wedge (t_0 \in \mathbb{R})$$

$$(1.3) \quad 0 < x \quad \blacksquare \quad 0 < x < 1+x \quad \blacksquare \quad t_0 = x/(1+x) > 0 \quad \blacksquare \quad t_0 > 0$$

$$(1.4) \quad 1 = (1+x)/(1+x) > x/(1+x) = t_0 \quad \blacksquare \quad 1 > t_0$$

$$(1.5) \quad (t_0 > 0) \wedge (1 > t_0) \quad \blacksquare \quad 0 < t_0 < 1$$

$$(1.6) \quad (0 < n \in \mathbb{Z}) \wedge (0 < t_0 < 1) \quad \blacksquare \quad t_0^n \leq t_0$$

$$(1.7) \quad 0 < x \quad \blacksquare \quad x > x/(1+x) = t_0 \quad \blacksquare \quad x > t_0$$

$$(1.8) \quad (t_0^n \leq t_0) \wedge (x > t_0) \quad \blacksquare \quad t_0^n < x$$

$$(1.9) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t_0 \in \mathbb{R}) \wedge (t_0 > 0) \wedge (t_0^n < x) \quad \blacksquare \quad t_0 \in E \quad \blacksquare \quad \emptyset \neq E$$

$$(1.10) \quad t_1 := \text{choice}(\{t \in \mathbb{R} : t > 1+x\}) \quad \blacksquare \quad (t_1 \in \mathbb{R}) \wedge (t_1 > 1+x)$$

$$(1.11) \quad x > 0 \quad \blacksquare \quad t_1 > 1+x > 1 \quad \blacksquare \quad t_1 > 1 \quad \blacksquare \quad t_1^n \geq t_1$$

$$(1.12) \quad (t_1^n \geq t_1) \wedge (t_1 > 1+x) \wedge (1 > 0) \quad \blacksquare \quad t_1^n \geq t_1 > 1+x > x \quad \blacksquare \quad t_1^n > x$$

$$(1.13) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t_1^n > x) \quad \blacksquare \quad t_1 \notin E \quad \blacksquare \quad E \subset \mathbb{R}$$

$$(1.14) \quad (\emptyset \neq E) \wedge (E \subset \mathbb{R}) \quad \blacksquare \quad \emptyset \neq E \subset \mathbb{R}$$

$$(1.15) \quad t \in E \implies \dots$$

$$(1.15.1) \quad (t \in E) \wedge (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \quad \blacksquare \quad t^n < x$$

$$(1.15.2) \quad (t_1^n > x) \wedge (t^n < x) \quad \blacksquare \quad t^n < x < t_1^n \quad \blacksquare \quad t < t_1$$

$$(1.16) \quad t \in E \implies t < t_1 \quad \blacksquare \quad \forall_{t \in E} (t \leq t_1) \quad \blacksquare \quad \text{Upper Bound}[t_1, E, \mathbb{R}, <] \quad \blacksquare \quad \text{Bounded Above}[E, \mathbb{R}, <]$$

$$(1.17) \quad \text{Completeness Of } \mathbb{R} \quad \blacksquare \quad \text{LUB Property}[\mathbb{R}, <]$$

$$(1.18) \quad (\text{LUB Property}[\mathbb{R}, <]) \wedge (\emptyset \neq E \subset \mathbb{R}) \wedge (\text{Bounded Above}[E, \mathbb{R}, <]) \quad \blacksquare \quad \exists_{y \in \mathbb{R}} (\text{LUB}[y, E, \mathbb{R}, <]) \quad \dots$$

$$(1.19) \quad \dots y_0 := \text{choice}(\{y \in \mathbb{R} : \text{LUB}[y, E, \mathbb{R}, <]\}) \quad \blacksquare \quad \text{LUB}[y_0, E, \mathbb{R}, <]$$

$$(1.20) \quad (\text{LUB}[y_0, E, \mathbb{R}, <]) \wedge (t_0 \in E) \wedge (t_0 > 0) \quad \blacksquare \quad 0 < t_0 \leq y_0 \in \mathbb{R} \quad \blacksquare \quad 0 < y_0 \in \mathbb{R}$$

$$(1.21) \quad y_0^n < x \implies \dots$$

$$(1.21.1) \quad k_0 := \frac{x - y_0^n}{n(y_0 + 1)^{n-1}} \quad \blacksquare \quad k_0 \in \mathbb{R}$$

$$(1.21.2) \quad y_0^n < x \quad \blacksquare \quad 0 < x - y_0^n$$

$$(1.21.3) \quad (n > 0) \wedge (y_0 > 0) \quad \blacksquare \quad 0 < n(y_0 + 1)^{n-1}$$

$$(1.21.4) \quad (0 < x - y_0^n) \wedge \left(0 < n(y_0 + 1)^{n-1} \right) \quad \blacksquare \quad 0 < \frac{x - y_0^n}{n(y_0 + 1)^{n-1}} = k_0 \quad \blacksquare \quad 0 < k_0$$

$$(1.21.5) \quad (0 < 1 \in \mathbb{R}) \wedge (0 < k_0 \in \mathbb{R}) \quad \blacksquare \quad 0 < \min(1, k_0) \in \mathbb{R}$$

$$(1.21.6) \quad \text{QDense In } \mathbb{R} \wedge (0, \min(1, k_0) \in \mathbb{R}) \wedge (0 < \min(1, k_0)) \quad \blacksquare \quad \exists_{h \in \mathbb{Q}} (0 < h < \min(1, k_0)) \quad \dots$$

$$(1.21.7) \quad \dots h_0 := \text{choice}(\{h \in \mathbb{Q} : 0 < h < \min(1, k_0)\}) \quad \blacksquare \quad (0 < h_0 < 1) \wedge \left(h_0 < k_0 = \frac{x - y_0^n}{n(y_0 + 1)^{n-1}} \right)$$

$$(1.21.8) \quad (y_0 > 0) \wedge (h_0 > 0) \quad \blacksquare \quad 0 < y_0 < y_0 + h_0$$

$$(1.21.9) \quad \text{Root Lemma} \wedge (0 < y_0 < y_0 + h_0) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < h_0 n(y_0 + h_0)^{n-1}$$

$$(1.21.10) \quad h_0 < 1 \quad \blacksquare \quad h_0 n(y_0 + h_0)^{n-1} < h_0 n(y_0 + 1)^{n-1}$$

$$(1.21.11) \quad \left((y_0 + h_0)^n - y_0^n < h_0 n(y_0 + h_0)^{n-1} \right) \wedge \left(h_0 n(y_0 + h_0)^{n-1} < h_0 n(y_0 + 1)^{n-1} \right) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < h_0 n(y_0 + 1)^{n-1}$$

$$(1.21.12) \quad \left(0 < n(y_0 + 1)^{n-1} \right) \wedge \left(h_0 < k_0 = \frac{x - y_0^n}{n(y_0 + 1)^{n-1}} \right) \quad \blacksquare \quad h_0 n(y_0 + 1)^{n-1} < x - y_0^n$$

$$(1.21.13) \quad \left((y_0 + h_0)^n - y_0^n < h_0 n(y_0 + 1)^{n-1} \right) \wedge \left(h_0 n(y_0 + 1)^{n-1} < x - y_0^n \right) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < x - y_0^n \quad \blacksquare \quad (y_0 + h_0)^n < x$$

$$(1.21.14) \quad (y_0 + h_0)^n - y_0^n < x - y_0^n \quad \blacksquare \quad (y_0 + h_0)^n < x$$

$$(1.21.15) \quad (0 < y_0 \in \mathbb{R}) \wedge (0 < h_0 \in \mathbb{R}) \quad \blacksquare \quad 0 < y_0 < y_0 + h_0 \in \mathbb{R}$$

$$(1.21.16) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge ((y_0 + h_0)^n < x) \wedge (0 < y_0 + h_0 \in \mathbb{R}) \quad \blacksquare \quad (y_0 + h_0)^n \in E$$

(1.21.17)	$((y_0 + h_0)^n \in E) \wedge (y_0 < y_0 + h_0) \blacksquare \exists_{e \in E} (y_0 < e)$
(1.21.18)	$\textcolor{blue}{LUB}[y_0, E, \mathbb{R}, <] \blacksquare \textcolor{blue}{UpperBound}[y_0, E, \mathbb{R}, <] \blacksquare \forall_{e \in E} (e \leq y_0) \blacksquare \neg \exists_{e \in E} (e > y_0)$
(1.21.19)	$(\exists_{e \in E} (e > y_0)) \wedge (\neg \exists_{e \in E} (e > y_0)) \blacksquare \perp$
(1.22)	$y_0^n < x \implies \perp \blacksquare y_0^n \geq x$
(1.23)	$y_0^n > x \implies \dots$
(1.23.1)	$k_1 := \frac{y_0^n - x}{ny_0^{n-1}} \blacksquare (k_1 \in \mathbb{R}) \wedge (k_1 ny_0^{n-1} = y_0^n - x)$
(1.23.2)	$(0 < x) \wedge (0 < n \in \mathbb{Z}) \blacksquare y_0^n - x < y_0^n \leq ny_0^n \blacksquare y_0^n - x < ny_0^n$
(1.23.3)	$y_0^n - x < ny_0^n \blacksquare k_1 = \frac{y_0^n - x}{ny_0^{n-1}} < \frac{ny_0^n}{ny_0^{n-1}} = y_0 \blacksquare k_1 < y_0$
(1.23.4)	$y_0^n > x \blacksquare 0 < y_0^n - x$
(1.23.5)	$(n > 0) \wedge (y_0 > 0) \blacksquare 0 < ny_0^{n-1}$
(1.23.6)	$(0 < y_0^n - x) \wedge 0 < (ny_0^{n-1}) \blacksquare 0 < \frac{y_0^n - x}{ny_0^{n-1}} = k_1 \blacksquare 0 < k_1$
(1.23.7)	$(k_1 < y_0) \wedge (0 < k_1) \blacksquare (0 < k_1 < y_0) \wedge (0 < y_0 - k_1 < y_0)$
(1.23.8)	$t \geq y_0 - k_1 \implies \dots$
(1.23.8.1)	$t \geq y_0 - k_1 \blacksquare t^n \geq (y_0 - k_1)^n \blacksquare -t^n \leq -(y_0 - k_1)^n \blacksquare y_0^n - t^n \leq y_0^n - (y_0 - k_1)^n$
(1.23.8.2)	$\textcolor{blue}{RootLemma} \wedge (0 < y_0 - k_1 < y_0) \blacksquare y_0^n - (y_0 - k_1)^n < k_1 ny_0^{n-1}$
(1.23.8.3)	$(y_0^n - t^n \leq y_0^n - (y_0 - k_1)^n) \wedge (y_0^n - (y_0 - k_1)^n < k_1 ny_0^{n-1}) \blacksquare y_0^n - t^n < k_1 ny_0^{n-1}$
(1.23.8.4)	$(k_1 ny_0^{n-1} = y_0^n - x) \wedge (y_0^n - t^n < k_1 ny_0^{n-1}) \blacksquare y_0^n - t^n < y_0^n - x \blacksquare -t^n < -x \blacksquare t^n > x$
(1.23.8.5)	$(t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t^n > x) \blacksquare t \notin E$
(1.23.9)	$t \geq y_0 - k_1 \implies t \notin E \blacksquare t \in E \implies t < y_0 - k_1 \blacksquare \forall_{t \in E} (t \leq y_0 - k_1) \blacksquare \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]$
(1.23.10)	$(\textcolor{blue}{LUB}[y_0, E, \mathbb{R}, <] \wedge (y_0 - k_1 < y_0)) \blacksquare \neg \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]$
(1.23.11)	$(\textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]) \wedge (\neg \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]) \blacksquare \perp$
(1.24)	$y_0^n > x \implies \perp \blacksquare y_0^n \leq x$
(1.25)	$\textcolor{blue}{Order}[\mathbb{R}, <] \blacksquare \textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]$
(1.26)	$(\textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]) \wedge (y_0^n \geq x) \wedge (y_0^n \leq x) \blacksquare y_0^n = x$
(1.27)	$(y_0^n = x) \wedge (y_0 \in \mathbb{R}) \blacksquare \exists_{y \in \mathbb{R}} (y^n = x)$
(1.28)	$y_1, y_2 := \textit{choice}(\{y \in \mathbb{R} : y^n = x\})$
(1.29)	$y_1 \neq y_2 \implies \dots$
(1.29.1)	$(\textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]) \wedge (y_1 \neq y_2) \blacksquare (y_1 < y_2) \vee (y_2 < y_1) \dots$
(1.29.2)	$\dots (x = y_1^n < y_2^n = x) \vee (x = y_2^n < y_1^n = x) \blacksquare (x < x) \vee (x > x) \blacksquare \perp \vee \perp \blacksquare \perp$
(1.30)	$y_1 \neq y_2 \implies \perp \blacksquare y_1 = y_2 \blacksquare \forall_{a,b \in \mathbb{R}} ((a^n = x \wedge b^n = x) \implies a = b)$
(1.31)	$(\exists_{y \in \mathbb{R}} (y^n = x)) \wedge (\forall_{a,b \in \mathbb{R}} ((a^n = x \wedge b^n = x) \implies a = b)) \blacksquare \exists!_{y \in \mathbb{R}} (y^n = x)$
(2)	$(0 < x \in \mathbb{R} \wedge 0 < n \in \mathbb{Z}) \implies \exists!_{y \in \mathbb{R}} (y^n = x) \blacksquare \forall_{0 < x \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} \exists!_{0 < y \in \mathbb{R}} (y_0^n = x)$

$$\textcolor{red}{RootExistenceInRCorollary} := \forall_{0 < a \in \mathbb{R}} \forall_{0 < b \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} \left((ab)^{1/n} = a^{1/n} b^{1/n} \right) \quad \text{---}$$

$$\textcolor{red}{ExtendedRealSystem}[\bar{\mathbb{R}}, +, *, <] := \left(\begin{array}{l} \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\} \quad \wedge \quad -\infty < x < \infty \quad \wedge \\ x + \infty = +\infty \quad \wedge \quad x - \infty = -\infty \quad \wedge \quad \frac{x}{+\infty} = \frac{x}{-\infty} = 0 \quad \wedge \\ (x > 0) \implies (x * (+\infty) = +\infty \wedge x * (-\infty) = -\infty) \quad \wedge \\ (x < 0) \implies (x * (+\infty) = -\infty \wedge x * (-\infty) = +\infty) \end{array} \right)$$

$$\mathbb{C} := \{\langle a, b \rangle \in \mathbb{R} \times \mathbb{R}\}$$

$$+_C[\langle a, b \rangle, \langle c, d \rangle] := \langle a +_{\mathbb{R}} c, b +_{\mathbb{R}} d \rangle$$

$$*_C[\langle a, b \rangle, \langle c, d \rangle] := \langle a *_R c - b *_R d, a *_R d + b *_R c \rangle$$

$$\textcolor{red}{FieldC} := \textcolor{blue}{Field}[\mathbb{C}, +_C, *_C] \quad \text{---}$$

$$\textcolor{red}{RSubfieldC} := \textcolor{blue}{Subfield}[\mathbb{R}, \mathbb{C}, +, *] \quad \text{---}$$

$$i := \langle 0, 1 \rangle \in \mathbb{C}$$

$$\textcolor{red}{iProperty} := i^2 = -1 \quad \text{---}$$

$$\textcolor{red}{CProperty} := (a, b \in \mathbb{R}) \implies (\langle a, b \rangle = a + bi) \quad \text{---}$$

Chapter 2

Abstract Algebra

2.1 Functions

$$Rel[r, X] := (X \neq \emptyset) \wedge (r \subseteq X)$$

$$Func[f, X, Y] := (Rel[f, X \times Y]) \wedge \left(\forall_{x \in X} \exists!_{y \in Y} (\langle x, y \rangle \in f) \right)$$

$$Comp[g \circ f, f, g, X, Y, Z] := (Func[f, X, Y]) \wedge (Func[g, Y, Z]) \wedge \left(g \circ f = \{ \langle x, g(f(x)) \rangle \in X \times Z \mid x \in X \} \right)$$

$$FuncComp := (Comp[g \circ f, f, g, X, Y, Z]) \implies (Func[g \circ f, X, Z])$$

(1) TODO

$$CompAssoc := ho(g \circ f) = (h \circ g) \circ f$$

(1) TODO

$$Domain[dom(f), f, X, Y] := (Func[f, X, Y]) \wedge (dom(f) = X)$$

$$Codomain[cod(f), f, X, Y] := (Func[f, X, Y]) \wedge (cod(f) = Y)$$

$$Image[im(A), A, f, X, Y] := (Func[f, X, Y]) \wedge (A \subseteq X) \wedge (im(A) = \{ f(a) \in Y \mid a \in A \})$$

$$Preimage[pim(B), B, f, X, Y] := (Func[f, X, Y]) \wedge (B \subseteq Y) \wedge (pim(B) = \{ a \in X \mid f(a) \in B \})$$

$$Range[rng(f), f, X, Y] := (Func[f, X, Y]) \wedge (Image[rng(f), dom(f), f, X, Y])$$

$$Inj[f, X, Y] := (Func[f, X, Y]) \wedge \left(\forall_{x_1, x_2 \in X} \left((f(x_1) = f(x_2)) \implies (x_1 = x_2) \right) \right)$$

$$Surj[f, X, Y] := (Func[f, X, Y]) \wedge \left(\forall_{y \in Y} \exists_{x \in X} (y = f(x)) \right)$$

$$Bij[f, X, Y] := (Inj[f, X, Y]) \wedge (Surj[f, X, Y])$$

$$Inv[f^{-1}, f, X, Y] := (Func[f, X, Y]) \wedge (Func[f^{-1}, Y, X]) \wedge (f \circ f^{-1} = I_Y) \wedge (f^{-1} \circ f = I_X)$$

$$SurjEquiv := (Surj[f, X, Y]) \iff (rng(f) = cod(f))$$

(1) TODO

$$BijEquiv := (Bij[f, X, Y]) \iff \left(\exists_{f^{-1}} (Inv[f^{-1}, f, X, Y]) \right)$$

(1) TODO

$$InjComp := ((Inj[f]) \wedge (Inj[g])) \implies (Inj[g \circ f])$$

(1) TODO

$$SurjComp := ((Surj[f]) \wedge (Surj[g])) \implies (Surj[g \circ f])$$

(1) TODO

2.2 Divisibility, Equivalence Relations, Partitions

$$\text{DivisionAlgorithm} := \forall_{b \in \mathbb{Z}} \forall_{a \in \mathbb{Z}^+} \exists!_{q, r \in \mathbb{Z}} ((b = aq + r) \wedge (0 \leq r < a))$$

(1) TODO

$$\text{Divides}[a, b] := (a, b \in \mathbb{Z}) \wedge (\exists_{c \in \mathbb{Z}} (b = ac))$$

$$\text{ComDiv}[a, b, c] := (\text{Divides}[a, b]) \wedge (\text{Divides}[a, c])$$

$$\text{GCD}[a, b, c] := (\text{ComDiv}[a, b, c]) \wedge \left(\forall_{d \in \mathbb{Z}} \left(((\text{Divides}[d, b]) \wedge (\text{Divides}[d, c])) \implies (\text{Divides}[d, a]) \right) \right)$$

$$\text{RelPrime}[a, b] := \text{GCD}[1, a, b]$$

$$\text{CongRel}[a, b, n] := \text{Divides}[n, a - b]$$

$$\text{Partition}[\mathcal{P}, S] := (\forall_{P \in \mathcal{P}} (P \neq \emptyset)) \wedge \left(S = \bigcup_{P \in \mathcal{P}} (P) \right) \wedge \left(\forall_{P_1, P_2 \in \mathcal{P}} ((P_1 \neq P_2) \implies (P_1 \cap P_2 = \emptyset)) \right)$$

$$\text{EqRel}[\sim, S] := (\text{Rel}[\sim, S]) \wedge (\forall_{a \in S} (a \sim a)) \wedge \left(\forall_{a, b \in S} ((a \sim b) \implies (b \sim a)) \right) \wedge \left(\forall_{a, b, c \in S} (((a \sim b) \wedge (b \sim c)) \implies (a \sim c)) \right)$$

$$\text{EqClass}[[s], s, \sim, S] := (\text{Rel}[\sim, S]) \wedge (s \in S) \wedge ([s] = \{x \in S \mid x \sim s\})$$

$$\text{PartitionInducesEqRel} := (\text{Partition}[\mathcal{P}, S]) \implies (\exists_{\sim} (\text{EqRel}[\sim, S]))$$

(1) TODO : $\sim = \{\langle a, b \rangle \in S \times S \mid (P \in \mathcal{P}) \wedge (a, b \in P)\}$

$$\text{EqRelInducesPartition} := (\text{EqRel}[\sim, S]) \implies (\exists_{\mathcal{P}} (\text{Partition}[\mathcal{P}, S]))$$

(1) TODO : $\text{Partition}[\text{EqClass}_1, \text{EqClass}_2, \dots]$

$$\text{EqRelCong} := \forall_{n \in \mathbb{Z}^+} (\text{EqRel}[\text{CongRel}, \mathbb{Z}])$$

(1) TODO

2.3 Groups

$$\text{Group}[G, *] := \left(\begin{array}{l} (\text{Function}[*, G, G]) \quad \wedge \\ \left(\forall_{a, b, c \in G} ((a * b) * c = a * (b * c)) \right) \wedge \\ \left(\exists_{e \in G} \forall_{a \in G} (a * e = a = e * a) \right) \quad \wedge \\ \left(\forall_{a \in G} \exists_{a^{-1} \in G} (a * a^{-1} = e = a^{-1} * a) \right) \end{array} \right)$$

$$\text{AbelianGroup}[G, *] := (\text{Group}[G, *]) \wedge (\forall_{a, b \in G} (a * b = b * a))$$

$$\text{CancelLaws} := \forall_G \left((\text{Group}[G, *]) \implies \left(\forall_{a, b, c \in G} \left(((a * b = a * c) \implies (b = c)) \wedge ((a * c = b * c) \implies (a = b)) \right) \right) \right)$$

(1) $(a * b = a * c) \implies \dots$

$$(1.1) \quad a \in G \quad \blacksquare \quad \exists_{a^{-1} \in G} (a * a^{-1} = e = a^{-1} * a)$$

$$(1.2) \quad \text{Function}[*, G, G] \quad \blacksquare \quad a^{-1} * a * b = a^{-1} * a * c$$

$$(1.3) \quad \left(\forall_{a, b, c \in G} ((a * b) * c = a * (b * c)) \right) \wedge \left(\forall_{a \in G} \exists_{a^{-1} \in G} (a * a^{-1} = e = a^{-1} * a) \right) \quad \blacksquare \quad b = c$$

(2) $(a * b = a * c) \implies (b = c)$

(3) $(a * c = b * c) \implies \dots$

(3.1) TODO

(4) $(a * c = b * c) \implies (a = b)$

(5) $((a * b = a * c) \implies (b = c)) \wedge ((a * c = b * c) \implies (a = b))$

$$\text{IdUniq} := \forall_G \left((\text{Group}[G, *]) \implies \left(\forall_{e_1, e_2 \in G} \forall_{a \in G} \left(((a * e_1 = a = e_1 * a) \wedge (a * e_2 = a = e_2 * a)) \implies (e_1 = e_2) \right) \right) \right)$$

(1) $(\text{CancelLaws}) \wedge \left(\forall_{a \in G} \exists_{a^{-1} \in G} (a * a^{-1} = e = a^{-1} * a) \right) \quad \blacksquare \quad a * e_1 = a = a * e_2 \quad \blacksquare \quad e_1 = e_2$

$$InvUniq := \forall_G \left((Group[G, *]) \implies \left(\forall_{a \in G} \forall_{a_1^{-1}, a_2^{-1} \in G} \left((a * a_1^{-1} = e = a_1^{-1} * a) \wedge (a * a_2^{-1} = e = a_2^{-1} * a) \implies (a_1^{-1} = a_2^{-1}) \right) \right) \right)$$

$$(1) \quad (CancelLaws) \wedge \left(\forall_{a \in G} \exists_{a^{-1} \in G} (a * a^{-1} = e = a^{-1} * a) \right) \blacksquare a * a_1^{-1} = e = a * a_2^{-1} \blacksquare a_1^{-1} = a_2^{-1}$$

$$InvProd := \forall_G \forall_{a, b \in G} \left((a * b)^{-1} = b^{-1} * a^{-1} \right)$$

$$(1) \quad (a * b) * (a * b)^{-1} = e$$

$$(2) \quad (a * b) * (b^{-1} * a^{-1}) = (a * (b * b^{-1}) * a^{-1}) = e$$

$$(3) \quad InvUniq \blacksquare (a * b)^{-1} = b^{-1} * a^{-1}$$

$$OrderEl[o(G), G, *] := (Group[G, *]) \wedge (o(G) = |G|)$$

$$gWitness[n, g, G, *] := (Group[G, *]) \wedge (n \in \mathbb{Z}^+) \wedge (g^n = e) \wedge (\forall_{m \in \mathbb{Z}^+} (m < n) \implies (g^m \neq e))$$

$$OrderEl[o(g), g, G, *] := (Group[G, *]) \wedge \left((\exists_n (gWitness[n, g, G, *])) \implies (o(g) = n) \right) \wedge \left((\neg \exists_n (gWitness[n, g, G, *])) \implies (o(g) = \infty) \right)$$

2.4 Subgroups

$$Subgroup[H, G, *] := (Group[G, *]) \wedge (H \subseteq G) \wedge (Group[H, *])$$

$$TrivSubgroup[H, G, *] := (H = \{e\}) \vee (H = G)$$

$$PropSubgroup[H, G, *] := (Subgroup[H, G, *]) \wedge (\neg TrivSubgroup[H, G, *])$$

$$SubgroupEquiv := \forall_{H, G} \left(\begin{array}{c} (Subgroup[H, G, *]) \\ \iff \\ \left((Group[G, *]) \wedge (\emptyset \neq H \subseteq G) \wedge (Function[*, H, H]) \wedge \left(\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a) \right) \right) \end{array} \right)$$

$$(1) \quad (Subgroup[H, G, *]) \implies \left((\emptyset \neq H \subseteq G) \wedge (Function[*, H, H]) \wedge \left(\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a) \right) \right)$$

$$(2) \quad \left((\emptyset \neq H \subseteq G) \wedge (Function[*, H, H]) \wedge \left(\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a) \right) \right) \implies \dots$$

$$(2.1) \quad Group[G, *] \blacksquare (a, b, c \in H) \implies (a, b, c \in G) \implies ((a * b) * c = a * (b * c)) \blacksquare \forall_{a, b, c \in H} ((a * b) * c = a * (b * c))$$

$$(2.2) \quad \emptyset \neq H \blacksquare \exists_h (h \in H)$$

$$(2.3) \quad h \in H \blacksquare \exists_{h^{-1} \in H} (h * h^{-1} = e = h^{-1} * h)$$

$$(2.4) \quad Function[*, H, H] \blacksquare e = h * h^{-1} \in H \blacksquare e \in H \blacksquare \exists_{e \in H} \forall_{a \in H} (a * e = a = e * a)$$

$$(2.5) \quad (Function[*, H, H]) \wedge \left(\forall_{a, b, c \in H} ((a * b) * c = a * (b * c)) \right) \wedge \left(\exists_{e \in H} \forall_{a \in H} (a * e = a = e * a) \wedge (\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a)) \right)$$

$$(2.6) \quad Group[H, *]$$

$$(2.7) \quad (Group[G, *]) \wedge (H \subseteq G) \wedge (Group[H, *]) \blacksquare Subgroup[H, G, *]$$

$$(3) \quad \left((\emptyset \neq H \subseteq G) \wedge (Function[*, H, H]) \wedge \left(\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a) \right) \right) \implies (Subgroup[H, G, *])$$

$$(4) \quad (Subgroup[H, G, *]) \iff \left((Group[G, *]) \wedge (\emptyset \neq H \subseteq G) \wedge (Function[*, H, H]) \wedge \left(\forall_{a \in H} \exists_{a^{-1} \in H} (a * a^{-1} = e = a^{-1} * a) \right) \right)$$

$$SubgroupEquivOST := \forall_{H, G} \left((Subgroup[H, G, *]) \iff \left((Group[G, *]) \wedge (\emptyset \neq H \subseteq G) \wedge \left(\forall_{a, b \in H} (a * b^{-1} \in H) \right) \right) \right)$$

$$(1) \quad \text{TODO}$$

$$SubgroupIntersection := \forall_{H_1, H_2, G} \left(((Subgroup[H_1, G, *]) \wedge (Subgroup[H_2, G, *])) \implies (Subgroup[H_1 \cap H_2, G, *]) \right)$$

$$(1) \quad Group[G, *]$$

$$(2) \quad (e \in H_1) \wedge (e \in H_2) \blacksquare e \in H_1 \cap H_2 \blacksquare \emptyset \neq H_1 \cap H_2$$

$$(3) \quad (H_1 \subseteq G) \wedge (H_2 \subseteq G) \blacksquare H_1 \cap H_2 \subseteq G$$

-
- (4) $\emptyset \neq H_1 \cap H_2 \subseteq G$
-
- (5) $(a, b \in H_1 \cap H_2) \implies \dots$
-
- (5.1) $a, b \in H_1 \blacksquare a * b \in H_1$
-
- (5.2) $a, b \in H_2 \blacksquare a * b \in H_2$
-
- (5.3) $a * b \in H_1 \cap H_2$
-
- (6) $(a, b \in H_1 \cap H_2) \implies (a * b \in H_1 \cap H_2) \blacksquare \text{Function}[* , H_1 \cap H_2, H_1 \cap H_2]$
-
- (7) $(a \in H_1 \cap H_2) \implies \dots$
-
- (7.1) $(a^{-1} \in H_1) \wedge (a^{-1} \in H_2) \blacksquare a^{-1} \in H_1 \cap H_2$
-
- (8) $(a \in H_1 \cap H_2) \implies (a^{-1} \in H_1 \cap H_2) \blacksquare \forall_{a \in H_1 \cap H_2} \exists_{a^{-1} \in H_1 \cap H_2} (a * a^{-1} = e = a^{-1} * a)$
-
- (9) $(\text{SubgroupEquiv}) \wedge (\text{Group}[G, *]) \wedge (\emptyset \neq H_1 \cap H_2 \subseteq G) \wedge (\text{Function}[* , H_1 \cap H_2, H_1 \cap H_2]) \wedge \dots$
-
- (10) $\dots \left(\forall_{a \in H_1 \cap H_2} \exists_{a^{-1} \in H_1 \cap H_2} (a * a^{-1} = e = a^{-1} * a) \right) \blacksquare \text{Subgroup}[H_1 \cap H_2, G, *]$
-

$$\text{Centralizer}[C(g), g, G, *] := (\text{Group}[G, *]) \wedge (g \in G) \wedge (C(g) = \{h \in G \mid g * h = h * g\})$$

$$\text{SubgroupCentralizer} := \forall_{g, G} \left((\text{Centralizer}[C(g), g, G, *]) \implies (\text{Subgroup}[C(g), G, *]) \right)$$

-
- (1) $e * g = g * e \blacksquare e \in C(g) \blacksquare C(g) \neq \emptyset$
-
- (2) $C(g) \subseteq G \blacksquare \emptyset \neq C(g) \subseteq G$
-
- (3) $(a, b \in C(g)) \implies \dots$
-
- (3.1) $(a * g = g * a) \wedge (b * g = g * b)$
-
- (3.2) $(a * b) * g = a * (b * g) = a * (g * b) = (a * g) * b = (g * a) * b = g * (a * b) \blacksquare a * b \in C(g)$
-
- (4) $(a, b \in C(g)) \implies (a * b \in C(g)) \blacksquare \forall_{a, b \in C(g)} (a * b \in C(g))$
-
- (5) $(a \in C(g)) \implies \dots$
-
- (5.1) $a * g = g * a$
-
- (5.2) $a^{-1} * (a * g) * a^{-1} = a^{-1} * (g * a) * a^{-1} \blacksquare g * a^{-1} = a^{-1} * g \blacksquare a^{-1} \in C(g)$
-
- (6) $(a \in C(g)) \implies (a^{-1} \in C(g)) \blacksquare \forall_{a \in C(g)} (a^{-1} \in C(g))$
-
- (7) $(\text{SubgroupEquiv}) \wedge (\emptyset \neq C(g) \subseteq G) \wedge \left(\forall_{a, b \in C(g)} (a * b \in C(g)) \right) \wedge \left(\forall_{a \in C(g)} (a^{-1} \in C(g)) \right) \blacksquare \text{Subgroup}[C(g), G, *]$
-

$$\text{Center}[Z(G), G, *] := (\text{Group}[G, *]) \wedge \left(Z(G) = \bigcap_{g \in G} (C(g)) \right)$$

$$\text{SubgroupCenter} := \forall_G \left((\text{Center}[Z(G), G, *]) \implies (\text{Subgroup}[Z(G), G, *]) \right)$$

-
- (1) $(\text{SubgroupCentralizer}) \wedge (\text{SubgroupIntersection}) \blacksquare \text{Subgroup}[Z(G), G, *]$
-

2.5 Special Groups

2.5.1 Cyclic Group

$$\text{CyclicSubgroup}[<g>, g, G, *] := (\text{Group}[G, *]) \wedge (g \in G) \wedge (<g> = \{g^n \mid n \in \mathbb{Z}\})$$

$$\text{Generator}[g, G, *] := \text{CyclicSubgroup}[G, g, G, *]$$

$$\text{CyclicGroup}[G, *] := \exists_{g \in G} (\text{Generator}[g, G, *])$$

$$\text{SubgroupOfCyclicGroupIsCyclic} := \forall_{G, H} \left(((\text{CyclicGroup}[G, *]) \wedge (\text{Subgroup}[H, G, *])) \implies (\text{CyclicGroup}[H, *]) \right)$$

-
- (1) $\exists_{g \in G} (\text{Generator}[g, G, *])$
-
- (2) $H \subseteq G \blacksquare \exists_{m \in \mathbb{Z}^+} \left((g^m \in H) \wedge \left(\forall_{k \in \mathbb{Z}^+} ((k < m) \implies (g^k \notin H)) \right) \right)$
-
- (3) $(b \in H) \implies \dots$
-
- (3.1) $H \subseteq G \blacksquare \exists_{n \in \mathbb{Z}^+} (b = g^n)$
-
- (3.2) $(\text{DivisionAlgorithm}) \wedge (n \in \mathbb{Z}) \wedge (m \in \mathbb{Z}^+) \blacksquare \exists!_{q, r \in \mathbb{Z}} ((n = mq + r) \wedge (0 \leq r < m))$
-

$$(3.3) \quad g^n = g^{mq+r} = g^{mq} * g^r \quad \blacksquare \quad g^r = (g^{mq})^{-1} * g^n$$

$$(3.4) \quad g^n, g^m \in H \quad \blacksquare \quad g^n, (g^{mq})^{-1} \in H \quad \blacksquare \quad g^r = g^{mq})^{-1} * g^n \in H \quad \blacksquare \quad g^r \in H$$

$$(3.5) \quad (g^r \in H) \wedge (0 \leq r < m) \wedge \left(\bigvee_{k \in \mathbb{Z}^+} ((k < m) \implies (g^k \notin H)) \right) \quad \blacksquare \quad r = 0$$

$$(3.6) \quad (r = 0) \wedge (g^n = g^{mq+r}) \wedge (b = g^n) \quad \blacksquare \quad b = g^n = g^{mq} \quad \blacksquare \quad b \in \langle g^m \rangle$$

$$(4) \quad (b \in H) \implies (b \in \langle g^m \rangle) \quad \blacksquare \quad H \subseteq \langle g^m \rangle$$

$$(5) \quad (b \in \langle g^m \rangle) \implies \dots$$

$$(5.1) \quad \exists_{k \in \mathbb{Z}} (b = g^{mk})$$

$$(5.2) \quad g^m \in H \quad \blacksquare \quad b = g^{mk} \in H \quad \blacksquare \quad b \in H$$

$$(6) \quad (b \in \langle g^m \rangle) \implies (b \in H) \quad \blacksquare \quad \langle g^m \rangle \subseteq H$$

$$(7) \quad (H \subseteq \langle g^m \rangle) \wedge (\langle g^m \rangle \subseteq H) \quad \blacksquare \quad H = \langle g^m \rangle \quad \blacksquare \quad \text{Generator}[g^m, H, *] \quad \blacksquare \quad \exists_{h \in G} (\text{Generator}[h, G, *]) \quad \blacksquare \quad \text{CyclicGroup}[H, *]$$

$$\text{ExpModOrder} := \forall_{G, g, n, s, t} \left(((\text{Group}[G, *]) \wedge (\text{OrderEl}[n, g, G, *])) \implies \left((g^s = g^t) \iff (s \equiv t \pmod{n}) \right) \right)$$

$$(1) \quad (s \equiv t \pmod{n}) \iff (\text{Divides}[n, s - t]) \iff (\exists_{k \in \mathbb{N}} (s - t = kn)) \iff \dots$$

$$(2) \quad \dots (\exists_{k \in \mathbb{N}} (s = kn + t)) \iff (g^s = g^{kn+t} = g^{kn} * g^t = e^k * g^t = g^t) \iff (g^s = g^t)$$

$$\text{ExpModOrderCorollary} := \forall_{G, g, n, s, t} \left(((\text{Group}[G, *]) \wedge (\text{OrderEl}[n, g, G, *])) \implies ((g^s = e) \iff (\text{Divides}[n, s])) \right)$$

$$(1) \quad \text{ExpModOrder} \quad \blacksquare \quad (g^s = e) \iff (g^s = g^0) \iff (s \equiv 0 \pmod{n}) \iff (\text{Divides}[n, s - 0]) \iff (\text{Divides}[n, s])$$

2.5.2 Symmetric and Alternating Groups

$$\text{SymmetricGroup}[S_n, n] := S_n = \{\text{permutation of a set with } n \text{ elements}\}$$

$$\text{SymmetricGroupOrder} := o(S_n) = n!$$

$$\text{SymmetricGroupAsDisjoinsCycles} := \forall_{\sigma \in S_n} \exists_{\Sigma \subseteq S_n} \left((\text{DisjointCycles}[\Sigma]) \wedge (\sigma = \prod(\sigma_i)) \right)$$

$$\text{SymmetricGroupAsTranspositions} := \forall_{\sigma \in S_n} \exists_{\Sigma \subseteq S_n} \left((\text{Transpositions}[\Sigma]) \wedge (\sigma = \prod(\sigma_i)) \right)$$

$$v\text{Function}[v(\sigma), \sigma, S_n] := v(\sigma) = n - |\text{DisjointFullCycles}[\Sigma]|$$

$$\text{signFunction}[\text{sign}(\sigma), \sigma, S_n] := \text{sign}(\sigma) = (-1)^{v(\sigma)}$$

$$\text{EvenPermutation}[\sigma, S_n] := \text{sign}(\sigma) = 1$$

$$\text{OddPermutation}[\sigma, S_n] := \text{sign}(\sigma) = -1$$

$$\text{TranspositionSigns} := \text{sign}(\tau\sigma) = -\text{sign}(\sigma)$$

$$\text{TranspositionSignsCorollary} := \text{sign}\left(\prod_{i=1}^r (\tau_i)\right) = (-1)^r$$

$$\text{SignProp} := \text{sign}(\sigma\pi) = \text{sign}(\sigma)\text{sign}(\pi)$$

$$\text{AlternatingGroup}[A_n, n] := A_n = \{\sigma \in S_n \mid \text{EvenPermutation}[\sigma, S_n]\}$$

$$\text{AlternatingGroupOrder} := o(A_n) = n!/2$$

2.5.3 Dihedral Group

$$\text{DihedralGroup}[D_n, *] := (D_n = \{a^r * b^s \mid (r \in \mathbb{N}_{0, n-1}) \wedge (s \in \mathbb{N}_{0, 1})\}) \wedge \left(\begin{array}{l} (a^p a^q = a^{(p+q)\%n}) \wedge \\ (a^p b a^q = a^{(p-q)\%n} b) \wedge \\ (a^p b a^q b = a^{(p-q)\%n}) \end{array} \right)$$

$$\text{DihedralGroupOrder} := o(D_n) = 2n$$

2.6 Lagrange's Theorem

$$\text{LeftCoset}[gH, g, H, G, *] := (\text{Subgroup}[H, G, *]) \wedge (g \in G) \wedge (gH = \{g * h \mid h \in H\})$$

$$\text{RightCoset}[Hg, g, H, G, *] := (\text{Subgroup}[H, G, *]) \wedge (g \in G) \wedge (Hg = \{h * g \mid h \in H\})$$

$$\text{CosetCardinality} := (\text{RightCoset}[Hg, g, H, G, *]) \implies (|H| = |Hg|)$$

$$(1) \text{ CancellationLaws} \blacksquare (h_1g = h_2g) \implies (h_1 = h_2) \blacksquare |H| = |Hg|$$

$$\text{CosetInduceEqRel} := \forall_{G,H} \left(((\text{Subgroup}[H, G, *]) \wedge (\sim = \{\langle a, b \rangle \mid a * b^{-1} \in H\})) \implies ((\text{EqRel}[\sim, G]) \wedge (\text{EqClass}[Ha, a, \sim, G])) \right)$$

$$(1) (a, b, c \in G) \implies \dots$$

$$(1.1) (\text{Subgroup}[H, G, *]) \implies (e \in H) \implies (a * a^{-1} \in H) \implies (a \sim a)$$

$$(1.2) (a \sim b) \implies (a * b^{-1} \in H) \implies (b * a^{-1} = (a * b^{-1})^{-1} \in H) \implies (b \sim a)$$

$$(1.3) ((a \sim b) \wedge (b \sim c)) \implies (a * b^{-1}, b * c^{-1} \in H) \implies (a * c^{-1} = (a * b^{-1}) * (b * c^{-1}) \in H) \blacksquare a \sim c$$

$$(2) \text{EqRel}[\sim, G]$$

$$(3) (a, x \in G) \implies \dots$$

$$(3.1) (x \sim a) \iff (x * a^{-1} \in H) \iff (\exists_{h \in H} (x * a^{-1} = h)) \iff (\exists_{h \in H} (x = h * a)) \iff (x \in Ha)$$

$$(4) [a] = \{x \in G \mid x \sim a\} = Ha$$

$$\text{CosetSet}[G : H, H, G, *] := (\text{Subgroup}[H, G, *]) \wedge (G : H = \{gH \mid g \in G\})$$

$$\text{IndexSubgroup}[|G : H|, H, G, *] := (\text{CosetSet}[G : H, H, G, *]) \wedge (|G : H| = |G : H|) \wedge (|G| = (|H|)(|G : H|))$$

$$\text{LagrangeTheorem} := \forall_{G,H} \left(((\text{Subgroup}[H, G, *]) \wedge (o(G), o(H) \in \mathbb{N})) \implies (o(G) = o(H)|G : H|) \wedge (\text{Divides}[o(H), o(G)]) \right)$$

$$(1) (\text{CosetInduceEqRel}) \wedge (\text{EqRelInducesPartition}) \wedge (\text{CosetCardinality}) \blacksquare (o(G) = o(H)|G : H|) \wedge (\text{Divides}[o(H), o(G)])$$

$$\text{OrderOrderElProp} := \forall_{g,G} \left(((\text{Order}[n, G, *]) \wedge (\text{OrderEl}[m, g, G, *])) \implies ((\text{Divides}[m, n]) \wedge (g^n = e)) \right)$$

$$(1) \text{CyclicSubgroup}[\langle g \rangle, g, G, *] \blacksquare \text{Order}[\langle g \rangle] = m$$

$$(2) (\text{LagrangeTheorem}) \wedge (\text{CyclicSubgroup}) \blacksquare \text{Divides}[\text{Order}[\langle g \rangle], \text{Order}[G]] \blacksquare \text{Divides}[m, n]$$

$$(3) g^n = g^{mk} = e^k = e$$

Any prime ordered cyclic group has no proper non-trivial subgroups and any non-identity element is a generator.

$$(1) \text{LagrangeTheorem} \blacksquare \text{Subgroups must have the order 1 or } p \blacksquare \text{Subgroups are trivial}$$

$$(2) \text{CyclicSubgroup of a non-identity element is } G \blacksquare \text{Non-identity elements generates } G$$

$$\left((\text{Subgroup}[H, G, *]) \wedge (\text{Subgroup}[K, G, *] \wedge (\text{RelPrime}(o(H), o(K))) \right) \implies (H \cap K = \{e\})$$

$$(1) (\text{LagrangeTheorem}) \wedge (\text{SubgroupIntersection}) \wedge (\text{RelPrime}(o(H), o(K))) \blacksquare H \cap K = \{e\}$$

2.7 Homomorphisms

$$\text{Homomorphism}[\phi, G, *, H, \diamond] := (\text{Function}[\phi, G, H]) \wedge \left(\forall_{a,b \in G} (\phi(a * b) = \phi(a) \diamond \phi(b)) \right)$$

$$\text{Monomorphism}[\phi, G, *, H, \diamond] := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (\text{Inj}[\phi, G, H])$$

$$\text{Epimorphism}[\phi, G, *, H, \diamond] := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (\text{Surj}[\phi, G, H])$$

$$\text{Isomorphism}[\phi, G, *, H, \diamond] := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (\text{Bij}[\phi, G, H])$$

$$\text{Isomorphic}[G, *, H, \diamond] := \exists_{\phi} (\text{Isomorphism}[\phi, G, *, H, \diamond]) \text{ ** Notation: } G \cong H \text{ **}$$

$$\text{Automorphism}[\phi, G, *] := \text{Isomorphism}[\phi, G, *, G, *]$$

$$\text{IdMapsId} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (\phi(e_G) = e_H)$$

$$(1) \phi(e_G) = \phi(e_G * e_G) = \phi(e_G) \diamond \phi(e_G) \blacksquare \phi(e_G) = \phi(e_G) \diamond \phi(e_G)$$

$$(2) e_H = \phi(e_G)^{-1} \diamond \phi(e_G) = \phi(e_G)^{-1} \diamond (\phi(e_G) \diamond \phi(e_G)) = \phi(e_G)$$

$$\text{InvMapsInv} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (\phi(g^{-1}) = \phi(g)^{-1})$$

$$(1) \quad IdMapsId \quad \blacksquare \quad e_H = \phi(e_G) = \phi(g * g^{-1}) = \phi(g) \diamond \phi(g^{-1}) \quad \blacksquare \quad e_H = \phi(g) \diamond \phi(g^{-1}) \quad \blacksquare \quad \phi(g^{-1}) = \phi(g)^{-1}$$

$$ExpMapsExp := (Homomorphism[\phi, G, *, H, \diamond]) \implies \left(\forall_{n \in \mathbb{N}^+} (\phi(g^n) = \phi(g)^n) \right)$$

$$(1) \quad \phi(g^1) = \phi(g) = \phi(g)^1 \quad \blacksquare \quad \phi(g^1) = \phi(g)^1$$

$$(2) \quad \left(\forall_{k \in \mathbb{N}^+} (\phi(g^k) = \phi(g)^k) \right) \implies \dots$$

$$(2.1) \quad \phi(g^{k+1}) = \phi(g^k * g) = \phi(g)^k \diamond \phi(g) = \phi(g)^{k+1} \quad \blacksquare \quad \phi(g^{k+1}) = \phi(g)^{k+1}$$

$$(3) \quad \left(\forall_{k \in \mathbb{N}^+} (\phi(g^k) = \phi(g)^k) \right) \implies \left(\phi(g^{k+1}) = \phi(g)^{k+1} \right)$$

$$(4) \quad \left(\phi(g^1) = \phi(g)^1 \right) \wedge \left(\left(\forall_{k \in \mathbb{N}^+} (\phi(g^k) = \phi(g)^k) \right) \implies \left(\phi(g^{k+1}) = \phi(g)^{k+1} \right) \right) \quad \blacksquare \quad \forall_{n \in \mathbb{N}^+} (\phi(g^n) = \phi(g)^n)$$

$$MapDivProp := ((Homomorphism[\phi, G, *, H, \diamond]) \wedge (Order[n, G, *])) \implies \left(\forall_{g \in G} \left((OrderEl[m, \phi(g), H, \diamond]) \implies (Divides[m, n]) \right) \right)$$

$$(1) \quad OrderOrderElProp \quad \blacksquare \quad g^n = e_G$$

$$(2) \quad (IdMapsId) \wedge (ExpMapsExp) \quad \blacksquare \quad e_G = \phi(g^n) = \phi(g)^n = e_H$$

$$(3) \quad OrderEl[m, \phi(g), H, \diamond] \quad \blacksquare \quad \phi(g)^m = e_H \quad \blacksquare \quad \phi(g)^m = e_H = e_H^k = \phi(g)^n$$

$$HomoCompInduceHomo := ((Homomorphism[\phi, G, *, H, \diamond]) \wedge (Homomorphism[\theta, H, \diamond, K, \square])) \implies (Homomorphism[\theta \circ \phi, G, *, K, \square])$$

$$(1) \quad FuncComp \quad \blacksquare \quad Func[\theta \circ \phi, G, K]$$

$$(2) \quad (g_1, g_2 \in G) \implies \dots$$

$$(2.1) \quad (Homomorphism[\phi, G, *, H, \diamond]) \wedge (Homomorphism[\theta, H, \diamond, K, \square]) \quad \blacksquare \quad \theta \circ \phi(g_1 * g_2) = \theta(\phi(g_1 * g_2)) = \dots$$

$$(2.2) \quad \dots \theta(\phi(g_1) \diamond \phi(g_2)) = \theta(\phi(g_1)) \square \theta(\phi(g_2)) = \theta \circ \phi(g_1) \square \theta \circ \phi(g_2) \quad \blacksquare \quad \theta \circ \phi(g_1 * g_2) = \theta \circ \phi(g_1) \square \theta \circ \phi(g_2)$$

$$(3) \quad (g_1, g_2 \in G) \implies (\theta \circ \phi(g_1 * g_2) = \theta \circ \phi(g_1) \square \theta \circ \phi(g_2)) \quad \blacksquare \quad \forall_{g_1, g_2 \in G} (\theta \circ \phi(g_1 * g_2) = \theta \circ \phi(g_1) \square \theta \circ \phi(g_2))$$

$$(4) \quad (Func[\theta \circ \phi, G, K]) \wedge \left(\forall_{g_1, g_2 \in G} (\theta \circ \phi(g_1 * g_2) = \theta \circ \phi(g_1) \square \theta \circ \phi(g_2)) \right) \quad \blacksquare \quad Homomorphism[\theta \circ \phi, G, *, K, \square]$$

$$IsoInvInduceIso := (Isomorphism[\phi, G, *, H, \diamond]) \implies (Isomorphism[\phi^{-1}, H, \diamond, G, *])$$

$$(1) \quad Isomorphism[\phi, G, *, H, \diamond] \quad \blacksquare \quad (Homomorphism[\phi, G, *, H, \diamond]) \wedge (Bij[\phi, G, H])$$

$$(2) \quad BijEquiv \quad \blacksquare \quad \exists_{\phi^{-1}} (Inv[\phi^{-1}, \phi, G, H])$$

$$(3) \quad \text{TODO continue}$$

$$KCycleGroupIsomorphic := \left(((CyclicGroup[G, *]) \wedge (CyclicGroup[H, \diamond]) \wedge (Order[n, G, *]) \wedge (Order[n, H, \diamond])) \implies \right. \\ \left. (Isomorphic[G, *, H, \diamond]) \right)$$

$$(1) \quad \left(\exists_{g \in G} (Generator[g, G, *]) \right) \wedge \left(\exists_{h \in H} (Generator[h, H, \diamond]) \right)$$

$$(2) \quad \phi := \{ \langle g^n, h^n \rangle \in (G \times H) \mid n \in \mathbb{Z} \}$$

$$(3) \quad (n_1, n_2 \in \mathbb{Z}) \implies \dots$$

$$(3.1) \quad (ExpModOrder) \wedge (Order[n, G, *]) \wedge (Order[n, H, \diamond]) \quad \blacksquare \quad (g^{n_1} = g^{n_2}) \iff (n_1 \equiv n_2 \pmod{n}) \iff (h^{n_1} = h^{n_2}) \iff \\ (\phi(g^{n_1}) = \phi(g^{n_2})) \quad \blacksquare \quad (g^{n_1} = g^{n_2}) \iff (\phi(g^{n_1}) = \phi(g^{n_2}))$$

$$(4) \quad (n_1, n_2 \in \mathbb{Z}) \implies \left((g^{n_1} = g^{n_2}) \iff (\phi(g^{n_1}) = \phi(g^{n_2})) \right) \dots$$

$$(5) \quad \dots (Func[\phi, G, H]) \wedge (Inj[\phi, G, H]) \wedge (Surj[\phi, G, H]) \quad \blacksquare \quad Bij[\phi, G, H]$$

$$(6) \quad \text{TODO: } Homomorphism[\phi, G, *, H, \diamond]$$

$$(7) \quad (Homomorphism[\phi, G, *, H, \diamond]) \wedge (Bij[\phi, G, H]) \quad \blacksquare \quad Isomorphism[\phi, G, *, H, \diamond]$$

$$(8) \quad \exists_{\phi} (Isomorphism[\phi, G, *, H, \diamond]) \quad \blacksquare \quad Isomorphic[G, *, H, \diamond]$$

2.8 Kernel and Image Homomorphisms

$$\text{Kernel}[ker_\phi, \phi, G, *, H, \diamond] := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge \left(ker_\phi = \{g \in G \mid \phi(g) = e_H\} \right)$$

$$\text{Image}[im_\phi, \phi, G, *, H, \diamond] := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge \left(im_\phi = \{\phi(g) \in H \mid g \in G\} \right)$$

$$\text{KernelSubgroupDomain} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (\text{Subgroup}[ker_\phi, G, *])$$

$$(1) \quad \text{Id Maps Id} \quad \blacksquare \phi(e_G) = e_H \quad \blacksquare e_G \in ker_\phi \quad \blacksquare ker_\phi \neq \emptyset$$

$$(2) \quad ker_\phi \subseteq G \quad \blacksquare \emptyset \neq ker_\phi \subseteq G$$

$$(3) \quad (a, b \in ker_\phi) \implies \dots$$

$$(3.1) \quad (\phi(a) = e_H) \wedge (\phi(b) = e_H) \quad \blacksquare \phi(a * b) = \phi(a) \diamond \phi(b) = e_H \diamond e_H = e_H \quad \blacksquare a * b \in ker_\phi$$

$$(4) \quad (a, b \in ker_\phi) \implies (a * b \in ker_\phi) \quad \blacksquare \forall_{a, b \in ker_\phi} (a * b \in ker_\phi)$$

$$(5) \quad (a \in ker_\phi) \implies \dots$$

$$(5.1) \quad \phi(a) = e_H$$

$$(5.2) \quad \text{Inv Maps Inv} \quad \blacksquare \phi(a^{-1}) = e_H^{-1} = e_H \quad \blacksquare a^{-1} \in ker_\phi$$

$$(6) \quad (a \in ker_\phi) \implies (a^{-1} \in ker_\phi) \quad \blacksquare \forall_{a \in ker_\phi} (a^{-1} \in ker_\phi)$$

$$(7) \quad (\text{SubgroupEquiv}) \wedge (\emptyset \neq ker_\phi \subseteq G) \wedge \left(\forall_{a, b \in ker_\phi} (a * b \in ker_\phi) \right) \wedge \left(\forall_{a \in ker_\phi} (a^{-1} \in ker_\phi) \right) \quad \blacksquare \text{Subgroup}[ker_\phi, G, *]$$

$$\text{ImageSubgroupCodomain} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (\text{Subgroup}[im_\phi, H, \diamond])$$

$$(1) \quad (\text{Id Maps Id}) \wedge (e_G \in G) \quad \blacksquare \phi(e_G) = e_H \in H \quad \blacksquare e_H \in im_\phi \quad \blacksquare \emptyset \neq im_\phi$$

$$(2) \quad im_\phi \subseteq H \quad \blacksquare \emptyset \neq im_\phi \subseteq H$$

$$(3) \quad (a, b \in im_\phi) \implies \dots$$

$$(3.1) \quad \left(\exists_{g_a \in G} (a = \phi(g_a)) \right) \wedge \left(\exists_{g_b \in G} (b = \phi(g_b)) \right)$$

$$(3.2) \quad (g_a * g_b \in G) \wedge (\phi(g_a * g_b) = \phi(g_a) * \phi(g_b) = a * b)$$

$$(3.3) \quad \exists_{g \in G} (a * b = \phi(g)) \quad \blacksquare a * b \in im_\phi$$

$$(4) \quad (a, b \in im_\phi) \implies (a * b \in im_\phi) \quad \blacksquare \forall_{a, b \in im_\phi} (a * b \in im_\phi)$$

$$(5) \quad (a \in im_\phi) \implies \dots$$

$$(5.1) \quad \exists_{g_a \in G} (a = \phi(g_a))$$

$$(5.2) \quad (g_a^{-1} \in G) \wedge (\text{Inv Maps Inv}) \quad \blacksquare \phi(g_a^{-1}) = \phi(g_a)^{-1} = a^{-1}$$

$$(5.3) \quad \exists_{g \in G} (a^{-1} = \phi(g)) \quad \blacksquare a^{-1} \in im_\phi$$

$$(6) \quad (a \in im_\phi) \implies (a^{-1} \in im_\phi) \quad \blacksquare \forall_{a \in im_\phi} (a^{-1} \in im_\phi)$$

$$(7) \quad (\text{SubgroupEquiv}) \wedge (\emptyset \neq im_\phi \subseteq H) \wedge \left(\forall_{a, b \in im_\phi} (a * b \in im_\phi) \right) \wedge \left(\forall_{a \in im_\phi} (a^{-1} \in im_\phi) \right) \quad \blacksquare \text{Subgroup}[im_\phi, H, \diamond]$$

$$\text{ImageCyclicIsCyclic} := ((\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (\text{CyclicGroup}[G, *])) \implies (\text{CyclicGroup}[im_\phi, \diamond])$$

$$(1) \quad \text{CyclicGroup}[G, *] \quad \blacksquare \exists_{g \in G} (\text{CyclicSubgroup}[G, g, G, *]) \quad \blacksquare \exists_{g_0 \in G} (G = \langle g_0 \rangle = \{g_0^n \mid n \in \mathbb{Z}\})$$

$$(2) \quad \text{Exp Maps Exp} \quad \blacksquare h \in im_\phi \iff \exists_{g \in G} (h = \phi(g)) \iff \exists_{n \in \mathbb{Z}} (h = \phi(g_0^n)) \iff \exists_{n \in \mathbb{Z}} (h = \phi(g_0)^n) \quad \blacksquare \text{Generator}[\phi(g_0), im_\phi, \diamond]$$

$$(3) \quad \exists_{h \in im_\phi} (\text{Generator}[h, im_\phi, \diamond]) \quad \blacksquare \text{CyclicGroup}[im_\phi, \diamond]$$

$$\text{MonomorphismEquiv} := (\text{Monomorphism}[\phi, G, *, H, \diamond]) \iff (ker_\phi = \{e_G\})$$

$$(1) \quad (\text{Monomorphism}[\phi, G, *, H, \diamond]) \implies \dots$$

$$(1.1) \quad \text{Id Maps Id} \quad \blacksquare \phi(e_G) = e_H \quad \blacksquare e_G \in ker_\phi \quad \blacksquare \{e_G\} \subseteq ker_\phi$$

$$(1.2) \quad (g \in ker_\phi) \implies \dots$$

$$(1.2.1) \quad (g \in ker_\phi) \wedge (\text{Id Maps Id}) \quad \blacksquare \phi(g) = e_H = \phi(e_G)$$

$$(1.2.2) \quad (\text{Injective}[\phi, G, H]) \wedge (\phi(g) = \phi(e_G)) \quad \blacksquare g = e_G \quad \blacksquare g \in \{e_G\}$$

$$(1.3) \quad (g \in ker_\phi) \implies (g \in \{e_G\}) \quad \blacksquare ker_\phi \subseteq \{e_G\}$$

$$(1.4) \quad (\{e_G\} \subseteq ker_\phi) \wedge (ker_\phi \subseteq \{e_G\}) \quad \blacksquare ker_\phi = \{e_G\}$$

$$(2) \text{ (Monomorphism}[\phi, G, *, H, \diamond]) \implies (ker_\phi = \{e_G\})$$

$$(3) (ker_\phi = \{e_G\}) \implies \dots$$

$$(3.1) \left((g_1, g_2 \in G) \wedge (\phi(g_1) = \phi(g_2)) \right) \implies \dots$$

$$(3.1.1) \text{ InvMapsInv } \blacksquare e_H = \phi(g_1) \diamond \phi(g_2)^{-1} = \phi(g_1) \diamond \phi(g_2^{-1}) = \phi(g_1 * g_2^{-1}) \blacksquare e_H = \phi(g_1 * g_2^{-1}) \blacksquare g_1 * g_2^{-1} \in ker_\phi$$

$$(3.1.2) (ker_\phi = \{e_G\}) \wedge (g_1 * g_2^{-1} \in ker_\phi) \blacksquare g_1 * g_2^{-1} = e_G \blacksquare g_1^{-1} = g_2^{-1}$$

$$(3.1.3) \text{ InvUniq } \blacksquare g_1 = g_2$$

$$(3.2) \left((g_1, g_2 \in G) \wedge (\phi(g_1) = \phi(g_2)) \right) \implies (g_1 = g_2) \blacksquare \text{Injective}[\phi, G, H] \blacksquare \text{Monomorphism}[\phi, G, *, H, \diamond]$$

$$(4) (ker_\phi = \{e_G\}) \implies (\text{Monomorphism}[\phi, G, *, H, \diamond])$$

$$(5) \left((\text{Monomorphism}[\phi, G, *, H, \diamond]) \implies (ker_\phi = \{e_G\}) \right) \wedge \left((ker_\phi = \{e_G\}) \implies (\text{Monomorphism}[\phi, G, *, H, \diamond]) \right)$$

$$(6) (\text{Monomorphism}[\phi, G, *, H, \diamond]) \iff (ker_\phi = \{e_G\})$$

$$\text{KerCountsMapSameEl} := ((\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (g \in G)) \implies ((ker_\phi)g = \{x \in G \mid \phi(x) = \phi(g)\})$$

$$(1) (x \in (ker_\phi)g) \implies \dots$$

$$(1.1) \exists_{K_x \in ker_\phi} (x = K_x * g) \blacksquare \phi(x) = \phi(K_x * g) = \phi(K_x) \diamond \phi(g) = e_H \diamond \phi(g) = \phi(g) \blacksquare \phi(x) = \phi(g)$$

$$(2) (x \in (ker_\phi)g) \implies (\phi(x) = \phi(g)) \blacksquare (ker_\phi)g \subseteq \{x \in G \mid \phi(x) = \phi(g)\}$$

$$(3) (\phi(x) = \phi(g)) \implies \dots$$

$$(3.1) e_H = \phi(x) \diamond \phi(g)^{-1} = \phi(x) \diamond \phi(g^{-1}) = \phi(x * g^{-1}) \blacksquare x * g^{-1} \in ker_\phi \blacksquare x \in (ker_\phi)g$$

$$(4) (\phi(x) = \phi(g)) \implies (x \in (ker_\phi)g) \blacksquare \{x \in G \mid \phi(x) = \phi(g)\} \subseteq (ker_\phi)g$$

$$(5) ((ker_\phi)g \subseteq \{x \in G \mid \phi(x) = \phi(g)\}) \wedge (\{x \in G \mid \phi(x) = \phi(g)\} \subseteq (ker_\phi)g) \blacksquare (ker_\phi)g = \{x \in G \mid \phi(x) = \phi(g)\}$$

$$\text{KerImPartitionsG} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (o(G) = o(ker_\phi)o(im_\phi))$$

$$(1) im_\phi \text{ forms equivalence classes of } G \text{ that maps to the same elements under } \phi$$

$$(2) (\text{KerCountsMapSameEl}) \wedge (\text{CosetCardinality}) \text{ counts the number of same element mappings / multiplicity for each pre-image class}$$

$$(3) o(G) = o(ker_\phi)o(im_\phi)$$

$$(4) \text{ TODO: formalize}$$

$$\text{ImageDividesGH} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies \left((\text{Divides}[o(im_\phi), o(G)]) \wedge (\text{Divides}[o(im_\phi), o(H)]) \right)$$

$$(1) \text{ KerImPartitionsG } \blacksquare \text{Divides}[r, o(G)]$$

$$(2) (\text{LagrangeTheorem}) \wedge (\text{ImageSubgroupCodomain}) \blacksquare \text{Divides}[r, o(H)]$$

2.9 Conjugacy

$$\text{Conjugate}[\sim^*, a, b, G, *] := (\text{Group}[G, *]) \wedge (a, b \in G) \wedge \left(\exists_{c \in G} (b = c^{-1} * a * c) \right)$$

$$\text{ConjugateEqRel} := \text{EqRel}[\sim^*, G]$$

$$(1) (a, b, c \in G) \implies \dots$$

$$(1.1) a = e^{-1} * a * e \blacksquare a \sim^* a$$

$$(1.2) (a \sim^* b) \implies (b = x_b^{-1} * a * x_b) \implies (x_b * b * x_b^{-1} = a) \implies (b \sim^* a)$$

$$(1.3) ((a \sim^* b) \wedge (b \sim^* c)) \implies ((b = x_b^{-1} * a * x_b) \wedge (c = x_c^{-1} * b * x_c)) \implies \dots$$

$$(1.4) \dots (c = x_c^{-1} * x_b^{-1} * a * x_b * x_c = (x_b * x_c)^{-1} * a * (x_b * x_c)) \blacksquare a \sim^* c$$

$$(2) \text{ EqRel}[\sim^*, G]$$

$$\text{ConjugacyClass}[C_g, g, G, *] := (\text{Group}[G, *]) \wedge (g \in G) \wedge (\text{EqClass}[C_g, g, \sim^*, G])$$

$$\text{ConjugacyClassEquiv} := (\text{ConjugacyClass}[C_g, g, G, *]) \iff \left(\forall_{x \in G} \left((x \in C_g) \iff \left(\exists_{c \in G} (x = c^{-1}gc) \right) \right) \right)$$

(1) TODO: by definition

$$\text{ConjugacyCenter} := (g \in G) \implies \left((C_g = \{g\}) \iff (g \in Z(G)) \right)$$

(1) $(C_g = \{g\}) \implies \dots$

(1.1) $(x \in G) \implies \dots$

$$(1.1.1) \quad (\text{ConjugacyClass}[C_g, g, G, *]) \wedge (\text{ConjugacyClassEquiv}) \wedge (x \in G) \quad \blacksquare \quad x^{-1}gx \in C_g$$

$$(1.1.2) \quad (C_g = \{g\}) \wedge (x^{-1}gx \in C_g) \quad \blacksquare \quad x^{-1}gx = g \quad \blacksquare \quad gx = xg$$

$$(1.2) \quad (x \in G) \implies (gx = xg) \quad \blacksquare \quad \forall_{x \in G} (gx = xg) \quad \blacksquare \quad g \in Z(G)$$

(2) $(C_g = \{g\}) \implies (g \in Z(G))$

(3) $(g \in Z(G)) \implies \dots$

$$(3.1) \quad (g \in Z(G)) \wedge (\text{Group}[G, *]) \quad \blacksquare \quad (\forall_{c \in G} (gc = cg)) \wedge (\exists_e (e \in G))$$

(3.2) $(x \in G) \implies \dots$

$$(3.2.1) \quad (\forall_{c \in G} (gc = cg)) \wedge (\exists_e (e \in G)) \quad \blacksquare \quad \left(\exists_{c \in G} (x = c^{-1}gc) \right) \iff \left(\exists_{c \in G} (x = c^{-1}gc = c^{-1}cg = g) \right) \iff (x = g) \iff (x \in \{g\})$$

$$(3.3) \quad (x \in G) \implies \left(\left(\exists_{c \in G} (x = c^{-1}gc) \right) \iff (x \in \{g\}) \right) \quad \blacksquare \quad \forall_{x \in G} \left((x \in \{g\}) \iff \left(\exists_{c \in G} (x = c^{-1}gc) \right) \right)$$

$$(3.4) \quad (\text{ConjugacyClassEquiv}) \wedge \left(\forall_{x \in G} \left((x \in \{g\}) \iff \left(\exists_{c \in G} (x = c^{-1}gc) \right) \right) \right) \quad \blacksquare \quad C_g = \{g\}$$

(4) $(g \in Z(G)) \implies (C_g = \{g\})$

(5) $(C_g = \{g\}) \iff (g \in Z(G))$

$$\text{ConjugacyAbelian} := \left(\forall_{g \in G} (C_g = \{g\}) \right) \iff (\text{AbelianGroup}[G, *])$$

$$(1) \quad \text{ConjugacyCenter} \quad \blacksquare \quad \left(\forall_{g \in G} (C_g = \{g\}) \right) \iff \left(\forall_{g \in G} (g \in Z(G)) \right) \iff (\text{AbelianGroup}[G, *])$$

$$\text{ConjugateExp} := \forall_{n \in \mathbb{N}^+} \left((x^{-1}gx)^n = x^{-1}g^n x \right)$$

(1) $(n = 1) \implies \dots$

$$(1.1) \quad (x^{-1}gx)^n = (x^{-1}gx)^1 = x^{-1}g^1 x = x^{-1}g^1 x \quad \blacksquare \quad (x^{-1}gx)^n = x^{-1}g^n x$$

(2) $(n = 1) \implies \left((x^{-1}gx)^n = x^{-1}g^n x \right)$

$$(3) \quad \left((n > 1) \wedge \left(\forall_{m \in \mathbb{N}^+} \left((m \leq n) \implies \left((x^{-1}gx)^m = x^{-1}g^m x \right) \right) \right) \right) \implies \dots$$

$$(3.1) \quad (x^{-1}gx)^{n+1} = (x^{-1}gx)^n * (x^{-1}gx) = (x^{-1}g^n x) * (x^{-1}gx) = x^{-1}g^{n+1} x \quad \blacksquare \quad (x^{-1}gx)^{n+1} = x^{-1}g^{n+1} x$$

$$(4) \quad \left((n > 1) \wedge \left(\forall_{m \in \mathbb{N}^+} \left((m \leq n) \implies \left((x^{-1}gx)^m = x^{-1}g^m x \right) \right) \right) \right) \implies \left((x^{-1}gx)^{n+1} = x^{-1}g^{n+1} x \right)$$

(5) $\forall_{n \in \mathbb{N}^+} \left((x^{-1}gx)^n = x^{-1}g^n x \right)$

$$\text{ConjugateOrder} := ((g_1, g_2 \in G) \wedge (g_1 \sim^* g_2)) \implies (o(g_1) = o(g_2))$$

(1) $\exists_{c \in G} (g_2 = c^{-1}g_1 c)$

$$(2) \quad \text{ConjugateExp} \quad \blacksquare \quad e = g_2^{o(g_2)} = (c^{-1}g_1 c)^{o(g_2)} = c^{-1}g_1^{o(g_2)} c \quad \blacksquare \quad e = c^{-1}g_1^{o(g_2)} c \quad \blacksquare \quad g_1^{o(g_2)} = e$$

(3) $\text{ExpModOrderCorollary} \quad \blacksquare \quad \text{Divides}[o(g_2), o(g_1)]$

$$(4) \quad \text{ConjugateExp} \quad \blacksquare \quad e = g_1^{o(g_1)} = (c g_2 c^{-1})^{o(g_1)} = c g_2^{o(g_1)} c^{-1} \quad \blacksquare \quad e = c g_2^{o(g_1)} c^{-1} \quad \blacksquare \quad g_2^{o(g_1)} = e$$

$$(5) \quad \text{ExpModOrderCorollary} \quad \blacksquare \quad \text{Divides}[o(g_1), o(g_2)]$$

$$(6) \quad (\text{Divides}[o(g_2), o(g_1)]) \wedge (\text{Divides}[o(g_1), o(g_2)]) \wedge (g_1, g_2 \in \mathbb{N}^+) \quad \blacksquare \quad o(g_1) = o(g_2)$$

$$(7) \quad \text{=====}$$

$$(8) \quad \exists_{c \in G} (g_2 = c^{-1} g_1 c) \quad \blacksquare \quad e = g_2^{o(g_2)} = (c^{-1} g_1 c)^{o(g_2)} = c^{-1} g_1^{o(g_2)} c \quad \blacksquare \quad e = c^{-1} g_1^{o(g_2)} c \quad \blacksquare \quad g_1^{o(g_2)} = e$$

$$(9) \quad (m \in \mathbb{Z}^+) \wedge (m < o(g_2)) \implies \dots$$

$$(9.1) \quad e \neq g_2^m = (c^{-1} g_1 c)^m = c^{-1} g_1^m c \quad \blacksquare \quad e \neq c^{-1} g_1^m c \quad \blacksquare \quad e = c * e * c^{-1} \neq g_1^m \quad \blacksquare \quad g_1^m \neq e$$

$$(10) \quad (m < o(g_2)) \implies (e \neq g_1^m) \quad \blacksquare \quad \forall_{m \in \mathbb{Z}^+} \left((m < o(g_2)) \implies (g_1^m \neq e) \right)$$

$$(11) \quad (g_1^{o(g_2)} = e) \wedge \left(\forall_{m \in \mathbb{Z}^+} \left((m < o(g_2)) \implies (g_1^m \neq e) \right) \right) \quad \blacksquare \quad o(g_1) = o(g_2)$$

$$\text{CentralizerConjugateCosets} := \forall_{c, g, h \in G} \left((h = c^{-1} g c) \implies (C(h) = c^{-1} C(g) c) \right)$$

$$(1) \quad (c^{-1} a c \in c^{-1} C(g) c) \implies \dots$$

$$(1.1) \quad a \in C(g) \quad \blacksquare \quad a g = g a$$

$$(1.2) \quad (c^{-1} a c) h = (c^{-1} a c) (c^{-1} g c) = c^{-1} a g c = c^{-1} g a c = c^{-1} g (c c^{-1}) a c = h (c^{-1} a c) \quad \blacksquare \quad (c^{-1} a c) h = h (c^{-1} a c) \quad \blacksquare \quad c^{-1} a c \in C(h)$$

$$(2) \quad (c^{-1} a c \in c^{-1} C(g) c) \implies (c^{-1} a c \in C(h)) \quad \blacksquare \quad c^{-1} C(g) c \subseteq C(h)$$

$$(3) \quad (a \in C(h)) \implies \dots$$

$$(3.1) \quad a \in C(h) \quad \blacksquare \quad a h = h a \quad \blacksquare \quad a (c^{-1} g c) = (c^{-1} g c) a$$

$$(3.2) \quad (c a c^{-1}) g = g (c a c^{-1}) \quad \blacksquare \quad c a c^{-1} \in C(g) \quad \blacksquare \quad a \in c^{-1} C(g) c$$

$$(4) \quad (a \in C(h)) \implies (a \in c^{-1} C(g) c) \quad \blacksquare \quad C(h) \subseteq c^{-1} C(g) c$$

$$(5) \quad (c^{-1} C(g) c \subseteq C(h)) \wedge (C(h) \subseteq c^{-1} C(g) c) \quad \blacksquare \quad C(h) = c^{-1} C(g) c$$

$$\text{ConjugatesMultiplicity} := (g \in G) \implies (o(G) = o(C(g)) | C_g|)$$

$$(1) \quad \phi := \{ \langle a^{-1} g a, C(g) a \rangle \in (C_g \times G : C(g)) \mid a \in G \}$$

$$(2) \quad (x, y \in G) \implies \dots$$

$$(2.1) \quad (x^{-1} g x = y^{-1} g y) \iff (g x = x y^{-1} g y) \iff (g (x y^{-1}) = (x y^{-1}) g) \iff \dots$$

$$(2.2) \quad \dots (x y^{-1} \in C(g)) \iff (C(g) (x y^{-1}) = C(g)) \iff (C(g) x = C(g) y)$$

$$(3) \quad (x, y \in G) \implies ((x^{-1} g x = y^{-1} g y) \iff (C(g) x = C(g) y)) \dots$$

$$(4) \quad \dots (Func[\phi, C_g, G : C(g)]) \wedge (Inj[\phi, C_g, G : C(g)]) \wedge (Surj[\phi, C_g, G : C(g)]) \quad \blacksquare \quad Bij[\phi, C_g, G : C(g)]$$

$$(5) \quad \exists_{\phi} (Bij[\phi, C_g, G : C(g)]) \quad \blacksquare \quad |C_g| = |G : C(g)|$$

$$(6) \quad (\text{LagrangeTheorem}) \wedge (\text{SubgroupCenter}) \wedge (|C_g| = |G : C(g)|) \quad \blacksquare \quad o(G) = o(C(g)) |G : C(g)| \quad \blacksquare \quad o(G) = o(C(g)) |C_g|$$

2.10 Normal Subgroups

$$\text{NormalSubgroup}[H, G, *] := (\text{Subgroup}[H, G, *]) \wedge \left(\forall_{h \in H} \forall_{g \in G} (g^{-1} h g \in H) \right)$$

$$\text{CenterNormalSubgroup} := \text{NormalSubgroup}[Z(G), G, *]$$

$$(1) \quad \text{SubgroupCenter} \quad \blacksquare \quad \text{Subgroup}[Z(G), G, *]$$

$$(2) \quad ((h \in Z(G)) \wedge (g \in G)) \implies \dots$$

$$(2.1) \quad h g = g h \quad \blacksquare \quad g^{-1} h g = h \in Z(G) \quad \blacksquare \quad g^{-1} h g \in Z(G)$$

$$(3) \quad \left((h \in Z(G)) \wedge (g \in G) \right) \implies \left(g^{-1}hg \in Z(G) \right) \quad \blacksquare \quad \forall_{h \in Z(G)} \forall_{g \in G} \left(g^{-1}hg \in Z(G) \right)$$

$$(4) \quad \left(\text{Subgroup}[Z(G), G, *] \wedge \left(\forall_{h \in Z(G)} \forall_{g \in G} \left(g^{-1}hg \in Z(G) \right) \right) \right) \quad \blacksquare \quad \text{NormalSubgroup}[Z(G), G, *]$$

$$\text{UnionConjugacyClassesNormalSubgroup} := (\text{NormalSubgroup}[H, G, *]) \implies \left(H = \bigcup_{z \in H} (C_z) \right)$$

$$(1) \quad (\text{NormalSubgroup}[H, G, *]) \implies \dots$$

$$(1.1) \quad \text{NormalSubgroup}[H, G, *] \quad \blacksquare \quad \forall_{x \in H} \forall_{g \in G} (g^{-1}xg \in H)$$

$$(1.2) \quad ((x \in H) \wedge (y \in C_x)) \implies \dots$$

$$(1.2.1) \quad \text{ConjugacyClassEquiv} \quad \blacksquare \quad \exists_{c \in G} (y = c^{-1}xc)$$

$$(1.2.2) \quad \left(\forall_{x \in H} \forall_{g \in G} (g^{-1}xg \in H) \right) \wedge (x \in H) \wedge (c \in G) \quad \blacksquare \quad y \in H$$

$$(1.3) \quad ((x \in H) \wedge (y \in C_x)) \implies (y \in H) \quad \blacksquare \quad \forall_{x \in H} (C_x \subseteq H)$$

$$(1.4) \quad \forall_{x \in H} (C_x \subseteq H) \quad \blacksquare \quad \forall_{x \in H} \forall_y (y \in C_x \implies y \in H) \quad \blacksquare \quad \forall_{x \in H} \forall_y (y \notin H \implies y \notin C_x)$$

$$(1.5) \quad (b \in H) \implies \left(b \in C_b \subseteq \bigcup_{z \in H} (C_z) \right) \quad \blacksquare \quad (b \in H) \implies \left(b \in \bigcup_{z \in H} (C_z) \right)$$

$$(1.6) \quad (b \notin H) \implies (\forall_{a \in H} (b \notin C_a)) \implies \left(b \notin \bigcup_{z \in H} (C_z) \right) \quad \blacksquare \quad (b \notin H) \implies \left(b \notin \bigcup_{z \in H} (C_z) \right)$$

$$(1.7) \quad \left((b \in H) \implies \left(b \in \bigcup_{z \in H} (C_z) \right) \right) \wedge \left((b \notin H) \implies \left(b \notin \bigcup_{z \in H} (C_z) \right) \right) \quad \blacksquare \quad (b \in H) \iff \left(b \in \bigcup_{z \in H} (C_z) \right)$$

$$(1.8) \quad \forall_b \left((b \in H) \iff \left(b \in \bigcup_{z \in H} (C_z) \right) \right) \quad \blacksquare \quad H = \bigcup_{z \in H} (C_z)$$

$$(2) \quad (\text{NormalSubgroup}[H, G, *]) \implies \left(H = \bigcup_{z \in H} (C_z) \right)$$

$$\text{NormalSubgroupCosetEquiv} := (\text{NormalSubgroup}[H, G, *]) \iff \left(\forall_{g \in G} (gH = Hg) \right)$$

$$(1) \quad \text{CosetCardinality} \quad \blacksquare \quad \forall_{g \in G} (|Hg| = |gH|) \quad \blacksquare \quad \left(\forall_{g \in G} ((Hg \subseteq gH) \iff (Hg = gH)) \right)$$

$$(2) \quad \left(\forall_{g \in G} ((Hg \subseteq gH) \iff (Hg = gH)) \right) \quad \blacksquare \quad (\text{NormalSubgroup}[H, G, *]) \iff \left(\forall_{h \in H} \forall_{g \in G} (g^{-1}hg \in H) \right) \iff \dots$$

$$(3) \quad \dots \left(\forall_{h \in H} \forall_{g \in G} (hg \in gH) \right) \iff \left(\forall_{g \in G} (Hg \subseteq gH) \right) \iff \left(\forall_{g \in G} (Hg = gH) \right)$$

$$\text{NormalSubgroupIndexEquiv} := (\text{NormalSubgroup}[H, G, *]) \iff (\text{IndexSubgroup}[2, H, G, *])$$

$$(1) \quad \text{NormalSubgroupCosetEquiv} \quad \blacksquare \quad (\text{IndexSubgroup}[2, H, G, *]) \iff \left(\forall_{g \in G} (gH = Hg) \right) \iff (\text{NormalSubgroup}[H, G, *])$$

$$\text{KerInduceNormalSubgroup} := (\text{Homomorphism}[\phi, G, *, H, \diamond]) \implies (\text{NormalSubgroup}[\ker \phi, G, *])$$

$$(1) \quad \text{KernelSubgroupDomain} \quad \blacksquare \quad \text{Subgroup}[\ker \phi, G, *]$$

$$(2) \quad ((h \in \ker \phi) \wedge (g \in G)) \implies \dots$$

$$(2.1) \quad h \in \ker \phi \quad \blacksquare \quad \phi(h) = e_H$$

$$(2.2) \quad (\text{Homomorphism}[\phi, G, *, H, \diamond]) \wedge (\text{InvMapsInv}) \quad \blacksquare \quad \phi(g^{-1} * h * g) = \phi(g^{-1}) \diamond \phi(h) \diamond \phi(g) = \phi(g)^{-1} \diamond e_H \diamond \phi(g) = e_H$$

$$(2.3) \quad \phi(g^{-1} * h * g) = e_H \quad \blacksquare \quad g^{-1}hg \in \ker \phi$$

$$(3) \quad ((h \in \ker \phi) \wedge (g \in G)) \implies (g^{-1}hg \in \ker \phi) \quad \blacksquare \quad \forall_{h \in \ker \phi} \forall_{g \in G} (g^{-1}hg \in \ker \phi)$$

$$(4) \quad (\text{Subgroup}[\ker \phi, G, *]) \wedge \left(\forall_{h \in \ker \phi} \forall_{g \in G} (g^{-1}hg \in \ker \phi) \right) \quad \blacksquare \quad \text{NormalSubgroup}[\ker \phi, G, *]$$

2.11 Quotient Groups

$$\text{QuotientSet}[G/H, H, G, *] := (\text{Subgroup}[H, G, *]) \wedge (G/H = \{Hg \mid g \in G\})$$

$$\text{CosetMul}[\bar{*}, H, G, *] := (\text{Subgroup}[H, G, *]) \wedge \left(\forall_{Hx, Hy \in G/H} (Hx \bar{*} Hy = \{h_1 x h_2 y \mid h_1, h_2 \in H\}) \right)$$

$$\text{SubsetMul}[\bar{\times}, G, *] := (\text{Group}[G, *]) \wedge \left(\forall_{A, B \subseteq G} (A \bar{\times} B = \{a * b \mid (a \in A) \wedge (b \in B)\}) \right)$$

$$\text{QuotientGroupLemma} := ((\text{NormalSubgroup}[H, G, *]) \wedge (x, y, z \in G)) \implies \left(\left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \iff \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \right)$$

$$(1) \quad \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \implies \dots$$

$$(1.1) \quad (\text{Group}[G, *]) \wedge (x \in G) \quad \blacksquare \quad x^{-1} \in G$$

$$(1.2) \quad (\text{NormalSubgroup}[H, G, *]) \wedge (x^{-1} \in G) \wedge (h_2 \in H) \quad \blacksquare \quad (x^{-1})^{-1} h_2 x^{-1} = x h_2 x^{-1} \in H$$

$$(1.3) \quad (\text{Group}[H, *]) \wedge (h_1, x h_2 x^{-1} \in H) \quad \blacksquare \quad h_1 x h_2 x^{-1} \in H$$

$$(1.4) \quad (h_1 x h_2 x^{-1})(xy) = h_1 x h_2 y = z \quad \blacksquare \quad (h_1 x h_2 x^{-1})(xy) = z$$

$$(1.5) \quad (h_1 x h_2 x^{-1} \in H) \wedge \left((h_1 x h_2 x^{-1})(xy) = z \right) \quad \blacksquare \quad \exists_{h_3 \in H} (z = h_3 x y)$$

$$(2) \quad \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \implies \left(\exists_{h_3 \in H} (z = h_3 x y) \right)$$

$$(3) \quad \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \implies \dots$$

$$(3.1) \quad (\text{NormalSubgroup}[H, G, *]) \wedge (x \in G) \wedge (h_3 \in H) \quad \blacksquare \quad x^{-1} h_3 x \in H$$

$$(3.2) \quad \text{Group}[H, *] \quad \blacksquare \quad e \in H$$

$$(3.3) \quad (e)x(x^{-1} h_3 x)y = h_3 x y = z \quad \blacksquare \quad (e)x(x^{-1} h_3 x)y = z$$

$$(3.4) \quad (x^{-1} h_3 x, e \in H) \wedge \left((e)x(x^{-1} h_3 x)y = h_3 x y = z \right) \quad \blacksquare \quad \exists_{h_1, h_2 \in H} (z = h_1 x h_2 y)$$

$$(4) \quad \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \implies \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right)$$

$$(5) \quad \left(\left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \implies \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \right) \wedge \left(\left(\exists_{h_3 \in H} (z = h_3 x y) \right) \implies \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \right)$$

$$(6) \quad \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \iff \left(\exists_{h_3 \in H} (z = h_3 x y) \right)$$

$$\text{QuotientGroupThm} := \left(\begin{array}{l} ((\text{NormalSubgroup}[H, G, *]) \wedge (\text{QuotientSet}[G/H, H, G, *]) \wedge (\text{CosetMul}[\bar{*}, x, y, H, G, *])) \implies \\ (\text{Group}[G/H, \bar{*}]) \end{array} \right)$$

$$(1) \quad (Hx, Hy \in G/H) \implies \dots$$

$$(1.1) \quad (\text{NormalSubgroup}[H, G, *]) \wedge (\text{QuotientGroupLemma}) \quad \blacksquare \quad \forall_{x, y, z \in G} \left(\left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \iff \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \right)$$

$$(1.2) \quad (z \in Hx \bar{*} Hy) \iff \left(\exists_{h_1, h_2 \in H} (z = h_1 x h_2 y) \right) \iff \left(\exists_{h_3 \in H} (z = h_3 x y) \right) \iff (z \in Hxy) \quad \blacksquare \quad Hx \bar{*} Hy = Hxy$$

$$(1.3) \quad (\text{Group}[G, *]) \wedge (x, y \in G) \quad \blacksquare \quad xy \in G \quad \blacksquare \quad Hxy \in G/H$$

$$(1.4) \quad (Hx \bar{*} Hy = Hxy) \wedge (Hxy \in G/H) \quad \blacksquare \quad \exists!_{Hxy \in G/H} (Hx \bar{*} Hy = Hxy)$$

$$(2) \quad (Hx, Hy \in G/H) \implies \left(\exists!_{Hxy \in G/H} (Hx \bar{*} Hy = Hxy) \right) \quad \blacksquare \quad \text{Func}[\bar{*}, G/H, G/H]$$

$$(3) \quad (Hx, Hy, Hz \in G/H) \implies \dots$$

$$(3.1) \quad (Hx \bar{*} Hy) \bar{*} Hz = Hxy \bar{*} Hz = Hxyz = Hx \bar{*} Hyz = Hx \bar{*} (Hy \bar{*} Hz) \quad \blacksquare \quad (Hx \bar{*} Hy) \bar{*} Hz = Hx \bar{*} (Hy \bar{*} Hz)$$

$$(4) \quad (Hx, Hy, Hz \in G/H) \implies \left((Hx \bar{*} Hy) \bar{*} Hz = Hx \bar{*} (Hy \bar{*} Hz) \right) \quad \blacksquare \quad \forall_{a, b, c \in G/H} ((a \bar{*} b) \bar{*} c = a \bar{*} (b \bar{*} c))$$

$$(5) \quad (He \in G/H) \wedge \left(\forall_{Hx \in G/H} (Hx \bar{*} He = Hxe = Hx = Hxe = He \bar{*} Hx) \right) \quad \blacksquare \quad \exists_{e \in G/H} \forall_{a \in G/H} (a \bar{*} e = a = e \bar{*} a)$$

$$(6) \quad (Hx \in G/H) \implies \dots$$

$$(6.1) \quad x \in G \quad \blacksquare \quad x^{-1} \in G \quad \blacksquare \quad Hx^{-1} \in G/H$$

$$(6.2) \quad Hx \bar{*} Hx^{-1} = Hxx^{-1} = He = Hx^{-1}x = Hx^{-1} \bar{*} Hx \quad \blacksquare \quad Hx \bar{*} Hx^{-1} = He = Hx^{-1} \bar{*} Hx$$

$$(6.3) \quad (Hx^{-1} \in G/H) \wedge (Hx \bar{*} Hx^{-1} = He = Hx^{-1} \bar{*} Hx) \quad \blacksquare \quad \exists_{Hx^{-1} \in G/H} (Hx \bar{*} Hx^{-1} = He = Hx^{-1} \bar{*} Hx)$$

$$(7) \quad (Hx \in G/H) \implies \left(\exists_{Hx^{-1} \in G/H} (Hx \bar{*} Hx^{-1} = He = Hx^{-1} \bar{*} Hx) \right) \blacksquare \forall_{a \in G/H} \exists_{a^{-1} \in G/H} (a \bar{*} a^{-1} = e = a^{-1} \bar{*} a)$$

$$(8) \quad (Func[\bar{*}, G/H, G/H]) \wedge \left(\forall_{a,b,c \in G/H} ((a \bar{*} b) \bar{*} c = a \bar{*} (b \bar{*} c)) \right) \wedge \left(\exists_{e \in G/H} \forall_{a \in G/H} (a \bar{*} e = a = e \bar{*} a) \right) \wedge \dots$$

$$(9) \quad \dots \left(\forall_{a \in G/H} \exists_{a^{-1} \in G/H} (a \bar{*} a^{-1} = e = a^{-1} \bar{*} a) \right) \blacksquare Group[G/H, \bar{*}]$$

$$QuotientMap[\bar{\phi}, H, G, *] := (NormalSubgroup[H, G, *]) \wedge (Func[\bar{\phi}, G, *, G/H, \bar{*}]) \wedge \left(\forall_{g \in G} (\bar{\phi}(g) = Hg) \right)$$

$$QuotientMapHomo := (QuotientMap[\bar{\phi}, H, G, *]) \implies (Homomorphism[\bar{\phi}, G, *, G/H, \bar{*}])$$

$$(1) \quad QuotientMap[\bar{\phi}, H, G, *] \blacksquare Func[\bar{\phi}, G, *, G/H, \bar{*}]$$

$$(2) \quad (x, y \in G) \implies \dots$$

$$(2.1) \quad \bar{\phi}(x * y) = Hxy = Hx \bar{*} Hy = \bar{\phi}(x) \bar{*} \bar{\phi}(y) \blacksquare \bar{\phi}(x * y) = \bar{\phi}(x) \bar{*} \bar{\phi}(y)$$

$$(3) \quad (x, y \in G) \implies (\bar{\phi}(x * y) = \bar{\phi}(x) \bar{*} \bar{\phi}(y)) \blacksquare \forall_{x,y \in G} (\bar{\phi}(x) \bar{*} \bar{\phi}(y))$$

$$(4) \quad (Func[\bar{\phi}, G, *, G/H, \bar{*}]) \wedge \left(\forall_{x,y \in G} (\bar{\phi}(x) \bar{*} \bar{\phi}(y)) \right) \blacksquare Homomorphism[\bar{\phi}, G, *, G/H, \bar{*}]$$

$$QuotientMapKer := (QuotientMap[\bar{\phi}, H, G, *]) \implies (ker_{\bar{\phi}} = H)$$

$$(1) \quad Group[H, *] \blacksquare ker_{\bar{\phi}} = \{x \in G \mid \bar{\phi}(x) = He\} = \{x \in G \mid Hx = H\} = H$$

$$FirstIsoThm := (Homomorphism[\phi, G, *, H, \diamond]) \implies (Isomorphic[G/ker_{\phi}, \bar{*}, im_{\phi}, \diamond])$$

$$(1) \quad \psi := \{\langle ker_{\phi}g, \phi(g) \rangle \in (G/ker_{\phi} \times im_{\phi}) \mid g \in G\}$$

$$(2) \quad (g, h \in G) \implies \dots$$

$$(2.1) \quad (ker_{\phi}g = ker_{\phi}h) \iff (ker_{\phi}gh^{-1} = ker_{\phi}) \iff (gh^{-1} \in ker_{\phi}) \iff (\phi(gh^{-1}) = e_H) \iff \dots$$

$$(2.2) \quad \dots \left(e_H = \phi(g) \diamond \phi(h^{-1}) = \phi(g) \diamond \phi(h)^{-1} \right) \iff (\phi(g) = \phi(h)) \blacksquare (ker_{\phi}g = ker_{\phi}h) \iff (\phi(g) = \phi(h))$$

$$(3) \quad (g, h \in G) \implies \left((ker_{\phi}g = ker_{\phi}h) \iff (\phi(g) = \phi(h)) \right) \dots$$

$$(4) \quad \dots (Func[\psi, G/ker_{\phi}, im_{\phi}]) \wedge (Inj[\psi, G/ker_{\phi}, im_{\phi}]) \wedge (Surj[\psi, G/ker_{\phi}, im_{\phi}]) \blacksquare Bij[\psi, G/ker_{\phi}, im_{\phi}]$$

$$(5) \quad (ker_{\phi}g, ker_{\phi}h \in G/ker_{\phi}) \implies \dots$$

$$(5.1) \quad \psi(ker_{\phi}g \bar{*} ker_{\phi}h) = \psi(ker_{\phi}gh) = \phi(g * h) = \phi(g) \diamond \phi(h) = \psi(ker_{\phi}g) \diamond \psi(ker_{\phi}h) \blacksquare \psi(ker_{\phi}g \bar{*} ker_{\phi}h) = \psi(ker_{\phi}g) \diamond \psi(ker_{\phi}h)$$

$$(6) \quad (ker_{\phi}g, ker_{\phi}h \in G/ker_{\phi}) \implies \left(\psi(ker_{\phi}g \bar{*} ker_{\phi}h) = \psi(ker_{\phi}g) \diamond \psi(ker_{\phi}h) \right) \dots$$

$$(7) \quad \dots \forall_{ker_{\phi}g, ker_{\phi}h \in G/ker_{\phi}} \left(\psi(ker_{\phi}g \bar{*} ker_{\phi}h) = \psi(ker_{\phi}g) \diamond \psi(ker_{\phi}h) \right)$$

$$(8) \quad (Bij[\psi, G/ker_{\phi}, im_{\phi}]) \wedge \left(\forall_{ker_{\phi}g, ker_{\phi}h \in G/ker_{\phi}} \left(\psi(ker_{\phi}g \bar{*} ker_{\phi}h) = \psi(ker_{\phi}g) \diamond \psi(ker_{\phi}h) \right) \right) \blacksquare Isomorphism[\psi, G/ker_{\phi}, \bar{*}, im_{\phi}, \diamond]$$

$$(9) \quad \exists_{\psi} (Isomorphism[\psi, G/ker_{\phi}, \bar{*}, im_{\phi}, \diamond]) \blacksquare Isomorphic[G/ker_{\phi}, \bar{*}, im_{\phi}, \diamond]$$

$$SecondIsoThm := ((Subgroup[H, G, *]) \wedge (NormalSubgroup[N, G, *])) \implies (Isomorphic[H/(H \cap N), \bar{*}, (HN)/N, \bar{*}])$$

$$(1) \quad (Group[H, *]) \wedge (Group[N, *]) \blacksquare (e \in H) \wedge (e \in N)$$

$$(2) \quad e = e * e \in HN \blacksquare \emptyset \neq HN \subseteq G$$

$$(3) \quad (h_1 n_1, h_2 n_2 \in HN) \implies \dots$$

$$(3.1) \quad h_2 \in G \blacksquare (h_2)^{-1} n_1 h_2 \in N$$

$$(3.2) \quad (h_1 n_1)(h_2 n_2) = h_1 \left(h_2 (h_2)^{-1} \right) n_1 h_2 n_2 = (h_1 h_2) \left((h_2)^{-1} n_1 h_2 n_2 \right) \blacksquare (h_1 n_1)(h_2 n_2) = (h_1 h_2) \left((h_2)^{-1} n_1 h_2 n_2 \right)$$

$$(3.3) \quad (Group[H, *]) \wedge (Group[N, *]) \blacksquare (h_1 h_2 \in H) \wedge \left((h_2)^{-1} n_1 h_2 n_2 \in N \right)$$

$$(3.4) \quad (h_1 n_1)(h_2 n_2) = (h_1 h_2) \left((h_2)^{-1} n_1 h_2 n_2 \in N \right) \blacksquare (h_1 n_1)(h_2 n_2) \in N$$

$$(4) \quad (h_1 n_1, h_2 n_2 \in HN) \implies \left((h_1 n_1)(h_2 n_2) \in N \right) \blacksquare \forall_{h_1 n_1, h_2 n_2 \in HN} \left((h_1 n_1)(h_2 n_2) \in N \right)$$

$$(5) \quad (hn \in HN) \implies \dots$$

$$(5.1) \quad (Subgroup[H, G, *]) \wedge (Group[N, *]) \blacksquare (h^{-1} \in G) \wedge (n^{-1} \in N)$$

$$(5.2) \quad (NormalSubgroup[N, G, *]) \wedge (h^{-1} \in G) \wedge (n^{-1} \in N) \blacksquare hn^{-1}h^{-1} \in N$$

-
- (5.3) $(hn)^{-1} = n^{-1}h^{-1} = (h^{-1}h)n^{-1}h^{-1} = h^{-1}(hn^{-1}h^{-1}) \in HN \quad \blacksquare (hn)^{-1} \in HN$
-
- (6) $(hn \in HN) \implies ((hn)^{-1} \in HN) \quad \blacksquare \forall_{hn \in HN} ((hn)^{-1} \in HN)$
-
- (7) $(\emptyset \neq HN \subseteq G) \wedge \left(\forall_{h_1n_1, h_2n_2 \in HN} ((h_1n_1)(h_2n_2) \in N) \right) \wedge \left(\forall_{hn \in HN} ((hn)^{-1} \in HN) \right) \quad \blacksquare \text{Subgroup}[HN, G, *] \quad \blacksquare \text{Group}[HN, *]$
-
- (8) $(N \subseteq HN) \wedge (\text{Group}[N, *]) \quad \blacksquare \text{Subgroup}[N, HN, *]$
-
- (9) $((n \in N) \wedge (h_1n_1 \in HN)) \implies \dots$
-
- (9.1) $(\text{NormalSubgroup}[N, G, *]) \wedge (h_1n_1 \in G) \quad \blacksquare (h_1n_1)^{-1}n(h_1n_1) \in N$
-
- (10) $((n \in N) \wedge (h_1n_1 \in HN)) \implies ((h_1n_1)^{-1}n(h_1n_1) \in N) \quad \blacksquare \forall_{n \in N} \forall_{h_1n_1 \in HN} ((h_1n_1)^{-1}n(h_1n_1) \in N)$
-
- (11) $(\text{Subgroup}[N, HN, *]) \wedge \left(\forall_{n \in N} \forall_{h_1n_1 \in HN} ((h_1n_1)^{-1}n(h_1n_1) \in N) \right) \quad \blacksquare \text{NormalSubgroup}[N, HN, *]$
-
- (12) $(\text{SubgroupIntersection}) \wedge (\text{Subgroup}[H, G, *]) \wedge (\text{Subgroup}[N, G, *]) \quad \blacksquare \text{Subgroup}[H \cap N, G, *] \quad \blacksquare \text{Group}[H \cap N, *]$
-
- (13) $(H \cap N \subseteq H) \wedge (\text{Group}[H \cap N, *]) \quad \blacksquare \text{Subgroup}[H \cap N, H, *]$
-
- (14) $((x \in H \cap N) \wedge (h \in H)) \implies \dots$
-
- (14.1) $x \in H \cap N \quad \blacksquare (x \in H) \wedge (x \in N)$
-
- (14.2) $(\text{Group}[H, *]) \wedge (h \in H) \quad \blacksquare h^{-1} \in H$
-
- (14.3) $(\text{Group}[H, *]) \wedge (x, h, h^{-1} \in H) \quad \blacksquare h^{-1}xh \in H$
-
- (14.4) $(\text{NormalSubgroup}[N, G, *]) \wedge (h \in G) \wedge (x \in N) \quad \blacksquare h^{-1}xh \in N$
-
- (14.5) $(h^{-1}xh \in H) \wedge (h^{-1}xh \in N) \quad \blacksquare h^{-1}xh \in H \cap N$
-
- (15) $((x \in H \cap N) \wedge (h \in H)) \implies (h^{-1}xh \in H \cap N) \quad \blacksquare \forall_{x \in H \cap N} \forall_{h \in H} (h^{-1}xh \in H \cap N)$
-
- (16) $(\text{Subgroup}[H \cap N, H, *]) \wedge \left(\forall_{x \in H \cap N} \forall_{h \in H} (h^{-1}xh \in H \cap N) \right) \quad \blacksquare \text{NormalSubgroup}[H \cap N, H, *]$
-
- (17) $(\text{Group}[HN, *]) \wedge (\text{NormalSubgroup}[N, HN, *]) \wedge (\text{Group}[H, *]) \wedge (\text{NormalSubgroup}[H \cap N, H, *])$
-
- (18) $\text{QuotientGroupThm} \quad \blacksquare (\text{Group}[(HN)/N, \bar{*}]) \wedge (\text{Group}[H/(H \cap N), \bar{*}])$
-
- (19) $\phi := \{ \langle h, hN \rangle \in (H \times (HN)/N) \mid h \in H \} \quad \blacksquare \text{Func}[\phi, H, (HN)/N]$
-
- (20) $(h_1, h_2 \in H) \implies \dots$
-
- (20.1) $\phi(h_1 * h_2) = (h_1 * h_2)N = (h_1N) \bar{*} (h_2N) = \phi(h_1) \bar{*} \phi(h_2) \quad \blacksquare \phi(h_1 * h_2) = \phi(h_1) \bar{*} \phi(h_2)$
-
- (21) $(h_1, h_2 \in H) \implies (\phi(h_1 * h_2) = \phi(h_1) \bar{*} \phi(h_2)) \quad \blacksquare \forall_{h_1, h_2 \in H} (\phi(h_1 * h_2) = \phi(h_1) \bar{*} \phi(h_2))$
-
- (22) $(\text{Func}[\phi, H, (HN)/N]) \wedge \left(\forall_{h_1, h_2 \in H} (\phi(h_1 * h_2) = \phi(h_1) \bar{*} \phi(h_2)) \right) \quad \blacksquare \text{Homomorphism}[\phi, H, *, (HN)/N, \bar{*}]$
-
- (23) $\text{im}_\phi = \{ \phi(h) \mid h \in H \} = \{ hN \mid h \in H \} = (HN)/N \quad \blacksquare \text{im}_\phi = (HN)/N$
-
- (24) $\ker_\phi = \{ h \in H \mid \phi(h) = e_{(HN)/N} \} = \{ h \in H \mid hN = N \} = \{ h \in H \mid h \in N \} = \{ h \mid (h \in H) \wedge (h \in N) \} = H \cap N \quad \blacksquare \ker_\phi = H \cap N$
-
- (25) $(\text{FirstIsoThm}) \wedge (\text{Homomorphism}[\phi, H, *, (HN)/N, \bar{*}]) \quad \blacksquare \text{Isomorphic}[H/\ker_\phi, \bar{*}, \text{im}_\phi, \bar{*}]$
-
- (26) $(\ker_\phi = H \cap N) \wedge (\text{im}_\phi = (HN)/N) \wedge (\text{Isomorphic}[H/\ker_\phi, \bar{*}, \text{im}_\phi, \bar{*}]) \quad \blacksquare \text{Isomorphic}[H/(H \cap N), \bar{*}, (HN)/N, \bar{*}]$
-

$$\text{ThirdIsoThm} := \left(\begin{array}{l} ((\text{NormalSubgroup}[K, G, *]) \wedge (\text{NormalSubgroup}[H, G, *]) \wedge (\text{Subgroup}[K, H, *])) \implies \\ (\text{Isomorphic}[G/H, \bar{*}, (G/K)/(H/K), \bar{*}]) \end{array} \right)$$

(1) TODO

Chapter 3

Linear Algebra

3.1 Matrix Operations and Special Matrices

$Matrix[A, m, n] := [a_{i,j}]_{m \times n} := m \text{ rows, } n \text{ columns of real numbers}$

$\mathcal{M}_{m,n} := \{A : Matrix[A, m, n]\}$

$O_{m,n} := (Matrix[O, m, n]) \wedge (a_{i,j} = 0)$

$Square[A, n] := Matrix[A, n, n]$

$UpperTriangular[A] := (Square[A]) \wedge (i > j \implies a_{i,j} = 0)$

$LowerTriangular[A] := (Square[A]) \wedge (i < j \implies a_{i,j} = 0)$

$Diagonal[A, n] := (Square[A, n]) \wedge (i \neq j \implies a_{i,j} = 0)$

$Scalar[A, n, k] := (Diagonal[A, n]) \wedge (a_{i,i} = k)$

$I_n := Scalar[I, n, 1]$

$+(A, B) := ((Matrix[A, m, n]) \wedge (Matrix[B, m, n])) \implies (A + B = [a_{i,j} + b_{i,j}]_{m \times n})$

$*(r, A) := ((r \in \mathbb{R}) \wedge (Matrix[A, m, n])) \implies (r * A = [ra_{i,j}]_{m \times n})$

$*(A, B) := ((Matrix[A, m, p]) \wedge (Matrix[B, p, n])) \implies \left(A * B = \left[\sum_{k=1}^p (a_{i,k} b_{k,j}) \right]_{m \times n} \right)$

$^T[A] := (Matrix[A, m, n]) \implies (A^T = [a_{j,i}]_{n \times m})$

$AddCom := \forall_{A,B \in \mathcal{M}} (A + B = B + A)$

(1) $A + B = [a_{i,j} + b_{i,j}] = [b_{i,j} + a_{i,j}] = B + A$

$AddAssoc := \forall_{A,B,C \in \mathcal{M}} ((A + B) + C = A + (B + C))$

(1) $(A + B) + C = [(a_{i,j} + b_{i,j}) + c_{i,j}] = [a_{i,j} + (b_{i,j} + c_{i,j})] = A + (B + C)$

$AddId := \forall_{A \in \mathcal{M}} \exists!_{O \in \mathcal{M}} (A + O = A = O + A)$

(1) $A + O = [a_{i,j} + 0] = A = [0 + a_{i,j}] = O + A$

(2) $A + O_1 = A = A + O_2 \quad \blacksquare \quad O_1 = O_2$

$AddInv := \forall_{A \in \mathcal{M}} \exists!_{(-A) \in \mathcal{M}} (A + (-A) = O = (-A) + A)$

(1) $A + (-A) = [a_{i,j} - a_{i,j}] = O = [-a_{i,j} + a_{i,j}] = (-A) + A$

(2) $A + (-A_1) = O = A + (-A_2) \quad \blacksquare \quad -A_1 = -A_2 \quad \blacksquare \quad A_1 = A_2$

$MulAssoc := \forall_{A,B,C \in \mathcal{M}} ((A * B) * C = A * (B * C))$

(1) $(A * B) * C = \left[\sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,j}) \right] * C = \left[\sum_{k_2=1}^{p_2} \left(\sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,k_2}) c_{k_2,j} \right) \right] = \left[\sum_{k_2=1}^{p_2} \sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,k_2} c_{k_2,j}) \right] = \dots$

(2) $\dots \left[\sum_{k_1=1}^{p_1} \sum_{k_2=1}^{p_2} (a_{i,k_1} b_{k_1,k_2} c_{k_2,j}) \right] = \left[\sum_{k_1=1}^{p_1} \left(a_{i,k_1} \sum_{k_2=1}^{p_2} (b_{k_1,k_2} c_{k_2,j}) \right) \right] = \dots = A * (B * C)$

$MulId := \forall_{A: Square[A,n]} (A * I_n = A = I_n * A)$

$$(1) \quad A * I_n = \left[\sum_{k=1}^n \left(a_{i,k} \begin{pmatrix} 1 & k=j \\ 0 & k \neq j \end{pmatrix} \right) \right] = [a_{i,j}] = A$$

$$(2) \quad \text{TODO} = A$$

$$\text{ScalAssoc} := \forall_{r,s \in \mathbb{R}} \forall_{A \in \mathcal{M}} (r(sA) = (rs)A = s(rA))$$

$$(1) \quad r(sA) = r[sa_{i,j}] = [rsa_{i,j}]$$

$$(2) \quad (rs)A = [rsa_{i,j}]$$

$$(3) \quad s(rA) = s[ra_{i,j}] = [sra_{i,j}] = [rsa_{i,j}]$$

$$\text{TransCancel} := \forall_{A \in \mathcal{M}} (A = (A^T)^T)$$

$$(1) \quad A = [a_{i,j}] = [a_{j,i}]^T = ([a_{i,j}]^T)^T = (A^T)^T$$

$$\text{ScalMulCom} := \forall_{r \in \mathbb{R}} \forall_{A,B \in \mathcal{M}} ((rA) * B = r(A * B) = A * (rB))$$

$$(1) \quad (rA) * B = [ra_{i,l}] * [b_{l,j}] = \left[\sum_{k=1}^p (ra_{i,k} b_{k,j}) \right] = r(A * B)$$

$$(2) \quad A * (rB) = [a_{i,l}] * [rb_{l,j}] = \left[\sum_{k=1}^p (a_{i,k} rb_{k,j}) \right] = \left[\sum_{k=1}^p (ra_{i,k} b_{k,j}) \right] = r(A * B)$$

$$\text{ScalDistLeft} := \forall_{r,s \in \mathbb{R}} \forall_{A \in \mathcal{M}} ((r+s)A = rA + sA)$$

$$(1) \quad \text{TODO}$$

$$\text{ScalDistRight} := \forall_{r \in \mathbb{R}} \forall_{A,B \in \mathcal{M}} (r(A+B) = rA + rB)$$

$$(1) \quad \text{TODO}$$

$$\text{MulDistRight} := \forall_{A,B,C \in \mathcal{M}} ((A+B) * C = A * C + B * C)$$

$$(1) \quad (A+B) * C = [a_{i,j} + b_{i,j}] * C = \left[\sum_{k=1}^p ((a_{i,k} + b_{i,k}) c_{k,j}) \right] = \dots$$

$$(2) \quad \dots \left[\sum_{k=1}^p (a_{i,k} c_{k,j} + b_{i,k} c_{k,j}) \right] = \left[\sum_{k=1}^p (a_{i,k} c_{k,j}) \right] + \left[\sum_{k=1}^p (b_{i,k} c_{k,j}) \right] = A * C + B * C$$

$$\text{MulDistLeft} := \forall_{A,B,C \in \mathcal{M}} (C * (A+B) = C * A + C * B)$$

$$(1) \quad \text{TODO}$$

$$\text{TransAddDist} := \forall_{A,B \in \mathcal{M}} ((A+B)^T = A^T + B^T)$$

$$(1) \quad \text{TODO}$$

$$\text{TransMulDist} := \forall_{A,B \in \mathcal{M}} ((A * B)^T = B^T * A^T)$$

$$(1) \quad (A * B)^T = \left[\sum_{k=1}^p (a_{i,k} b_{k,j}) \right]^T = \left[\sum_{k=1}^p (a_{j,k} b_{k,i}) \right] = \left[\sum_{k=1}^p (b_{k,i} a_{j,k}) \right] = \left[\sum_{k=1}^p (b_{i,k}^T a_{k,j}^T) \right] = B^T * A^T$$

$$\text{Sym}[A] := A = A^T$$

$$\text{SkewSym}[A] := A = -A^T$$

$$\text{Invertible}[A] := (\text{Square}[A, n]) \wedge \left(\exists_{A^{-1} \in \mathcal{M}} (A * A^{-1} = I_n = A^{-1} * A) \right)$$

$$\text{SymGen} := \forall_{A \in \mathcal{M}} (\text{Sym}[A + A^T])$$

$$(1) \quad (A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T$$

$$\text{SkewSymGen} := \forall_{A \in \mathcal{M}} (\text{SkewSym}[A - A^T])$$

$$(1) \quad -(A - A^T)^T = -(A^T - (A^T)^T) = -(A^T - A) = (A - A^T)$$

$$SymDecomp := \forall_{A \in \mathcal{M}} \exists! B : Sym[B] \exists! C : SkewSym[C] (A = B + C)$$

$$(1) \quad B := (1/2) * (A + A^T) ; C := (1/2) * (A - A^T)$$

$$(2) \quad SymGen[B] \wedge SkewSymGen[C]$$

$$(3) \quad A = (1/2) * (A + A^T) + (1/2) * (A - A^T) = B + C$$

$$(4) \quad (1/2) * (A_1 + A_1^T) = (1/2) * (A_2 + A_2^T) \quad \blacksquare \quad A_1 = A_2$$

$$(5) \quad (1/2) * (A_3 - A_3^T) = (1/2) * (A_4 - A_4^T) \quad \blacksquare \quad A_3 = A_4$$

$$InvId := \forall_{A : Invertible[A]} \left(\exists! A^{-1} \in \mathcal{M} (A * A^{-1} = I_n = A^{-1} * A) \right)$$

$$(1) \quad A^{-1}_1 = A^{-1}_1 * I_n = A^{-1}_1 * (A * A^{-1}_2) = (A^{-1}_1 * A) * A^{-1}_2 = I_n * A^{-1}_2 = A^{-1}_2$$

$$InvCancel := \forall_{A : Invertible[A]} \left((A^{-1})^{-1} = A \right)$$

$$(1) \quad (A * A^{-1})^{-1} = I_n^{-1} = I_n$$

$$(2) \quad (A^{-1})^{-1} * A^{-1} = I_n \quad \blacksquare \quad A^{-1})^{-1} = I_n * A = A$$

$$InvDist := \forall_{A : Invertible[A]} \forall_{B : Invertible[B]} \left((A * B)^{-1} = B^{-1} * A^{-1} \right)$$

$$(1) \quad (A * B) * (A * B)^{-1} = I \quad \blacksquare \quad B * (A * B)^{-1} = A^{-1} \quad \blacksquare \quad (A * B)^{-1} = B^{-1} * A^{-1}$$

$$InvTrans := \forall_{A : Invertible[A]} \left((A^T)^{-1} = (A^{-1})^T \right) \quad \blacksquare \quad \Leftarrow$$

$$(1) \quad A^T * (A^{-1})^T = (A^{-1} * A)^T = I^T = I \quad \blacksquare \quad (A^{-1})^T = (A^T)^{-1}$$

3.2 Elementary Matrices on Invertibility and Systems of Linear Equations

$$Sys[A, B] := (Matrix[A, m, n]) \wedge (Matrix[B, m, 1])$$

$$Sol[X, A, B] := (Sys[A, B]) \wedge (Matrix[X, n, 1]) \wedge (A * X = B)$$

$$ConsistentSys[A, B] := (Sys[A, B]) \wedge \exists_X (Sol[X, A, B])$$

$$TrivSol[X, A] := (Sol[X, A, O]) \wedge (X = O)$$

$$NonTrivSol[X, A] := (Sol[X, A, O]) \wedge (X \neq O)$$

$$HomoSysProps := (Sys[A, O]) \implies \dots$$

$$(1) \quad u_0 := O ; u_1 := choice(\{X \in \mathcal{M} \mid X \neq O\}) ; k := choice(\mathbb{R})$$

$$(2) \quad TrivSol[u_0, A]$$

$$(3) \quad (NonTrivSol[u_1, A]) \implies (Sol[u_1 + ku_0])$$

$$(4) \quad (TrivSol[\vec{X}, A]) \implies (TrivSol[LC(\vec{X}), A])$$

$$ElemMat[E] := (E = Swap[I_n, i, j]) \vee (Scale_*(I_n, i, c)) \vee (Combine_*(I_n, i, c, j))$$

$$ElemMatProd[E^*] := \exists_{\langle E \rangle} \left(\forall_{E_i \in E^*} (ElemMat[E_i]) \wedge \left(E^* = \prod_{E_i \in E^*} (E_i) \right) \right)$$

$$RowEquiv[A, B] := \exists_{E^*} ((ElemMatProd[E^*]) \wedge (B = E^* * A))$$

$$ElemMatInv := \forall_{E \in \mathcal{M}} ((ElemMat[E]) \implies (Invertible[E]))$$

$$(1) \quad E - RowSwap[E] \implies TODO ; E - RowScale_*(E) \implies TODO ; E - RowCombine_*(E) \implies TODO$$

$$ElemMatProdInv := \forall_{E^*} ((ElemMatProd[E^*]) \implies (Invertible[E^*]))$$

$$(1) \quad TODO$$

$$RowEquivSys := \forall_{A, B, C, D, X \in \mathcal{M}} \left(((Sys[A, B]) \wedge (Sys[C, D]) \wedge (RowEquiv[[AB], [CD]])) \implies (Sol[X, A, B] \iff Sol[X, C, D]) \right)$$

$$(1) \quad \exists_{E^* : ElemMatProd[E^*]} ([CD] = E^* * [AB])$$

-
- (2) $(E^* * A = C) \wedge (E^* * B = D)$
-
- (3) $Sol[Y, A, B] \implies \dots$
-
- (3.1) $A * Y = B$
-
- (3.2) $C * Y = (E^* * A) * Y = E^* * (A * Y) = E^* * B = D \quad \blacksquare \quad Sol[Y, C, D]$
-
- (4) $Sol[Y, A, B] \implies Sol[Y, C, D]$
-
- (5) $\left(A = (E^*)^{-1} * C \right) \wedge \left(B = (E^*)^{-1} * D \right)$
-
- (6) $Sol[Z, C, D] \implies \dots$
-
- (6.1) $C * Z = D$
-
- (6.2) $A * Z = \left((E^*)^{-1} * C \right) * Z = (E^*)^{-1} * (C * Z) = (E^*)^{-1} * D = B$
-
- (7) $Sol[Z, C, D] \implies Sol[Z, A, B]$
-
- (8) $Sol[X, A, B] \iff Sol[X, C, D]$
-

$$RowEquivHomoSysSol := \forall_{A, C, X \in \mathcal{M}} \left((RowEquiv[A, C]) \implies ((Sol[X, A, O]) \iff (Sol[X, C, O])) \right)$$

-
- (1) Set $B = D = O$
-

$$RREF[A] := (A \in \mathcal{M}) \wedge \left(\begin{array}{l} \text{All zero rows are at the bottom of the matrix.} \\ \text{The leading entry after the first occurs to the right of the leading entry of the previous row.} \\ \text{The leading entry in any nonzero row is 1.} \\ \text{All entries in the column above and below a leading 1 are zero.} \end{array} \right) \wedge$$

$$GaussJordanElim := \forall_{A \in \mathcal{M}} \exists!_{B \in \mathcal{M}} ((RREF[B]) \wedge (RowEquiv[A, B]))$$

-
- (1) Hit A with $ElemMat$'s until it becomes B
-
- (2) $(B = E^* * A) \wedge (RREF[B])$
-

$$HasZero[A] := (Matrix(A, m, n)) \wedge (\exists_{i \leq m} (A_{i,:} = O))$$

$$HasZeroNonInvertible := \forall_{A \in \mathcal{M}} ((HasZero[A]) \implies (\neg Invertible[A]))$$

-
- (1) $i := choice(\{i \leq m \mid A_{i,:} = O\})$
-
- (2) $(B \in \mathcal{M}) \implies \dots$
-
- (2.1) $(A * B)_{i,:} = O \neq I_{n_i,:} \quad \blacksquare \quad A * B \neq I_n$
-
- (3) $(B \in \mathcal{M}) \implies (A * B \neq I_n) \quad \blacksquare \quad \forall_{B \in \mathcal{M}} (A * B \neq I_n) \quad \blacksquare \quad \neg Invertible[A]$
-

$$InvIf f RowEquivI := \forall_{A \in \mathcal{M}} ((Invertible[A]) \iff (RowEquiv[A, I_n]))$$

-
- (1) $(Invertible[A]) \implies \dots$
-
- (1.1) $(RREF[B]) \wedge (RowEquiv[A, B])$
-
- (1.2) $B = E^* * A$
-
- (1.3) $(Invertible[E^*]) \wedge (Invertible[A]) \quad \blacksquare \quad Invertible[B]$
-
- (1.4) $Invertible[B] \quad \blacksquare \quad \neg HasZero[B]$
-
- (1.5) $(RREF[B]) \wedge (\neg HasZero[B]) \quad \blacksquare \quad B = I_n$
-
- (1.6) $RowEquiv[A, I_n]$
-
- (2) $(Invertible[A]) \implies (RowEquiv[A, I_n])$
-
- (3) $(RowEquiv[A, I_n]) \implies \dots$
-
- (3.1) $I_n = E^* * A \quad \blacksquare \quad (E^*)^{-1} = A$
-
- (3.2) $A^{-1} = E^*_{DescSort} \quad \blacksquare \quad Invertible[A]$
-
- (4) $(RowEquiv[A, I_n]) \implies (Invertible[A])$
-
- (5) $(Invertible[A]) \iff (RowEquiv[A, I_n])$
-

$$RowEquivIf f TrivSol := \forall_{A \in \mathcal{M}} \left((RowEquiv[A, I_n]) \iff \left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right) \right)$$

-
- (1) $(RowEquiv[A, I_n]) \implies \dots$
-
- (1.1) $RowEquiv[A, I_n] \quad \blacksquare \quad Invertible[A]$
-

(1.2)	$(Sol[X, A, O]) \implies \dots$
(1.2.1)	$A * X = O \quad \blacksquare \quad X = A^{-1} * O = O \quad \blacksquare \quad X = O$
(1.3)	$(Sol[X, A, O]) \implies (X = O)$
(1.4)	$(X = O) \implies (Sol[X, A, O])$
(1.5)	$(X = O) \iff (Sol[X, A, O]) \quad \blacksquare \quad \forall_X ((X = O) \iff (Sol[X, A, O]))$
(2)	$(RowEquiv[A, I_n]) \implies \left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right)$
(3)	$\left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right) \implies \dots$
(3.1)	$(RREF[B]) \wedge (RowEquiv[A, B])$
(3.2)	$Sol[X, B, O]$
(3.3)	$(B \neq I_n) \implies \dots$
(3.3.1)	$\left(\exists_{Y \neq X} (Sol[Y, B, O]) \right)$
(3.3.2)	$Sol[Y, A, O] \quad \blacksquare \quad Y = X$
(3.3.3)	$(Y \neq X) \wedge (Y = X) \quad \blacksquare \quad \perp$
(3.4)	$(B \neq I_n) \implies \perp \quad \blacksquare \quad B = I_n$
(3.5)	$(RowEquiv[A, B]) \wedge (B = I_n) \quad \blacksquare \quad RowEquiv[A, I_n]$
(4)	$\left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right) \implies (RowEquiv[A, I_n])$
(5)	$(RowEquiv[A, I_n]) \iff \left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right)$

$$InvIf fUniqSol := \forall_{A \in \mathcal{M}} \left((Invertible[A]) \iff \left(\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B]) \right) \right)$$

(1)	$(Invertible[A] \wedge B \in \mathcal{M}) \implies \dots$
(1.1)	$(Invertible[A]) \wedge (Sys[A, B])$
(1.2)	$(X = A^{-1} * B) \iff (Sol[X, A, B]) \quad \blacksquare \quad \exists!_{X \in \mathcal{M}} (Sol[X, A, B])$
(2)	$\left(\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B]) \right) \implies \dots$
(2.1)	$X_i := choice(\{X_i \mid Sol[X_i, A, I_{n:,i}]\})$
(2.2)	$A * [X_1 \dots X_n] = [(A * X_1) \dots (A * X_n)] = [I_{n:,1} \dots I_{n:,n}] = I_n$
(2.3)	$A^{-1} = [X_1 \dots X_n]$
(3)	$\left(\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B]) \right) \implies (Invertible[A])$

$$SquareTheorems_4 := \forall_{A \in \mathcal{M}} \left(\begin{array}{ccc} (Invertible[A]) & \iff & \\ (RowEquiv[A, I_n]) & \iff & \\ \left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right) & \iff & \\ \left(\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B]) \right) & & \end{array} \right)$$

3.3 Vector Spaces

$$VectorSpace[V, +, *] := \exists_{O \in V} \forall_{\alpha, \beta \in \mathbb{R}} \forall_{u, v, w \in V} \left(\begin{array}{l} (u + v \in V) \wedge (u + v = v + u) \wedge ((u + v) + w = u + (v + w)) \wedge \\ (u + O = u) \wedge \left(\exists_{-u \in V} (u + (-u) = O) \right) \wedge \\ (\alpha * u \in V) \wedge (\alpha * (\beta * u) = (\alpha\beta) * u) \wedge (1 * u = u) \wedge \\ (\alpha * (u + v) = (\alpha * u) + (\alpha * v)) \wedge ((\alpha + \beta) * u = (\alpha * u) + (\beta * u)) \end{array} \right)$$

$$ZeroVectorUniq := \forall_{O', v \in V} ((v + O' = v) \implies (O' = O))$$

(1)	$O' = O' + O = O + O' = O \quad \blacksquare \quad O' = O$
-----	------------------------------------------------------------

$$AddInvUniq := \forall_{-v', v \in V} ((v + -v' = O) \implies (-v' = -v))$$

(1)	$-v' = -v' + O = -v' + (v + -v) = (-v' + v) + -v = (v + -v') + -v = O + -v = -v \quad \blacksquare \quad -v' = -v$
-----	--------------------------------------------------------------------------------------------------------------------

$$AddInvGen := \forall_{v \in V} ((-1) * v = -v)$$

$$(1) \quad v + (-1) * v = (1 - 1) * v = 0 * v = O \quad \blacksquare \quad (-1) * v = -v$$

$$ZeroVectorGenLeft := \forall_{v \in V} (0 * v = O)$$

$$(1) \quad 0 * v = (0 + 0) * v = (0 * v) + (0 * v) \quad \blacksquare \quad O = 0 * v$$

$$ZeroVectorGenRight := \forall_{r \in \mathbb{R}} (r * O = O)$$

$$(1) \quad r * O = r * (O + O) = (r * O) + (r * O) \quad \blacksquare \quad O = r * O$$

$$ZeroVectorEquiv := \forall_{r \in \mathbb{R}} \forall_{v \in V} \left((r * v = O) \iff ((v = O) \vee (r = 0)) \right)$$

$$(1) \quad (ZeroVectorGenLeft) \wedge (ZeroVectorGenRight) \quad \blacksquare \quad ((v = O) \vee (r = 0)) \implies (r * v = O)$$

$$(2) \quad (r * v = O) \implies \dots$$

$$(2.1) \quad (r \neq 0) \implies \dots$$

$$(2.1.1) \quad r \neq 0 \quad \blacksquare \quad r^{-1} \in \mathbb{R}$$

$$(2.1.2) \quad ZeroVectorGenRight \quad \blacksquare \quad O = r^{-1} * O = r^{-1} * (r * v) = (r^{-1}r) * v = 1 * v = v \quad \blacksquare \quad O = v$$

$$(2.2) \quad (r \neq 0) \implies (v = O) \quad \blacksquare \quad (r = 0) \vee (v = O)$$

$$(3) \quad (r * v = O) \implies ((r = 0) \vee (v = O))$$

$$(4) \quad (r * v = O) \iff ((r = 0) \vee (v = O))$$

3.4 Subspaces and Special Subspaces

$$Subspace[S, V, +, *] := (VectorSpace[V, +, *]) \wedge (S \subseteq V) \wedge (VectorSpace[S, +, *])$$

$$SubspaceEquiv := \forall_{V, S} \left(\begin{array}{c} (VectorSpace[V, +, *]) \\ \left((Subspace[S, V, +, *]) \iff ((\emptyset \neq S \subseteq V) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S))) \right) \end{array} \implies \right)$$

$$(1) \quad (Subspace[S, V, +, *]) \implies \dots$$

$$(1.1) \quad Subspace[S, V, +, *] \quad \blacksquare \quad S \subseteq V$$

$$(1.2) \quad VectorSpace[S, V, +, *] \quad \blacksquare \quad \exists_{O \in V} \forall_{v \in V} (v + O = v) \quad \blacksquare \quad O \in S \quad \blacksquare \quad \emptyset \neq S$$

$$(1.3) \quad (\emptyset \neq S) \wedge (S \subseteq V) \quad \blacksquare \quad \emptyset \neq S \subseteq V$$

$$(1.4) \quad VectorSpace[S, V, +, *] \quad \blacksquare \quad (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S))$$

$$(1.5) \quad (\emptyset \neq S \subseteq V) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S))$$

$$(2) \quad (Subspace[S, V, +, *]) \implies ((\emptyset \neq S \subseteq V) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S)))$$

$$(3) \quad ((\emptyset \neq S \subseteq V) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S))) \implies \dots$$

$$(3.1) \quad ((\emptyset \neq S) \wedge (\alpha, \beta \in \mathbb{R}) \wedge (u, v, w \in S)) \implies \dots$$

$$(3.1.1) \quad \emptyset \neq S \quad \blacksquare \quad \exists_x (x \in V)$$

$$(3.1.2) \quad (ZeroVectorGenLeft) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S)) \wedge (x \in V) \quad \blacksquare \quad O = 0 * x \in S \quad \blacksquare \quad O \in S$$

$$(3.1.3) \quad u, v \in V \quad \blacksquare \quad u + v = v + u$$

$$(3.1.4) \quad u, v, w \in V \quad \blacksquare \quad (u + v) + w = u + (v + w)$$

$$(3.1.5) \quad u \in V \quad \blacksquare \quad u + O = u$$

$$(3.1.6) \quad (AddInvGen) \wedge (u \in S) \quad \blacksquare \quad (-1) * u = -u \in S$$

$$(3.1.7) \quad u \in V \quad \blacksquare \quad \alpha * (\beta * u) = (\alpha\beta) * u$$

$$(3.1.8) \quad u \in V \quad \blacksquare \quad 1 * u = u$$

$$(3.1.9) \quad u, v \in V \quad \blacksquare \quad \alpha * (u + v) = (\alpha * u) + (\alpha * v)$$

$$(3.1.10) \quad u \in V \quad \blacksquare \quad (\alpha + \beta) * u = (\alpha * u) + (\beta * u)$$

$$(4) \quad ((\emptyset \neq S) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S))) \implies (Subspace[S, V, +, *])$$

$$(5) \quad (Subspace[S, V, +, *]) \iff ((\emptyset \neq S) \wedge (\forall_{r, s \in S} (r + s \in S)) \wedge (\forall_{\alpha \in \mathbb{R}} \forall_{s \in S} (\alpha * s \in S)))$$

$$\text{SetSum}[A + B, A, B, V, +, *] := (\text{VectorSpace}[V, +, *]) \wedge (A, B \subseteq V) \wedge (A + B = \{a + b \mid (a \in A) \wedge (b \in B)\})$$

$$\text{SumSubContains} := \forall_{A,B,V} \left(\left((\text{Subspace}[A, V, +, *]) \wedge (\text{Subspace}[B, V, +, *]) \wedge (\text{SetSum}[A + B, A, B, V, +, *]) \right) \implies \left((\text{Subspace}[A + B, V, +, *]) \wedge (A, B \subseteq A + B) \right) \right)$$

$$(1) \quad (\text{Subspace}[A, V, +, *]) \wedge (\text{Subspace}[B, V, +, *]) \quad \blacksquare \quad (O \in A) \wedge (O \in B)$$

$$(2) \quad (\text{SetSum}[A + B, A, B, V, +, *]) \wedge (O \in A) \wedge (O \in B) \quad \blacksquare \quad O = O + O \in A + B \quad \blacksquare \quad \emptyset \neq A + B$$

$$(3) \quad (v \in A + B) \implies \dots$$

$$(3.1) \quad \exists_{a \in A} \exists_{b \in B} (v = a + b)$$

$$(3.2) \quad (A \subseteq V) \wedge (B \subseteq V) \quad \blacksquare \quad a, b \in V$$

$$(3.3) \quad \text{VectorSpace}[V, +, *] \quad \blacksquare \quad v = a + b \in V$$

$$(4) \quad (v \in A + B) \implies (v \in V) \quad \blacksquare \quad A + B \subseteq V$$

$$(5) \quad (\emptyset \neq A + B) \wedge (A + B \subseteq V) \quad \blacksquare \quad \emptyset \neq A + B \subseteq V$$

$$(6) \quad (u, v \in A + B) \implies \dots$$

$$(6.1) \quad \left(\exists_{a_1 \in A} \exists_{b_1 \in B} (u = a_1 + b_1) \right) \wedge \left(\exists_{a_2 \in A} \exists_{b_2 \in B} (v = a_2 + b_2) \right)$$

$$(6.2) \quad u + v = (a_1 + b_1) + (a_2 + b_2) = (a_1 + a_2) + (b_1 + b_2)$$

$$(6.3) \quad (a_1 + a_2 \in A) \wedge (b_1 + b_2 \in B) \quad \blacksquare \quad u + v \in A + B$$

$$(7) \quad (u, v \in A + B) \implies (u + v \in A + B) \quad \blacksquare \quad \forall_{u,v \in A+B} (u + v \in A + B)$$

$$(8) \quad ((r \in \mathbb{R}) \wedge (v \in A + B)) \implies \dots$$

$$(8.1) \quad \exists_{a \in A} \exists_{b \in B} (v = a + b)$$

$$(8.2) \quad r * v = r * (a + b) = r * a + r * b$$

$$(8.3) \quad (r * a \in A) \wedge (r * b \in B) \quad \blacksquare \quad r * v \in A + B$$

$$(9) \quad ((r \in \mathbb{R}) \wedge (v \in A + B)) \implies (r * v \in A + B) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{v \in A+B} (r * v \in A + B)$$

$$(10) \quad (\text{SubspaceEquiv}) \wedge (\emptyset \neq A + B \subseteq V) \wedge (\forall_{u,v \in A+B} (u + v \in A + B)) \wedge (\forall_{r \in \mathbb{R}} \forall_{v \in A+B} (r * v \in A + B)) \quad \blacksquare \quad \text{Subspace}[A + B, V, +, *]$$

$$(11) \quad (O \in B) \wedge (\forall_{a \in A} (a + O) = a) \quad \blacksquare \quad A \subseteq A + B$$

$$(12) \quad (O \in A) \wedge (\forall_{b \in B} (b + O) = b) \quad \blacksquare \quad B \subseteq A + B$$

$$(13) \quad (A \subseteq A + B) \wedge (B \subseteq A + B) \quad \blacksquare \quad A, B \subseteq A + B$$

$$(14) \quad (\text{Subspace}[A + B, V, +, *]) \wedge (A, B \subseteq A + B)$$

$$\text{SumSubMinContains} := \forall_{A,B,V} \left(\left((\text{Subspace}[A, V, +, *]) \wedge (\text{Subspace}[B, V, +, *]) \wedge (\text{SetSum}[A + B, A, B, V, +, *]) \right) \implies \left(\forall_C ((\text{Subspace}[C, V, +, *]) \wedge (A, B \subseteq C)) \implies (A + B \subseteq C) \right) \right)$$

$$(1) \quad \text{SumSub} \quad \blacksquare \quad (A, B \subseteq A + B) \wedge (\text{Subspace}[A + B, V, +, *])$$

$$(2) \quad ((\text{Subspace}[C, V, +, *]) \wedge (A, B \subseteq C)) \implies \dots$$

$$(2.1) \quad (s \in A + B) \implies \dots$$

$$(2.1.1) \quad \exists_{a \in A} \exists_{b \in B} (s = a + b)$$

$$(2.1.2) \quad (A, B \subseteq C) \quad \blacksquare \quad a, b \in C$$

$$(2.1.3) \quad (\text{VectorSpace}[C, V, +, *]) \wedge (a, b \in C) \quad \blacksquare \quad s = a + b \in C$$

$$(2.2) \quad (s \in A + B) \implies (s \in C) \quad \blacksquare \quad A + B \subseteq C$$

$$(3) \quad ((\text{Subspace}[C, V, +, *]) \wedge (A, B \subseteq C)) \implies (A + B \subseteq C)$$

$$\text{DirSum}[A \oplus B, A, B, V, +, *] := \left(\begin{array}{c} (\text{Subspace}[A, V, +, *]) \quad \wedge \quad (\text{Subspace}[B, V, +, *]) \\ (\text{SetSum}[A + B, A, B, V, +, *]) \wedge \left(\forall_{s \in A+B} \exists!_{\langle a,b \rangle \in A \times B} (s = a + b) \right) \end{array} \right)$$

$$\text{DirSumEquiv} := \forall_{A,B,V} \left(\left((\text{Subspace}[A, V, +, *]) \wedge (\text{Subspace}[B, V, +, *]) \wedge (\text{SetSum}[A + B, A, B, V, +, *]) \right) \implies \left((\text{DirSum}[A \oplus B, A, B, V, +, *]) \iff \left(\exists!_{\langle a,b \rangle \in A \times B} (O = a + b) \right) \right) \right)$$

$$(1) \quad (\text{DirSum}[A \oplus B, A, B, V, +, *]) \implies \dots$$

$$(1.1) \quad (\text{Subspace}[A, V, +, *]) \wedge (\text{Subspace}[B, V, +, *]) \quad \blacksquare \quad (O \in A) \wedge (O \in B)$$

$$(1.2) \quad (SubSum[A \oplus B, A, B, V, +, *]) \wedge (O \in A) \wedge (O \in B) \quad \blacksquare \quad O = O + O \in A \oplus B$$

$$(1.3) \quad (DirSum[A \oplus B, A, B, V, +, *]) \wedge (O \in A \oplus B) \quad \blacksquare \quad \exists!_{\langle a, b \rangle \in A \times B} (O = a + b)$$

$$(2) \quad (DirSum[A \oplus B, A, B, V, +, *]) \implies \left(\exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right)$$

$$(3) \quad \left(\exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right) \implies \dots$$

$$(3.1) \quad (s \in A \oplus B) \implies \dots$$

$$(3.1.1) \quad \left(\exists_{\langle a, b \rangle \in A \times B} (s = a + b) \right)$$

$$(3.1.2) \quad ((s = a_1 + b_1) \wedge (s = a_2 + b_2)) \implies \dots$$

$$(3.1.2.1) \quad O = s - s = (a_1 + b_1) - (a_2 + b_2) = (a_1 - a_2) + (b_1 - b_2)$$

$$(3.1.2.2) \quad (Subspace[A, V, +, *]) \wedge (Subspace[B, V, +, *]) \quad \blacksquare \quad (a_1 - a_2 \in A) \wedge (b_1 - b_2 \in B)$$

$$(3.1.2.3) \quad ((a_1 - a_2 \neq O) \vee (b_1 - b_2 \neq O)) \implies \left(\neg \exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right) \implies \perp$$

$$(3.1.2.4) \quad (a_1 - a_2 = O) \wedge (b_1 - b_2 = O) \quad \blacksquare \quad \langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle$$

$$(3.1.3) \quad ((s = a_1 + b_1) \wedge (s = a_2 + b_2)) \implies \langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle$$

$$(3.1.4) \quad \forall_{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B} \left(((s = a_1 + b_1) \wedge (s = a_2 + b_2)) \implies (\langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle) \right)$$

$$(3.1.5) \quad \exists_{\langle a, b \rangle \in A \times B} (s = a + b) \wedge \forall_{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B} \left(((s = a_1 + b_1) \wedge (s = a_2 + b_2)) \implies (\langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle) \right) \quad \blacksquare \quad \exists!_{\langle a, b \rangle \in A \times B} (s = a + b)$$

$$(3.2) \quad (s \in A + B) \implies \exists!_{\langle a, b \rangle \in A \times B} (s = a + b) \quad \blacksquare \quad \forall_{s \in A + B} \exists!_{\langle a, b \rangle \in A \times B} (s = a + b) \quad \blacksquare \quad DirSum[A \oplus B, A, B, V, +, *]$$

$$(4) \quad \left(\exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right) \implies (DirSum[A \oplus B, A, B, V, +, *])$$

$$(5) \quad (DirSum[A \oplus B, A, B, V, +, *]) \iff \left(\exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right)$$

$$DirSumSubspace := \forall_{A, B, V} \left(\begin{array}{l} ((Subspace[A, V, +, *]) \wedge (Subspace[B, V, +, *]) \wedge (SetSum[A + B, A, B, V, +, *])) \implies \\ ((DirSum[A \oplus B, A, B, V, +, *]) \iff (A \cap B = \{O\})) \end{array} \right)$$

$$(1) \quad (DirSum[A \oplus B, A, B, V, +, *]) \implies \dots$$

$$(1.1) \quad (v \in A \cap B) \implies \dots$$

$$(1.1.1) \quad (v \in A \cap B) \wedge (VectorSpace[B, +, *]) \quad \blacksquare \quad (v \in A) \wedge (v \in B) \quad \blacksquare \quad (v \in A) \wedge (-v \in B)$$

$$(1.1.2) \quad (v \in A) \wedge (-v \in B) \quad \blacksquare \quad v + (-v) = O \in A + B$$

$$(1.1.3) \quad DirSum[A \oplus B, A, B, V, +, *] \quad \blacksquare \quad \exists!_{\langle a, b \rangle \in A \times B} (O = a + b)$$

$$(1.1.4) \quad (v \neq O) \implies \left(\neg \exists!_{\langle a, b \rangle \in A \times B} (O = a + b) \right) \implies \perp \quad \blacksquare \quad v = O$$

$$(1.2) \quad (v \in A \cap B) \implies (v = O) \quad \blacksquare \quad A + B \subseteq \{O\}$$

$$(1.3) \quad (v = O) \implies \dots$$

$$(1.3.1) \quad (Subspace[A, V, +, *]) \wedge (Subspace[B, V, +, *]) \quad \blacksquare \quad (O \in A) \wedge (O \in B) \quad \blacksquare \quad v = O \in A \cup B$$

$$(1.4) \quad (v = O) \implies (v \in A \cap B) \quad \blacksquare \quad \{O\} \subseteq A \cap B$$

$$(1.5) \quad (A + B \subseteq \{O\}) \wedge (\{O\} \subseteq A \cap B) \quad \blacksquare \quad A \cap B = \{O\}$$

$$(2) \quad (DirSum[A \oplus B, A, B, V, +, *]) \implies (A \cap B = \{O\})$$

$$(3) \quad (A \cap B = \{O\}) \implies \dots$$

$$(3.1) \quad (O \in A) \wedge (O \in B) \wedge (O = O + O \in A + B) \quad \blacksquare \quad \exists_{\langle a, b \rangle \in A \times B} (O = a + b)$$

$$(3.2) \quad ((\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B) \wedge (O = a_1 + b_1) \wedge (O = a_2 + b_2)) \implies \dots$$

$$(3.2.1) \quad (O = a_1 + b_1) \wedge (O = a_2 + b_2) \quad \blacksquare \quad (a_1 = -b_1) \wedge (a_2 = -b_2)$$

$$(3.2.2) \quad VectorSpace[B, +, *] \quad \blacksquare \quad -b_1, -b_2 \in B$$

$$(3.2.3) \quad (a_1 \in A) \wedge (a_1 = -b_1 \in B) \quad \blacksquare \quad a_1 \in A \cap B \quad \blacksquare \quad a_1 = O \quad \blacksquare \quad a_1 = b_1 = O$$

$$(3.2.4) \quad (a_2 \in A) \wedge (a_2 = -b_2 \in B) \quad \blacksquare \quad a_2 \in A \cap B \quad \blacksquare \quad a_2 = O \quad \blacksquare \quad a_2 = b_2 = O$$

$$(3.2.5) \quad \langle a_1, b_1 \rangle = \langle O, O \rangle = \langle a_2, b_2 \rangle$$

$$(3.3) \quad ((\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B) \wedge (O = a_1 + b_1) \wedge (O = a_2 + b_2)) \implies (\langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle)$$

$$(3.4) \quad \forall_{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B} \left(((O = a_1 + b_1) \wedge (O = a_2 + b_2)) \implies (\langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle) \right)$$

$$(3.5) \quad \left(\exists_{\langle a, b \rangle \in A \times B} (O = a + b) \right) \wedge \left(\forall_{\langle a_1, b_1 \rangle, \langle a_2, b_2 \rangle \in A \times B} \left(((O = a_1 + b_1) \wedge (O = a_2 + b_2)) \implies (\langle a_1, b_1 \rangle = \langle a_2, b_2 \rangle) \right) \right)$$

$$(3.6) \quad \left(\exists!_{\langle a,b \rangle \in A \times B} (O = a + b) \right) \wedge (DirSumEquiv) \quad \blacksquare \quad DirSum[A \oplus B, A, B, V, +, *]$$

$$(4) \quad (A \cap B = \{O\}) \implies (DirSum[A \oplus B, A, B, V, +, *])$$

$$(5) \quad (DirSum[A \oplus B, A, B, V, +, *]) \iff (A \cap B = \{O\})$$

$$NullSpace[N, A, m, n] := (Matrix[A, m, n]) \wedge (N = \{x \in \mathbb{R}^n \mid A * x = O\})$$

$$RowSpace[R, A, m, n] := (Matrix[A, m, n]) \wedge (R = \{x^T * A \in \mathbb{R}^n \mid x \in \mathbb{R}^m\})$$

$$ColSpace[C, A, m, n] := (Matrix[A, m, n]) \wedge (C = \{A * x \in \mathbb{R}^m \mid x \in \mathbb{R}^n\})$$

$$NullSubspace := (NullSpace[N, A, m, n]) \implies (Subspace[N, \mathbb{R}^n, +, *])$$

(1) TODO

$$RowSubspace := (RowSpace[R, A, m, n]) \implies (Subspace[R, \mathbb{R}^n, +, *])$$

(1) TODO

$$ColSubspace := (ColSpace[C, A, m, n]) \implies (Subspace[C, \mathbb{R}^m, +, *])$$

(1) TODO

3.5 Linear Combination, Linear Span, Linear Independence

$$LinComb[c, U, K, V, +, *] := (VectorSpace[V, +, *]) \wedge (n \in \mathbb{N}) \wedge (U \in V^n) \wedge (K \in \mathbb{R}^n) \wedge (c = \sum_{i=1}^n (k_i * u_i))$$

$$LinSpan[S', S, V, +, *] := \left(\begin{array}{l} (VectorSpace[V, +, *]) \wedge (S \in V^n) \wedge ((S = \emptyset) \implies (S' = \{O\})) \quad \wedge \\ ((S \neq \emptyset) \implies (S' = \{c \in V \mid (K \in \mathbb{R}^n) \wedge (LinComb[c, S, K, V, +, *])\})) \end{array} \right)$$

$$LinSpanSubContains := \forall_{S', S, V} \left((LinSpan[S', S, V, +, *]) \implies ((Subspace[S', V, +, *]) \wedge (S \subseteq S')) \right)$$

(1) $(S = \emptyset) \implies \dots$

$$(1.1) \quad LinSpan[S', S, V, +, *] \quad \blacksquare \quad S' = \{O\}$$

$$(1.2) \quad Subspace[\{O\}, V, +, *] \quad \blacksquare \quad Subspace[S', V, +, *]$$

$$(1.3) \quad S = \emptyset \subseteq \{O\} = S' \quad \blacksquare \quad S \subseteq S'$$

$$(1.4) \quad (Subspace[S', V, +, *]) \wedge (S \subseteq S')$$

(2) $(S = \emptyset) \implies ((Subspace[S', V, +, *]) \wedge (S \subseteq S'))$

(3) $(S \neq \emptyset) \implies \dots$

$$(3.1) \quad LinSpan[S', S, V, +, *] \quad \blacksquare \quad S' = \{c \in V \mid (K \in \mathbb{R}^n) \wedge (LinComb[c, S, K, V, +, *])\} \quad \blacksquare \quad S' \subseteq V$$

$$(3.2) \quad (\{0\}^n \subseteq \mathbb{R}^n) \wedge (LinComb[O, S, \{0\}^n, V, +, *]) \quad \blacksquare \quad O \in S' \quad \blacksquare \quad \emptyset \neq S'$$

$$(3.3) \quad (S' \subseteq V) \wedge (\emptyset \neq S') \quad \blacksquare \quad \emptyset \neq S' \subseteq V$$

(3.4) $(a, b \in S') \implies \dots$

$$(3.4.1) \quad \left(\exists_{K_a \in \mathbb{R}^n} (LinComb[a, S, K_a, V, +, *]) \right) \wedge \left(\exists_{K_b \in \mathbb{R}^n} (LinComb[b, S, K_b, V, +, *]) \right) \quad \blacksquare \quad (a = \sum_{i=1}^n (k_{ai} * s_i)) \wedge (b = \sum_{i=1}^n (k_{bi} * s_i))$$

$$(3.4.2) \quad a + b = \sum_{i=1}^n (k_{ai} * s_i) + \sum_{i=1}^n (k_{bi} * s_i) = \sum_{i=1}^n ((k_{ai} + k_{bi}) * s_i) \quad \blacksquare \quad a + b = \sum_{i=1}^n ((k_{ai} + k_{bi}) * s_i)$$

$$(3.4.3) \quad \langle k_{ai} + k_{bi} \mid i \in \mathbb{N}_{1,n} \rangle \in \mathbb{R}^n$$

$$(3.4.4) \quad \left(a + b = \sum_{i=1}^n ((k_{ai} + k_{bi}) * s_i) \right) \wedge (\langle k_{ai} + k_{bi} \mid i \in \mathbb{N}_{1,n} \rangle \in \mathbb{R}^n) \quad \dots$$

$$(3.4.5) \quad \dots \exists_{M \in \mathbb{R}^n} (a + b = \sum_{i=1}^n (m_i * s_i)) \quad \blacksquare \quad \exists_{M \in \mathbb{R}^n} (LinComb[a + b, S, M, V, +, *]) \quad \blacksquare \quad a + b \in S'$$

$$(3.5) \quad (a, b \in S') \implies (a + b \in S') \quad \blacksquare \quad \forall_{a,b \in S'} (a + b \in S')$$

(3.6) $((r \in \mathbb{R}) \wedge (u \in S')) \implies \dots$

$$(3.6.1) \quad \exists_{K \in \mathbb{R}^n} (LinComb[u, S, K, V, +, *]) \quad \blacksquare \quad u = \sum_{i=1}^n (k_i * s_i)$$

$$(3.6.2) \quad r * u = r * \sum_{i=1}^n (k_i * s_i) = \sum_{i=1}^n (r * (k_i * s_i)) = \sum_{i=1}^n (rk_i) * s_i \quad \blacksquare \quad r * u = \sum_{i=1}^n (rk_i) * s_i$$

$$(3.6.3) \quad \langle rk_i \in \mathbb{R} \mid i \in \mathbb{N}_{1,n} \rangle \in \mathbb{R}^n$$

$$(3.6.4) \quad (r * u = \sum_{i=1}^n (rk_i) * s_i) \wedge (\langle rk_i \in \mathbb{R} \mid i \in \mathbb{N}_{1,n} \rangle \in \mathbb{R}^n) \quad \blacksquare \quad \exists_{M \in \mathbb{R}^n} (r * u = \sum_{i=1}^n (m_i * s_i))$$

$$(3.6.5) \quad \exists_{M \in \mathbb{R}^n} (LinComb[r * u, S, M, V, +, *]) \quad \blacksquare \quad r * u \in S'$$

$$(3.7) \quad ((r \in \mathbb{R}) \wedge (u \in S')) \implies (r * u \in S') \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{u \in S'} (r * u \in S')$$

$$(3.8) \quad (SubspaceEquiv) \wedge (\emptyset \neq S' \subseteq V) \wedge (\forall_{a,b \in S'} (a + b \in S')) \wedge (\forall_{r \in \mathbb{R}} \forall_{u \in S'} (r * u \in S')) \quad \blacksquare \quad Subspace[S', V, +, *]$$

$$(3.9) \quad (s_i \in S) \implies \dots$$

$$(3.9.1) \quad K_s := \left\langle \begin{cases} 1 & j = i \\ 0 & j \neq i \end{cases} \mid j \in \mathbb{N}_{1,n} \right\rangle \quad \blacksquare \quad (K_s \in \mathbb{R}^n) \wedge \left(\sum_{j=1}^n (k_{sj} * s_j) = s_i \right)$$

$$(3.9.2) \quad \exists_{K \in \mathbb{R}^n} (LinComb[s_j, S, K, V, +, *]) \quad \blacksquare \quad s_j \in S'$$

$$(3.10) \quad (s_i \in S) \implies (s_i \in S') \quad \blacksquare \quad S \subseteq S'$$

$$(3.11) \quad (Subspace[S', V, +, *]) \wedge (S \subseteq S')$$

$$(4) \quad (S \neq \emptyset) \implies ((Subspace[S', V, +, *]) \wedge (S \subseteq S'))$$

$$(5) \quad \left((S = \emptyset) \implies ((Subspace[S', V, +, *]) \wedge (S \subseteq S')) \right) \wedge \left((S \neq \emptyset) \implies ((Subspace[S', V, +, *]) \wedge (S \subseteq S')) \right) \dots$$

$$(6) \quad \dots (Subspace[S', V, +, *]) \wedge (S \subseteq S')$$

$$LinSpanSubMinContains := \forall_{S', S, V, +, *} \left((LinSpan[S', S, V, +, *]) \implies \left(\forall_W (((Subspace[W, V, +, *]) \wedge (S \subseteq W)) \implies (S' \subseteq W)) \right) \right)$$

$$(1) \quad (s' \in S') \implies \dots$$

$$(1.1) \quad \exists_{K \in \mathbb{R}^n} (LinComb[s', S, K, V, +, *]) \quad \blacksquare \quad s' = \sum_{i=1}^n (k_i * s_i)$$

$$(1.2) \quad (S \subseteq W) \wedge (VectorSpace[W, V, +, *]) \quad \blacksquare \quad s' = \sum_{i=1}^n (k_i * s_i) \in W \quad \blacksquare \quad s' \in W$$

$$(2) \quad (s' \in S') \implies (s' \in W) \quad \blacksquare \quad S' \subseteq W$$

$$Spans[S, V, +, *] := LinSpan[V, S, V, +, *]$$

$$FinDim[V, +, *] := \exists_{S \in V^n} (Spans[S, V, +, *])$$

$$LinInd[S, V, +, *] := (VectorSpace[V, +, *]) \wedge (S \in V^n) \wedge \left((S \neq \emptyset) \implies \left(\forall_{K \in \mathbb{R}^n} ((LinComb[O, S, K, V, +, *]) \implies (K = \{0\}^n)) \right) \right)$$

$$ZeroDependent := (O \in S) \implies (\neg LinInd[S, V, +, *])$$

$$(1) \quad O \in S \quad \blacksquare \quad \exists_{u_i \in S} (u_i = O) \quad \blacksquare \quad K := \left\langle \begin{cases} 1 & u_i = O \\ 0 & u_i \neq O \end{cases} \mid i \in \mathbb{N}_{1,n} \right\rangle \quad \blacksquare \quad \{O\}^n \neq K \in \mathbb{R}^n$$

$$(2) \quad O = \sum_{i=1}^n (k_i * s_i) \quad \blacksquare \quad LinComb[O, S, K, V, +, *]$$

$$(3) \quad (LinComb[O, S, K, V, +, *]) \wedge (\{O\}^n \neq K \in \mathbb{R}^n) \quad \blacksquare \quad \exists_{K \in \mathbb{R}^n} ((LinComb[O, S, K, V, +, *]) \wedge (K \neq \{0\}^n)) \quad \blacksquare \quad \neg LinInd[S, V, +, *]$$

$$SingletonNonZeroIndependent := (v \neq O) \implies (LinInd[\langle v \rangle, V, +, *])$$

$$(1) \quad \left((\langle r \rangle \in \mathbb{R}^1) \wedge (LinComb[O, \langle v \rangle, \langle r \rangle, V, +, *]) \right) \implies \dots$$

$$(1.1) \quad (ZeroVectorEquiv) \wedge (r * v = O) \quad \blacksquare \quad (r * v = O) \iff ((r = 0) \vee (v \neq O))$$

$$(1.2) \quad v \neq O \quad \blacksquare \quad r = 0$$

$$(2) \quad \left((\langle r \rangle \in \mathbb{R}^1) \wedge (LinComb[O, \langle v \rangle, \langle r \rangle, V, +, *]) \right) \implies (r = 0) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} ((LinComb[O, \langle v \rangle, \langle r \rangle, V, +, *]) \implies (r = 0))$$

$$(3) \quad LinInd[\langle v \rangle, V, +, *]$$

$$SubIndependent := \forall_{V, A, B} \left(\begin{aligned} & ((VectorSpace[V, +, *]) \wedge (A \subseteq B) \wedge (A \in V^n) \wedge (B \in V^m)) \implies \\ & ((LinInd[B, V, +, *]) \implies (LinInd[A, V, +, *])) \end{aligned} \right)$$

$$(1) \quad ((K \in \mathbb{R}^n) \wedge (LinComb[O, A, K, V, +, *])) \implies \dots$$

$$(1.1) \quad n \leq m \quad \blacksquare \quad L := \left\langle \left\{ \begin{pmatrix} k_j & j \leq n \\ 0 & j > n \end{pmatrix} \mid j \in \mathbb{N}_{1,m} \right\rangle \mid L \in \mathbb{R}^m \right.$$

$$(1.2) \quad A \subseteq B \quad \blacksquare \quad \forall_{j \in \mathbb{N}_{1,n}} (a_j = b_j) \quad \blacksquare \quad \sum_{i=1}^n (k_i * a_i) = \sum_{j=1}^m (l_j * b_j))$$

$$(1.3) \quad \text{LinComb}[O, A, K, V, +, *] \quad \blacksquare \quad O = \sum_{i=1}^n (k_i * a_i) = \sum_{j=1}^m (l_j * b_j) \quad \blacksquare \quad \text{LinComb}[O, B, L, V, +, *]$$

$$(1.4) \quad (\text{LinInd}[B, V, +, *]) \wedge (\text{LinComb}[O, B, L, V, +, *]) \quad \blacksquare \quad L = \{0\}^m \quad \blacksquare \quad K = \{0\}^n$$

$$(2) \quad ((K \in \mathbb{R}^n) \wedge (\text{LinComb}[O, A, K, V, +, *])) \implies (K = \{0\}^n) \quad \blacksquare \quad \text{LinInd}[A, V, +, *]$$

$$\text{SuperDependent} := \forall_{V,A,B} \left(((\text{VectorSpace}[V, +, *]) \wedge (A \subseteq B \subseteq V)) \implies ((\neg \text{LinInd}[A, V, +, *]) \implies (\neg \text{LinInd}[B, V, +, *])) \right)$$

$$(1) \quad \neg \text{LinInd}[A, V, +, *] \quad \blacksquare \quad \exists_K ((\text{LinComb}[O, A, K, V, +, *]) \wedge (K \neq \{0\}^n))$$

$$(2) \quad n \leq m \quad \blacksquare \quad L := \left\langle \left\{ \begin{pmatrix} k_j & j \leq n \\ 0 & j > n \end{pmatrix} \mid j \in \mathbb{N}_{1,m} \right\rangle \mid L \in \mathbb{R}^m \right.$$

$$(3) \quad A \subseteq B \quad \blacksquare \quad \forall_{j \in \mathbb{N}_{1,n}} (a_j = b_j) \quad \blacksquare \quad \sum_{i=1}^n (k_i * a_i) = \sum_{j=1}^m (l_j * b_j))$$

$$(4) \quad \text{LinComb}[O, A, K, V, +, *] \quad \blacksquare \quad \text{LinComb}[O, B, L, V, +, *]$$

$$(5) \quad K \neq \{0\}^n \quad \blacksquare \quad L \neq \{0\}^m$$

$$(6) \quad \exists_L ((\text{LinComb}[O, B, L, V, +, *]) \wedge (L \neq \{0\}^m)) \quad \blacksquare \quad \neg \text{LinInd}[B, V, +, *]$$

$$\text{LinDepProp} := \forall_{S,V} \left((\neg \text{LinInd}[S, V, +, *]) \implies \left(\exists_{s_j \in S} \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \right)$$

$$(1) \quad \neg \text{LinInd}[S, V, +, *] \quad \blacksquare \quad \exists_{K \in \mathbb{R}^n} ((\text{LinComb}[O, S, K, V, +, *]) \wedge (K \neq \{0\}^n))$$

$$(2) \quad K \neq \{0\}^n \quad \blacksquare \quad \exists_{j \in \mathbb{N}_{1,n}} \left((k_j \neq 0) \wedge \left(\forall_{i \in \mathbb{N}_{j+1,n}} (k_i = 0) \right) \right) \quad \dots$$

$$(3) \quad \dots \sum_{i=1}^n (k_i * s_i) = \sum_{i=1}^j (k_i * s_i) = \sum_{i=1}^{j-1} (k_i * s_i) + k_j * s_j \quad \blacksquare \quad \sum_{i=1}^n (k_i * s_i) = \sum_{i=1}^{j-1} (k_i * s_i) + k_j * s_j$$

$$(4) \quad (\text{LinComb}[O, S, K, V, +, *]) \wedge \left(\sum_{i=1}^n (k_i * s_i) = \sum_{i=1}^{j-1} (k_i * s_i) + k_j * s_j \right) \quad \blacksquare \quad O = \sum_{i=1}^n (k_i * s_i) = \sum_{i=1}^{j-1} (k_i * s_i) + k_j * s_j$$

$$(5) \quad s_j = (-1/k_j) \sum_{i=1}^{j-1} (k_i * s_i) = \sum_{i=1}^{j-1} ((-k_i/k_j) * s_i) \quad \blacksquare \quad s_j = \sum_{i=1}^{j-1} ((-k_i/k_j) * s_i)$$

$$(6) \quad \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *])$$

$$\text{LinDepPropCorollary} := \forall_{P,S,V} \left(((\neg \text{LinInd}[S, V, +, *]) \wedge (\text{LinSpan}[P, S, V, +, *])) \implies \left(\exists_{s_j \in S} (\text{LinSpan}[P, S \setminus \{s_j\}, V, +, *]) \right) \right)$$

$$(1) \quad \text{LinDepProp} \quad \blacksquare \quad \exists_{s_j \in S} \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *])$$

$$(2) \quad \forall_{u \in P} \left(\left(\exists_{K_1} (\text{LinComb}[u, S, K_1, V, +, *]) \right) \implies \left(\exists_{K_2} (\text{LinComb}[u, S \setminus \{s_j\}, K_2, V, +, *]) \right) \right) \quad \blacksquare \quad \text{LinSpan}[P, S \setminus \{s_j\}, V, +, *]$$

$$\text{LinIndEquiv} := \forall_{S,V} \left((\text{LinInd}[S, V, +, *]) \iff \left(\forall_{s_j \in S} \forall_{K \in \mathbb{R}^{n-1}} (\neg \text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \right)$$

$$(1) \quad \text{LinDepProp} \quad \blacksquare \quad (\neg \text{LinInd}[S, V, +, *]) \implies \left(\exists_{s_j \in S} \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \quad \dots$$

$$(2) \quad \dots \left(\forall_{s_j \in S} \forall_{K \in \mathbb{R}^{n-1}} (\neg \text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \implies (\text{LinInd}[S, V, +, *])$$

$$(3) \quad \left(\exists_{s_j \in S} \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \implies \dots$$

$$(3.1) \quad L := \left\langle \left\{ \begin{pmatrix} k_i & i \neq j \\ -1 & i = j \end{pmatrix} \mid i \in \mathbb{N}_{1,n} \right\rangle \mid (L \in \mathbb{R}^n) \wedge (L \neq \{0\}^n) \right.$$

$$(3.2) \quad \text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *] \quad \blacksquare \quad \dots \quad \sum_{i=1}^{j-1} (k_i * s_i) + k_j * s_j = \sum_{i=1}^{j-1} (k_i * s_i) + - \sum_{i=1}^{j-1} (k_i * s_i) = O \quad \dots$$

$$(3.3) \quad \dots \text{LinComb}[O, S, L, V, +, *]$$

$$(3.4) \quad (\text{LinComb}[O, S, L, V, +, *]) \wedge (L \neq \{0\}^n) \quad \blacksquare \quad \exists_{L \in \mathbb{R}^n} ((\text{LinComb}[O, S, L, V, +, *]) \wedge (L \neq \{0\}^n)) \quad \blacksquare \quad (\neg \text{LinInd}[S, V, +, *])$$

$$(4) \quad \left(\exists_{s_j \in S} \exists_{K \in \mathbb{R}^{n-1}} (\text{LinComb}[s_j, S \setminus \{s_j\}, K, V, +, *]) \right) \implies (\neg \text{LinInd}[S, V, +, *])$$

$$(5) \quad (LinInd[S, V, +, *]) \implies \left(\forall_{s_j \in S} \forall_{K \in \mathbb{R}^{n-1}} (\neg LinComb[s_j, S \setminus \{s_j\}, K, V, +, *]) \right)$$

$$(6) \quad (LinInd[S, V, +, *]) \iff \left(\forall_{s_j \in S} \forall_{K \in \mathbb{R}^{n-1}} (\neg LinComb[s_j, S \setminus \{s_j\}, K, V, +, *]) \right)$$

$$LinIndSuperspace := \forall_{U, V} \left((Subspace[U, V]) \implies \left(\forall_W ((LinInd[W, U, +, *]) \implies (LinInd[W, V, +, *])) \right) \right)$$

$$(1) \quad (\neg LinInd[W, V, +, *]) \implies \dots$$

$$(1.1) \quad \exists_{j \in W} (LinComb[j, W \setminus \{j\}, +, *]) \quad \blacksquare \quad \neg LinInd[W, U, +, *]$$

$$(1.2) \quad (LinInd[W, U, +, *]) \wedge (\neg LinInd[W, V, +, *]) \quad \blacksquare \quad \perp$$

$$(2) \quad (\neg LinInd[W, V, +, *]) \implies \perp \quad \blacksquare \quad LinInd[W, V, +, *]$$

3.6 Bases and Dimensions

$$Basis[S, V, +, *] := (Spans[S, V, +, *]) \wedge (LinInd[S, V, +, *])$$

$$BasisEquiv := \forall_{S, V} ((Basis[S, V, +, *]) \iff (\forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *])))$$

$$(1) \quad (Basis[S, V, +, *]) \implies \dots$$

$$(1.1) \quad (v \in V) \implies \dots$$

$$(1.1.1) \quad Basis[S, V, +, *] \quad \blacksquare \quad Spans[V, S, +, *] \quad \blacksquare \quad \exists_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *])$$

$$(1.1.2) \quad ((K_1, K_2 \in \mathbb{R}^n) \wedge (LinComb[v, S, K_1, V, +, *]) \wedge (LinComb[v, S, K_2, V, +, *])) \implies \dots$$

$$(1.1.2.1) \quad (v = \sum(k_{1i} * s_i)) \wedge (v = \sum(k_{2i} * s_i))$$

$$(1.1.2.2) \quad 0 = v - v = \sum(k_{1i} * s_i) - \sum(k_{2i} * s_i) = \sum((k_{1i} - k_{2i}) * s_i)$$

$$(1.1.2.3) \quad L := \langle k_{1i} - k_{2i} \mid i \in \mathbb{N}_{i=1}^n \rangle \in \mathbb{R}^n$$

$$(1.1.2.4) \quad (LinInd[S, V, +, *]) \wedge (LinComb[0, S, L, V, +, *]) \quad \blacksquare \quad L = \{0\}^n \quad \blacksquare \quad K_2 = K_1$$

$$(1.1.3) \quad ((K_1, K_2 \in \mathbb{R}^n) \wedge (LinComb[v, S, K_1, V, +, *]) \wedge (LinComb[v, S, K_2, V, +, *])) \implies (K_1 = K_2)$$

$$(1.1.4) \quad \forall_{K_1, K_2 \in \mathbb{R}^n} ((LinComb[v, S, K_1, V, +, *]) \wedge (LinComb[v, S, K_2, V, +, *]) \implies (K_1 = K_2))$$

$$(1.1.5) \quad \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *])$$

$$(1.2) \quad (v \in V) \implies (\exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *]))$$

$$(2) \quad (Basis[S, V, +, *]) \implies (\forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *]))$$

$$(3) \quad (\forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *])) \implies \dots$$

$$(3.1) \quad \forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *]) \quad \blacksquare \quad \forall_{v \in V} \exists_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *]) \quad \blacksquare \quad Spans[S, V, +, *]$$

$$(3.2) \quad 0 \in V \quad \blacksquare \quad \exists!_{K \in \mathbb{R}^n} (LinComb[0, S, K, V, +, *])$$

$$(3.3) \quad (K \neq \{0\}^n) \implies (\neg \exists!_{K \in \mathbb{R}^n} (LinComb[0, S, K, V, +, *])) \implies \perp \quad \blacksquare \quad K = \{0\}^n$$

$$(3.4) \quad (\exists!_{K \in \mathbb{R}^n} (LinComb[0, S, K, V, +, *])) \wedge (K = \{0\}^n) \quad \blacksquare \quad LinInd[S, V, +, *]$$

$$(3.5) \quad (Spans[S, V, +, *]) \wedge (LinInd[S, V, +, *]) \quad \blacksquare \quad Basis[S, V, +, *]$$

$$(4) \quad (\forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, S, K, V, +, *])) \implies (Basis[S, V, +, *])$$

$$SpanReduceBasis := \forall_{S, V} \left((Spans[S, V, +, *]) \implies \left(\exists_B ((B \subseteq S) \wedge (Basis[B, V, +, *])) \right) \right)$$

$$(1) \quad LinDepPropCorollary \quad \blacksquare \quad \exists_B ((B \subseteq S) \wedge (LinInd[B, V, +, *]) \wedge (Spans[B, V, +, *])) \quad \blacksquare \quad \exists_B ((B \subseteq S) \wedge (Basis[B, V, +, *]))$$

$$(2) \quad \text{TODO - formalize removing latter entries first}$$

$$FinDimBasis := \forall_V \left((FinDim[V, +, *]) \implies (\exists_B (Basis[B, V, +, *])) \right)$$

$$(1) \quad FinDim[V, +, *] \quad \blacksquare \quad \exists_{S \in V^n} (Spans[S, V, +, *])$$

$$(2) \quad (SpanReduceBasis) \wedge (Spans[S, V, +, *]) \quad \blacksquare \quad \exists_B (Basis[B, V, +, *])$$

$$LinIndExpandBasis := \forall_{L, V} \left((LinInd[L, V, +, *]) \implies \left(\exists_B ((L \subseteq B) \wedge (Basis[B, V, +, *])) \right) \right)$$

-
- (1) $FinDimBasis \dashv \exists_C (Basis[C, V, +, *])$
-
- (2) $S := L \cup C$
-
- (3) $Basis[C, V, +, *] \dashv Spans[C, V, +, *] \dashv Spans[S, V, +, *]$
-
- (4) $SpanReduceBasis \dashv \left(\exists_B ((B \subseteq S) \wedge (Basis[B, V, +, *])) \right) \wedge (L \subseteq B)$
-

$$SpanLinIndLength := \forall_{S,T,V} \left(((Span[S, V, +, *]) \wedge (LinInd[T, V, +, *])) \implies (|T| \leq |S|) \right)$$

-
- (1) $((Span[S, V, +, *]) \wedge (|T| > |S|)) \implies \dots$
-
- (1.1) $Span[S, V, +, *] \dashv \forall_{i \in \mathbb{N}_{1,|H|}} \exists_{K_i \mathbb{R}^{|S|}} (LinComb[t_i, S, K_i V, +, *])$
-
- (1.2) $|H| > |S| \dashv \exists_{L \in \mathbb{R}^{|H|-1}} (LinComb[t_{|H|}, T \setminus \{t_{|H|}\}, L, V, +, *])$
-
- (1.3) $L = -1 * K \dashv (\sum (K + L) = O) \wedge (K + L \neq \{0\}^{|T|}) \dashv \neg LinInd[T, V, +, *]$
-
- (1.4) TODO - tidy up
-
- (2) $((Span[S, V, +, *]) \wedge (|T| > |S|)) \implies (\neg LinInd[T, V, +, *]) \dashv ((Span[S, V, +, *]) \wedge (LinInd[T, V, +, *])) \implies (|T| \leq |S|)$
-

$$BasisLength := \forall_{S,T,V} \left(((Basis[S, V, +, *]) \wedge (Basis[T, V, +, *])) \implies (|T| = |S|) \right)$$

-
- (1) $(Span[T, V, +, *]) \wedge (LinInd[S, V, +, *]) \dashv |S| \leq |T|$
-
- (2) $(Span[S, V, +, *]) \wedge (LinInd[T, V, +, *]) \dashv |T| \leq |S|$
-
- (3) $(|S| \leq |T|) \wedge (|T| \leq |S|) \dashv |T| = |S|$
-

$$Dim[d, V, +, *] := ((V = \{O\}) \implies (d = 0)) \wedge ((V \neq \{O\}) \implies ((\exists_B (Basis[B, V, +, *])) \wedge (d = |B|)))$$

$$LinIndLengthDim := \forall_{U,V} \left(((LinInd[U, V, +, *]) \wedge (Dim[|U|, V, +, *])) \implies (Basis[U, V, +, *]) \right)$$

-
- (1) $(LinIndExpandBasis) \wedge (LinInd[U, V, +, *]) \dashv \exists_B ((U \subseteq B) \wedge (Basis[B, V, +, *]))$
-
- (2) $(BasisLength) \wedge (Dim[|U|, V, +, *]) \wedge (Basis[B, V, +, *]) \dashv |B| = |U| \dashv B = U \dashv Basis[U, V, +, *]$
-

$$SpanLengthDim := \forall_{U,V} \left(((Spans[U, V, +, *]) \wedge (Dim[|U|, V, +, *])) \implies (Basis[U, V, +, *]) \right)$$

-
- (1) $(SpanReduceBasis) \wedge (Spans[U, V, +, *]) \dashv \exists_B ((B \subseteq U) \wedge (Basis[B, V, +, *]))$
-
- (2) $(BasisLength) \wedge (Dim[|U|, V, +, *]) \wedge (Basis[B, V, +, *]) \dashv |B| = |U| \dashv B = U \dashv Basis[U, V, +, *]$
-

$$LinDepLengthDim := \forall_{U,V} \left(((U \subseteq V) \wedge (|U| > Dim[V])) \implies (\neg LinInd[U, V, +, *]) \right)$$

-
- (1) Contrapositive of $BasisLinearIndCard$
-
- (2) TODO - cleanup
-

$$NonSpanLengthDim := \forall_{U,V} \left(((U \subseteq V) \wedge (|U| < Dim[V])) \implies (\neg Spans[U, V, +, *]) \right)$$

-
- (1) Suppose $Spans[U, V, +, *]$, $B = SpanReduceBasis[U]$ to form a basis, $(|B| \leq |U| < Dim[V]) \wedge |B| = Dim[V] \dashv \perp$
-
- (2) $\neg Spans[U, V, +, *]$
-
- (3) TODO - cleanup
-

3.7 Rank

$$Nullity[n, A] := (NullSpace[N, A]) \wedge (Dim[n, N, +, *])$$

$$Rank[r, A, m, n] := (Matrix[A, m, n]) \wedge (RowSpace[R, A, m, n]) \wedge (Dim[r, R, A, +, *])$$

$$RowRankEqColRank := \forall_A (TODO)$$

-
- (1) TODO
-

$$RankNullity := \forall_A ((Matrix[A, m, n]) \implies (Rank[A] + Nullity[A] = n))$$

(1) TODO

$$RankInv := \forall_A \left((Matrix[A, m, n]) \implies ((Rank[A] = n) \iff (Inv[A])) \right)$$

(1) TODO

$$RankNonTrivialSol := \left(\exists_X ((A * X = O) \wedge (X \neq O)) \right) \iff (Rank[A] < n)$$

(1) TODO

$$RankUniqueSol := (\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B])) \iff (Rank[A] = n)$$

(1) TODO

$$SquareTheorems_8 := \forall_{A \in \mathcal{M}} \left(\begin{array}{l} (Invertible[A]) \\ (RowEquiv[A, I_n]) \\ \left(\forall_X ((X = O) \iff (Sol[X, A, O])) \right) \\ (\forall_{B \in \mathcal{M}} \exists!_{X \in \mathcal{M}} (Sol[X, A, B])) \\ (Rank[A] = n) \\ (Nullity[A] = 0) \\ (\text{The rows form a linearly independent set of vectors (to get full rank)}) \\ (\text{The columns form a linearly independent set of vectors (to get full rank)}) \end{array} \right)$$

3.8 Linear Transformations

$$LinTrans[L, V, +_v, *_v, W, +_w, *_w] := \left(\begin{array}{l} (Function[f, V, W]) \wedge (VectorSpace[V, +_v, *_v]) \wedge (VectorSpace[W, +_w, *_w]) \wedge \\ \left(\forall_{\alpha, \beta \in V} (L(\alpha +_v \beta) = L(\alpha) +_w L(\beta)) \right) \wedge \left(\forall_{r \in \mathbb{R}} \forall_{\alpha \in V} (L(r *_v \alpha) = r *_w L(\alpha)) \right) \end{array} \right)$$

$$LinOp[L, V, +_v, *_v] := LinTrans[L, V, +_v, *_v, V, +_v, *_v]$$

$$\mathcal{L}[V, W] := \{L \mid LinTrans[L, V, +_v, *_v, W, +_w, *_w]\}$$

$$ZeroMapsToZero := \forall_{L, V, W} \left((LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \implies (L(O_v) = O_w) \right)$$

(1) $L(O_v) = L(O_v +_v O_v) = L(O_v) +_w L(O_v)$

(2) $O_w = L(O_v) - L(O_v) = L(O_v)$

$$SplitAddInv := \forall_{L, V, W} \left((LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \implies \left(\forall_{\alpha, \beta \in V} (L(\alpha -_v \beta) = L(\alpha) -_w L(\beta)) \right) \right)$$

(1) $L(\alpha - \beta) = L(\alpha + (-\beta)) = L(\alpha) + L(-\beta) = L(\alpha) + (-1) * L(\beta) = L(\alpha) - L(\beta)$

$$UniqBasisLT := \forall_{V, W} \left(\begin{array}{l} ((VectorSpace[V, +_v, *_v]) \wedge (VectorSpace[W, +_w, *_w]) \wedge (Basis[A, V, +_v, *_v]) \wedge (Basis[B, W, +_w, *_w])) \implies \\ \left(\exists!_T \left((LinTrans[T, V, +_v, *_v, W, +_w, *_w]) \wedge \left(\forall_{i \in \mathbb{N}_{1,n}} (T(a_i) = b_i) \right) \right) \right) \end{array} \right)$$

(1) $T(\sum_{i=1}^n (k_i * a_i)) := \sum_{i=1}^n (k_i * b_i)$

(2) $(i \in \mathbb{N}_{1,n}) \implies \dots$

$$(2.1) \quad L := \left\langle \left\{ \begin{array}{ll} 1 & j = i \\ 0 & j \neq i \end{array} \right\} \mid j \in \mathbb{N}_{1,n} \right\rangle \quad \blacksquare \quad L \in \mathbb{R}^n$$

$$(2.2) \quad T(a_i) = T(\sum_{i=1}^n (l_i * a_i)) = \sum_{i=1}^n (l_i * b_i) = b_i \quad \blacksquare \quad T(a_i) = b_i$$

(3) $(i \in \mathbb{N}_{1,n}) \implies (T(a_i) = b_i) \quad \blacksquare \quad \forall_{i \in \mathbb{N}_{1,n}} (T(a_i) = b_i)$

(4) $(BasisEquiv) \wedge (Basis[A, V, +_v, *_v]) \quad \blacksquare \quad \forall_{v \in V} \exists!_{K \in \mathbb{R}^n} (LinComb[v, A, K, V, +, *]) \quad \dots$

(5) $\dots \forall_{v_1, v_2 \in V} \left((v_1 = v_2) \implies (T(v_1) = T(v_2)) \right) \quad \blacksquare \quad Function[T, V, W]$

(6) $(\alpha, \beta \in V) \implies \dots$

$$(6.1) \quad \left(\exists_{K_\alpha} (LinComb[\alpha, A, K_\alpha, V, +_v, *_v]) \right) \wedge \left(\exists_{K_\beta} (LinComb[\beta, A, K_\beta, V, +_v, *_v]) \right) \quad \blacksquare \quad \left(\alpha = \sum_{i=1}^n (k_{\alpha_i} * a_i) \right) \wedge \left(\beta = \sum_{i=1}^n (k_{\beta_i} * a_i) \right)$$

$$(6.2) \quad T(\alpha + \beta) = T\left(\sum_{i=1}^n (k_{\alpha_i} * a_i) + \sum_{i=1}^n (k_{\beta_i} * a_i)\right) = T\left(\sum_{i=1}^n ((k_{\alpha_i} + k_{\beta_i}) * a_i)\right) = \sum_{i=1}^n ((k_{\alpha_i} + k_{\beta_i}) * b_i) = \dots$$

$$(6.3) \quad \dots \sum_{i=1}^n (k_{\alpha_i} * b_i) + \sum_{i=1}^n (k_{\beta_i} * b_i) = T\left(\sum_{i=1}^n (k_{\alpha_i} * a_i)\right) + T\left(\sum_{i=1}^n (k_{\beta_i} * a_i)\right) = T(\alpha) + T(\beta)$$

$$(7) \quad (\alpha, \beta \in V) \implies (L(\alpha +_v \beta) = L(\alpha) +_w L(\beta)) \quad \blacksquare \quad \forall_{\alpha, \beta \in V} (L(\alpha +_v \beta) = L(\alpha) +_w L(\beta))$$

$$(8) \quad ((r \in \mathbb{R}) \wedge (\alpha \in V)) \implies \dots$$

$$(8.1) \quad \exists_K (LinComb[\alpha, A, K, V, +_v, *_v]) \quad \blacksquare \quad \alpha = \sum_{i=1}^n (k_i * a_i)$$

$$(8.2) \quad L(r *_v \alpha) = L(r *_v \sum_{i=1}^n (k_i * a_i)) = L\left(\sum_{i=1}^n ((rk_i) *_v a_i)\right) = \dots$$

$$(8.3) \quad \dots \sum_{i=1}^n ((rk_i) *_w b_i) = r *_w \sum_{i=1}^n (k_i *_w b_i) = r *_w L\left(\sum_{i=1}^n (k_i *_v a_i)\right) = r *_w L(\alpha)$$

$$(9) \quad ((r \in \mathbb{R}) \wedge (\alpha \in V)) \implies (L(r *_v \alpha) = r *_w L(\alpha)) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{\alpha \in V} (L(r *_v \alpha) = r *_w L(\alpha))$$

$$(10) \quad \left(\forall_{i \in \mathbb{N}_{1,n}} (T(a_i) = b_i)\right) \wedge (Function[T, V, W]) \wedge \left(\forall_{\alpha, \beta \in V} (L(\alpha +_v \beta) = L(\alpha) +_w L(\beta))\right) \wedge \left(\forall_{r \in \mathbb{R}} \forall_{\alpha \in V} (L(r *_v \alpha) = r *_w L(\alpha))\right) \wedge \dots$$

$$(11) \quad \dots (VectorSpace[V, +_v, *_v]) \wedge (VectorSpace[W, +_w, *_w]) \quad \blacksquare \quad \left(\forall_{i \in \mathbb{N}_{1,n}} (T(a_i) = b_i)\right) \wedge (LinTrans[T, V, +_v, *_v, W, +_w, *_w])$$

$$(12) \quad \left(\left(\forall_{i \in \mathbb{N}_{1,n}} (T_2(a_i) = b_i)\right) \wedge (LinTrans[T_2, V, +_v, *_v, W, +_w, *_w])\right) \implies \dots$$

$$(12.1) \quad \forall_{i \in \mathbb{N}_{1,n}} (T_2(a_i) = b_i) \quad \blacksquare \quad \forall_{i \in \mathbb{N}_{1,n}} (T_2(c_i * a_i) = c_i * b_i) \quad \blacksquare \quad T_2\left(\sum_{i=1}^n (c_i * a_i)\right) = \sum_{i=1}^n (c_i * b_i) \quad \blacksquare \quad T_2 = T$$

$$(13) \quad \left(\left(\forall_{i \in \mathbb{N}_{1,n}} (T_2(a_i) = b_i)\right) \wedge (LinTrans[T_2, V, +_v, *_v, W, +_w, *_w])\right) \implies (T_2 = T)$$

$$+_L[S + T, S, T] := (S + T)(v) = S(v) + T(v)$$

$$*_L[r * T, r, T] := (r * T)(v) = r * (T(v))$$

$$LTVectorSpace := \forall_{V, W} (VectorSpace[\mathcal{L}[V, W], +_L, *_L])$$

(1) TODO

$$*_L[S * T, S, T] := (S * T)(v) = S(T(v))$$

$$LTProdProperties := (associativity) \wedge (identity) \wedge (distributive)$$

(1) TODO

$$Ker[ker_L, L, V, +_v, *_v, W, +_w, *_w] := (LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge (ker_L = \{\alpha \in V \mid L(\alpha) = O_w\})$$

$$KerSubspace := \forall_{L, V, W} ((Ker[ker_L, L, V, +_v, *_v, W, +_w, *_w]) \implies (Subspace[ker_L, V, +_v, *_v]))$$

$$(1) \quad ZeroMapsToZero \quad \blacksquare \quad L(O_v) = O_w \quad \blacksquare \quad O_v \in ker_L \quad \blacksquare \quad \emptyset \neq ker_L \quad \blacksquare \quad \emptyset \neq ker_L \subseteq V$$

$$(2) \quad (\alpha, \beta \in ker_L) \implies \dots$$

$$(2.1) \quad (L(\alpha) = O_w) \wedge (L(\beta) = O_w)$$

$$(2.2) \quad L(\alpha + \beta) = L(\alpha) + L(\beta) = O_w + O_w = O_w \quad \blacksquare \quad L(\alpha + \beta) \in ker_L$$

$$(3) \quad (\alpha, \beta \in ker_L) \implies (\alpha + \beta \in ker_L) \quad \blacksquare \quad \forall_{\alpha, \beta \in ker_L} (\alpha + \beta \in ker_L)$$

$$(4) \quad ((r \in \mathbb{R}) \wedge (\alpha \in ker_L)) \implies \dots$$

$$(4.1) \quad L(\alpha) = O_w \quad \blacksquare \quad L(r * \alpha) = r * L(\alpha) = r * O_w = O_w \quad \blacksquare \quad r * \alpha \in ker_L$$

$$(5) \quad ((r \in \mathbb{R}) \wedge (\alpha \in ker_L)) \implies (r * \alpha \in ker_L) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{\alpha \in ker_L} (r * \alpha \in ker_L)$$

$$(6) \quad (SubspaceEquiv) \wedge (\emptyset \neq ker_L \subseteq V) \wedge \left(\forall_{\alpha, \beta \in ker_L} (\alpha + \beta \in ker_L)\right) \wedge \left(\forall_{r \in \mathbb{R}} \forall_{\alpha \in ker_L} (r * \alpha \in ker_L)\right) \quad \blacksquare \quad Subspace[ker_L, V, +_v, *_v]$$

$$Rng[rng_L, L, V, +_v, *_v, W, +_w, *_w] := (LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge (rng_L = \{\beta \in W \mid \exists_{\alpha \in V} (\beta = L(\alpha))\})$$

$$RangeSubspace := \forall_{L, V, W} ((Ran[rng_L, L, V, +_v, *_v, W, +_w, *_w]) \implies (Subspace[rng_L, W, +_w, *_w]))$$

$$(1) \quad ZeroMapsToZero \quad \blacksquare \quad O_w = L(O_v) \quad \blacksquare \quad \exists_{\alpha \in V} (O_w = L(\alpha)) \quad \blacksquare \quad O_w \in rng_L \quad \blacksquare \quad \emptyset \neq rng_L \quad \blacksquare \quad \emptyset \neq rng_L \subseteq W$$

$$(2) \quad (\alpha, \beta \in rng_L) \implies \dots$$

-
- (2.1) $\left(\exists_{u \in V} (\alpha = L(u)) \right) \wedge \left(\exists_{v \in V} (\beta = L(v)) \right)$
-
- (2.2) $\alpha + \beta = L(u) + L(v) = L(u + v) \quad \blacksquare \quad \exists_{w \in V} (\alpha + \beta = L(w)) \quad \blacksquare \quad \alpha + \beta \in \text{rng}_L$
-
- (3) $(\alpha, \beta \in \text{rng}_L) \implies (\alpha + \beta \in \text{rng}_L) \quad \blacksquare \quad \forall_{\alpha, \beta \in \text{rng}_L} (\alpha + \beta \in \text{rng}_L)$
-
- (4) $((r \in \mathbb{R}) \wedge (\alpha \in \text{rng}_L)) \implies \dots$
-
- (4.1) $\exists_{v \in V} (\alpha = L(v)) \quad \blacksquare \quad L(r * v) = r * L(v) = r * \alpha \quad \blacksquare \quad \exists_{w \in V} (r * \alpha = L(w)) \quad \blacksquare \quad r * \alpha \in \text{rng}_L$
-
- (5) $((r \in \mathbb{R}) \wedge (\alpha \in \text{rng}_L)) \implies (r * \alpha \in \text{rng}_L) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{\alpha \in \text{rng}_L} (r * \alpha \in \text{rng}_L)$
-
- (6) $(\text{SubspaceEquiv}) \wedge (\emptyset \neq \text{rng}_L \subseteq W) \wedge \left(\forall_{\alpha, \beta \in \text{rng}_L} (\alpha + \beta \in \text{rng}_L) \right) \wedge \left(\forall_{r \in \mathbb{R}} \forall_{\alpha \in \text{rng}_L} (r * \alpha \in \text{rng}_L) \right) \quad \blacksquare \quad \text{Subspace}[\text{rng}_L, W, +_w, *_w]$
-

$$\text{KerInjective} := \forall_{L, V, W} \left((\text{Ker}[\text{ker}_L, L, V, +_v, *_v, W, +_w, *_w]) \implies ((\text{Injective}[L, V, W]) \iff (\text{ker}_L = \{O_v\})) \right)$$

-
- (1) $(\text{Injective}[L, V, W]) \implies \dots$
-
- (1.1) $\text{ZeroMapsToZero} \quad \blacksquare \quad L(O_v) = O_w$
-
- (1.2) $O_v \in \text{ker}_L \quad \blacksquare \quad \{O_v\} \subseteq \text{ker}_L$
-
- (1.3) $(v \in \text{ker}_L) \implies \dots$
-
- (1.3.1) $L(v) = O_w$
-
- (1.3.2) $(\text{Injective}[L, V, W]) \wedge (L(O_v) = O_w) \quad \blacksquare \quad O_v = v$
-
- (1.4) $(v \in \text{ker}_L) \implies (v = O_v) \quad \blacksquare \quad \text{ker}_L \subseteq \{O_v\}$
-
- (1.5) $(\{O_v\} \subseteq \text{ker}_L) \wedge (\text{ker}_L \subseteq \{O_v\}) \quad \blacksquare \quad \text{ker}_L = \{O_v\}$
-
- (2) $(\text{Injective}[L, V, W]) \implies (\text{ker}_L = \{O_v\})$
-
- (3) $(\text{ker}_L = \{O_v\}) \implies \dots$
-
- (3.1) $\left((u, v \in V) \wedge (L(u) = L(v)) \right) \implies \dots$
-
- (3.1.1) $O_w = L(u) - L(v) = L(u - v) \quad \blacksquare \quad u - v \in \text{ker}_L$
-
- (3.1.2) $\text{ker}_L = \{O_v\} \quad \blacksquare \quad u - v = O_v \quad \blacksquare \quad u = v$
-
- (3.2) $\left((u, v \in V) \wedge (L(u) = L(v)) \right) \implies (u = v) \quad \blacksquare \quad \forall_{u, v \in V} \left((L(u) = L(v)) \implies (u = v) \right) \quad \blacksquare \quad \text{Injective}[L, V, W]$
-
- (4) $(\text{ker}_L = \{O_v\}) \implies (\text{Injective}[L, V, W])$
-
- (5) $(\text{Injective}[L, V, W]) \iff (\text{ker}_L = \{O_v\})$
-

$$\text{RngSurjective} := \forall_{L, V, W} \left((\text{Ran}[\text{rng}_L, L, V, +_v, *_v, W, +_w, *_w]) \implies ((\text{Surjective}[L, V, W]) \iff (\text{rng}_L = W)) \right)$$

-
- (1) $(\text{SurjEquiv}) \wedge (\text{rng}(L) = \text{rng}_L) \quad \blacksquare \quad (\text{Surjective}[L, V, W]) \iff (\text{rng}_L = W)$
-

$$\text{RankNullityLT} := \forall_{L, V, W} \left((\text{LinTrans}[L, V, +_v, *_v, W, +_w, *_w]) \implies (\text{Dim}[V] = \text{Dim}[\text{ker}_L] + \text{Dim}[\text{rng}_L]) \right)$$

-
- (1) $\text{KerSubspace} \quad \blacksquare \quad (\exists_U (\text{Basis}[U, \text{ker}_L, +_v, *_v])) \wedge (\text{Dim}[\text{ker}_L] = |U|)$
-
- (2) $(\text{LinIndSuperspace}) \wedge (\text{LinInd}[U, \text{ker}_L, +_v, *_v]) \quad \blacksquare \quad \text{LinInd}[U, V, +_v, *_v]$
-
- (3) $(\text{LinIndExpandBasis}) \wedge (\text{LinInd}[U, V, +_v, *_v]) \quad \blacksquare \quad \left(\exists_B ((U \subseteq B) \wedge (\text{Basis}[B, V, +_v, *_v])) \right) \wedge (\text{Dim}[V] = |B|)$
-
- (4) $U \subseteq B \quad \blacksquare \quad \exists_T (B = U \cup T)$
-
- (5) $(w \in \text{rng}_L) \implies \dots$
-
- (5.1) $\exists_{v \in V} (w = L(v))$
-
- (5.2) $(\text{Basis}[B, V, +_v, *_v]) \wedge (B = U \cup T) \quad \blacksquare \quad \exists_{K \in \mathbb{R}^{|B|}} \left(v = \sum_{i=1}^{|B|} (k_i * b_i) = \sum_{i=1}^{|U|} (k_i * u_i) + \sum_{i=1}^{|T|} (k_{|U|+i} * t_i) \right)$
-
- (5.3) $w = L(v) = L \left(\sum_{i=1}^{|U|} (k_i * u_i) + \sum_{i=1}^{|T|} (k_{|U|+i} * t_i) \right) = L \left(\sum_{i=1}^{|U|} (k_i * u_i) \right) + L \left(\sum_{i=1}^{|T|} (k_{|U|+i} * t_i) \right) = \dots$
-
- (5.4) $O + L \left(\sum_{i=1}^{|T|} (k_{|U|+i} * t_i) \right) = \sum_{i=1}^{|T|} (L(k_{|U|+i} * t_i)) = \sum_{i=1}^{|T|} (k_{|U|+i} * L(t_i)) \quad \blacksquare \quad \exists_K (\text{LinComb}[w, L(T), K, W, +, *])$
-
- (6) $(w \in \text{rng}_L) \implies \left(\exists_L (\text{LinComb}[w, L(T), L, W, +, *]) \right) \quad \blacksquare \quad \text{Spans}[L(T), \text{rng}_L, W, +, *]$
-
- (7) $\left((K \in \mathbb{R}^n) \wedge (\text{LinComb}[O_w, L(T), K, W, +_w, *_w]) \right) \implies \dots$
-
- (7.1) $O_w = \sum_{i=1}^n (k_i * L(t_i)) = L \left(\sum_{i=1}^n (k_i * t_i) \right) \quad \blacksquare \quad \sum_{i=1}^n (k_i * t_i) \in \text{ker}_L$
-

$$(7.2) \quad (Basis[U, ker_L, +_v, *_v]) \wedge (\sum_{i=1}^n (k_i * t_i) \in ker_L) \quad \blacksquare \quad \exists_{D \in \mathbb{R}^m} (\sum_{i=1}^n (k_i * t_i) = \sum_{i=1}^m (d_i * u_i))$$

$$(7.3) \quad Basis[B] \quad \blacksquare \quad LinInd[B] \quad \blacksquare \quad LinInd[U \cup T] \quad \blacksquare \quad \forall_{s_j \in U \cup T} \forall_{K \in \mathbb{R}^{n-1}} (\neg LinComb[s_j, U \cup T \setminus \{s_j\}, K, V, +, *])$$

$$(7.4) \quad (\sum_{i=1}^n (k_i * t_i) = \sum_{i=1}^m (d_i * u_i)) \wedge (\forall_{s_j \in U \cup T} \forall_{K \in \mathbb{R}^{n-1}} (\neg LinComb[s_j, U \cup T \setminus \{s_j\}, K, V, +, *])) \quad \blacksquare \quad (D = \{O\}) \wedge (K = \{O\})$$

$$(8) \quad ((K \in \mathbb{R}^n) \wedge (LinComb[O_w, L(T), K, W, +_w, *_w])) \implies (K = \{O\}) \quad \blacksquare \quad LinInd[L(T), W, +_w, *_w]$$

$$(9) \quad (SubIndependent) \wedge (LinInd[L(T), W, +_w, *_w]) \quad \blacksquare \quad LinInd[L(T), rng_L, +_w, *_w]$$

$$(10) \quad (Spans[L(T), rng_L, W, +, *]) \wedge (LinInd[L(T), rng_L, +_w, *_w]) \quad \blacksquare \quad Basis[L(T), rng_L, +_w, *_w] \quad \blacksquare \quad Dim[rng_L] = |L(T)| = |T|$$

$$(11) \quad B = U \cup T \quad \blacksquare \quad |B| = |U| + |T| \quad \blacksquare \quad Dim[V] = Dim[ker_L] + Dim[rng_L]$$

$$InjectiveSurjectiveEqualDim := \forall_{T,V,W} \left(\begin{array}{l} ((LinTrans[T, V, +_v, *_v, W, +_w, *_w]) \wedge (Dim[V] = Dim[W]) \wedge (Injective[T, V, W])) \implies \\ (Surjective[T, V, W]) \end{array} \right)$$

$$(1) \quad (KerInjective) \wedge (Injective[T, V, W]) \quad \blacksquare \quad ker_T = \{O\} \quad \blacksquare \quad Dim[ker_T] = 0$$

$$(2) \quad (RankNullityLT) \wedge (Dim[ker_T] = 0) \quad \blacksquare \quad Dim[V] = Dim[ker_T] + Dim[rng_T] = Dim[rng_T] \quad \blacksquare \quad Dim[V] = Dim[rng_T]$$

$$(3) \quad (Dim[V] = Dim[W]) \wedge (Dim[V] = Dim[rng_T]) \quad \blacksquare \quad Dim[W] = Dim[rng_T]$$

$$(4) \quad RangeSubspace \quad \blacksquare \quad Subspace[rng_T, W, +_w, *_w]$$

$$(5) \quad (Subspace[rng_T, W, +_w, *_w]) \wedge (Dim[W] = Dim[rng_T]) \quad \blacksquare \quad \exists_B ((Basis[B, W, +_w, *_w]) \wedge (Basis[B, rng_T, +_w, *_w]))$$

$$(6) \quad (Spans[W] = Spans[rng_T]) \quad \blacksquare \quad W = rng_T \quad \blacksquare \quad Surjective[T, V, W]$$

$$SurjectiveInjectiveEqualDim := \forall_{T,V,W} \left(\begin{array}{l} ((LinTrans[T, V, +_v, *_v, W, +_w, *_w]) \wedge (Dim[V] = Dim[W]) \wedge (Surjective[T, V, W])) \implies \\ (Injective[T, V, W]) \end{array} \right)$$

$$(1) \quad RankNullityLT \quad \blacksquare \quad Dim[V] = Dim[ker_T] + Dim[rng_T]$$

$$(2) \quad Surjective[T, V, W] \quad \blacksquare \quad rng_T = W \quad \blacksquare \quad Dim[rng_T] = Dim[W]$$

$$(3) \quad (Dim[V] = Dim[W]) \wedge (Dim[V] = Dim[ker_T] + Dim[rng_T]) \wedge (Dim[rng_T] = Dim[W]) \quad \blacksquare \quad Dim[ker_T] + Dim[rng_T] = Dim[rng_T] \\ \blacksquare \quad Dim[ker_T] = 0 \quad \blacksquare \quad ker_T = \{O\}$$

$$(4) \quad (KerInjective) \wedge (ker_T = \{O\}) \quad \blacksquare \quad Injective[T, V, W]$$

$$SmallerMapNotInjective := \forall_{T,V,W} \left(((LinTrans[T, V, +_v, *_v, W, +_w, *_w]) \wedge (Dim[V] > Dim[W])) \implies (\neg Injective[T, V, W]) \right)$$

$$(1) \quad (RankNullityLT) \wedge (Dim[W] \geq Dim[rng_T]) \quad \blacksquare \quad Dim[ker_T] = Dim[V] - Dim[rng_T] \geq Dim[V] - Dim[W] > 0 \quad \blacksquare \quad Dim[ker_T] \neq 0$$

$$(2) \quad (KerInjective) \wedge (Dim[ker_T] \neq 0) \quad \blacksquare \quad \neg Injective[T, V, W]$$

$$LargerMapNotSurjective := \forall_{T,V,W} \left(((LinTrans[T, V, +_v, *_v, W, +_w, *_w]) \wedge (Dim[V] < Dim[W])) \implies (\neg Surjective[T, V, W]) \right)$$

$$(1) \quad RankNullityLT \quad \blacksquare \quad Dim[rng_T] = Dim[V] - Dim[ker_T] \leq Dim[V] < Dim[W]$$

$$(2) \quad Dim[rng_T] < Dim[W] \quad \blacksquare \quad Dim[rng_T] \neq Dim[W] \quad \blacksquare \quad \neg Surjective[T, V, W]$$

A linear transformation $L : V \rightarrow W$ is one-to-one if and only if the image of every linearly independent set of vectors in V is linearly independent set of vectors in W .

(1) TODO

A homogeneous system of linear equations with more variables than equations has nonzero solutions.

(1) TODO

An inhomogeneous system of linear equations with more equations than variables has no solution for some choice of the constant terms.

(1) TODO

$$LTInv[L^{-1}, L, V, +_v, *_v, W, +_w, *_w] := \left(\begin{array}{l} (LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge (LinTrans[L^{-1}, W, +_w, *_w, V, +_v, *_v]) \wedge \\ (L^{-1} \circ L = 1_v) \end{array} \right) \quad \wedge \quad (L \circ L^{-1} = 1_w)$$

$$LTInvUniq := \forall_{L_1^{-1}, L_2^{-1}} \left(((LTInv[L_1^{-1}, L, V, +_v, *_v, W, +_w, *_w]) \wedge (LTInv[L_2^{-1}, L, V, +_v, *_v, W, +_w, *_w])) \implies (L_1^{-1} = L_2^{-1}) \right)$$

$$(1) \quad L_1^{-1} = L_1^{-1} \circ 1_w = L_1^{-1} \circ (L \circ L_2^{-1}) = (L_1^{-1} \circ L) \circ L_2^{-1} = 1_v \circ L_2^{-1} = L_2^{-1} \quad \blacksquare \quad L_1^{-1} = L_2^{-1}$$

$$LTInvertible[L, V, +_v, *_v, W, +_w, *_w] := \exists_{L^{-1}}(LTInv[L^{-1}, L, V, +_v, *_v, W, +_w, *_w])$$

$$InvertibleBijectiveEquiv := \forall_L \left((LTInvertible[L, V, +_v, *_v, W, +_w, *_w]) \iff ((Injective[L, V, W]) \wedge (Surjective[L, V, W])) \right)$$

$$(1) \quad (LTInvertible[L, V, +_v, *_v, W, +_w, *_w]) \implies \dots$$

$$(1.1) \quad \exists_{L^{-1}}(LTInv[L^{-1}, L, V, +_v, *_v, W, +_w, *_w])$$

$$(1.2) \quad (L(u) = L(w)) \implies \dots$$

$$(1.2.1) \quad u = L^{-1}(L(u)) = L^{-1}(L(w)) = w \quad \blacksquare \quad u = w$$

$$(1.3) \quad (L(u) = L(w)) \implies (u = w) \quad \blacksquare \quad \forall_{u,w} \left((L(u) = L(w)) \implies (u = w) \right) \quad \blacksquare \quad Injective[L, V, W]$$

$$(1.4) \quad (w \in W) \implies \dots$$

$$(1.4.1) \quad L^{-1}(w) \in V$$

$$(1.4.2) \quad L \circ L^{-1} = 1_w \quad \blacksquare \quad L(L^{-1}(w) = w)$$

$$(1.4.3) \quad (L^{-1}(w) \in V) \wedge (L(L^{-1}(w)) = w) \quad \blacksquare \quad \exists_{v \in V} (w = L(v))$$

$$(1.5) \quad (w \in W) \implies \left(\exists_{v \in V} (w = L(v)) \right) \quad \blacksquare \quad \forall_{w \in W} \exists_{v \in V} (w = L(v)) \quad \blacksquare \quad Surjective[L, V, W]$$

$$(1.6) \quad (Injective[L, V, W]) \wedge (Surjective[L, V, W])$$

$$(2) \quad (LTInvertible[L, V, +_v, *_v, W, +_w, *_w]) \implies ((Injective[L, V, W]) \wedge (Surjective[L, V, W]))$$

$$(3) \quad ((Injective[L, V, W]) \wedge (Surjective[L, V, W])) \implies \dots$$

$$(3.1) \quad (Injective[L, V, W]) \wedge (Surjective[L, V, W]) \quad \blacksquare \quad \forall_{w \in W} \exists!_{v \in V} (w = L(v))$$

$$(3.2) \quad S := \{(w, v) \in W \times V \mid w = L(v)\}$$

$$(3.3) \quad \left(\forall_{w \in W} \exists!_{v \in V} (w = L(v)) \right) \wedge (S = \{(w, v) \in W \times V \mid w = L(v)\}) \quad \blacksquare \quad Function[S, W, V]$$

$$(3.4) \quad \left(\forall_{v \in V} (S(L(v)) = v) \right) \wedge \left(\forall_{w \in W} (L(S(w)) = w) \right)$$

$$(3.5) \quad (w_1, w_2 \implies W) \implies \dots$$

$$(3.5.1) \quad (LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge \left(\forall_{w \in W} (L(S(w)) = w) \right) \quad \blacksquare \quad L(S(w_1) + S(w_2)) = L(S(w_1)) + L(S(w_2)) = w_1 + w_2$$

$$(3.5.2) \quad \left(\forall_{w \in W} (L(S(w)) = w) \right) \wedge (w_1 + w_2 \in W) \quad \blacksquare \quad L(S(w_1 + w_2)) = w_1 + w_2$$

$$(3.5.3) \quad L(S(w_1) + S(w_2)) = w_1 + w_2 = L(S(w_1 + w_2)) \quad \blacksquare \quad L(S(w_1) + S(w_2)) = L(S(w_1 + w_2))$$

$$(3.5.4) \quad (Injective[L, V, W]) \wedge (L(S(w_1) + S(w_2)) = L(S(w_1 + w_2))) \quad \blacksquare \quad S(w_1) + S(w_2) = S(w_1 + w_2)$$

$$(3.6) \quad (w_1, w_2 \implies W) \implies (S(w_1 + w_2) = S(w_1) + S(w_2)) \quad \blacksquare \quad \forall_{w_1, w_2 \in W} (S(w_1 + w_2) = S(w_1) + S(w_2))$$

$$(3.7) \quad (r \in \mathbb{R}) \wedge (w \in W) \implies \dots$$

$$(3.7.1) \quad (LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge \left(\forall_{w \in W} (L(S(w)) = w) \right) \quad \blacksquare \quad L(r * S(w)) = r * L(S(w)) = r * w$$

$$(3.7.2) \quad \left(\forall_{w \in W} (L(S(w)) = w) \right) \wedge (r * w \in W) \quad \blacksquare \quad L(S(r * w)) = r * w$$

$$(3.7.3) \quad L(r * S(w)) = r * w = L(S(r * w)) \quad \blacksquare \quad L(r * S(w)) = L(S(r * w))$$

$$(3.7.4) \quad (Injective[L, V, W]) \wedge (L(r * S(w)) = L(S(r * w))) \quad \blacksquare \quad r * S(w) = S(r * w)$$

$$(3.8) \quad (r \in \mathbb{R}) \wedge (w \in W) \implies (r * S(w) = S(r * w)) \quad \blacksquare \quad \forall_{r \in \mathbb{R}} \forall_{w \in W} (S(r * w) = r * S(w))$$

$$(3.9) \quad (Function[S, W, V]) \wedge \left(\forall_{w_1, w_2 \in W} (S(w_1 + w_2) = S(w_1) + S(w_2)) \right) \wedge \left(\forall_{r \in \mathbb{R}} \forall_{w \in W} (S(r * w) = r * S(w)) \right)$$

$$(3.10) \quad LinTrans[S, W, +_w, *_w, V, +_v, *_v]$$

$$(3.11) \quad \forall_{v \in V} \left((S(L(v)) = v) \right) \quad \blacksquare \quad S \circ L = 1_v$$

$$(3.12) \quad \forall_{w \in W} (L(S(w)) = w) \quad \blacksquare \quad L \circ S = 1_w$$

$$(3.13) \quad (LinTrans[S, W, +_w, *_w, V, +_v, *_v]) \wedge (S \circ L = 1_v) \wedge (L \circ S = 1_w) \quad \blacksquare \quad L T Inv[S, L, V, +_v, *_v, W, +_w, *_w]$$

$$(3.14) \quad \exists_{L^{-1}} (L T Inv[L^{-1}, L, V, +_v, *_v, W, +_w, *_w]) \quad \blacksquare \quad L T Invertible[L, V, +_v, *_v, W, +_w, *_w]$$

$$(4) \quad ((Injective[L, V, W]) \wedge (Surjective[L, V, W])) \implies (L T Invertible[L, V, +_v, *_v, W, +_w, *_w])$$

$$(5) \quad (L T Invertible[L, V, +_v, *_v, W, +_w, *_w]) \iff ((Injective[L, V, W]) \wedge (Surjective[L, V, W]))$$

TODO: some corollary of InjectiveSurjectiveEqualDim + SurjectiveInjectiveEqualDim + InvertibleBijjectiveEquiv

$$Isomorphism[L, V, +_v, *_v, W, +_w, *_w] := L T Invertible[L, V, +_v, *_v, W, +_w, *_w]$$

$$Isomorphic[V, +_v, *_v, W, +_w, *_w] := \exists_L (Isomorphism[L, V, +_v, *_v, W, +_w, *_w])$$

3.9 Matrix of a Linear Transform

$$CoordVec[[\alpha]_S, \alpha, S, V, +, *] := (Basis[S, V, +, *]) \wedge (S * [\alpha]_S = \alpha \in V)$$

$$L T Matrix := \forall_{L, V, W} \left(\begin{array}{l} ((LinTrans[L, V, +_v, *_v, W, +_w, *_w]) \wedge (Basis[A, V, +_v, *_v]) \wedge (Basis[B, W, +_w, *_w])) \implies \\ (\forall_{v \in V} (CoordVec[[L(v)]_B, L(v), B, W, +_w, *_w] = \langle [L(a_i)]_B \mid a_i \in A \rangle * CoordVec[[v]_A, v, A, V, +_v, *_v])) \end{array} \right)$$

$$(1) \quad Basis[A, V, +_v, *_v] \quad \blacksquare \quad \exists_{K \in \mathbb{R}^n} (v = \sum_{i=1}^n (k_i * a_i)) \quad \blacksquare \quad K^T = CoordVec[[v]_A, v, A, V, +, *]$$

$$(2) \quad [L(v)]_B = [L(\sum_{i=1}^n (k_i * a_i))]_B = [\sum_{i=1}^n (L(k_i * a_i))]_B = \sum_{i=1}^n ([L(k_i * a_i)]_B) = \sum_{i=1}^n ([k_i * L(a_i)]_B) = \sum_{i=1}^n (k_i * [L(a_i)]_B) = \dots$$

$$(3) \quad \dots \langle [L(a)]_B \mid a \in A \rangle * K^T = \langle [L(a)]_B \mid a \in A \rangle * [v]_A \quad \blacksquare \quad [L(v)]_B = \langle [L(a)]_B \mid a \in A \rangle * [v]_A$$

Note: Shorthand is to RREF the augmented matrix [Columns of B | Columns of A] into [I | M], thus M is the transition matrix

$$TransitionMatrix := \forall_{L, V} \left(\begin{array}{l} ((Basis[A, V, +, *]) \wedge (Basis[B, V, +, *])) \implies \\ (\forall_{v \in V} (CoordVec[[v]_B, v, B, W, +_w, *_w] = \langle [a]_B \mid a \in A \rangle * CoordVec[[v]_A, v, A, V, +_v, *_v])) \end{array} \right)$$

$$(1) \quad (L T Matrix) \wedge (LinTrans[I, V, +, *, V, +, *]) \quad \blacksquare \quad [I(v)]_B = \langle [I(a)]_B \mid a \in A \rangle * [v]_A \quad \blacksquare \quad [v]_B = \langle [a]_B \mid a \in A \rangle * [v]_A$$

$$L T OverTransition := \left(([L(a)]_T = A * [a]_S) \wedge (P * [a]_{S'} = [a]_S) \wedge (Q * [L(a)]_{T'} = [L(a)]_T) \right) \implies ([L(a)]_{T'} = (Q^{-1} * A * P) * [a]_{S'})$$

$$(1) \quad [L(a)]_{T'} = Q^{-1} * [L(a)]_T = Q^{-1} * A * [a]_S = Q^{-1} * A * P * [a]_{S'} \quad \blacksquare \quad [L(a)]_{T'} = (Q^{-1} * A * P) * [a]_{S'}$$

$$L O OverTransition := \left(([L(a)]_S = A * [a]_S) \wedge (P * [a]_{S'} = [a]_S) \right) \implies ([L(a)]_{S'} = (P^{-1} * A * P) * [a]_{S'})$$

$$(1) \quad P * [a]_{S'} = [a]_S \quad \blacksquare \quad P * [L(a)]_{S'} = [L(a)]_S$$

$$(2) \quad L T OverTransition \quad \blacksquare \quad [L(a)]_{S'} = P^{-1} * [L(a)]_S = P^{-1} * A * [a]_S = P^{-1} * A * P * [a]_{S'} \quad \blacksquare \quad [L(a)]_{S'} = (P^{-1} * A * P) * [a]_{S'}$$

$$RankNullityRelation := (Rank[A] \equiv Dim[rng_L]) \wedge (Nullity[A] \equiv Dim[ker_L]) \wedge (RankNullity \equiv RankNullityLT)$$

$$(1) \quad \text{TODO}$$

$$SimMatrix[A, B] := \exists_P (B = P * A * P^{-1})$$

$$SimMatrixEquiv := (SimMatrix[A, B]) \iff \left(\exists_{S, T, S', T'} ([L(a)]_T = A * [a]_S) \wedge ([L(a)]_{T'} = B * [a]_{S'}) \right)$$

$$(1) \quad \text{TODO}$$

$$SimRank := (SimMatrix[A, B]) \implies (Rank[A] = Rank[B])$$

$$(1) \quad \text{TODO}$$

3.10 Determinants

$$\text{Perm}[\sigma, S] := \text{Bij}[\sigma, S, S]$$

$$\text{IntPermSet}[S_n, n] := S_n = \{\sigma \mid \text{Perm}[\sigma, \mathbb{N}_{1,n}]\}$$

$$\text{IntPermSetCard} := (\text{IntPermSet}[S_n, n]) \implies (|S_n| = n!)$$

(1) TODO: Combinatorics / induction on N

$$\text{IntPermGroup} := \text{Group}[S_n, \circ]$$

$$(1) \text{ Perm}[I_n, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}] \blacksquare I_n \in S_n$$

$$(2) (\sigma, \tau, v \in S_n) \implies \dots$$

$$(2.1) (\text{Bij}[\sigma, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}]) \wedge (\text{Bij}[\tau, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}]) \blacksquare \text{Bij}[\sigma \circ \tau, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}] \blacksquare \sigma \circ \tau \in S_n$$

$$(2.2) (\text{Bij}[\sigma, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}]) \wedge (\text{Bij}[\tau, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}]) \wedge (\text{Bij}[v, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}]) \blacksquare (\sigma \circ \tau) \circ v = \sigma \circ (\tau \circ v)$$

$$(2.3) \sigma \circ I_n = \sigma = I_n \circ \sigma$$

$$(2.4) \text{Bij}[\sigma, \mathbb{N}_{1,n}, \mathbb{N}_{1,n}] \blacksquare \sigma \circ \sigma^{-1} = I_n = \sigma^{-1} \circ \sigma$$

$$(3) \text{Group}[S_n, \circ]$$

$$\text{IntPermSetDecomp} := (\text{IntPermSet}[S_n, n]) \wedge (\text{Perm}[\tau, \mathbb{N}_{1,n}]) \implies (S_n = \{\tau \circ \sigma \mid \sigma \in S_n\} = \{\sigma \circ \tau \mid \sigma \in S_n\})$$

$$(1) (\sigma \in S_n) \iff ()$$