

# Contents



# Chapter 1

## Real Analysis

(1.5)

*OrderTrichotomy* $[\prec, S] := \forall_{x,y \in S} (x \prec y \vee x = y \vee y \prec x)$

*OrderTransitivity* $[\prec, S] := \forall_{x,y,z \in S} ((x \prec y \wedge y \prec z) \implies x \prec z)$

*Order* $[\prec, S] := (\textcolor{teal}{OrderTrichotomy}[\prec, S]) \wedge (\textcolor{teal}{OrderTransitivity}[\prec, S])$

(1.7)

*Bounded Above* $[E, S, \prec] := (\textcolor{teal}{Order}[\prec, S]) \wedge (E \subset S) \wedge (\exists_{\beta \in S} \forall_{x \in E} (x \leq \beta))$

*Bounded Below* $[E, S, \prec] := (\textcolor{teal}{Order}[\prec, S]) \wedge (E \subset S) \wedge (\exists_{\beta \in S} \forall_{x \in E} (\beta \leq x))$

*Upper Bound* $[\beta, E, S, \prec] := (\textcolor{teal}{Order}[\prec, S]) \wedge (E \subset S) \wedge (\beta \in S \wedge \forall_{x \in E} (x \leq \beta))$

*Lower Bound* $[\beta, E, S, \prec] := (\textcolor{teal}{Order}[\prec, S]) \wedge (E \subset S) \wedge (\beta \in S \wedge \forall_{x \in E} (\beta \leq x))$

(1.8)

*LU B* $[\alpha, E, S, \prec] := (\textcolor{teal}{Upper Bound}[\alpha, E, S, \prec]) \wedge (\forall_{\gamma} (\gamma < \alpha \implies \neg \textcolor{teal}{Upper Bound}[\gamma, E, S, \prec]))$

*GLB* $[\alpha, E, S, \prec] := (\textcolor{teal}{Lower Bound}[\alpha, E, S, \prec]) \wedge (\forall_{\beta} (\alpha < \beta \implies \neg \textcolor{teal}{Lower Bound}[\beta, E, S, \prec]))$

(1.10)

*LU BProperty* $[S, \prec] := \forall_E (((\emptyset \neq E \subset S) \wedge (\textcolor{teal}{Bounded Above}[E, S, \prec]) \implies \exists_{\alpha \in S} (\textcolor{teal}{LU B}[\alpha, E, S, \prec])))$

*GLBProperty* $[S, \prec] := \forall_E (((\emptyset \neq E \subset S) \wedge (\textcolor{teal}{Bounded Below}[E, S, \prec]) \implies \exists_{\alpha \in S} (\textcolor{teal}{GLB}[\alpha, E, S, \prec])))$

(1.11)

*LU BPropertyImpliesGLBProperty*  $:= \textcolor{teal}{LU BProperty}[S, \prec] \implies \textcolor{teal}{GLBProperty}[S, \prec]$

(1)  $\textcolor{teal}{LU BProperty}[S, \prec] \implies \dots$

wts: 2

(1.1)  $(\emptyset \neq B \subset S \wedge \textcolor{teal}{Bounded Below}[B, S, \prec]) \implies \dots$

wts: 1.2

from: *Bounded Below*, 1.1

(1.1.1)  $\textcolor{teal}{Order}[\prec, S] \wedge \exists_{\delta' \in S} (\textcolor{teal}{Lower Bound}[\delta', B, S, \prec])$

(1.1.2)  $|B| = 1 \implies \dots$

wts: 1.1.3

from: 1.1.2

(1.1.2.1)  $\exists_{u'} (u' \in B) \blacksquare u := \textit{choice}(\{u' : u' \in B\}) \blacksquare B = \{u\}$

(1.1.2.2)  $\textcolor{teal}{GLB}[u, B, S, \prec] \blacksquare \exists_{\epsilon_0 \in S} (\textcolor{teal}{GLB}[\epsilon_0, B, S, \prec])$

(1.1.3)  $|B| = 1 \implies \exists_{\epsilon_0 \in S} (\textcolor{teal}{GLB}[\epsilon_0, B, S, \prec])$

(1.1.4)  $|B| \neq 1 \implies \dots$

wts: 1.1.5

from: *LU BProperty*, 1

(1.1.4.1)  $\forall_E (((\emptyset \neq E \subset S \wedge \textcolor{teal}{Bounded Above}[E, S, \prec]) \implies \exists_{\alpha \in S} (\textcolor{teal}{LU B}[\alpha, E, S, \prec])))$

(1.1.4.2)  $L := \{s \in S : \textcolor{teal}{Lower Bound}[s, B, S, \prec]\}$

(1.1.4.3)  $|B| > 1 \wedge \textcolor{teal}{OrderTrichotomy}[\prec, S] \blacksquare \exists_{b_1' \in B} \exists_{b_0' \in B} (b_0' < b_1')$

from: *Order*, 1.1.1  
wts: 1.1.4.7

(1.1.4.4)  $b_1 := \textit{choice}(\{b_1' \in B : \exists_{b_0' \in B} (b_0' < b_1')\}) \blacksquare \neg \textcolor{teal}{Lower Bound}[b_1, B, S, \prec]$

from: 1.1.4.2

(1.1.4.5)  $b_1 \notin L \blacksquare L \subset S$

(1.1.4.6)  $\delta := \textit{choice}(\{\delta' \in S : \textcolor{teal}{Lower Bound}[\delta', B, S, \prec]\}) \blacksquare \delta \in L \blacksquare \emptyset \neq L$

from: 1.1.1

(1.1.4.7)  $\emptyset \neq L \subset S$

from: 1.1.4.5, 1.1.4.6

(1.1.4.8)  $\forall_{y \in L} (\textcolor{teal}{Lower Bound}[y_0, B, S, \prec]) \blacksquare \forall_{y \in L} \forall_{x \in B} (y_0 \leq x)$

from: *Lower Bound*, 1.1.4.2  
wts: 1.1.4.10

(1.1.4.9)  $\forall_{x \in B} (x \in S \wedge \forall_{y \in L} (y_0 \leq x)) \blacksquare \forall_{x \in B} (\textcolor{teal}{Upper Bound}[x, L, S, \prec])$

from: *Upper Bound*

(1.1.4.10)  $\exists_{x \in S} (\textcolor{teal}{Upper Bound}[x, L, S, \prec]) \blacksquare \textcolor{teal}{Bounded Above}[L, S, \prec]$

(1.1.4.11)	$\emptyset \neq L \subset S \wedge \text{BoundedAbove}[L, S, <]$	from: 1.1.4.7, 1.1.4.10
(1.1.4.12)	$\exists_{\alpha' \in S}(\text{LUB}[\alpha', L, S, <]) \blacksquare \alpha := \text{choice}(\{\alpha' \in S : (\text{LUB}[\alpha', L, S, <])\})$	from: 1.1.4.1 wts: 1.1.4.21
(1.1.4.13)	$\forall_x(x \in B \implies \text{UpperBound}[x, L, S, <])$	from: 1.1.4.9 wts: 1.1.4.17
(1.1.4.14)	$\forall_x(\neg \text{UpperBound}[x, L, S, <] \implies x \notin B)$	
(1.1.4.15)	$\gamma < \alpha \implies \dots$	wts: 1.1.4.16
(1.1.4.15.1)	$\neg \text{UpperBound}[\gamma, L, S, <] \blacksquare \gamma \notin B$	from: LUB, 1.1.4.12, 1.1.4.14
(1.1.4.16)	$\gamma < \alpha \implies \gamma \notin B \blacksquare \gamma \in B \implies \gamma \geq \alpha$	
(1.1.4.17)	$\forall_{\gamma \in B}(\alpha \leq \gamma) \blacksquare \text{LowerBound}[\alpha, B, S, <]$	from: LowerBound
(1.1.4.18)	$\alpha < \beta \implies \dots$	wts: 1.1.4.19
(1.1.4.18.1)	$\forall_{y \in L}(y_0 \leq \alpha < \beta) \blacksquare \forall_{y \in L}(y_0 \neq \beta)$	from: LUB, 1.1.4.12, 1.1.4.18
(1.1.4.18.2)	$\beta \notin L \blacksquare \neg \text{LowerBound}[\beta, B, S, <]$	from: 1.1.4.2
(1.1.4.19)	$\alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <] \blacksquare \forall_{\beta \in S}(\alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <])$	
(1.1.4.20)	$\text{LowerBound}[\alpha, B, S, <] \wedge \forall_{\beta \in S}(\alpha < \beta \implies \neg \text{LowerBound}[\beta, B, S, <])$	from: 1.1.4.17, 1.1.4.19
(1.1.4.21)	$\text{GLB}[\alpha, B, S, <] \blacksquare \exists_{\epsilon_1 \in S}(\text{GLB}[\epsilon_1, B, S, <])$	
(1.1.5)	$ B  \neq 1 \implies \exists_{\epsilon_1 \in S}(\text{GLB}[\epsilon_1, B, S, <])$	
(1.1.6)	$( B  = 1 \implies \exists_{\epsilon_0 \in S}(\text{GLB}[\epsilon_0, B, S, <])) \wedge ( B  \neq 1 \implies \exists_{\epsilon_1 \in S}(\text{GLB}[\epsilon_1, B, S, <]))$	from: 1.1.3, 1.1.5
(1.1.7)	$( B  = 1 \vee  B  \neq 1) \implies \exists_{\epsilon \in S}(\text{GLB}[\epsilon, B, S, <]) \blacksquare \exists_{\epsilon \in S}(\text{GLB}[\epsilon, B, S, <])$	
(1.2)	$(\emptyset \neq B \subset S \wedge \text{BoundedBelow}[B, S, <]) \implies \exists_{\epsilon \in S}(\text{GLB}[\epsilon, B, S, <])$	
(1.3)	$\forall_B((\emptyset \neq B \subset S \wedge \text{BoundedBelow}[B, S, <]) \implies \exists_{\epsilon \in S}(\text{GLB}[\epsilon, B, S, <]))$	
(1.4)	$\text{GLBProperty}[S, <]$	
(2)	$\text{LUBProperty}[S, <] \implies \text{GLBProperty}[S, <]$	

(1.12)

$$\text{Field}[F, +, *] := \exists_{0, 1 \in F} \forall_{x, y, z \in F} \left( \begin{array}{llll} x + y \in F & \wedge & x * y \in F & \wedge \\ x + y = y + x & \wedge & x * y = y * x & \wedge \\ (x + y) + z = x + (y + z) & \wedge & (x * y) * z = x * (y * z) & \wedge \\ 1 \neq 0 & \wedge & x * (y + z) = (x * y) + (x * z) & \wedge \\ 0 + x = x & \wedge & 1 * x = x & \wedge \\ \exists_{-x \in F}(x + (-x) = 0) \wedge (x \neq 0 \implies \exists_{1/x \in F}(x * (1/x) = 1)) \end{array} \right)$$

\*\*\*\*\* (Field[F, +, \*]  $\wedge$   $x, y, z \in F$ )  $\implies$  ... \*\*\*\*\*

(1.14)

$$\text{AdditiveCancellation} := (x + y = x + z) \implies y = z$$

(1)	$y = 0 + y = (x + (-x)) + y = ((-x) + x) + y = (-x) + (x + y) = \dots$	from: Field
-----	--	-------------

(2)	$(-x) + (x + z) = ((-x) + x) + z = (x + (-x)) + z = 0 + z = z$	from: Field
-----	--	-------------

$$\text{AdditiveIdentityUniqueness} := (x + y = x) \implies y = 0$$

(1)	$x + y = x = 0 + x = x + 0$	from: Field
-----	-----------------------------	-------------

(2)	$y = 0$	from: AdditiveCancellation
-----	---------	----------------------------

$$\text{AdditiveInverseUniqueness} := (x + y = 0) \implies y = -x$$

(1)	$x + y = 0 = x + (-x)$	from: Field
-----	------------------------	-------------

(2)	$y = -x$	from: AdditiveCancellation
-----	----------	----------------------------

$$\text{DoubleNegative} := x = -(-x)$$

(1)	$0 = x + (-x) = (-x) + x \blacksquare 0 = (-x) + x$	from: Field
-----	---	-------------

(2)	$x = -(-x)$	from: AdditiveInverseUniqueness
-----	-------------	---------------------------------

(1.15)

$$\begin{aligned}
\textcolor{red}{\textit{MultiplicativeCancellation}} &:= (x \neq 0 \wedge x * y = x * z) \implies y = z & \text{---} \\
\textcolor{red}{\textit{MultiplicativeIdentityUniqueness}} &:= (x \neq 0 \wedge x * y = x) \implies y = 1 & \text{---} \\
\textcolor{red}{\textit{MultiplicativeInverseUniqueness}} &:= (x \neq 0 \wedge x * y = 1) \implies y = 1/x & \text{---} \\
\textcolor{red}{\textit{DoubleReciprocal}} &:= (x \neq 0) \implies x = 1/(1/x) & \text{---}
\end{aligned}$$

(1.16)

$$\textcolor{red}{\textit{Domination}} := 0 * x = 0$$

$$(1) \quad 0 * x = (0 + 0) * x = 0 * x + 0 * x \quad \blacksquare \quad 0 * x = 0 * x + 0 * x$$

from: [Field](#)

$$(2) \quad 0 * x = 0$$

from: [AdditiveIdentityUniqueness](#)

$$\textcolor{red}{\textit{NonDomination}} := (x \neq 0 \wedge y \neq 0) \implies x * y \neq 0$$

$$(1) \quad (x \neq 0 \wedge y \neq 0) \implies \dots$$

$$(1.1) \quad (x * y = 0) \implies \dots$$

$$(1.1.1) \quad 1 = 1 * 1 = (x * (1/x)) * (y * (1/y)) = (x * y) * ((1/x) * (1/y)) = 0 * ((1/x) * (1/y)) = 0$$

from: [Field](#), [Domination](#), 1, 1.1

$$(1.1.2) \quad 1 = 0 \wedge 1 \neq 0 \quad \blacksquare \quad \perp$$

from: [Field](#)

$$(1.2) \quad (x * y = 0) \implies \perp \quad \blacksquare \quad x * y \neq 0$$

$$(2) \quad (x \neq 0 \wedge y \neq 0) \implies x * y \neq 0$$

$$\textcolor{red}{\textit{NegationCommutativity}} := (-x) * y = -(x * y) = x * (-y)$$

$$(1) \quad x * y + (-x) * y = (x + -x) * y = 0 * y = 0 \quad \blacksquare \quad x * y + (-x) * y = 0$$

from: [Field](#), [Domination](#)  
wts: 2

$$(2) \quad (-x) * y = -(x * y)$$

from: [AdditiveInverseUniqueness](#)

$$(3) \quad x * y + x * (-y) = x * (y_0 + -y) = x * 0 = 0 \quad \blacksquare \quad x * y + x * (-y) = 0$$

from: [Field](#), [Domination](#)  
wts: 4

$$(4) \quad x * (-y) = -(x * y)$$

from: [AdditiveInverseUniqueness](#)

$$(5) \quad (-x) * y = -(x * y) = x * (-y)$$

from: 2, 4

$$\textcolor{red}{\textit{NegativeMultiplication}} := (-x) * (-y) = x * y$$

$$(1) \quad (-x) * (-y) = -(x * (-y)) = -(-(x * y)) = x * y$$

from: [NegationCommutativity](#), [DoubleNegative](#)

(1.17)

$$\textcolor{red}{\textit{OrderedField}}[F, +, *, <] := \left( \begin{array}{l} \textcolor{blue}{\textit{Field}}[F, +, *] \quad \wedge \quad \textcolor{blue}{\textit{Order}}[<, F] \quad \wedge \\ \forall_{x,y,z \in F} (y_0 < z \implies x + y < x + z) \quad \wedge \\ \forall_{x,y \in F} ((x > 0 \wedge y > 0) \implies x * y > 0) \end{array} \right)$$

$$***** \textcolor{blue}{(\textit{OrderedField}[F, +, *, <] \wedge x, y, z \in F)} \implies \dots *****$$

(1.18)

$$\textcolor{red}{\textit{NegationOnOrder}} := x > 0 \iff -x < 0$$

$$(1) \quad x > 0 \implies \dots$$

$$(1.1) \quad 0 = (-x) + x > (-x) + 0 = -x \quad \blacksquare \quad 0 > -x \quad \blacksquare \quad -x < 0$$

from: [OrderedField](#)

$$(2) \quad x > 0 \implies -x < 0$$

$$(3) \quad -x < 0 \implies \dots$$

$$(3.1) \quad 0 = x + (-x) < x + 0 = x \quad \blacksquare \quad 0 < x \quad \blacksquare \quad x > 0$$

from: [OrderedField](#)

$$(4) \quad -x < 0 \implies x > 0$$

$$(5) \quad x > 0 \implies -x < 0 \wedge -x < 0 \implies x > 0 \quad \blacksquare \quad x > 0 \iff -x < 0$$

from: 2, 4

$$\textcolor{red}{\textit{PositiveFactorPreservesOrder}} := (x > 0 \wedge y < z) \implies x * y < x * z$$

$$(1) \quad (x > 0 \wedge y < z) \implies \dots$$

$$(1.1) \quad (-y) + z > (-y) + y = 0 \quad \blacksquare \quad z + (-y) = 0$$

from: [OrderedField](#)

$$(1.2) \quad x * (z + (-y)) > 0 \quad \blacksquare \quad x * z + x * (-y) > 0$$

from: [OrderedField](#)

$$(1.3) \quad x * z = 0 + x * z = (x * y + -(x * y)) + x * z = (x * y + x * (-y)) + x * z = \dots$$

from: [Field](#), [NegationCommutativity](#)

$$(1.4) \quad x * y + (x * z + x * (-y)) > x * y + 0 = x * y$$

from: [Field](#), 1.2

$$(1.5) \quad x * z > x * y$$

from: 1.3, 1.4

$$(2) \quad (x > 0 \wedge y < z) \implies x * z > x * y$$

$$\textcolor{red}{NegativeFactorFlipsOrder} := (x < 0 \wedge y < z) \implies x * y > x * z$$

$$(1) \quad (x < 0 \wedge y < z) \implies \dots$$

$$(1.1) \quad -x > 0$$

from: [NegationOnOrder](#)

$$(1.2) \quad (-x) * y < (-x) * z \quad \blacksquare \quad 0 = x * y + (-x) * y < x * y + (-x) * z \quad \blacksquare \quad 0 < x * y + (-x) * z$$

from: [PositiveFactorPreservesOrder](#)

$$(1.3) \quad 0 < (-x) * (-y + z) \quad \blacksquare \quad 0 > x * (-y + z) \quad \blacksquare \quad 0 > -(x * y) + x * z$$

from: [NegationOnOrder](#)

$$(1.4) \quad x * y > x * z$$

$$(2) \quad (x < 0 \wedge y < z) \implies x * y > x * z$$

$$\textcolor{red}{SquareIsPositive} := (x \neq 0) \implies x * x > 0$$

$$(1) \quad (x > 0) \implies x * x > 0$$

from: [OrderedField](#)

$$(2) \quad (x < 0) \implies \dots$$

$$(2.1) \quad -x > 0 \quad \blacksquare \quad x * x = (-x) * (-x) > 0 \quad \blacksquare \quad x * x > 0$$

from: [NegationOnOrder](#), [OrderedField](#), [NegativeMultiplication](#)

$$(3) \quad (x < 0) \implies x * x > 0$$

$$(4) \quad x \neq 0 \implies (x > 0 \vee x < 0) \implies x * x > 0 \quad \blacksquare \quad x \neq 0 \implies x * x > 0$$

from: [OrderTrichotomy](#), 1, 3

$$\textcolor{red}{OneIsPositive} := 1 > 0$$

$$(1) \quad 1 \neq 0 \quad \blacksquare \quad 1 = 1 * 1 > 0$$

from: [Field](#), [SquareIsPositive](#)

$$\textcolor{red}{ReciprocationOnOrder} := (0 < x < y) \implies 0 < 1/y < 1/x$$

$$(1) \quad (0 < x < y) \implies \dots$$

$$(1.1) \quad x * (1/x) = 1 > 0 \quad \blacksquare \quad x * (1/x) > 0$$

from: [Field](#), [OneIsPositive](#)

$$(1.2) \quad 1/x < 0 \implies x * (1/x) < 0 \wedge x * (1/x) > 0 \implies \perp \quad \blacksquare \quad 1/x > 0$$

from: [NegativeFactorFlipsOrder](#), 1

$$(1.3) \quad y * (1/y) = 1 > 0 \quad \blacksquare \quad y * (1/y) > 0$$

from: [Field](#), [OneIsPositive](#)

$$(1.4) \quad 1/y < 0 \implies y * (1/y) < 0 \wedge y * (1/y) > 0 \implies \perp \quad \blacksquare \quad 1/y > 0$$

from: [NegativeFactorFlipsOrder](#), 1

$$(1.5) \quad (1/x) * (1/y) > 0$$

from: [OrderedField](#)

$$(1.6) \quad 0 < 1/y = ((1/x) * (1/y)) * x < ((1/x) * (1/y)) * y = 1/x$$

from: [OrderedField](#), 1, 1.4, 1.5

$$(1.19)$$

$$\textcolor{red}{OrderedFieldQ} := \textcolor{blue}{OrderedField}[\mathbb{Q}, +, *, <] \quad \text{---}$$

$$\textcolor{blue}{Subfield}[K, F, +, *] := \textcolor{blue}{Field}[F, +, *] \wedge K \subset F \wedge \textcolor{blue}{Field}[K, +, *]$$

$$\textcolor{blue}{OrderedSubfield}[K, F, +, *, <] := \textcolor{blue}{OrderedField}[F, +, *, <] \wedge K \subset F \wedge \textcolor{blue}{OrderedField}[K, +, *, <]$$

$$\textcolor{blue}{CutI}[\alpha] := \emptyset \neq \alpha \subset \mathbb{Q}$$

$$\textcolor{blue}{CutII}[\alpha] := \forall_{p \in \alpha} \forall_{q \in \mathbb{Q}} (q < p \implies q \in \alpha)$$

$$\textcolor{blue}{CutIII}[\alpha] := \forall_{p \in \alpha} \exists_{r \in \alpha} (p < r)$$

$$\mathbb{R} := \{\alpha \in \mathbb{Q} : \textcolor{blue}{CutI}[\alpha] \wedge \textcolor{blue}{CutII}[\alpha] \wedge \textcolor{blue}{CutIII}[\alpha]\}$$

$$\textcolor{red}{CutCorollaryI} := (\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies p < q$$

$$(1) \quad (\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies \dots$$

$$(1.1) \quad \forall_{p' \in \alpha} \forall_{q' \in \mathbb{Q}} (q' < p' \implies q' \in \alpha)$$

from: [CutII](#), 1

$$(1.2) \quad q < p \implies q \in \alpha \quad \blacksquare \quad q \notin \alpha \implies q \geq p$$

from: 1

$$(1.3) \quad (q \notin \alpha) \implies \dots$$

$$(1.3.1) \quad q \geq p$$

from: 1.2

$$(1.3.2) \quad (q = p) \implies (p \in \alpha \wedge p \notin \alpha) \implies \perp \quad \blacksquare \quad q \neq p$$

from: 1, 1.3

$$(1.3.3) \quad q \geq p \wedge q \neq p \quad \blacksquare \quad p < q$$

$$(1.4) \quad q \notin \alpha \implies p < q \quad \blacksquare \quad p < q$$

from: 1

$$(2) \quad (\alpha \in \mathbb{R} \wedge p \in \alpha \wedge q \in \mathbb{Q} \wedge q \notin \alpha) \implies p < q$$

---

**CutCorollaryI1**  $:= (\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies s \notin \alpha$

---

(1)  $(\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies \dots$

---

(1.1)  $\forall_{s' \in \alpha} \forall_{r' \in \mathbb{Q}} (r' < s' \implies r' \in \alpha)$

---

from: [CutI1](#), 1

(1.2)  $s \in \alpha \implies (r \in \mathbb{Q} \implies (r < s \implies r \in \alpha)) \blacksquare s \in \alpha \implies r \in \alpha$

---

from: 1, 1.1

(1.3)  $r \notin \alpha \implies s \notin \alpha \blacksquare s \notin \alpha$

---

from: 1, 1.2

(2)  $(\alpha \in \mathbb{R} \wedge r, s \in \mathbb{Q} \wedge r < s \wedge r \notin \alpha) \implies s \notin \alpha$

---

$<_{\mathbb{R}}[\alpha, \beta] := \alpha, \beta \in \mathbb{R} \wedge \alpha < \beta$

**OrderTrichotomyOfR**  $:= \text{OrderTrichotomy}[\mathbb{R}, <_{\mathbb{R}}]$

---

(1)  $(\alpha, \beta \in \mathbb{R}) \implies \dots$

---

(1.1)  $\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \implies \dots$

---

(1.1.1)  $\alpha \not\subset \beta \wedge \alpha \neq \beta$

---

from:  $<_{\mathbb{R}}$ , 1.1

(1.1.2)  $\exists_{p'} (p' \in \alpha \wedge p' \notin \beta) \blacksquare p := \text{choice}(\{p' : p' \in \alpha \wedge p' \notin \beta\})$

---

(1.1.3)  $q \in \beta \implies \dots$

---

(1.1.3.1)  $p, q \in \mathbb{Q}$

---

(1.1.3.2)  $q < p$

---

from: [CutCorollaryI](#)

(1.1.3.3)  $q \in \alpha$

---

from: [CutI1](#)

(1.1.4)  $q \in \beta \implies q \in \alpha$

---

(1.1.5)  $\forall_{q \in \beta} (q \in \alpha) \blacksquare \beta \subseteq \alpha$

---

(1.1.6)  $\beta \subset \alpha \blacksquare \beta <_{\mathbb{R}} \alpha$

---

(1.2)  $\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \implies \beta <_{\mathbb{R}} \alpha$

---

(1.3)  $\neg(\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \vee (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta) \blacksquare (\beta <_{\mathbb{R}} \alpha) \vee (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta)$

---

(1.4)  $\alpha = \beta \implies \neg(\alpha <_{\mathbb{R}} \beta \vee \beta <_{\mathbb{R}} \alpha)$

---

(1.5)  $\alpha <_{\mathbb{R}} \beta \implies \neg(\alpha = \beta \vee \beta <_{\mathbb{R}} \alpha)$

---

(1.6)  $\beta <_{\mathbb{R}} \alpha \implies \neg(\alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$

---

(1.7)  $\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta$

---

(2)  $(\alpha, \beta \in \mathbb{R}) \implies (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$

---

(3)  $\forall_{\alpha, \beta \in \mathbb{R}} (\alpha <_{\mathbb{R}} \beta \vee \alpha = \beta \vee \alpha <_{\mathbb{R}} \beta)$

---

(4) [OrderTrichotomy](#) $[\mathbb{R}, <_{\mathbb{R}}]$

---

**OrderTransitivityOfR**  $:= \text{OrderTransitivity}[\mathbb{R}, <_{\mathbb{R}}]$

---

(1)  $(\alpha, \beta, \gamma \in \mathbb{R}) \implies \dots$

---

(1.1)  $(\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \dots$

---

(1.1.1)  $\alpha \subset \beta \wedge \beta \subset \gamma$

---

(1.1.2)  $\forall_{a \in \alpha} (a \in \beta) \wedge \forall_{b \in \beta} (b \in \gamma)$

---

(1.1.3)  $\forall_{a \in \alpha} (\alpha \in \gamma) \blacksquare \alpha \subset \gamma \blacksquare \alpha <_{\mathbb{R}} \gamma$

---

(1.2)  $(\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma$

---

(2)  $(\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma)$

---

(3)  $\forall_{\alpha, \beta, \gamma \in \mathbb{R}} ((\alpha <_{\mathbb{R}} \beta \wedge \beta <_{\mathbb{R}} \gamma) \implies \alpha <_{\mathbb{R}} \gamma)$

---

(4) [OrderTransitivity](#) $[\mathbb{R}, <_{\mathbb{R}}]$

---

**OrderOfR**  $:= \text{Order}[<_{\mathbb{R}}, \mathbb{R}]$

---

from: [OrderTrichotomyR](#), [OrderTransitivityR](#)  
wts:

**LUBPropertyOfR**  $:= \text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}]$

---

(1)  $(\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \dots$

---

(1.1)  $\gamma := \{p \in \mathbb{Q} : \exists_{\alpha \in A} (p \in \alpha)\}$

---

(1.2)  $A \neq \emptyset \blacksquare \exists_{\alpha} (\alpha \in A) \blacksquare \alpha_0 := \text{choice}(\{\alpha : \alpha \in A\})$

---

(1.3)  $\alpha_0 \neq \emptyset \blacksquare \exists_a (a \in \alpha_0) \blacksquare a_0 := \text{choice}(\{a : a \in \alpha_0\}) \blacksquare a_0 \in \gamma \blacksquare \gamma \neq \emptyset$

---

(1.4) [BoundedAbove](#) $[A, \mathbb{R}, <_{\mathbb{R}}] \blacksquare \exists_{\beta} (\text{UpperBound}[\beta, A, \mathbb{R}, <_{\mathbb{R}}])$

---

(1.5)	$\beta_0 := \text{choice}(\{\beta : \text{UpperBound}[\beta, A, \mathbb{R}, <_{\mathbb{R}}]\})$
(1.6)	$\text{UpperBound}[\beta_0, A, \mathbb{R}, <_{\mathbb{R}}] \blacksquare \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \beta_0) \blacksquare \forall_{\alpha \in A}(\alpha \subseteq \beta_0) \blacksquare \forall_{\alpha \in A} \forall_{a \in \alpha}(a \in \beta_0)$
(1.7)	$(\alpha \in A \wedge a \in \alpha) \iff a \in \gamma \blacksquare \forall_{a \in \gamma}(a \in \beta_0) \blacksquare \gamma \subseteq \beta_0$
(1.8)	$\beta_0 \subset \mathbb{Q} \blacksquare \gamma \subseteq \beta_0 \subset \mathbb{Q} \blacksquare \gamma \subset \mathbb{Q}$
(1.9)	$\emptyset \neq \gamma \subset \mathbb{Q} \blacksquare \text{CutI}[\gamma]$
(1.10)	$(p \in \gamma \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$
(1.10.1)	$p \in \gamma \blacksquare \exists_{\alpha \in A}(p \in \alpha) \blacksquare \alpha_1 := \text{choice}(\{\alpha \in A : p \in \alpha\})$
(1.10.2)	$p \in \alpha_1 \wedge q \in \mathbb{Q} \wedge q < p \blacksquare q \in \alpha_1 \blacksquare q \in \gamma$
(1.11)	$(p \in \gamma \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \gamma \blacksquare \forall_{p \in \gamma} \forall_{q \in \mathbb{Q}}(q < p \implies q \in \gamma) \blacksquare \text{CutII}[\gamma]$
(1.12)	$p \in \gamma \implies \dots$
(1.12.1)	$\exists_{\alpha \in A}(p \in \alpha) \blacksquare \alpha_2 := \text{choice}(\{\alpha \in A : p \in \alpha\})$
(1.12.2)	$\alpha_2 \in \mathbb{R} \blacksquare \text{CutII}[\alpha_2] \blacksquare \exists_{r \in \alpha_2}(p < r) \blacksquare r_0 := \text{choice}(\{r \in \alpha_2 : p < r\})$
(1.12.3)	$r_0 \in \alpha_2 \blacksquare r_0 \in \gamma$
(1.12.4)	$p < r_0 \blacksquare p < r_0 \wedge r_0 \in \gamma \blacksquare \exists_{r \in \gamma}(p < r)$
(1.13)	$p \in \gamma \implies \exists_{r \in \gamma}(p < r) \blacksquare \forall_{p \in \gamma} \exists_{r \in \gamma}(p < r) \blacksquare \text{CutIII}[\gamma]$
(1.14)	$\text{CutI}[\gamma] \wedge \text{CutII}[\gamma] \wedge \text{CutIII}[\gamma] \blacksquare \gamma \in \mathbb{R}$
(1.15)	$\forall_{\alpha \in A}(\alpha \subseteq \gamma) \blacksquare \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \gamma)$
(1.16)	$\forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \gamma) \wedge \gamma \in \mathbb{R} \blacksquare \text{UpperBound}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}]$
(1.17)	$\delta <_{\mathbb{R}} \gamma \implies \dots$
(1.17.1)	$\delta \subset \gamma \blacksquare \exists_s(s \in \gamma \wedge s \notin \delta) \blacksquare s_0 := \text{choice}(\{s \in \mathbb{Q} : s \in \gamma \wedge s \notin \delta\})$
(1.17.2)	$s_0 \in \gamma \blacksquare \exists_{\alpha \in A}(s_0 \in \alpha) \blacksquare \alpha_3 := \text{choice}(\{\alpha \in A : s_0 \in \alpha\})$
(1.17.3)	$s_0 \in \alpha_3 \wedge s_0 \notin \delta \blacksquare \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta)$
(1.17.4)	$\delta \geq_{\mathbb{R}} \alpha_3 \implies \dots$
(1.17.4.1)	$\alpha_3 \subseteq \delta \blacksquare \forall_{s \in \mathbb{Q}}(s \in \alpha_3 \implies s \in \delta) \blacksquare \neg \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta)$
(1.17.4.2)	$\neg \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta) \wedge \exists_{s \in \mathbb{Q}}(s \in \alpha_3 \wedge s \notin \delta) \blacksquare \perp$
(1.17.5)	$\delta \geq_{\mathbb{R}} \alpha_3 \implies \perp \blacksquare \delta <_{\mathbb{R}} \alpha_3 \blacksquare \exists_{\alpha \in A}(\delta <_{\mathbb{R}} \alpha) \blacksquare \exists_{\alpha \in A}(\neg(\alpha \leq_{\mathbb{R}} \delta))$
(1.17.6)	$\neg \forall_{\alpha \in A}(\alpha \leq_{\mathbb{R}} \delta) \blacksquare \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}]$
(1.18)	$\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}] \blacksquare \forall_{\delta}(\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}])$
(1.19)	$\text{UpperBound}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}] \wedge \forall_{\delta}(\delta <_{\mathbb{R}} \gamma \implies \neg \text{UpperBound}[\delta, A, \mathbb{R}, <_{\mathbb{R}}])$
(1.20)	$\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}] \blacksquare \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}])$
(2)	$(\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}])$
(3)	$\forall_A((\emptyset \neq A \subset \mathbb{R} \wedge \text{BoundedAbove}[A, \mathbb{R}, <_{\mathbb{R}}]) \implies \exists_{\gamma \in S}(\text{LUB}[\gamma, A, \mathbb{R}, <_{\mathbb{R}}])) \blacksquare \text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}]$

$$+_{\mathbb{R}}[\alpha, \beta] := \alpha, \beta \in \mathbb{R} \wedge (\alpha +_{\mathbb{R}} \beta) = \{r + s : r \in \alpha \wedge s \in \beta\}$$

$$0_{\mathbb{R}} := \{x \in \mathbb{Q} : x < 0\}$$

$$0InR := 0_{\mathbb{R}} \in \mathbb{R}$$

(1)	$-1 \in 0_{\mathbb{R}} \wedge 1 \notin 0_{\mathbb{R}} \blacksquare \emptyset \neq 0_{\mathbb{R}} \subseteq \mathbb{Q} \blacksquare \text{CutI}[0_{\mathbb{R}}]$
(2)	$(x \in 0_{\mathbb{R}} \wedge y \in \mathbb{Q} \wedge y < x) \implies y < x < 0 \implies y < 0 \implies y \in 0_{\mathbb{R}} \blacksquare \forall_{x \in 0_{\mathbb{R}}} \forall_{y \in \mathbb{Q}}(y_0 < x \implies y \in 0_{\mathbb{R}}) \blacksquare \text{CutII}[0_{\mathbb{R}}]$
(3)	$y := x/2 \blacksquare (x \in 0_{\mathbb{R}}) \implies (x < y < 0) \implies \exists_{y \in 0_{\mathbb{R}}}(x < y) \blacksquare \forall_{x \in 0_{\mathbb{R}}} \exists_{y \in 0_{\mathbb{R}}}(x < y) \blacksquare \text{CutIII}[0_{\mathbb{R}}]$
(4)	$\text{CutI}[0_{\mathbb{R}}] \wedge \text{CutII}[0_{\mathbb{R}}] \wedge \text{CutIII}[0_{\mathbb{R}}] \blacksquare 0_{\mathbb{R}} \in \mathbb{R}$

$$\text{FieldAdditionClosureOf } R := (\alpha, \beta \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) \in \mathbb{R})$$

(1)	$(\alpha, \beta \in \mathbb{R}) \implies \dots$
(1.1)	$(\alpha +_{\mathbb{R}} \beta) = \{r + s : r \in \alpha \wedge s \in \beta\}$
(1.2)	$\emptyset \neq \alpha \subset \mathbb{Q} \wedge \emptyset \neq \beta \subset \mathbb{Q}$
(1.3)	$\exists_a(a \in \alpha) ; \exists_b(b \in \beta) \blacksquare a_0 := \text{choice}(\{a : a \in \alpha\}) ; b_0 := \text{choice}(\{b : b \in \beta\}) \blacksquare a_0 + b_0 \in \alpha +_{\mathbb{R}} \beta$
(1.4)	$\exists_x(x \notin \alpha) ; \exists_y(y_0 \notin \beta) \blacksquare x_0 := \text{choice}(\{x : x \notin \alpha\}) ; y_0 := \text{choice}(\{y : y \notin \beta\})$
(1.5)	$\forall_{r \in \alpha}(r < x_0) ; \forall_{s \in \beta}(s < y_0) \blacksquare \forall_{r \in \alpha} \forall_{s \in \beta}(r + s < x_0 + y_0) \blacksquare x_0 + y_0 \notin \alpha +_{\mathbb{R}} \beta$
(1.6)	$\emptyset \neq \alpha +_{\mathbb{R}} \beta \subset \mathbb{Q} \blacksquare \text{CutI}[\alpha +_{\mathbb{R}} \beta]$



- (1.7)  $(p \in \alpha +_{\mathbb{R}} \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$
- (1.7.1)  $\exists_{r \in \alpha} \exists_{s \in \beta} (p = r + s) \blacksquare (r_0, s_0) := \text{choice}((r, s) \in \alpha \times \beta : p = r + s)$
- (1.7.2)  $q < p = r_0 + s_0 \blacksquare (q - s_0) < r_0 \blacksquare (q - s_0) \in \alpha$
- (1.7.3)  $s_0 \in \beta \blacksquare q = (q - s_0) + s_0 \in \alpha +_{\mathbb{R}} \beta \blacksquare q \in \alpha +_{\mathbb{R}} \beta$
- (1.8)  $(p \in \alpha +_{\mathbb{R}} \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \alpha +_{\mathbb{R}} \beta \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} \beta} \forall_{q \in \mathbb{Q}} (q < p \implies q \in \alpha +_{\mathbb{R}} \beta) \blacksquare \text{CutII}[\alpha +_{\mathbb{R}} \beta]$
- (1.9)  $p \in \alpha \implies \dots$
- (1.9.1)  $\exists_{r \in \alpha} \exists_{s \in \beta} (p = r + s) \blacksquare (r_1, s_1) := \text{choice}(\{(r, s) \in \alpha \times \beta : p = r + s\})$
- (1.9.2)  $r_1 \in \alpha \blacksquare \exists_{t \in \alpha} (r_1 < t) \blacksquare t_0 := \text{choice}(\{t \in \alpha : r_1 < t\})$
- (1.9.3)  $s_1 \in \beta \blacksquare t + s_1 \in \alpha +_{\mathbb{R}} \beta \wedge p = r_1 + s_1 < t + s_1 \blacksquare \exists_{r \in \alpha +_{\mathbb{R}} \beta} (p < r)$
- (1.10)  $p \in \alpha \implies \exists_{r \in \alpha +_{\mathbb{R}} \beta} (p < r) \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} \beta} \exists_{r \in \alpha +_{\mathbb{R}} \beta} (p < r) \blacksquare \text{CutIII}[\alpha +_{\mathbb{R}} \beta]$
- (1.11)  $\text{CutI}[\alpha +_{\mathbb{R}} \beta] \wedge \text{CutII}[\alpha +_{\mathbb{R}} \beta] \wedge \text{CutIII}[\alpha +_{\mathbb{R}} \beta] \blacksquare \alpha +_{\mathbb{R}} \beta \in \mathbb{R}$
- (2)  $(\alpha, \beta \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) \in \mathbb{R})$

**FieldAdditionCommutativityOfR** :=  $(\alpha, \beta \in \mathbb{R}) \implies (\alpha +_{\mathbb{R}} \beta = \beta +_{\mathbb{R}} \alpha)$

- (1)  $\alpha +_{\mathbb{R}} \beta = \{r + s : r \in \alpha \wedge s \in \beta\} = \{s + r : s \in \beta \wedge r \in \alpha\} = \beta +_{\mathbb{R}} \alpha$

**FieldAdditionAssociativityOfR** :=  $(\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma))$

- (1)  $(\alpha, \beta, \gamma \in \mathbb{R}) \implies \dots$
- (1.1)  $(\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \{(a + b) + c : a \in \alpha \wedge b \in \beta \wedge c \in \gamma\} = \dots$
- (1.2)  $\{a + (b + c) : a \in \alpha \wedge b \in \beta \wedge c \in \gamma\} = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma)$
- (2)  $(\alpha, \beta, \gamma \in \mathbb{R}) \implies (\alpha +_{\mathbb{R}} \beta) +_{\mathbb{R}} \gamma = \alpha +_{\mathbb{R}} (\beta +_{\mathbb{R}} \gamma)$

**FieldAdditionIdentityOfR** :=  $(\alpha \in \mathbb{R}) \implies 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$

- (1)  $\alpha \in \mathbb{R} \implies \dots$
- (1.1)  $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \implies \dots$
- (1.1.1)  $s < 0 \blacksquare r + s < r + 0 = r \blacksquare r + s < r \blacksquare r + s \in \alpha$
- (1.2)  $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \implies r + s \in \alpha \blacksquare \forall_{r \in \alpha} \forall_{s \in 0_{\mathbb{R}}} (r + s \in \alpha)$
- (1.3)  $(r \in \alpha \wedge s \in 0_{\mathbb{R}}) \iff (r + s \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}) \blacksquare \forall_{p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}} (p \in \alpha) \blacksquare \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \subseteq \alpha$
- (1.4)  $p \in \alpha \implies \dots$
- (1.4.1)  $\exists_{r \in \alpha} (p < r) \blacksquare r_2 := \text{choice}(\{r \in \alpha : p < r\})$
- (1.4.2)  $p < r_2 \blacksquare p - r_2 < r_2 - r_2 = 0 \blacksquare (p - r_2) < 0 \blacksquare (p - r_2) \in 0_{\mathbb{R}}$
- (1.4.3)  $r_2 \in \alpha \blacksquare p = r_2 + (p - r_2) \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}$
- (1.5)  $p \in \alpha \implies p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare \forall_{p \in \alpha} (p \in \alpha +_{\mathbb{R}} 0_{\mathbb{R}}) \blacksquare \alpha \subseteq \alpha +_{\mathbb{R}} 0_{\mathbb{R}}$
- (1.6)  $\alpha +_{\mathbb{R}} 0_{\mathbb{R}} \subseteq \alpha \wedge \alpha \subseteq \alpha +_{\mathbb{R}} 0_{\mathbb{R}} \blacksquare 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$
- (2)  $\alpha \in \mathbb{R} \implies 0_{\mathbb{R}} +_{\mathbb{R}} \alpha = \alpha$

**FieldAdditionInverseOfR** :=  $(\alpha \in \mathbb{R}) \implies \exists_{-\alpha \in \mathbb{R}} (\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$

- (1)  $\alpha \in \mathbb{R} \implies \dots$
- (1.1)  $\beta := \{p \in \mathbb{Q} : \exists_{r > 0} (-p - r \notin \alpha)\}$
- (1.2)  $\alpha \subset \mathbb{Q} \blacksquare \exists_{s \in \mathbb{Q}} (s \notin \alpha) \blacksquare s_0 := \text{choice}(\{s : s \notin \alpha\}) \blacksquare p_0 := -s_0 - 1$
- (1.3)  $-p_0 - 1 = -(-s_0 - 1) - 1 = s_0 \notin \alpha \blacksquare -p_0 - 1 \notin \alpha \blacksquare \exists_{r > 0} (-p_0 - r \notin \alpha) \blacksquare p_0 \in \beta$
- (1.4)  $\emptyset \neq \alpha \blacksquare \exists_{q \in \alpha} \blacksquare q_0 := \text{choice}(\{q \in \mathbb{Q} : q \in \alpha\})$
- (1.5)  $r > 0 \implies \dots$
- (1.5.1)  $q_0 \in \alpha \blacksquare -(-q_0) - r = q_0 - r < q_0 \blacksquare -(-q_0) - r < q_0 \blacksquare -(-q_0) - r \in \alpha$
- (1.6)  $\forall_{r > 0} (-(-q_0) - r \in \alpha) \blacksquare \neg \exists_{r > 0} (-(-q_0) - r \notin \alpha) \blacksquare -q_0 \notin \beta$
- (1.7)  $\emptyset \neq \beta \subset \mathbb{Q} \blacksquare \text{CutI}[\beta]$
- (1.8)  $(p \in \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies \dots$
- (1.8.1)  $p \in \beta \blacksquare \exists_{r > 0} (-p - r \notin \alpha) \blacksquare r_0 := \text{choice}(\{r > 0 : -p - r \notin \alpha\})$
- (1.8.2)  $q < p \blacksquare -p - r < -q - r$
- (1.8.3)  $-q - r \notin \alpha \blacksquare q \in \beta$
- (1.9)  $(p \in \beta \wedge q \in \mathbb{Q} \wedge q < p) \implies q \in \beta \blacksquare \forall_{p \in \beta} \forall_{q \in \mathbb{Q}} (q < p \implies q \in \beta) \blacksquare \text{CutII}[\beta]$

(1.10)	$p \in \beta \implies \dots$	
(1.10.1)	$p \in \beta \implies \exists_{r>0}(-p-r \notin \alpha) \blacksquare r_1 := \text{choice}(\{r > 0 : -p-r \notin \alpha\})$	
(1.10.2)	$t_0 := p + (r_1/2)$	
(1.10.3)	$r_1 > 0 \blacksquare r_1/2 > 0$	
(1.10.4)	$t_0 > t_0 - (r_1/2) = p \blacksquare t_0 > p$	
(1.10.5)	$-t_0 - (r_1/2) = -(p + (r_1/2)) - (r_1/2) = -p - r_1$	
(1.10.6)	$-p - r_1 \notin \alpha \blacksquare -t_0 - (r_1/2) \notin \alpha \blacksquare \exists_{r>0}(-t_0 - r \notin \alpha) \blacksquare t_0 \in \beta$	
(1.10.7)	$t_0 > p \wedge t_0 \in \beta \blacksquare \exists_{t \in \beta}(p < t)$	
(1.11)	$p \in \beta \implies \exists_{t \in \beta}(p < t) \blacksquare \forall_{p \in \beta} \exists_{t \in \beta}(p < t) \blacksquare \text{CutIII}[\beta]$	
(1.12)	$\text{CutI}[\beta] \wedge \text{CutII}[\beta] \wedge \text{CutIII}[\beta] \blacksquare \beta \in \mathbb{R}$	
(1.13)	$(r \in \alpha \wedge s \in \beta) \implies \dots$	
(1.13.1)	$s \in \beta \blacksquare \exists_{t>0}(-s-t \notin \alpha) \blacksquare t_1 := \text{choice}(\{t > 0 : -s-t \notin \alpha\}) \blacksquare -s - t_1 < -s$	
(1.13.2)	$\alpha \in \mathbb{R} \wedge s, t_1 \in \mathbb{Q} \wedge -s - t_1 < -s \wedge -s - t_1 \notin \alpha \blacksquare -s \notin \alpha$	
(1.13.3)	$\alpha \in \mathbb{R} \wedge r \in \alpha \wedge -s \notin \alpha \blacksquare r < -s \blacksquare r + s < 0 \blacksquare r + s \in 0_{\mathbb{R}}$	
(1.14)	$(r \in \alpha \wedge s \in \beta) \implies r + s \in 0_{\mathbb{R}} \blacksquare \forall_{(r,s) \in \alpha \times \beta}(r + s \in 0_{\mathbb{R}}) \blacksquare \alpha +_{\mathbb{R}} \beta \subseteq 0_{\mathbb{R}}$	
(1.15)	$v \in 0_{\mathbb{R}} \implies \dots$	
(1.15.1)	$v < 0 \blacksquare w_0 := -v/2 \blacksquare w > 0$	
(1.15.2)	$\exists_{n \in \mathbb{Z}}(nw_0 \in \alpha \wedge (n+1)w_0 \notin \alpha) \blacksquare n_0 := \text{choice}(\{n \in \mathbb{Z} : nw_0 \in \alpha \wedge (n+1)w_0 \notin \alpha\})$	from: ARCHIMEDEANPROPERTYOFQ + LUB???
(1.15.3)	$p_0 := -(n_0 + 2)w_0 \blacksquare -p_0 - w_0 = (n_0 + 2)w_0 - w_0 = (n_0 + 1)w_0 \notin \alpha \blacksquare -p_0 - w_0 \notin \alpha \blacksquare p_0 \in \beta$	
(1.15.4)	$n_0 w_0 \in \alpha \wedge p_0 \in \beta \blacksquare n_0 w_0 + p_0 = n_0(-v/2) + -(n_0 + 2) - v/2 = v \in \alpha +_{\mathbb{R}} \beta$	
(1.16)	$v \in 0_{\mathbb{R}} \implies v \in \alpha +_{\mathbb{R}} \beta \blacksquare \forall_{v \in 0_{\mathbb{R}}}(v \in \alpha +_{\mathbb{R}} \beta) \blacksquare 0_{\mathbb{R}} \subseteq \alpha +_{\mathbb{R}} \beta$	
(1.17)	$\alpha +_{\mathbb{R}} \beta \subseteq 0_{\mathbb{R}} \wedge 0_{\mathbb{R}} \subseteq \alpha +_{\mathbb{R}} \beta \blacksquare \alpha +_{\mathbb{R}} \beta = 0_{\mathbb{R}}$	
(1.18)	$\beta \in \mathbb{R} \wedge \alpha +_{\mathbb{R}} \beta = 0_{\mathbb{R}} \blacksquare \exists_{-\alpha \in \mathbb{R}}(\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$	
(2)	$\alpha \in \mathbb{R} \implies \exists_{-\alpha \in \mathbb{R}}(\alpha +_{\mathbb{R}} (-\alpha) = 0_{\mathbb{R}})$	

$*_{\mathbb{R}}[\alpha, \beta] := \text{---}$

$1_{\mathbb{R}} := \{x \in \mathbb{Q} : x < 1\}$

$11sNot0 := 0_{\mathbb{R}} \neq 1_{\mathbb{R}} \text{ ---}$

$11nR := 1_{\mathbb{R}} \in \mathbb{R} \text{ ---}$

$\text{FieldMultiplicationClosureOf } \mathbb{R} := (\alpha, \beta \in \mathbb{R}) \implies ((\alpha *_{\mathbb{R}} \beta) \in \mathbb{R}) \text{ ---}$

$\text{FieldMultiplicationCommutativityOf } \mathbb{R} := (\alpha, \beta \in \mathbb{R}) \implies (\alpha *_{\mathbb{R}} \beta = \beta *_{\mathbb{R}} \alpha) \text{ ---}$

$\text{FieldMultiplicationAssociativityOf } \mathbb{R} := (\alpha, \beta, \gamma \in \mathbb{R}) \implies ((\alpha *_{\mathbb{R}} \beta) *_{\mathbb{R}} \gamma = \alpha *_{\mathbb{R}} (\beta *_{\mathbb{R}} \gamma)) \text{ ---}$

$\text{FieldMultiplicationIdentityOf } \mathbb{R} := (\alpha \in \mathbb{R}) \implies 1_{\mathbb{R}} *_{\mathbb{R}} \alpha = \alpha \text{ ---}$

$\text{FieldMultiplicationInverseOf } \mathbb{R} := (\alpha \in \mathbb{R}) \implies \exists_{1/\alpha \in \mathbb{R}}(\alpha *_{\mathbb{R}} (1/\alpha) = 1_{\mathbb{R}}) \text{ ---}$

$\text{FieldDistributivityOf } \mathbb{R} := (\alpha, \beta, \gamma \in \mathbb{R}) \implies \gamma *_{\mathbb{R}} (\alpha +_{\mathbb{R}} \beta) = \gamma *_{\mathbb{R}} \alpha + \gamma *_{\mathbb{R}} \beta \text{ ---}$

$\text{FieldWith } \mathbb{R} := \text{Field}[\mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}] \text{ ---}$

$\text{OrderedFieldWith } \mathbb{R} := \text{OrderedField}[\mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}] \text{ ---}$

$\mathbb{Q}_{\mathbb{R}} := \{\{r \in \mathbb{Q} : r < q\} : q \in \mathbb{Q}\}$

$\text{QROrderedSubfieldOf } \mathbb{R} := \text{OrderedSubfield}[\mathbb{Q}_{\mathbb{R}}, \mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}] \text{ ---}$

$\text{QIsomorphicToQR} := \mathbb{Q}_{\mathbb{R}} \simeq \mathbb{Q} \text{ ---}$

$\text{CompletenessOf } \mathbb{R} := \exists_{\mathbb{R}}(\text{LUBProperty}[\mathbb{R}, <_{\mathbb{R}}] \wedge \text{OrderedSubfield}[\mathbb{Q}, \mathbb{R}, +_{\mathbb{R}}, *_{\mathbb{R}}, <_{\mathbb{R}}]) \text{ ---}$

(1.20)

$\text{ArchimedeanPropertyOf } \mathbb{R} := \forall_{x,y \in \mathbb{R}}(x > 0 \implies \exists_{n \in \mathbb{N}^+}(nx > y))$

(1)  $(x, y \in \mathbb{R} \wedge x > 0) \implies \dots$

(1.1)  $A := \{nx : n \in \mathbb{N}^+\} \blacksquare (\emptyset \neq A \subset \mathbb{R}) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a))$

(1.2)  $\neg \exists_{n \in \mathbb{N}^+}(nx > y) \implies \dots$

(1.2.1)  $\neg \exists_{n \in \mathbb{N}^+}(nx > y) \blacksquare \forall_{n \in \mathbb{N}^+}(nx \leq y) \blacksquare \text{UpperBound}[y_0, A, \mathbb{R}, <] \blacksquare \text{BoundedAbove}[A, \mathbb{R}, <]$

(1.2.2)  $\text{CompletenessOf } \mathbb{R} \blacksquare \text{LUBProperty}[\mathbb{R}, <]$

(1.2.3)  $(\text{LUBProperty}[\mathbb{R}, <]) \wedge (\emptyset \neq A \subset \mathbb{R}) \wedge (\text{BoundedAbove}[A, \mathbb{R}, <]) \blacksquare \exists_{\alpha \in \mathbb{R}}(\text{LUB}[\alpha, A, \mathbb{R}, <]) \dots$

(1.2.4)	$\dots \alpha_0 := \text{choice}(\{\alpha \in \mathbb{R} : \textcolor{teal}{LUB}[\alpha, A, \mathbb{R}, <]\}) \quad \blacksquare \quad \textcolor{teal}{LUB}[\alpha_0, A, \mathbb{R}, <]$
(1.2.5)	$x > 0 \quad \blacksquare \quad \alpha_0 - x < \alpha_0$
(1.2.6)	$(\alpha_0 - x < \alpha_0) \wedge (\textcolor{teal}{LUB}[\alpha_0, A, \mathbb{R}, <]) \quad \blacksquare \quad \neg \textcolor{teal}{UpperBound}[\alpha_0 - x, A, \mathbb{R}, <]$
(1.2.7)	$\neg \textcolor{teal}{UpperBound}[\alpha_0 - x, A, \mathbb{R}, <] \quad \blacksquare \quad \exists_{c \in A}(\alpha_0 - x < c) \quad \dots$
(1.2.8)	$\dots c_0 := \text{choice}(\{c \in A : \alpha_0 - x < c\}) \quad \blacksquare \quad (c_0 \in A) \wedge (\alpha_0 - x < c_0)$
(1.2.9)	$(c_0 \in A) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a)) \quad \blacksquare \quad \exists_{m \in \mathbb{N}^+}(mx = c_0) \quad \dots$
(1.2.10)	$\dots m_0 := \text{choice}(\{m \in \mathbb{N}^+ : mx = c_0\}) \quad \blacksquare \quad (m_0 \in \mathbb{N}^+) \wedge (m_0 x = c_0)$
(1.2.11)	$(\alpha_0 - x < c_0) \wedge (m_0 x = c_0) \quad \blacksquare \quad \alpha_0 - x < c_0 = m_0 x \quad \blacksquare \quad \alpha_0 < m_0 x + x \quad \blacksquare \quad \alpha_0 < (m_0 + 1)x$
(1.2.12)	$m_0 \in \mathbb{N}^+ \quad \blacksquare \quad m_0 + 1 \in \mathbb{N}^+$
(1.2.13)	$(m_0 + 1 \in \mathbb{N}^+) \wedge (a \in A \iff \exists_{m \in \mathbb{N}^+}(mx = a)) \quad \blacksquare \quad (m_0 + 1)x \in A$
(1.2.14)	$(\alpha_0 < (m_0 + 1)x) \wedge ((m_0 + 1)x \in A) \quad \blacksquare \quad \exists_{c \in A}(\alpha_0 < c)$
(1.2.15)	$\textcolor{teal}{LUB}[\alpha_0, A, \mathbb{R}, <] \quad \blacksquare \quad \textcolor{teal}{UpperBound}[\alpha_0, A, \mathbb{R}, <] \quad \blacksquare \quad \forall_{c \in A}(c \leq \alpha_0) \quad \blacksquare \quad \neg \exists_{c \in A}(c > \alpha_0) \quad \blacksquare \quad \neg \exists_{c \in A}(\alpha_0 < c)$
(1.2.16)	$(\exists_{c \in A}(\alpha_0 < c)) \wedge (\neg \exists_{c \in A}(\alpha_0 < c)) \quad \blacksquare \quad \perp$
(1.3)	$\neg \exists_{n \in \mathbb{N}^+}(nx > y) \implies \perp \quad \blacksquare \quad \exists_{n \in \mathbb{N}^+}(nx > y)$
(2)	$(x, y \in \mathbb{R} \wedge x > 0) \implies \exists_{n \in \mathbb{N}^+}(nx > y) \quad \blacksquare \quad \forall_{x, y \in \mathbb{R}}(x > 0 \implies \exists_{n \in \mathbb{N}^+}(nx > y))$

***QDenseInR*** :=  $\forall_{x, y \in \mathbb{R}}(x < y \implies \exists_{p \in \mathbb{Q}}(x < p < y))$

(1)	$(x, y \in \mathbb{R} \wedge x < y) \implies \dots$
(1.1)	$x < y \quad \blacksquare \quad (0 < y - x) \wedge (y - x \in \mathbb{R})$
(1.2)	$\textcolor{teal}{ArchimedeanPropertyOfR} \wedge (0 < y - x) \wedge (y - x, 1 \in \mathbb{R}) \quad \blacksquare \quad \exists_{n \in \mathbb{N}^+}(n(y - x) > 1) \quad \dots$
(1.3)	$\dots n_0 := \text{choice}(\{n \in \mathbb{N}^+ : n(y - x) > 1\}) \quad \blacksquare \quad (n_0 \in \mathbb{N}^+) \wedge (n_0(y - x) > 1)$
(1.4)	$(n_0 \in \mathbb{N}^+) \wedge (x \in \mathbb{R}) \quad \blacksquare \quad n_0 x, -n_0 x \in \mathbb{R}$
(1.5)	$\textcolor{teal}{ArchimedeanPropertyOfR} \wedge (1 > 0) \wedge (n_0 x, 1 \in \mathbb{R}) \quad \blacksquare \quad \exists_{m \in \mathbb{N}^+}(m(1) > n_0 x) \quad \dots$
(1.6)	$\dots m_1 := \text{choice}(\{m \in \mathbb{N}^+ : m(1) > n_0 x\}) \quad \blacksquare \quad (m_1 \in \mathbb{N}^+) \wedge (m_1 > n_0 x)$
(1.7)	$\textcolor{teal}{ArchimedeanPropertyOfR} \wedge (1 > 0) \wedge (-n_0 x, 1 \in \mathbb{R}) \quad \blacksquare \quad \exists_{m \in \mathbb{N}^+}(m(1) > -n_0 x) \quad \dots$
(1.8)	$\dots m_2 := \text{choice}(\{m \in \mathbb{N}^+ : m(1) > -n_0 x\}) \quad \blacksquare \quad (m_2 \in \mathbb{N}^+) \wedge (m_2 > -n_0 x)$
(1.9)	$(m_1 > n_0 x) \wedge (m_2 > -n_0 x) \quad \blacksquare \quad -m_2 < n_0 x < m_1$
(1.10)	$m_1, m_2 \in \mathbb{N}^+ \quad \blacksquare \quad  m_1 - (-m_2)  \geq 2$
(1.11)	$(-m_2 < n_0 x < m_1) \wedge ( m_1 - (-m_2)  \geq 2) \quad \blacksquare \quad \exists_{m \in \mathbb{Z}}((-m_2 < m < m_1) \wedge (m - 1 \leq n_0 x < m)) \quad \dots$
(1.12)	$\dots m_0 := \text{choice}(\{m \in \mathbb{Z} : (-m_2 < m < m_1) \wedge (m - 1 \leq n_0 x < m)\}) \quad \blacksquare \quad (-m_2 < m_0 < m_1) \wedge (m_0 - 1 \leq n_0 x < m_0)$
(1.13)	$(n_0(y - x) > 1) \wedge (m_0 - 1 \leq n_0 x < m_0) \quad \blacksquare \quad n_0 x < m_0 \leq 1 + n_0 x < n_0 y \quad \blacksquare \quad n_0 x < m_0 < n_0 y$
(1.14)	$(n_0 \in \mathbb{N}^+) \wedge (n_0 x < m_0 < n_0 y) \quad \blacksquare \quad x < m_0/n_0 < y$
(1.15)	$m_0, n_0 \in \mathbb{Z} \quad \blacksquare \quad m_0/n_0 \in \mathbb{Q}$
(1.16)	$(m_0/n_0 \in \mathbb{Q}) \wedge (x < m_0/n_0 < y) \quad \blacksquare \quad \exists_{p \in \mathbb{Q}}(x < p < y)$
(2)	$(x, y \in \mathbb{R} \wedge x < y) \implies \exists_{p \in \mathbb{Q}}(x < p < y) \quad \blacksquare \quad \forall_{x, y \in \mathbb{R}}(x < y \implies \exists_{p \in \mathbb{Q}}(x < p < y))$

(1.21)

***Root Lemma*** :=  $(0 < a < b) \implies (b^n - a^n \leq (b - a)nb^{n-1})$

(1)	$(0 < a < b) \implies \dots$
(1.1)	$b^n - a^n = (b - a) \sum_{i=1}^n (b^{n-i} a^{i-1})$
(1.2)	$0 < a < b \quad \blacksquare \quad b/a > 1$
(1.3)	$b/a > 1 \quad \blacksquare \quad \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq \sum_{i=1}^n (b^{n-i} a^{i-1} (b/a)^{i-1}) = \sum_{i=1}^n (b^{n-1}) = nb^{n-1} \quad \blacksquare \quad \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq \sum_{i=1}^n (b^{n-1}) = nb^{n-1}$
(1.4)	$b^n - a^n = (b - a) \sum_{i=1}^n (b^{n-i} a^{i-1}) \leq (b - a)nb^{n-1} \quad \blacksquare \quad b^n - a^n \leq (b - a)nb^{n-1}$
(2)	$(0 < a < b) \implies (b^n - a^n \leq (b - a)nb^{n-1})$

***Root Existence In R*** :=  $\forall_{0 < x \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} \exists!_{0 < y \in \mathbb{R}} (y_0^n = x)$

(1)	$(0 < x \in \mathbb{R} \wedge 0 < n \in \mathbb{Z}) \implies \dots$
(1.1)	$E := \{t \in \mathbb{R} : t > 0 \wedge t^n < x\} \quad \blacksquare \quad t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)$
(1.2)	$t_0 := x/(1 + x) \quad \blacksquare \quad (t_0 = x/(1 + x)) \wedge (t_0 \in \mathbb{R})$
(1.3)	$0 < x \quad \blacksquare \quad 0 < x < 1 + x \quad \blacksquare \quad t_0 = x/(1 + x) > 0 \quad \blacksquare \quad t_0 > 0$

$$(1.4) \quad 1 = (1+x)/(1+x) > x/(1+x) = t_0 \quad \blacksquare \quad 1 > t_0$$

$$(1.5) \quad (t_0 > 0) \wedge (1 > t_0) \quad \blacksquare \quad 0 < t_0 < 1$$

$$(1.6) \quad (0 < n \in \mathbb{Z}) \wedge (0 < t_0 < 1) \quad \blacksquare \quad t_0^n \leq t_0$$

$$(1.7) \quad 0 < x \quad \blacksquare \quad x > x/(1+x) = t_0 \quad \blacksquare \quad x > t_0$$

$$(1.8) \quad (t_0^n \leq t_0) \wedge (x > t_0) \quad \blacksquare \quad t_0^n < x$$

$$(1.9) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t_0 \in \mathbb{R}) \wedge (t_0 > 0) \wedge (t_0^n < x) \quad \blacksquare \quad t_0 \in E \quad \blacksquare \quad \emptyset \neq E$$

$$(1.10) \quad t_1 := \text{choice}(\{t \in \mathbb{R} : t > 1+x\}) \quad \blacksquare \quad (t_1 \in \mathbb{R}) \wedge (t_1 > 1+x)$$

$$(1.11) \quad x > 0 \quad \blacksquare \quad t_1 > 1+x > 1 \quad \blacksquare \quad t_1 > 1 \quad \blacksquare \quad t_1^n \geq t_1$$

$$(1.12) \quad (t_1^n \geq t_1) \wedge (t_1 > 1+x) \wedge (1 > 0) \quad \blacksquare \quad t_1^n \geq t_1 > 1+x > x \quad \blacksquare \quad t_1^n > x$$

$$(1.13) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t_1^n > x) \quad \blacksquare \quad t_1 \notin E \quad \blacksquare \quad E \subset \mathbb{R}$$

$$(1.14) \quad (\emptyset \neq E) \wedge (E \subset \mathbb{R}) \quad \blacksquare \quad \emptyset \neq E \subset \mathbb{R}$$

$$(1.15) \quad t \in E \implies \dots$$

$$(1.15.1) \quad (t \in E) \wedge (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \quad \blacksquare \quad t^n < x$$

$$(1.15.2) \quad (t_1^n > x) \wedge (t^n < x) \quad \blacksquare \quad t^n < x < t_1^n \quad \blacksquare \quad t < t_1$$

$$(1.16) \quad t \in E \implies t < t_1 \quad \blacksquare \quad \forall_{t \in E} (t \leq t_1) \quad \blacksquare \quad \text{UpperBound}[t_1, E, \mathbb{R}, <] \quad \blacksquare \quad \text{BoundedAbove}[E, \mathbb{R}, <]$$

$$(1.17) \quad \text{CompletenessOfR} \quad \blacksquare \quad \text{LUBProperty}[\mathbb{R}, <]$$

$$(1.18) \quad (\text{LUBProperty}[\mathbb{R}, <]) \wedge (\emptyset \neq E \subset \mathbb{R}) \wedge (\text{BoundedAbove}[E, \mathbb{R}, <]) \quad \blacksquare \quad \exists_{y \in \mathbb{R}} (\text{LUB}[y, E, \mathbb{R}, <]) \quad \dots$$

$$(1.19) \quad \dots y_0 := \text{choice}(\{y \in \mathbb{R} : \text{LUB}[y, E, \mathbb{R}, <]\}) \quad \blacksquare \quad \text{LUB}[y_0, E, \mathbb{R}, <]$$

$$(1.20) \quad (\text{LUB}[y_0, E, \mathbb{R}, <]) \wedge (t_0 \in E) \wedge (t_0 > 0) \quad \blacksquare \quad 0 < t_0 \leq y_0 \in \mathbb{R} \quad \blacksquare \quad 0 < y_0 \in \mathbb{R}$$

$$(1.21) \quad y_0^n < x \implies \dots$$

$$(1.21.1) \quad k_0 := \frac{x-y_0^n}{n(y_0+1)^{n-1}} \quad \blacksquare \quad k_0 \in \mathbb{R}$$

$$(1.21.2) \quad y_0^n < x \quad \blacksquare \quad 0 < x - y_0^n$$

$$(1.21.3) \quad (n > 0) \wedge (y_0 > 0) \quad \blacksquare \quad 0 < n(y_0 + 1)^{n-1}$$

$$(1.21.4) \quad (0 < x - y_0^n) \wedge (0 < n(y_0 + 1)^{n-1}) \quad \blacksquare \quad 0 < \frac{x-y_0^n}{n(y_0+1)^{n-1}} = k_0 \quad \blacksquare \quad 0 < k_0$$

$$(1.21.5) \quad (0 < 1 \in \mathbb{R}) \wedge (0 < k_0 \in \mathbb{R}) \quad \blacksquare \quad 0 < \min(1, k_0) \in \mathbb{R}$$

$$(1.21.6) \quad \text{QDenseInR} \wedge (0, \min(1, k_0) \in \mathbb{R}) \wedge (0 < \min(1, k_0)) \quad \blacksquare \quad \exists_{h \in \mathbb{Q}} (0 < h < \min(1, k_0)) \quad \dots$$

$$(1.21.7) \quad \dots h_0 := \text{choice}(\{h \in \mathbb{Q} : 0 < h < \min(1, k_0)\}) \quad \blacksquare \quad (0 < h_0 < 1) \wedge (h_0 < k_0 = \frac{x-y_0^n}{n(y_0+1)^{n-1}})$$

$$(1.21.8) \quad (y_0 > 0) \wedge (h_0 > 0) \quad \blacksquare \quad 0 < y_0 < y_0 + h_0$$

$$(1.21.9) \quad \text{RootLemma} \wedge (0 < y_0 < y_0 + h_0) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < h_0 n (y_0 + h_0)^{n-1}$$

$$(1.21.10) \quad h_0 < 1 \quad \blacksquare \quad h_0 n (y_0 + h_0)^{n-1} < h_0 n (y_0 + 1)^{n-1}$$

$$(1.21.11) \quad ((y_0 + h_0)^n - y_0^n < h_0 n (y_0 + h_0)^{n-1}) \wedge (h_0 n (y_0 + h_0)^{n-1} < h_0 n (y_0 + 1)^{n-1}) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < h_0 n (y_0 + 1)^{n-1}$$

$$(1.21.12) \quad (0 < n(y_0 + 1)^{n-1}) \wedge (h_0 < k_0 = \frac{x-y_0^n}{n(y_0+1)^{n-1}}) \quad \blacksquare \quad h_0 n (y_0 + 1)^{n-1} < x - y_0^n$$

$$(1.21.13) \quad ((y_0 + h_0)^n - y_0^n < h_0 n (y_0 + 1)^{n-1}) \wedge (h_0 n (y_0 + 1)^{n-1} < x - y_0^n) \quad \blacksquare \quad (y_0 + h_0)^n - y_0^n < x - y_0^n \quad \blacksquare \quad (y_0 + h_0)^n < x$$

$$(1.21.14) \quad (y_0 + h_0)^n - y_0^n < x - y_0^n \quad \blacksquare \quad (y_0 + h_0)^n < x$$

$$(1.21.15) \quad (0 < y_0 \in \mathbb{R}) \wedge (0 < h_0 \in \mathbb{R}) \quad \blacksquare \quad 0 < y_0 < y_0 + h_0 \in \mathbb{R}$$

$$(1.21.16) \quad (t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge ((y_0 + h_0)^n < x) \wedge (0 < y_0 + h_0 \in \mathbb{R}) \quad \blacksquare \quad (y_0 + h_0)^n \in E$$

$$(1.21.17) \quad ((y_0 + h_0)^n \in E) \wedge (y_0 < y_0 + h_0) \quad \blacksquare \quad \exists_{e \in E} (y_0 < e)$$

$$(1.21.18) \quad \text{LUB}[y_0, E, \mathbb{R}, <] \quad \blacksquare \quad \text{UpperBound}[y_0, E, \mathbb{R}, <] \quad \blacksquare \quad \forall_{e \in E} (e \leq y_0) \quad \blacksquare \quad \neg \exists_{e \in E} (e > y_0)$$

$$(1.21.19) \quad (\exists_{e \in E} (e > y_0)) \wedge (\neg \exists_{e \in E} (e > y_0)) \quad \blacksquare \quad \perp$$

$$(1.22) \quad y_0^n < x \implies \perp \quad \blacksquare \quad y_0^n \geq x$$

$$(1.23) \quad y_0^n > x \implies \dots$$

$$(1.23.1) \quad k_1 := \frac{y_0^n - x}{n y_0^{n-1}} \quad \blacksquare \quad (k_1 \in \mathbb{R}) \wedge (k_1 n y_0^{n-1} = y_0^n - x)$$

$$(1.23.2) \quad (0 < x) \wedge (0 < n \in \mathbb{Z}) \quad \blacksquare \quad y_0^n - x < y_0^n \leq n y_0^n \quad \blacksquare \quad y_0^n - x < n y_0^n$$

$$(1.23.3) \quad y_0^n - x < n y_0^n \quad \blacksquare \quad k_1 = \frac{y_0^n - x}{n y_0^{n-1}} < \frac{n y_0^n}{n y_0^{n-1}} = y_0 \quad \blacksquare \quad k_1 < y_0$$

$$(1.23.4) \quad y_0^n > x \quad \blacksquare \quad 0 < y_0^n - x$$

$$(1.23.5) \quad (n > 0) \wedge (y_0 > 0) \quad \blacksquare \quad 0 < n y_0^{n-1}$$

$$(1.23.6) \quad (0 < y_0^n - x) \wedge 0 < (n y_0^{n-1}) \quad \blacksquare \quad 0 < \frac{y_0^n - x}{n y_0^{n-1}} = k_1 \quad \blacksquare \quad 0 < k_1$$

(1.23.7)	$(k_1 < y_0) \wedge (0 < k_1) \quad \blacksquare \quad (0 < k_1 < y_0) \wedge (0 < y_0 - k_1 < y_0)$	
(1.23.8)	$t \geq y_0 - k_1 \implies \dots$	
(1.23.8.1)	$t \geq y_0 - k_1 \quad \blacksquare \quad t^n \geq (y_0 - k_1)^n \quad \blacksquare \quad -t^n \leq -(y_0 - k_1)^n \quad \blacksquare \quad y_0^n - t^n \leq y_0^n - (y_0 - k_1)^n$	
(1.23.8.2)	$\textcolor{blue}{RootLemma} \wedge (0 < y_0 - k_1 < y_0) \quad \blacksquare \quad y_0^n - (y_0 - k_1)^n < k_1 n y_0^{n-1}$	
(1.23.8.3)	$(y_0^n - t^n \leq y_0^n - (y_0 - k_1)^n) \wedge (y_0^n - t^n < k_1 n y_0^{n-1}) \quad \blacksquare \quad y_0^n - t^n < k_1 n y_0^{n-1}$	
(1.23.8.4)	$(k_1 n y_0^{n-1} = y_0^n - x) \wedge (y_0^n - t^n < k_1 n y_0^{n-1}) \quad \blacksquare \quad y_0^n - t^n < y_0^n - x \quad \blacksquare \quad -t^n < -x \quad \blacksquare \quad t^n > x$	
(1.23.8.5)	$(t \in E \iff (t \in \mathbb{R} \wedge t > 0 \wedge t^n < x)) \wedge (t^n > x) \quad \blacksquare \quad t \notin E$	
(1.23.9)	$t \geq y_0 - k_1 \implies t \notin E \quad \blacksquare \quad t \in E \implies t < y_0 - k_1 \quad \blacksquare \quad \forall_{t \in E} (t \leq y_0 - k_1) \quad \blacksquare \quad \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]$	
(1.23.10)	$(\textcolor{blue}{LUB}[y_0, E, \mathbb{R}, <] \wedge (y_0 - k_1 < y_0)) \quad \blacksquare \quad \neg \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]$	
(1.23.11)	$(\textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]) \wedge (\neg \textcolor{blue}{UpperBound}[y_0 - k_1, E, \mathbb{R}, <]) \quad \blacksquare \quad \perp$	
(1.24)	$y_0^n > x \implies \perp \quad \blacksquare \quad y_0^n \leq x$	
(1.25)	$\textcolor{blue}{Order}[\mathbb{R}, <] \quad \blacksquare \quad \textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]$	
(1.26)	$(\textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]) \wedge (y_0^n \geq x) \wedge (y_0^n \leq x) \quad \blacksquare \quad y_0^n = x$	
(1.27)	$(y_0^n = x) \wedge (y_0 \in \mathbb{R}) \quad \blacksquare \quad \exists_{y \in \mathbb{R}} (y^n = x)$	
(1.28)	$y_1, y_2 := \textit{choice}(\{y \in \mathbb{R} : y^n = x\})$	
(1.29)	$y_1 \neq y_2 \implies \dots$	
(1.29.1)	$(\textcolor{blue}{OrderTrichotomy}[\mathbb{R}, <]) \wedge (y_1 \neq y_2) \quad \blacksquare \quad (y_1 < y_2) \vee (y_2 < y_1) \quad \dots$	
(1.29.2)	$\dots (x = y_1^n < y_2^n = x) \vee (x = y_2^n < y_1^n = x) \quad \blacksquare \quad (x < x) \vee (x > x) \quad \blacksquare \quad \perp \vee \perp \quad \blacksquare \quad \perp$	
(1.30)	$y_1 \neq y_2 \implies \perp \quad \blacksquare \quad y_1 = y_2 \quad \blacksquare \quad \forall_{a,b \in \mathbb{R}} ((a^n = x \wedge b^n = x) \implies a = b)$	
(1.31)	$(\exists_{y \in \mathbb{R}} (y^n = x)) \wedge (\forall_{a,b \in \mathbb{R}} ((a^n = x \wedge b^n = x) \implies a = b)) \quad \blacksquare \quad \exists!_{y \in \mathbb{R}} (y^n = x)$	
(2)	$(0 < x \in \mathbb{R} \wedge 0 < n \in \mathbb{Z}) \implies \exists!_{y \in \mathbb{R}} (y^n = x) \quad \blacksquare \quad \forall_{0 < x \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} \exists!_{0 < y \in \mathbb{R}} (y_0^n = x)$	

$$\textcolor{red}{RootExistenceInRCorollary} := \forall_{0 < a \in \mathbb{R}} \forall_{0 < b \in \mathbb{R}} \forall_{0 < n \in \mathbb{Z}} ((ab)^{1/n} = a^{1/n} b^{1/n}) \quad \text{---}$$

$$\textcolor{red}{ExtendedRealSystem}[\bar{\mathbb{R}}, +, *, <] := \left( \begin{array}{l} \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\} \quad \wedge \quad -\infty < x < \infty \quad \wedge \\ x + \infty = +\infty \quad \wedge \quad x - \infty = -\infty \quad \wedge \quad \frac{x}{+\infty} = \frac{x}{-\infty} = 0 \quad \wedge \\ (x > 0) \implies (x * (+\infty) = +\infty \wedge x * (-\infty) = -\infty) \wedge \\ (x < 0) \implies (x * (+\infty) = -\infty \wedge x * (-\infty) = +\infty) \end{array} \right)$$

$$\mathbb{C} := \{\langle a, b \rangle \in \mathbb{R} \times \mathbb{R}\}$$

$$+_C[\langle a, b \rangle, \langle c, d \rangle] := \langle a +_{\mathbb{R}} c, b +_{\mathbb{R}} d \rangle$$

$$*_C[\langle a, b \rangle, \langle c, d \rangle] := \langle a *_R c - b *_R d, a *_R d + b *_R c \rangle$$

$$\textcolor{red}{FieldC} := \textcolor{blue}{Field}[\mathbb{C}, +_C, *_C] \quad \text{---}$$

$$\textcolor{red}{RSubfieldC} := \textcolor{blue}{Subfield}[\mathbb{R}, \mathbb{C}, +, *] \quad \text{---}$$

$$i := \langle 0, 1 \rangle \in \mathbb{C}$$

$$\textcolor{red}{iProperty} := i^2 = -1 \quad \text{---}$$

$$\textcolor{red}{CProperty} := (a, b \in \mathbb{R}) \implies (\langle a, b \rangle = a + bi) \quad \text{---}$$

$$\textcolor{red}{Conjugate}[\overline{a + bi}] := a - bi$$

$$\textcolor{red}{ConjugateProperties} := (w, z \in \mathbb{C}) \implies \dots \quad \text{---}$$

$$(1) \quad \overline{z + w} = \bar{z} + \bar{w}$$

$$(2) \quad \overline{z * w} = \bar{z} * \bar{w}$$

$$(3) \quad \textit{Re}(z) = (1/2)(z + \bar{z}) \wedge \textit{Im}(z) = (1/2)(z - \bar{z})$$

$$(4) \quad 0 \leq z * \bar{z} \in \mathbb{R}$$

$$\textcolor{red}{AbsoluteValueC}[|z|] = (z * \bar{z})^{1/2}$$

$$\textcolor{red}{AbsoluteValueProperties} := (z, w \in \mathbb{C}) \implies \dots \quad \text{---}$$

$$(1) \quad 123123$$



# Chapter 2

## Abstract Algebra

$Relation(f, X) := f \subseteq X$   
 $Function(f, X, Y) := X \neq \emptyset \neq Y \wedge Relation(f, X \times Y) \wedge \forall_{x \in X} \exists!_{y \in Y} ((x, y) \in f)$

$(Function(f, X, Y) \wedge A \subseteq X \wedge B \subseteq Y) \implies \dots$

(1)  $Domain(f) := X; Codomain(f) := Y$

(2)  $Image(f, A) := \{f(a) : a \in A\}; Preimage(f, B) := \{a : f(a) \in B\}$

(3)  $Range(f) := Image(Domain(f))$

$Injective(f, X, Y) := Function(f, X, Y) \wedge \forall_{x_1, x_2 \in X} (x_1 \neq x_2 \implies f(x_1) \neq f(x_2))$

$Surjective(f, X, Y) := Function(f, X, Y) \wedge \forall_{y \in Y} \exists_{x \in X} (y = f(x))$

$Bijjective(f, X, Y) := Injective(f, X, Y) \wedge Surjective(f, X, Y)$

**Surjective Equivalent**  $:= (Range(f) = Codomain(f)) \implies Surjective(f)$

$(Function(f, X, Y) \wedge Function(g, Y, Z)) \implies (f \circ g)(x) := f(g(x)); Function(f \circ g, X, Z)$

**Properties of Functions**  $:= (Function(f, A, B) \wedge Function(g, B, C) \wedge Function(h, C, D)) \implies \dots$

(1)  $h \circ (g \circ f) = (h \circ g) \circ f$

(2)  $(Injective(f) \wedge Injective(g)) \implies Injective(g \circ f)$

(3)  $(Surjective(f) \wedge Surjective(g)) \implies Surjective(g \circ f)$

(4)  $(Bijjective(f, A, B)) \implies \exists_{f^{-1}} (Function(f^{-1}, B, A) \wedge \forall_{a \in A} (f^{-1}(f(a)) = a) \wedge \forall_{b \in B} (f(f^{-1}(b)) = b))$

$|(a, b) := a, b \in \mathbb{Z} \wedge a \neq 0 \wedge \exists_{c \in \mathbb{Z}} (b = ac)$

**Divisibility Theorems**  $:= (a, b, c, m, x, y \in \mathbb{Z}) \implies \dots$

(1)  $(a|b) \implies a|bc$

(2)  $(a|b \wedge b|c) \implies a|c$

(3)  $(a|b \wedge b|c) \implies a|(bx + cy)$

(4)  $(a|b \wedge b|a) \implies a = \pm b$

(5)  $(a|b \wedge a > 0 \wedge b > 0) \implies (a \leq b)$

(6)  $(a|b) \iff (m \neq 0 \wedge ma|mb)$

**Division Algorithm**  $:= (a, b \in \mathbb{Z} \wedge a > 0) \implies \exists!_{q, r \in \mathbb{Z}} (b = aq + r)$

**CD** $(a, b, c) := a, b, c \in \mathbb{Z} \wedge a : b \wedge a : c$

**GCD** $(a, b, c) := CD(a, b, c) \wedge \forall_d ((d : b \wedge d : c) \implies d : a)$

**GCD Equivalent**  $:= 123123$





# Chapter 3

## Linear Algebra

BASE ON ADVANCED BOOK

$Matrix(A, m, n) := [a_{i,j}]_{m \times n} := m \text{ rows, } n \text{ columns of real numbers}$

$\mathcal{M}_{m,n} := \{A : Matrix(A, m, n)\}$

$O_{m,n} := (Matrix(O, m, n)) \wedge (a_{i,j} = 0)$

$Square(A, n) := Matrix(A, n, n)$

$UpperTriangular(A) := (Square(A)) \wedge (i > j \implies a_{i,j} = 0)$

$LowerTriangular(A) := (Square(A)) \wedge (i < j \implies a_{i,j} = 0)$

$Diagonal(A, n) := (Square(A, n)) \wedge (i \neq j \implies a_{i,j} = 0)$

$Scalar(A, n, k) := (Diagonal(A, n)) \wedge (a_{i,i} = k)$

$I_n := Scalar(I, n, 1)$

$+(A, B) := ((Matrix(A, m, n)) \wedge (Matrix(B, m, n))) \implies (A + B = [a_{i,j} + b_{i,j}]_{m \times n})$

$*(r, A) := ((r \in \mathbb{R}) \wedge (Matrix(A, m, n))) \implies (r * A = [ra_{i,j}]_{m \times n})$

$*(A, B) := ((Matrix(A, m, p)) \wedge (Matrix(B, p, n))) \implies (A * B = \left[ \sum_{k=1}^p (a_{i,k} b_{k,j}) \right]_{m \times n})$

$T(A) := (Matrix(A, m, n)) \implies (A^T = [a_{j,i}]_{n \times m})$

$AddCom := \forall_{A,B \in \mathcal{M}} (A + B = B + A) \quad \blacksquare \Leftarrow A + B = [a_{i,j} + b_{i,j}] = [b_{i,j} + a_{i,j}] = B + A$

$AddAssoc := \forall_{A,B,C \in \mathcal{M}} ((A + B) + C = A + (B + C)) \quad \blacksquare \Leftarrow (A + B) + C = [(a_{i,j} + b_{i,j}) + c_{i,j}] = [a_{i,j} + (b_{i,j} + c_{i,j})] = A + (B + C)$

$AddId := \forall_{A \in \mathcal{M}} \exists!_{O \in \mathcal{M}} (A + O = A = O + A) \quad \text{---}$

$AddInv := \forall_{A \in \mathcal{M}} \exists!_{(-A) \in \mathcal{M}} (A + (-A) = O = (-A) + A) \quad \text{---}$

$MulAssoc := \forall_{A,B,C \in \mathcal{M}} ((A * B) * C = A * (B * C)) \quad \blacksquare \Leftarrow$

---


$$(1) \quad (A * B) * C = \left[ \sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,j}) \right] * C = \left[ \sum_{k_2=1}^{p_2} \left( \sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,k_2}) c_{k_2,j} \right) \right] = \left[ \sum_{k_2=1}^{p_2} \sum_{k_1=1}^{p_1} (a_{i,k_1} b_{k_1,k_2} c_{k_2,j}) \right] = \dots$$


---

$$(2) \quad \dots \left[ \sum_{k_1=1}^{p_1} \sum_{k_2=1}^{p_2} (a_{i,k_1} b_{k_1,k_2} c_{k_2,j}) \right] = \left[ \sum_{k_1=1}^{p_1} (a_{i,k_1} \sum_{k_2=1}^{p_2} (b_{k_1,k_2} c_{k_2,j})) \right] = \dots = A * (B * C)$$


---

$MulId := \forall_{A: Square(A,n)} (A * I_n = A = I_n * A) \quad \text{---}$

$LeftDist := \forall_{A,B,C \in \mathcal{M}} ((A + B) * C = A * C + B * C) \quad \blacksquare \Leftarrow$

---


$$(1) \quad (A + B) * C = [a_{i,j} + b_{i,j}] * C = \left[ \sum_{k=1}^p ((a_{i,k} + b_{i,k}) c_{k,j}) \right] = \dots$$


---

$$(2) \quad \dots \left[ \sum_{k=1}^p (a_{i,k} c_{k,j} + b_{i,k} c_{k,j}) \right] = \left[ \sum_{k=1}^p (a_{i,k} c_{k,j}) \right] + \left[ \sum_{k=1}^p (b_{i,k} c_{k,j}) \right] = A * C + B * C$$


---

$RightDist := \forall_{A,B,C \in \mathcal{M}} (C * (A + B) * C = C * A + C * B) \quad \text{---}$

$Scalar1 := \forall_{r,s \in \mathbb{R}} \forall_{A \in \mathcal{M}} (r(sA) = (rs)A = s(rA)) \quad \text{---}$

$Scalar2 := \forall_{r \in \mathbb{R}} \forall_{A,B \in \mathcal{M}} (A(rB) = r(A * B)) \quad \text{---}$

$Scalar3 := \forall_{r,s \in \mathbb{R}} \forall_{A \in \mathcal{M}} ((r + s)A = rA + sA) \quad \text{---}$

$Scalar4 := \forall_{r \in \mathbb{R}} \forall_{A,B \in \mathcal{M}} (r(A + B) = rA + rB) \quad \text{---}$

$Trans1 := \forall_{A \in \mathcal{M}} (A = (A^T)^T) \quad \blacksquare \Leftarrow A = [a_{i,j}] = [a_{j,i}]^T = ([a_{i,j}]^T)^T = (A^T)^T$

$Trans2 := \forall_{A,B \in \mathcal{M}} ((A + B)^T = A^T + B^T) \quad \text{---}$

$Trans3 := \forall_{A,B \in \mathcal{M}} ((A * B)^T = B^T * A^T) \quad \blacksquare \Leftarrow$

---


$$(1) \quad (A * B)^T = \left[ \sum_{k=1}^p (a_{i,k} b_{k,j}) \right]^T = \left[ \sum_{k=1}^p (a_{j,k} b_{k,i}) \right] = \left[ \sum_{k=1}^p (b_{k,i} a_{j,k}) \right] = \left[ \sum_{k=1}^p (b_{i,k}^T a_{k,j}^T) \right] = B^T * A^T$$


---

$$\begin{aligned}
\text{Sym}(A) &:= A = A^T \\
\text{SkewSym}(A) &:= A = -A^T \\
\text{SymGen} &:= \forall_{A \in \mathcal{M}} (\text{Sym}(A + A^T)) \quad \blacksquare \Leftarrow (A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T \\
\text{SkewSymGem} &:= \forall_{A \in \mathcal{M}} (\text{SkewSym}(A - A^T)) \quad \blacksquare \Leftarrow -(A - A^T)^T = -(A^T - (A^T)^T) = -(A^T - A) = (A - A^T) \\
\text{SymDec} &:= \forall_{A \in \mathcal{M}} \exists!_{B: \text{Sym}(B)} \exists!_{C: \text{SkewSym}(C)} (A = B + C = (1/2) * (A + A^T) + (1/2) * (A - A^T)) \quad \text{---}
\end{aligned}$$

$$\begin{aligned}
\text{Invertible}(A) &:= (\text{Square}(A, n)) \wedge (\exists_{A^{-1} \in \mathcal{M}} (A * A^{-1} = I_n = A^{-1} * A)) \\
\text{Inv1} &:= \forall_{A: \text{Invertible}(A)} (\exists!_{A^{-1} \in \mathcal{M}} (A * A^{-1} = I_n = A^{-1} * A)) \quad \text{---} \\
\text{Inv2} &:= \forall_{A: \text{Invertible}(A)} \forall_{B: \text{Invertible}(B)} ((A * B)^{-1} = B^{-1} * A^{-1}) \quad \blacksquare \Leftarrow
\end{aligned}$$

$$(1) \quad (A * B) * (A * B)^{-1} = I \quad \blacksquare \quad B * (A * B)^{-1} = A^{-1} \quad \blacksquare \quad (A * B)^{-1} = B^{-1} * A^{-1}$$

$$\begin{aligned}
\text{Inv3} &:= \forall_{A: \text{Invertible}(A)} ((A^{-1})^{-1} = A) \quad \text{---} \\
\text{InvTrans} &:= \forall_{A: \text{Invertible}(A)} ((A^T)^{-1} = (A^{-1})^T) \quad \blacksquare \Leftarrow
\end{aligned}$$

$$(1) \quad A^T * (A^{-1})^T = (A^{-1} * A)^T = I^T = I \quad \blacksquare \quad (A^{-1})^T = (A^T)^{-1}$$

$$\begin{aligned}
\text{RowOp}(\phi) &:= (\text{Definition1.19}) \\
\text{RowEquiv}(A, B) &:= \exists_{\Phi: |\Phi| \in \mathbb{N}} ((\forall_{\phi \in \Phi} (\text{RowOp}(\phi))) \wedge (\Phi(A) = B)) \\
\text{RREF}(A) &:= (\text{Definition1.18}) \\
\text{GaussJordan} &:= \forall_A \exists!_B (\text{RREF}(B) \wedge \text{RowEquiv}(A, B)) \quad \text{---}
\end{aligned}$$

TRASH CAN

$$\begin{aligned}
\text{Sys}(A, X, B) &:= (A * X = B) \wedge (\text{Matrix}(A, m, n)) \wedge (\text{Matrix}(X, n, 1)) \wedge (\text{Matrix}(B, m, 1)) \\
\text{Sol}(X, A, B) &:= \text{Sys}(A, X, B) \\
\text{TrivSol}(X, A, B) &:= (\text{Sol}(X, A, B)) \wedge (x_{i,1} = 0) \\
\text{SysAlgebra} &:= \forall_{A, B, C, D, X} ((\text{RowEquiv}([A|B], [C|D])) \implies (\text{Sys}(A, X, B) \iff \text{Sys}(C, X, D))) \quad \text{---} \\
\text{SysAlgebraCorollary} &:= \forall_{A, B, X} ((\text{RowEquiv}(A, B)) \implies (\text{Sys}(A, X, O) \iff \text{Sys}(B, X, O))) \quad \text{---} \\
\text{TrivSolLemma} &:= (m < n) \implies (\exists_X (\neg \text{TrivSol}(X, A, O))) \quad \blacksquare \quad (\forall_X (\text{TrivSol}(X, A, O)) \implies (m \geq n)) \quad \blacksquare \Leftarrow
\end{aligned}$$

$$(1) \quad B = \text{GaussJordan}(A) \quad \blacksquare \quad \text{RowEquiv}(A, B) \quad \blacksquare \quad \text{Sys}(A, X, O) \iff \text{Sys}(B, X, O)$$

$$(2) \quad ((m < n) \wedge \text{Sys}(B, X, O)) \implies X \text{ has at least } n - m \text{ free variables}$$

$$(3) \quad \text{Let } B = \text{GaussJordan}(A) \text{ and let } r \text{ be the number of non-zero rows of } B \quad \blacksquare \quad r \leq m$$

$$(4) \quad (r \leq m) \wedge (m < n) \quad \blacksquare \quad r < n \quad \blacksquare \quad 0 < n - r$$

$$(5) \quad \text{The solution of } B \text{ will have } r \text{ fixed variables and } n - r \text{ free variables which is also more than zero}$$

$$(6) \quad \text{The solution of } B \text{ will have at least one free variable that can be non-zero, i.e., non-trivial}$$

TRASH CAN

$$\begin{aligned}
\text{Elem}(A, n) &:= \text{RowEquiv}(A, I_n) \\
\text{RowOpAsElem} &:= \forall_{\phi: \text{RowOp}(\phi)} ((E = \phi(I)) \implies (\phi(A) = E * A)) \\
\text{RowEquiv2} &:= \text{RowEquiv}(A, B) \iff \exists_{E: |E| \in \mathbb{N}} (A = \Pi_{E_i \in E} (E_i) * B) \\
(\text{Elem}_i(A)) &\implies (\text{Invertible}(A) \wedge \text{Elem}_i(A^{-1})) \\
(\text{RowEquiv}(A, B)) &\iff (B = \Pi(E_i) * A) \\
(\text{RowEquiv}(I, B)) &\iff (B = \Pi(E_i))
\end{aligned}$$

$$(\forall_X (\text{TrivSol}(X, A, O)) \implies (\text{RowEquiv}(A, I))) \quad \blacksquare \Leftarrow$$

$$(1) \quad \text{Let } B = \text{GaussJordan}(A) \quad \blacksquare \quad \text{RowEquiv}(A, B)$$

$$(2) \quad \forall_X (\text{TrivSol}(X, B, O) \quad \blacksquare \quad n \geq n \quad \blacksquare \quad n = n \quad \blacksquare \quad B = I)$$

$$(3) \quad \text{RowEquiv}(A, B) \quad \blacksquare \quad \text{RowEquiv}(A, I)$$

$$(\text{Invertible}(A)) \iff (A = \Pi(E_i)) \quad \blacksquare \Leftarrow$$

$$(1) \quad \text{Invertible}(A) \implies \dots$$

$$(1.1) \quad AX = O \quad \blacksquare \quad X = A^{-1}O = O \quad \blacksquare \quad \forall_X (\text{TrivSol}(X, A, O))$$

$$(1.2) \quad \forall_X (\text{TrivSol}(X, A, O) \quad \blacksquare \quad \text{RowEquiv}(A, I) \quad \blacksquare \quad A = \Pi(E_i))$$

$$(2) \quad A = \Pi(E_i) \implies \dots$$

$$(2.1) \quad \Pi(E_i^{-1}) = A^{-1} \quad \blacksquare \quad \text{Invertible}(A)$$

$$(3) \quad (\text{Invertible}(A)) \iff (A = \Pi(E_i))$$

---


$$(Invertible(A)) \iff (RowEquiv(I, A)) \blacksquare \Leftarrow (RowEquiv(I, B) \iff B = \Pi(E_i)) \wedge ((Invertible(A)) \iff (A = \Pi(E_i)))$$

$$(\forall_X(TrivSol(X, A, O))) \implies (Invertible(A)) \blacksquare \Leftarrow$$

---


$$(1) \quad ((\forall_X(TrivSol(X, A, O)) \implies (RowEquiv(A, I))) \wedge ((Invertible(A)) \iff (RowEquiv(I, A))))$$


---

$$(Invertible(A)) \implies (\forall_X(TrivSol(X, A, O))) \blacksquare \Leftarrow AX = O \blacksquare X = A^{-1}O = O \blacksquare \forall_X(TrivSol(X, A, O))$$