

MT621M-Elementary Number Theory

Mathematical Preliminaries

Francis Joseph Campena, PhD
Mathematics and Statistics Department
De La Salle University-Manila

Introduction

In this chapter, we recall some mathematical tools needed to prove theorems and results in number theory. Most statements in mathematics are stated in a form of propositions, theorems or implications which are often made up of the **HYPOTHESIS** and the **CONCLUSION**.

These statements are often phrased in the following manner:

- **Conditional:** $P \Rightarrow Q$ which is read as " P implies Q " or IF P , then Q .
- **Equivalence:** $P \Leftrightarrow Q$ which is read as " P if and only if Q ". This means that P is both necessary and sufficient condition for Q .

Introduction

In this chapter, we recall some mathematical tools needed to prove theorems and results in number theory. Most statements in mathematics are stated in a form of propositions, theorems or implications which are often made up of the **HYPOTHESIS** and the **CONCLUSION**.

These statements are often phrased in the following manner:

- **Conditional:** $P \Rightarrow Q$ which is read as " P implies Q " or IF P , then Q .
- **Equivalence:** $P \Leftrightarrow Q$ which is read as " P if and only if Q ". This means that P is both necessary and sufficient condition for Q .

Introduction

In this chapter, we recall some mathematical tools needed to prove theorems and results in number theory. Most statements in mathematics are stated in a form of propositions, theorems or implications which are often made up of the **HYPOTHESIS** and the **CONCLUSION**.

These statements are often phrased in the following manner:

- **Conditional:** $P \Rightarrow Q$ which is read as " P implies Q " or IF P , then Q .
- **Equivalence:** $P \Leftrightarrow Q$ which is read as " P if and only if Q ". This means that P is both necessary and sufficient condition for Q .

Introduction

In this chapter, we recall some mathematical tools needed to prove theorems and results in number theory. Most statements in mathematics are stated in a form of propositions, theorems or implications which are often made up of the **HYPOTHESIS** and the **CONCLUSION**.

These statements are often phrased in the following manner:

- **Conditional:** $P \Rightarrow Q$ which is read as " P implies Q " or IF P , then Q .
- **Equivalence:** $P \Leftrightarrow Q$ which is read as " P if and only if Q ". This means that P is both necessary and sufficient condition for Q .

Methods of Proof

There are four main methods of proving propositions.

1. Direct Proof
2. Indirect Proof
3. Proof by Contradiction
4. Proof by Mathematical Induction

Methods of Proof

There are four main methods of proving propositions.

1. Direct Proof
2. Indirect Proof
3. Proof by Contradiction
4. Proof by Mathematical Induction

Methods of Proof

There are four main methods of proving propositions.

1. Direct Proof
2. Indirect Proof
3. Proof by Contradiction
4. Proof by Mathematical Induction

Methods of Proof

There are four main methods of proving propositions.

1. Direct Proof
2. Indirect Proof
3. Proof by Contradiction
4. Proof by Mathematical Induction

Methods of Proof

There are four main methods of proving propositions.

1. Direct Proof
2. Indirect Proof
3. Proof by Contradiction
4. Proof by Mathematical Induction

Direct Proof

Direct Proof: In this method, we begin with the hypothesis and progress logically using known results or facts and definitions to arrive at the conclusion.

Example

If x, y are odd integers, then $x + y$ is even.

Proof.

Let $x, y \in \mathbb{Z}$. Since, x, y are odd integers, there exists integers a, b such that $x = 2a + 1, y = 2b + 1$. This gives us,

$$\begin{aligned}x + y &= 2a + 1 + 2b + 1 \\ &= 2(a + b + 1)\end{aligned}$$

Since $x + y = 2(a + b + 1)$, the conclusion follows. □

Direct Proof

Direct Proof: In this method, we begin with the hypothesis and progress logically using known results or facts and definitions to arrive at the conclusion.

Example

If x, y are odd integers, then $x + y$ is even.

Proof.

Let $x, y \in \mathbb{Z}$. Since, x, y are odd integers, there exists integers a, b such that $x = 2a + 1, y = 2b + 1$. This gives us,

$$\begin{aligned}x + y &= 2a + 1 + 2b + 1 \\ &= 2(a + b + 1)\end{aligned}$$

Since $x + y = 2(a + b + 1)$, the conclusion follows. □

Indirect Proof

Indirect Proof: In this method, we use the fact that the conditional statement $P \Rightarrow Q$ is equivalent to its contrapositive that is $\neg Q \Rightarrow \neg P$

Example

If x, y are odd integers, then $x + y$ is even.

Show that if $x + y$ is not even, then exactly one of x or y must be even.

Indirect Proof Example

Proof by Contradiction

Proof by Contradiction: In this method, we use the negation of the conclusion together with our hypothesis to obtain a false statement or contradiction.

Example

Show that there is no largest even integer.

Proof.

Suppose now. That is, suppose there is a largest even integer, say k . Since k is even, there exist $n \in \mathbb{Z}$ such that $k = 2n$. Now, consider $k + 2$. $k + 2 = (2n) + 2 = 2(n + 1)$. So $k + 2$ is even. But $k + 2$ is larger than k . This contradicts our assumption that k was the largest even integer. So our original claim must have been true. \square

If x, y are odd integers, then $x + y$ is even: Proof By Contradiction

Proof by Mathematical Induction

Theorem (The Well-Ordering Property)

Every non-empty set of positive integers has a least element.

Mathematical Induction : This method of proof is used when the statement to be proven contains a function of an integer variable.

Definition (First Principle of Mathematical Induction)

Let S_m, S_{m+1}, \dots be a sequence of statements, where m is a positive integer. Suppose that the following conditions are true:

1. S_m is true
2. If an arbitrary statement S_k is true, then the next statement S_{k+1} is also true.

Proof by Mathematical Induction

Theorem (The Well-Ordering Property)

Every non-empty set of positive integers has a least element.

Mathematical Induction : This method of proof is used when the statement to be proven contains a function of an integer variable.

Definition (First Principle of Mathematical Induction)

Let S_m, S_{m+1}, \dots be a sequence of statements, where m is a positive integer. Suppose that the following conditions are true:

1. S_m is true
2. If an arbitrary statement S_k is true, then the next statement S_{k+1} is also true.

Proof by Mathematical Induction

Definition (Second Principle of Mathematical Induction)

A set of positive integers which contains the integer 1, and which has the property that if it contains all the positive integers $1, 2, \dots, k$, then it also contains the integer $k + 1$, must be the set of all positive integers.

An alternative definition for the Second Principle of Mathematical Induction is given below:

Definition (Second Principle of Mathematical Induction)

Suppose that a property $P(n)$ is true for $n = 1$. If $P(n)$ is also true for all $n < k$, where k is an arbitrary integer greater than or equal to 1 implies $P(n)$ is true for $n = k$ then $P(n)$ must be true for all $n \in \mathbb{N}$.

EXERCISES Part 1

1. Show that the sum of two consecutive integers is odd.
2. Prove that if an integer k is odd then k^2 is odd.
3. Use Mathematical induction to prove the following statements
 - a) Show that $4^n - 1$ is divisible by 3 for all positive integer n .
 - b) Show that for $n \in \mathbb{Z}^+$, the sum of the first n consecutive integers is $\frac{n(n+1)}{2}$.
 - c) Show that the sum of the first n odd natural numbers is n^2 .
 - d) Show that for all positive integers n , $3^n > 3n - 1$.
 - e) Show that for any natural number n , $n^3 - n$ is divisible by 3.

EXERCISES Part 2

4. A triangular number is a number that represents the number of dots that can be arranged evenly in an equilateral triangle. Equivalently, the k^{th} triangular number is given by $t_k = \frac{k(k+1)}{2}$.
- (a) List the first 6 triangular numbers.
 - (b) Prove that an integer n is a triangular number if and only if $8n + 1$ is a perfect square.
 - (c) Let n be a triangular number. Prove that $(2k + 1)2n + t_k$ is also a triangular number.