


Summary of Suicide Deaths in Nebraska


284 Total Deaths (14.6 per 100,000 Population)

Demographic Information




In 2021, male suicide victims were predominantly white males aged 40-49 with a high school education. American Indian/Alaska Native non-Hispanic and Asian/Pacific Islander non-Hispanic males were disproportionately affected by suicide in Nebraska. The average age of male deaths was 45, while for females, it was 42.

Cause of Death




In 2021, firearms were the most common method of suicide among males in Nebraska, with an average age of 48. Strangulation had an average age of 40, and poisoning (including drug overdose) averaged 41. Among females, strangulation was the most common method, with an average age of 37. Firearms had an average age of 42, and poisoning (including drug overdose) averaged 46.

Suicide Deaths by Month



Suicide Death Rate by Census Tracts



Due to small numbers, rates may be unstable.

Other Decedent Characteristics

Statistics are based on 253 cases. Categories are not mutually exclusive.



73.9%

of total suicide deaths took place in a residence.

9.5%

of total suicide victims experienced homelessness or housing instability.

52.6%

of total suicide victims had previous mental health problems.

7.9%

of total suicide victims were diagnosed with depression.



7.5%

of total suicide victims were diagnosed with anxiety disorder.

24.9%

of total suicide victims had history of mental health treatment.

17.0%

of total suicide victims had current mental health treatment.

57.7%

of total suicide victims were perceived by others to be depressed at the time of death.

34.0%

of total suicide victims had a history of suicidal ideation.

14.6%

of total suicide victims had a history of suicide attempt.



26.5%

of total suicide victims had recently disclosed their suicidal intent.

11.5%

of total suicide victims had disclosed their suicidal intent to their intimate partner.

32.8%

of total suicide victims left suicide notes.



22.5%

of total suicide victims had a history of alcohol abuse.

22.5%

of total suicide victims had a history of substance abuse other than alcohol.

22.9%

of total suicide victims had a history of physical health problems.

11.5%

of total suicide victims had criminal or legal problems.

26.1%

of total suicide victims had intimate partner problems.

7.9%

of total suicide victims had family or other relationship problems.



16.6%

of total suicide victims had known recent argument.

41.1%

of the suicide deaths were among never married individuals.

26.9%

of the suicide deaths were among married individuals.

The reduced death count in this section is due to incomplete law enforcement or coroner reports.