



# Office Hours!

Instructor:

Peter M. Garfield, [garfield@math.ucsb.edu](mailto:garfield@math.ucsb.edu)

Office Hours:

Mondays 2–3PM

Tuesdays 10:30–11:30AM

Thursdays 1–2PM

or by appointment

Office:


South Hall 6510

© 2017 Daryl Cooper, Peter M. Garfield

# Sketching some simple graphs

It's useful to be able to sketch...

## (1) Quadratics



$$y = 2x^2 + 4x - 44$$


$$y = -2x^2 + 4x + 38$$

- $y = ax^2 + bx + c$
- Bowl-shaped:
  - ★ Opens up if  $a > 0$
  - ★ Opens down if  $a < 0$
- Model curve:  $y = x^2$

# Sketching some simple graphs

It's useful to be able to sketch...

## (1) Quadratics



$$y = 2x^2 + 4x - 44$$


$$y = -2x^2 + 4x + 38$$

- $y = ax^2 + bx + c$
- Bowl-shaped:
  - ★ Opens up if  $a > 0$
  - ★ Opens down if  $a < 0$
- Model curve:  $y = x^2$   
Shown here!

# Sketching some simple graphs

It's useful to be able to sketch...

## (2) Cubics



$$y = 2x^3 - 6x - 15$$


$$y = -2x^3 + 3x^2 + 12x - 10$$

- $y = ax^3 + bx^2 + cx + d$
- “S”-shaped:
  - ★ Goes to  $+\infty$  if  $a > 0$
  - ★ Goes to  $-\infty$  if  $a < 0$
- Model curve:  $y = x^3$

# Sketching some simple graphs

It's useful to be able to sketch...

## (2) Cubics



$$y = 2x^3 - 6x - 15$$

$$y = -2x^3 + 3x^2 + 12x - 10$$


- $y = ax^3 + bx^2 + cx + d$
- “S”-shaped:
  - ★ Goes to  $+\infty$  if  $a > 0$
  - ★ Goes to  $-\infty$  if  $a < 0$
- Model curve:  $y = x^3$

Shown here!

# Sketching some simple graphs

It's useful to be able to sketch...

## (2) Cubics



$$y = 2x^3 - 6x - 15$$

$$y = -2x^3 + 3x^2 + 12x - 10$$


- $y = ax^3 + bx^2 + cx + d$
- “S”-shaped:
  - ★ Goes to  $+\infty$  if  $a > 0$
  - ★ Goes to  $-\infty$  if  $a < 0$
- Model curve:  $y = x^3$

Shown here!


For a polynomial, the **highest power** of  $x$  **dominates** when  $x$  is big

# The Derivatives of Simple Functions

The derivative of a constant is...?



# The Derivatives of Simple Functions




The derivative of a constant is zero because:

- derivative = rate of change
- constants don't change




# The Derivatives of Simple Functions



The derivative of a constant is zero because:

- derivative = rate of change
- constants don't change
- derivative = slope
- slope = 0

# The Derivatives of Simple Functions




The derivative of a constant is zero because:

- derivative = rate of change
- constants don't change
- derivative = slope
- slope = 0

So  $\frac{d}{dx}(5) = 0$

# The Derivatives of Simple Functions




The derivative of a constant is zero because:

- derivative = rate of change
- constants don't change
- derivative = slope
- slope = 0

$$\text{So } \frac{d}{dx}(5) = 0$$

The derivative of a straight line is...?

# The Derivatives of Simple Functions



The derivative of a constant is zero because:


- derivative = rate of change
- constants don't change
- derivative = slope
- slope = 0

$$\text{So } \frac{d}{dx}(5) = 0$$

The derivative of a straight line is its slope because

- derivative = slope

# The Derivatives of Simple Functions



The derivative of a constant is zero because:

- derivative = rate of change
- constants don't change
- derivative = slope
- slope = 0

$$\text{So } \frac{d}{dx}(5) = 0$$

The derivative of a straight line is its slope because

- derivative = slope


$$\text{So } \frac{d}{dx}(2x) = 2$$

# Meaning of Derivatives

$$\frac{d}{dx}(x^2) = 2x$$

What this means

The slope of the graph  
of  $y = x^2$  at  $x = a$  is  $2a$




# Meaning of Derivatives

$$\frac{d}{dx}(x^2) = 2x$$

What this means

The **slope** of the graph  
of  $y = x^2$  at  $x = a$  is  $2a$




# Meaning of Derivatives

$$\frac{d}{dx}(x^2) = 2x$$

What this means

The **slope** of the graph  
of  $y = x^2$  at  $x = a$  is  $2a$



at  $x = 0$ , slope is  $2(0) = 0$




# Meaning of Derivatives

$$\frac{d}{dx}(x^2) = 2x$$

What this means

The **slope** of the graph  
of  $y = x^2$  at  $x = a$  is  $2a$




at  $x = 2$ , slope is  $2(2) = 4$

# Meaning of Derivatives

$$\frac{d}{dx}(x^2) = 2x$$

What this means

The slope of the graph  
of  $y = x^2$  at  $x = a$  is  $2a$



at  $x = 2$ , slope is  $2(2) = 4$

derivative = rate of change = slope of graph = slope of tangent line

# General Rule:

$$\frac{d}{dx} (x^2) = 2x$$

$$\frac{d}{dx} (x^3) = 3x^2$$

$$\frac{d}{dx} (x^4) = 4x^3$$

# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

The **exponent** comes out front. Then **subtract** one from exponent.

# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

The **exponent** comes out front. Then **subtract** one from exponent.

**Examples:**

# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

The **exponent** comes out front. Then **subtract** one from exponent.

**Examples:**

**1.**  $\frac{d}{dx}(x^7) =$

$$A = 7x^7 \quad B = 6x^6 \quad C = 6x^7 \quad D = 7x^6 \quad E = 0$$

# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

The **exponent** comes out front. Then **subtract** one from exponent.

**Examples:**

**1.**  $\frac{d}{dx}(x^7) =$

A =  $7x^7$     B =  $6x^6$     C =  $6x^7$     D =  $7x^6$     E = 0    D

**2.**  $\frac{d}{dx}(x^{-3}) =$

A =  $3x^{-2}$     B =  $-3x^{-2}$     C =  $-2x^{-4}$     D =  $-3x^{-4}$



# General Rule:

$$\frac{d}{dx}(x^2) = 2x$$

$$\frac{d}{dx}(x^3) = 3x^2$$

$$\frac{d}{dx}(x^4) = 4x^3$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

The **exponent** comes out front. Then **subtract** one from exponent.

**Examples:**

**1.**  $\frac{d}{dx}(x^7) =$

A =  $7x^7$     B =  $6x^6$     C =  $6x^7$     D =  $7x^6$     E = 0    D

**2.**  $\frac{d}{dx}(x^{-3}) =$

A =  $3x^{-2}$     B =  $-3x^{-2}$     C =  $-2x^{-4}$     D =  $-3x^{-4}$     D

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

$$A = \frac{1}{2}x^{1/2} \quad B = -\frac{1}{2}x^{-1/2} \quad C = \frac{1}{2}x^{-1/2}$$

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

A =  $\frac{1}{2}x^{1/2}$     B =  $-\frac{1}{2}x^{-1/2}$     C =  $\frac{1}{2}x^{-1/2}$     C

# More Examples

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx} (x^{1/2}) =$

$$A = \frac{1}{2}x^{1/2} \quad B = -\frac{1}{2}x^{-1/2} \quad C = \frac{1}{2}x^{-1/2} \quad \boxed{C}$$

**Rule:** ALWAYS rewrite the thing you want derivative of as  $x^n$

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

$$A = \frac{1}{2}x^{1/2} \quad B = -\frac{1}{2}x^{-1/2} \quad C = \frac{1}{2}x^{-1/2} \quad \boxed{C}$$

**Rule:** ALWAYS rewrite the thing you want derivative of as  $x^n$

**4.**  $\frac{d}{dx}\left(\frac{1}{x^3}\right) =$

$$A = \frac{1}{3x^2} \quad B = -3x^{-2} \quad C = -3x^{-4}$$

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

$$A = \frac{1}{2}x^{1/2} \quad B = -\frac{1}{2}x^{-1/2} \quad C = \frac{1}{2}x^{-1/2} \quad \boxed{C}$$

**Rule:** ALWAYS rewrite the thing you want derivative of as  $x^n$

**4.**  $\frac{d}{dx}\left(\frac{1}{x^3}\right) =$

$$A = \frac{1}{3x^2} \quad B = -3x^{-2} \quad C = -3x^{-4} \quad \boxed{C}$$

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

$$A = \frac{1}{2}x^{1/2} \quad B = -\frac{1}{2}x^{-1/2} \quad C = \frac{1}{2}x^{-1/2} \quad \boxed{C}$$

**Rule:** ALWAYS rewrite the thing you want derivative of as  $x^n$

**4.**  $\frac{d}{dx}\left(\frac{1}{x^3}\right) =$

$$A = \frac{1}{3x^2} \quad B = -3x^{-2} \quad C = -3x^{-4} \quad \boxed{C}$$

**5.**  $\frac{d}{dx}(\sqrt{x}) =$

$$A = -\frac{1}{2}\sqrt{x} \quad B = \frac{1}{2}x^{-1/2} \quad C = -\frac{1}{2}x^{-1/2}$$

# More Examples

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**3.**  $\frac{d}{dx}(x^{1/2}) =$

A =  $\frac{1}{2}x^{1/2}$     B =  $-\frac{1}{2}x^{-1/2}$     C =  $\frac{1}{2}x^{-1/2}$     C

**Rule:** ALWAYS rewrite the thing you want derivative of as  $x^n$

**4.**  $\frac{d}{dx}\left(\frac{1}{x^3}\right) =$

A =  $\frac{1}{3x^2}$     B =  $-3x^{-2}$     C =  $-3x^{-4}$     C

**5.**  $\frac{d}{dx}(\sqrt{x}) =$

A =  $-\frac{1}{2}\sqrt{x}$     B =  $\frac{1}{2}x^{-1/2}$     C =  $-\frac{1}{2}x^{-1/2}$     B



# Polynomials

$$\frac{d}{dx} (4x^5 + 7x^2 - 5x + 7) = 4(5)x^4 + 7(2)x^1 - 5 + 0$$

# Polynomials

$$\frac{d}{dx} (4x^5 + 7x^2 - 5x + 7) = 4(5)x^4 + 7(2)x^1 - 5 + 0$$

Special cases

- $\frac{d}{dx} (-5x) = -5$

# Polynomials

$$\frac{d}{dx} (4x^5 + 7x^2 - 5x + 7) = 4(5)x^4 + 7(2)x^1 - 5 + 0$$

## Special cases

- $\frac{d}{dx} (-5x) = -5$
- $\frac{d}{dx} (7) = 0$

# Polynomials

$$\frac{d}{dx} (4x^5 + 7x^2 - 5x + 7) = 4(5)x^4 + 7(2)x^1 - 5 + 0$$

## Special cases

- $\frac{d}{dx} (-5x) = -5$
- $\frac{d}{dx} (7) = 0$

**6.**  $\frac{d}{dx} (3x^4 + 9x^3 + 7) = ?$

A= I have an answer

B= I am working on it

C= Help!

# Polynomials

$$\frac{d}{dx} (4x^5 + 7x^2 - 5x + 7) = 4(5)x^4 + 7(2)x^1 - 5 + 0$$

## Special cases

- $\frac{d}{dx} (-5x) = -5$
- $\frac{d}{dx} (7) = 0$

**6.**  $\frac{d}{dx} (3x^4 + 9x^3 + 7) = ?$

A= I have an answer

B= I am working on it

C= Help!

**Answer:**  $12x^3 + 27x^2$

A= I got it

B= I nearly got it

C= I want my mommy!

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.



# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.
- The **slope of the tangent line** to the graph at  $x = 2$  is **7**.

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.
- The **slope of the tangent line** to the graph at  $x = 2$  is **7**.
- The **instantaneous rate of change** of  $f(x)$  at  $x = 2$  is **7**.

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.
- The **slope of the tangent line** to the graph at  $x = 2$  is **7**.
- The **instantaneous rate of change** of  $f(x)$  at  $x = 2$  is **7**.
- At  $x = 2$  the output (value of  $f(x)$ ) changes **7** times as fast as the **input** (value of  $x$ ).

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.
- The **slope of the tangent line** to the graph at  $x = 2$  is **7**.
- The **instantaneous rate of change** of  $f(x)$  at  $x = 2$  is **7**.
- At  $x = 2$  the output (value of  $f(x)$ ) changes **7** times as fast as the **input** (value of  $x$ ).
- $\Delta f \approx 7\Delta x$  near  $x = 2$ .

# The Meanings of Derivatives

The derivative of  $f(x) = x^2 + 3x + 1$  is  $f'(x) = \frac{df}{dx} = 2x + 3$ . This means:

- This is the **slope** of the graph  $y = x^2 + 3x + 1$  at the point  $x$
- It is the **instantaneous rate of change** of  $f(x)$  at  $x$ .

That  $f'(2) = 7$  means:

- The **slope** of the graph  $y = f(x)$  at  $x = 2$  is **7**.
- The **slope of the tangent line** to the graph at  $x = 2$  is **7**.
- The **instantaneous rate of change** of  $f(x)$  at  $x = 2$  is **7**.
- At  $x = 2$  the output (value of  $f(x)$ ) changes **7** times as fast as the **input** (value of  $x$ ).
- $\Delta f \approx 7\Delta x$  near  $x = 2$ .
- $f(2 + \Delta x) \approx f(2) + 7\Delta x$ .

# Applications

**7.** What is the slope of the graph  $y = 3x^2 - 7x + 5$  at  $x = 1$ ?

A =  $-2$     B =  $-1$     C =  $0$     D =  $1$     E =  $2$

# Applications

**7.** What is the slope of the graph  $y = 3x^2 - 7x + 5$  at  $x = 1$ ?

A = -2    B = -1    C = 0    D = 1    E = 2    B

**8.** What is the instantaneous rate of change of  $f(x) = x^3 - 2x + 3$  at  $x = 1$ ?

A = -2    B = -1    C = 0    D = 1    E = 2

# Applications

7. What is the slope of the graph  $y = 3x^2 - 7x + 5$  at  $x = 1$ ?

A = -2   B = -1   C = 0   D = 1   E = 2   B

8. What is the instantaneous rate of change of  $f(x) = x^3 - 2x + 3$  at  $x = 1$ ?

A = -2   B = -1   C = 0   D = 1   E = 2   D

9. After  $t$  seconds a hamster on a skate board is  $4t^2 + 2t$  cm from the origin on the  $x$ -axis. What is the exact speed of the hamster (in cm/sec) after 2 seconds?

A = 10   B = 16   C = 18   D = 20   E = 14



# Applications

7. What is the slope of the graph  $y = 3x^2 - 7x + 5$  at  $x = 1$ ?

A = -2    B = -1    C = 0    D = 1    E = 2    B

8. What is the instantaneous rate of change of  $f(x) = x^3 - 2x + 3$  at  $x = 1$ ?

A = -2    B = -1    C = 0    D = 1    E = 2    D

9. After  $t$  seconds a hamster on a skate board is  $4t^2 + 2t$  cm from the origin on the  $x$ -axis. What is the exact speed of the hamster (in cm/sec) after 2 seconds?

A = 10    B = 16    C = 18    D = 20    E = 14    C

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + \Delta x$  then take limit as  $\Delta x \rightarrow 0$ .

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

$$\left( \begin{array}{c} \text{average rate} \\ \text{of change between} \\ x \text{ and } x + h \end{array} \right)$$

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

$$\left( \begin{array}{c} \text{average rate} \\ \text{of change between} \\ x \text{ and } x + h \end{array} \right) = \frac{(x + h)^3 - x^3}{(x + h) - x}$$

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

$$\begin{aligned} \left( \begin{array}{c} \text{average rate} \\ \text{of change between} \\ x \text{ and } x + h \end{array} \right) &= \frac{(x + h)^3 - x^3}{(x + h) - x} \\ &= 3x^2 + 3xh + h^2 \end{aligned}$$

# Why This Works (§8.9)

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

**Example:  $n = 3$ :** Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

$$\begin{aligned} \left( \begin{array}{c} \text{average rate} \\ \text{of change between} \\ x \text{ and } x + h \end{array} \right) &= \frac{(x + h)^3 - x^3}{(x + h) - x} \\ &= 3x^2 + 3xh + h^2 \end{aligned}$$

Limit as  $h \rightarrow 0$  is  $3x^2$



# Why This Works (§8.9)

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

**Example:**  $n = 3$ : Calculate the average rate of change of  $x^3$  between  $x$  and  $x + h$  then take limit as  $h \rightarrow 0$ .

$$\begin{aligned} \left( \begin{array}{c} \text{average rate} \\ \text{of change between} \\ x \text{ and } x + h \end{array} \right) &= \frac{(x + h)^3 - x^3}{(x + h) - x} \\ &= 3x^2 + 3xh + h^2 \end{aligned}$$

Limit as  $h \rightarrow 0$  is  $3x^2$

A similar calculation works for  $x^n$  for any  $n$ .

# More Applications

**10.** What is the equation of the tangent line at  $x = 1$  to the graph of  $y = x^3 - x + 4$ ? The tangent line is  $y = \dots$ ?

A =  $x + 3$     B =  $3x + 1$     C =  $2x - 2$     D =  $2x + 2$     E =  $6x - 2$

# More Applications

**10.** What is the equation of the tangent line at  $x = 1$  to the graph of  $y = x^3 - x + 4$ ? The tangent line is  $y = \dots$ ?

A =  $x + 3$     B =  $3x + 1$     C =  $2x - 2$     D =  $2x + 2$     E =  $6x - 2$

Answer: D


# More Applications

**10.** What is the equation of the tangent line at  $x = 1$  to the graph of  $y = x^3 - x + 4$ ? The tangent line is  $y = \dots$ ?

A =  $x + 3$     B =  $3x + 1$     C =  $2x - 2$     D =  $2x + 2$     E =  $6x - 2$

Answer: D

Here's a picture:



# Another Example

- 11.** The temperature in an oven after  $t$  minutes is  $50 + t^3$  °F. How quickly is the temperature rising after 2 minutes?

$$A = 58 \quad B = 3 \quad C = 12 \quad D = 50 \quad E = 8$$

# Another Example

- 11.** The temperature in an oven after  $t$  minutes is  $50 + t^3$  °F. How quickly is the temperature rising after 2 minutes?

$$A = 58 \quad B = 3 \quad C = 12 \quad D = 50 \quad E = 8$$

Answer: C

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:**  $5x^4 = \frac{d}{dx} (x^5) = \frac{d}{dx} (x^2 \cdot x^3) \neq (2x)(3x^2) = 6x^3$



# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:**  $5x^4 = \frac{d}{dx} (x^5) = \frac{d}{dx} (x^2 \cdot x^3) \neq (2x)(3x^2) = 6x^3$

**Example:** Find the derivative of  $(x + 1)(2x + 3)$

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:**  $5x^4 = \frac{d}{dx} (x^5) = \frac{d}{dx} (x^2 \cdot x^3) \neq (2x)(3x^2) = 6x^3$

**Example:** Find the derivative of  $(x+1)(2x+3)$

**12.**  $\frac{d}{dx} ((x^2+1)(x^3+1)) = ?$

A =  $6x^3$     B =  $5x^4 + 3x^2 + 2x$     C =  $x^5 + x^3 + x^2 + 1$     D = Other

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:**  $5x^4 = \frac{d}{dx} (x^5) = \frac{d}{dx} (x^2 \cdot x^3) \neq (2x)(3x^2) = 6x^3$

**Example:** Find the derivative of  $(x+1)(2x+3)$

**12.**  $\frac{d}{dx} ((x^2+1)(x^3+1)) = ?$

A =  $6x^3$     B =  $5x^4 + 3x^2 + 2x$     C =  $x^5 + x^3 + x^2 + 1$     D = Other

**Answer:** B

# Once upon a time...

There was a happy math professor and he told his happy students:

# Once upon a time...

There was a happy math professor and he told his happy students:

“When you work out **derivatives** **ALWAYS** write the  $\frac{d}{dx}$  part so you write something like

$$\frac{d}{dx} (3x^2 + 5x + 2) = 6x + 5$$

and you never-ever-ever write

$$3x^2 + 5x + 2 \quad 6x + 5 \quad \text{or even worse}$$

$$3x^2 + 5x + 2 = 6x + 5.$$

Because if you don't do as I say I will become a sad math professor and you will repeat this class.”