

Office Hours!

Instructor:

Peter M. Garfield, garfield@math.ucsb.edu
South Hall 6510

Office Hours:

Monday: TBA
Tuesday: TBA

Final Exam:

Wednesday: 4:00–7:00PM

© 2017 Daryl Cooper, Peter M. Garfield

HW 23 Problem #4

A commuter railway has 800 passengers per day and charges each one two dollars per day. For each 4 cents that the fare is increased, 5 fewer people will go by train. What is the greatest profit that can be earned?

HW 23 Problem #4

A commuter railway has 800 passengers per day and charges each one two dollars per day. For each 4 cents that the fare is increased, 5 fewer people will go by train. What is the greatest profit that can be earned?

“Profit” in this case must mean “Revenue”

W '14 Problem #9

Carol's chocolate cookies cost \$2 each and she sells 2200 at this price. For each cent she raises the price she sells 5 fewer cookies. The ingredients for 10 cookies cost \$2.

If Carol increases the price of a cookie by x cents:

- (a) How many cookies will she sell?
- (b) How many dollars profit does she make on each cookie?
- (c) Express the total profit (in \$) in terms of x .
- (d) What should x be to make the most profit?
- (e) What should the price in \$ of one cookie be to make the most profit?

W '15 Problem #8

Let $f(x) = 20\sqrt{x}$.

(a) Find $f'(4)$.

[Simplify your answer to something like $7/3$.]

(b) Find the tangent line approximation to $y = f(x)$ at $x = 4$.

(c) Use this to approximate the value of $20\sqrt{5}$.

W '15 Problem #3(c)

Compute $\frac{d}{dx}((3x^2 + 5)/x^k)$.

[Here k is a constant.]

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 C

(b) What is the instantaneous rate of change of $f(x)$ at $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 C

(b) What is the instantaneous rate of change of $f(x)$ at $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 E

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 C

(b) What is the instantaneous rate of change of $f(x)$ at $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 E

(2) The table to the right shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

(a) On average, how many people were treated per day during the first week?

A = 56 B = 38 C = 81 D = 8

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 C

(b) What is the instantaneous rate of change of $f(x)$ at $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 E

(2) The table to the right shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

(a) On average, how many people were treated per day during the first week?

A = 56 B = 38 C = 81 D = 8 D

(b) Which period had the greatest average number of cases per day?

A = 0 – 3 B = 3 – 7 C = 7 – 9

Review: Rates of Change

(1) Suppose $f(x) = x^2 - x$.

(a) What is the average rate of change of $f(x)$ between $x = 1$ and $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 C

(b) What is the instantaneous rate of change of $f(x)$ at $x = 3$?

A = 1 B = 2 C = 3 D = 4 E = 5 E

(2) The table to the right shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

(a) On average, how many people were treated per day during the first week?

A = 56 B = 38 C = 81 D = 8 D

(b) Which period had the greatest average number of cases per day?

A = 0 – 3 B = 3 – 7 C = 7 – 9 C

Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.

(1) Which of the following equations is true?

A $J + M = 270$ B $2J = 4M$ C $J - M = 270$

D $2J + 4M = 270$ E $2J = 270 + 4M$

Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.

(1) Which of the following equations is true?

A $J + M = 270$ B $2J = 4M$ C $J - M = 270$

D $2J + 4M = 270$ E $2J = 270 + 4M$ D

Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.

(1) Which of the following equations is true?

A $J + M = 270$ B $2J = 4M$ C $J - M = 270$

D $2J + 4M = 270$ E $2J = 270 + 4M$ D

- At 3pm, they are 100 miles apart.

(2) Which of the following equations is true?

A $J + M = 100$ B $2J = 4M$ C $J - M = 100$

D $2J + 4M = 100$ E $2J = 100 + 4M$

Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.

(1) Which of the following equations is true?

A $J + M = 270$ B $2J = 4M$ C $J - M = 270$

D $2J + 4M = 270$ E $2J = 270 + 4M$ D

- At 3pm, they are 100 miles apart.

(2) Which of the following equations is true?

A $J + M = 100$ B $2J = 4M$ C $J - M = 100$

D $2J + 4M = 100$ E $2J = 100 + 4M$ A

Jason & Marie (continued)

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.
- At 3pm, they are 100 miles apart.

(3) What was Jason's speed ?

$$A = 35 \quad B = 45 \quad C = 55 \quad D = 65 \quad E = 75$$

Jason & Marie (continued)

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed M starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed J .
- They meet at 4pm.
- At 3pm, they are 100 miles apart.

(3) What was Jason's speed ?

$A = 35$


$B = 45$

$C = 55$

$D = 65$


$E = 75$

D



Use the graph of $y = 10^x$ to find:

- (A) $10^{3.65}$
- (B) Solve $10^x = 73$
- (C) The slope of the graph at $x = 0.65$
- (D) The average rate of change of 10^x between $x = 0.1$ and $x = 0.6$



Use the graph of $y = 10^x$ to find:


(A) $10^{3.65}$

Answer: 4500

(B) Solve $10^x = 73$

(C) The slope of the graph at $x = 0.65$

(D) The average rate of change of 10^x between $x = 0.1$ and $x = 0.6$



Use the graph of $y = 10^x$ to find:

(A) $10^{3.65}$


Answer: 4500

(B) Solve $10^x = 73$

Answer: 1.86

(C) The slope of the graph at $x = 0.65$

(D) The average rate of change of 10^x between $x = 0.1$ and $x = 0.6$



Use the graph of $y = 10^x$ to find:

(A) $10^{3.65}$

Answer: 4500


(B) Solve $10^x = 73$

Answer: 1.86

(C) The slope of the graph at $x = 0.65$

Answer: 10

(D) The average rate of change of 10^x between $x = 0.1$ and $x = 0.6$



Use the graph of $y = 10^x$ to find:

(A) $10^{3.65}$

Answer: 4500

(B) Solve $10^x = 73$

Answer: 1.86

(C) The slope of the graph at $x = 0.65$

Answer: 10

(D) The average rate of change of 10^x between $x = 0.1$ and $x = 0.6$

Answer: 5

Review: Lines!

- 1.** Find the equation of the line with slope 3 that contains the point $(2, 5)$.

A $y = 3x + 5$ B $y = 3x - 1$ C $y = 3x + 2$

Review: Lines!

- 1.** Find the equation of the line with slope 3 that contains the point $(2, 5)$.

A $y = 3x + 5$ B $y = 3x - 1$ C $y = 3x + 2$ B

- 2.** What is the x -coordinate of the point where the two lines

$$y = 3x + 2 \quad \text{and} \quad y - 4x + 1 = 0$$

cross?

A $x = -3$ B $x = -1$ C $x = 1$ D $x = 3$ E $x = 4$

Review: Lines!

- 1.** Find the equation of the line with slope 3 that contains the point $(2, 5)$.

A $y = 3x + 5$ B $y = 3x - 1$ C $y = 3x + 2$ B

- 2.** What is the x -coordinate of the point where the two lines

$$y = 3x + 2 \quad \text{and} \quad y - 4x + 1 = 0$$

cross?

A $x = -3$ B $x = -1$ C $x = 1$ D $x = 3$ E $x = 4$ D

Review: Logs!

3. Solve $3^x = 7$.

A $x = 7/3$ B $x = \log(7/3)$

C $x = \log(7)/\log(3)$ D $x = \log(7) - \log(3)$

Review: Logs!

3. Solve $3^x = 7$.

A $x = 7/3$ B $x = \log(7/3)$

C $x = \log(7)/\log(3)$ D $x = \log(7) - \log(3)$ C

Remember half-life:

• Half-life = K years

• Initial amount = A

• Amount after t years is $= A \times 2^{-t/K}$

4. Let's start with 8 grams of an element with half-life of 5 years.

(a) How many grams remain after 10 years?

A = 0 B = 2 C = 4 D = 8

Review: Logs!

3. Solve $3^x = 7$.

A $x = 7/3$ B $x = \log(7/3)$

C $x = \log(7)/\log(3)$ D $x = \log(7) - \log(3)$ C

Remember half-life:

• Half-life = K years

• Initial amount = A

• Amount after t years is $= A \times 2^{-t/K}$

4. Let's start with 8 grams of an element with half-life of 5 years.

(a) How many grams remain after 10 years?

A = 0 B = 2 C = 4 D = 8 B

(b) How many years until 3 grams remain?

A = $8/3$ B = $-5 \log(3/8)/\log(2)$

C = $-5 \log(3/16)$ D = $\log(3/8) - \log(2)$

Review: Logs!

3. Solve $3^x = 7$.

A $x = 7/3$ B $x = \log(7/3)$

C $x = \log(7)/\log(3)$ D $x = \log(7) - \log(3)$ C

Remember half-life:

• Half-life = K years

• Initial amount = A

• Amount after t years is $= A \times 2^{-t/K}$

4. Let's start with 8 grams of an element with half-life of 5 years.

(a) How many grams remain after 10 years?

A = 0 B = 2 C = 4 D = 8 B

(b) How many years until 3 grams remain?

A = $8/3$ B = $-5 \log(3/8)/\log(2)$

C = $-5 \log(3/16)$ D = $\log(3/8) - \log(2)$ B