



# Office Hours:

**Instructor:**

Peter M. Garfield  
garfield@math.ucsb.edu  
South Hall 6510

To Be Announced  
For Exam Week

**TAs:**

Christine Alar  
christine@math.ucsb.edu

Justin Rogers  
justin\_rogers@math.ucsb.edu

Abe Schulte  
aschulte@math.ucsb.edu

© 2019 Daryl Cooper, Peter M. Garfield  
Please do not distribute outside of this course.

# Review: Rates of Change

**1.** Suppose  $f(x) = x^2 - x$ .

(a) What is the average rate of change of  $f(x)$  between  $x = 1$  and  $x = 3$ ?

(A) 1

(B) 2

(C) 3

(D) 4

(E) 5

# Review: Rates of Change

**1.** Suppose  $f(x) = x^2 - x$ .

(a) What is the average rate of change of  $f(x)$  between  $x = 1$  and  $x = 3$ ?

(A) 1

(B) 2

(C) 3

(D) 4

(E) 5

Answer: **C**

(b) What is the instantaneous rate of change of  $f(x)$  at  $x = 3$ ?

(A) 1

(B) 2

(C) 3

(D) 4

(E) 5

# Review: Rates of Change

1. Suppose  $f(x) = x^2 - x$ .

(a) What is the average rate of change of  $f(x)$  between  $x = 1$  and  $x = 3$ ?

(A) 1

(B) 2

(C) 3

(D) 4

(E) 5

Answer: C

(b) What is the instantaneous rate of change of  $f(x)$  at  $x = 3$ ?

(A) 1

(B) 2

(C) 3

(D) 4

(E) 5

Answer: E

# Review: Rates of Change (cont'd)

- 2.** The table below shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

- (a) On average, how many people were treated per day during the first week?

(A) 56

(B) 38

(C) 81

(D) 8

# Review: Rates of Change (cont'd)

2. The table below shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

- (a) On average, how many people were treated per day during the first week?

(A) 56

(B) 38

(C) 81

(D) 8

D

- (b) Which period had the greatest average number of cases per day?

(A) 0 – 3

(B) 3 – 7

(C) 7 – 9

# Review: Rates of Change (cont'd)

2. The table below shows the number total number of people treated in a hospital up to and including the day shown during a flu outbreak.

days	0	3	7	9
cases	0	18	56	81

- (a) On average, how many people were treated per day during the first week?

(A) 56

(B) 38

(C) 81

(D) 8

D

- (b) Which period had the greatest average number of cases per day?

(A) 0 – 3

(B) 3 – 7

(C) 7 – 9

C

# Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.

**3.** Which of the following equations is true?

(A)  $J + M = 270$

(B)  $2J = 4M$

(C)  $J - M = 270$

(D)  $2J + 4M = 270$

(E)  $2J = 270 + 4M$



# Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.

**3.** Which of the following equations is true?

(A)  $J + M = 270$

(B)  $2J = 4M$

(C)  $J - M = 270$

(D)  $2J + 4M = 270$

(E)  $2J = 270 + 4M$

**D**

# Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.

**3.** Which of the following equations is true?

(A)  $J + M = 270$

(B)  $2J = 4M$

(C)  $J - M = 270$

(D)  $2J + 4M = 270$

(E)  $2J = 270 + 4M$

**D**

- At 3pm, they are 100 miles apart.

# Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.

**3.** Which of the following equations is true?

(A)  $J + M = 270$

(B)  $2J = 4M$

(C)  $J - M = 270$

(D)  $2J + 4M = 270$

(E)  $2J = 270 + 4M$

**D**

- At 3pm, they are 100 miles apart.

**4.** Which of the following equations is true?

(A)  $J + M = 100$

(B)  $2J = 4M$

(C)  $J - M = 100$

(D)  $2J + 4M = 100$

(E)  $2J = 100 + 4M$

# Jason & Marie

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.

**3.** Which of the following equations is true?

(A)  $J + M = 270$

(B)  $2J = 4M$

(C)  $J - M = 270$

(D)  $2J + 4M = 270$

(E)  $2J = 270 + 4M$

**A**

- At 3pm, they are 100 miles apart.

**4.** Which of the following equations is true?

(A)  $J + M = 100$

(B)  $2J = 4M$

(C)  $J - M = 100$

(D)  $2J + 4M = 100$

(E)  $2J = 100 + 4M$

**A**

# Jason & Marie (continued)

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.
- At 3pm, they are 100 miles apart.

**5.** What was Jason's speed?

(A) 35

(B) 45

(C) 55

(D) 65

(E) 75

# Jason & Marie (continued)

- Jason Bourne and Marie Kreutz are 270 miles apart at noon.
- Marie drives towards Jason at constant speed  $M$  starting at noon.
- Jason sets out at 2pm driving towards Marie at constant speed  $J$ .
- They meet at 4pm.
- At 3pm, they are 100 miles apart.

**5.** What was Jason's speed?

(A) 35


(B) 45

(C) 55

(D) 65

(E) 75

**D**




**6.** Use the graph of  $y = 10^x$  to find:

(A)  $10^{3.65}$

(B) Solve  $10^x = 73$

(C) The slope of the graph at  $x = 0.65$

(D) The average rate of change of  $10^x$  between  $x = 0.1$  and  $x = 0.6$



**6.** Use the graph of  $y = 10^x$  to find:

(A)  $10^{3.65}$


**Answer:** 4500

(B) Solve  $10^x = 73$

(C) The slope of the graph at  $x = 0.65$

(D) The average rate of change of  $10^x$  between  $x = 0.1$  and  $x = 0.6$





**6.** Use the graph of  $y = 10^x$  to find:

(A)  $10^{3.65}$


**Answer:** 4500

(B) Solve  $10^x = 73$

**Answer:** 1.86

(C) The slope of the graph at  $x = 0.65$

(D) The average rate of change of  $10^x$  between  $x = 0.1$  and  $x = 0.6$



**6.** Use the graph of  $y = 10^x$  to find:

(A)  $10^{3.65}$

**Answer:** 4500


(B) Solve  $10^x = 73$

**Answer:** 1.86

(C) The slope of the graph at  $x = 0.65$

**Answer:** 10 (10.285)

(D) The average rate of change of  $10^x$  between  $x = 0.1$  and  $x = 0.6$



**6.** Use the graph of  $y = 10^x$  to find:

(A)  $10^{3.65}$

**Answer:** 4500

(B) Solve  $10^x = 73$

**Answer:** 1.86

(C) The slope of the graph at  $x = 0.65$

**Answer:** 10 (10.285)

(D) The average rate of change of  $10^x$  between  $x = 0.1$  and  $x = 0.6$

**Answer:** 5 (5.444)

# HW 23 Problem #4

A commuter railway has 800 passengers per day and charges each one two dollars per day. For each 4 cents that the fare is increased, 5 fewer people will go by train. What is the greatest profit that can be earned?

# HW 23 Problem #4

A commuter railway has 800 passengers per day and charges each one two dollars per day. For each 4 cents that the fare is increased, 5 fewer people will go by train. What is the greatest profit that can be earned?

“Profit” in this case must mean “Revenue”

# Old Final #4 Problem #10

Bob's House of Index Cards sells note cards for exams. If the price is 5 cents, he sells 350 cards. For each cent he increases the price, he sells 10 fewer cards.


- (a) If he sells each card for  $(5 + x)$  cents, how many are sold?
- (b) What is the total amount of money Bob gets by selling cards for  $5 + x$  cents each. *Simplify your answer.*
- (c) What price for the card gives Bob the most money?
- (d) How many cards does he sell to make the most money?

# Old Final #4 Problem #8

Let  $f(x) = 2e^{3x} - 5x$ .

- (a) Find  $f'(0)$ .
- (b) Find the tangent line approximation to  $y = f(x)$  at  $x = 0$ .
- (c) Use this to approximate the value of  $f(-0.1)$ .


# Old Final #4 Problem #7



(a) For which value(s) of  $x$  is  $f'(x) = 0$ ?




# Old Final #4 Problem #7




(b) For which values of  $x$  is  $f''(x) < 0$ ?

# Old Final #4 Problem #7




(c) Find a single value of  $x$  with  $f'(x) = -2$ .

# Old Final #4 Problem #7




(c) Find a single value of  $x$  with  $f'(x) = -2$ .

# Old Final #4 Problem #7




(c) Find a single value of  $x$  with  $f'(x) = -2$ .

# Old Final #4 Problem #7



(c) Find a single value of  $x$  with  $f'(x) = -2$ .

# Old Final #4 Problem #7



(c) Find a single value of  $x$  with  $f'(x) = -2$ .

# Old Final #4 Problem #3(c)

Compute  $\frac{d}{dx} \left( (2x^k + 5)/x^2 \right)$ .

[Here  $k$  is a constant.]