



# Office Hours!

Instructor:

Peter M. Garfield, [garfield@math.ucsb.edu](mailto:garfield@math.ucsb.edu)

Office Hours:

Mondays 2–3PM

Tuesdays 10:30–11:30AM

Thursdays 1–2PM

or by appointment

Office:

South Hall 6510

© 2017 Daryl Cooper, Peter M. Garfield

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate *puppy* =  $17^{3.1}$

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate  $puppy = 17^{3.1}$

(i) **doggy** = **log**(*puppy*)

(ii) rules of **logs** to expand **doggy**

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate  $puppy = 17^{3.1}$

(i) **doggy** = **log**(*puppy*)

(ii) rules of **logs** to expand **doggy**

(iii) look up **logs** of individual terms in **doggy**. Move decimal point trick.

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate  $puppy = 17^{3.1}$

(i) **doggy** = **log**(*puppy*)

(ii) rules of **logs** to expand **doggy**

(iii) look up **logs** of individual terms in **doggy**. Move decimal point trick.

(iv) Now have numerical value for **doggy**.

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate  $puppy = 17^{3.1}$

(i)  $doggy = \log(puppy)$

(ii) rules of **logs** to expand **doggy**

(iii) look up **logs** of individual terms in **doggy**. Move decimal point trick.

(iv) Now have numerical value for **doggy**.

(v) so  $puppy = \text{antilog}(doggy)$  is the answer.

# Summary of calculations with logs

[Courtesy of Daryl Cooper]

Calculate the **log** of the thing you want then take **antilog** of the result.

Example: To calculate  $puppy = 17^{3.1}$

(i) **doggy** = **log**(*puppy*)

(ii) rules of **logs** to expand **doggy**

(iii) look up **logs** of individual terms in **doggy**. Move decimal point trick.

(iv) Now have numerical value for **doggy**.

(v) so  $puppy = \text{antilog}(\text{doggy})$  is the answer.

Make sure you never jot down a number on its own. It should always be part of an equation like  $\log(945 \times 32) \approx 4.48$  This way one can read and understand what is written. Otherwise you get **gibberish**

**Write math the way I do.** With **words** and **equations**. One should be able to **read and understand** what is on the paper **without being telepathic**. Imagine it is a **report** for your employer. In reality you are **explaining it to yourself**.

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10)$$



## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

Why didn't I use antilog?

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

Why didn't I use antilog?

Answer: I found  $x$ , so why would I?

I have **written** equations so I can **see** what each thing I write **means**. I can **see** that I've found  $x$  and so don't need to take antilog.

## §7.7: Solving Exponential Eq'ns

1. Find  $x$  by solving  $10^x = 5$ .

$$\begin{array}{llll} A = 5 & B = 0.5 & C = \log(5) & D = \log(0.5) \\ & E = \log(5) - \log(10) & \boxed{C} & \end{array}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

Why didn't I use antilog?

Answer: I found  $x$ , so why would I?

I have **written** equations so I can **see** what each thing I write **means**. I can **see** that I've found  $x$  and so don't need to take antilog.

How do I know when to take antilog?

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

Why didn't I use antilog?

Answer: I found  $x$ , so why would I?

I have **written** equations so I can **see** what each thing I write **means**. I can **see** that I've found  $x$  and so don't need to take antilog.

How do I know when to take antilog?

How do **you**

know?

## §7.7: Solving Exponential Eq'ns

**1.** Find  $x$  by solving  $10^x = 5$ .

$$A = 5 \quad B = 0.5 \quad C = \log(5) \quad D = \log(0.5)$$

$$E = \log(5) - \log(10) \quad \boxed{C}$$

Look how I write the answer!

$$\log(10^x) = \log(5) \quad \text{Take logs of both sides}$$

$$x = \log(10^x) = \log(5) \quad \text{Using } \log(a^p) = p \log(a) \text{ and } \log(10) = 1$$

Why didn't I use antilog?

Answer: I found  $x$ , so why would I?

I have **written** equations so I can **see** what each thing I write **means**. I can **see** that I've found  $x$  and so don't need to take antilog.

How do I know when to take antilog? How do **you** know? My answer: If you write the problem the way I do, so it makes sense, you can **see** what to do.

# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

**2.** Solve  $3^x = 7$

$$A = \log(7/3) \quad B = \log(7) - \log(3) \quad C = \log(7) + \log(3)$$

$$D = \log(3)/\log(7) \quad E = \log(7)/\log(3)$$



# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

**2.** Solve  $3^x = 7$

$$A = \log(7/3) \quad B = \log(7) - \log(3) \quad C = \log(7) + \log(3)$$

$$D = \log(3)/\log(7) \quad E = \log(7)/\log(3) \quad \boxed{E}$$

Look how I write the answer:

$$\log(3^x) = \log(7)$$

Take logs of both sides

$$x \log(3) = \log(3^x) = \log(7)$$

Using  $\log(a^p) = p \log(a)$

$$\text{So:} \quad x = \log(7)/\log(3)$$

# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

**3.** Solve  $7^{x+2} = 30$ .

$$A = \frac{\log(30) - 2\log(7)}{\log(7)} \quad B = \frac{\log(30)}{\log(7)} - 2 \quad C = \frac{\log(30) - \log(49)}{\log(7)}$$

$$D = \frac{\log(30/49)}{\log(7)} \quad E \approx -0.25213$$

# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

**3.** Solve  $7^{x+2} = 30$ .

$$A = \frac{\log(30) - 2\log(7)}{\log(7)} \quad B = \frac{\log(30)}{\log(7)} - 2 \quad C = \frac{\log(30) - \log(49)}{\log(7)}$$

$$D = \frac{\log(30/49)}{\log(7)} \quad E \approx -0.25213$$

All are correct!

# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

**4.** Solve  $7 \times 3^y = 2^{4y+3}$

$$A = \frac{3 \log(2) - \log(7)}{\log(3) - 4 \log(2)} \quad B = \frac{3 \log(2)}{7 \log(3)} \quad C = \frac{3 \log(2)}{7 \log(3) - 4 \log(2)}$$

$$D = \frac{7 \log(3) - 4 \log(2)}{3 \log(2)}$$

E = none of the above

# Examples:

Use the Fourth Law:

$$\log(a^x) = x \log(a)$$

Slogan: Logs bring exponents down to ground level.

4. Solve  $7 \times 3^y = 2^{4y+3}$

$$A = \frac{3 \log(2) - \log(7)}{\log(3) - 4 \log(2)} \quad B = \frac{3 \log(2)}{7 \log(3)} \quad C = \frac{3 \log(2)}{7 \log(3) - 4 \log(2)}$$

$$D = \frac{7 \log(3) - 4 \log(2)}{3 \log(2)}$$

E = none of the above

A

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

After 1 year:

$$\$10,000 \times 1.07 = \$10,700$$



# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

After 1 year:  $\$10,000 \times 1.07 = \$10,700$

After 2 years:  $\$10,700 \times 1.07 = \$10,000 \times 1.07 \times 1.07 = \$11,449$

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

After 1 year:  $\$10,000 \times 1.07 = \$10,700$

After 2 years:  $\$10,700 \times 1.07 = \$10,000 \times 1.07 \times 1.07 = \$11,449$

After 3 years:  $\$11,449 \times 1.07 = \$10,000 \times (1.07)^3 = \$12,250.40$

Each year **what you had before** is **multiplied** by 1.07. Thus **compound interest**.

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

After 1 year:  $\$10,000 \times 1.07 = \$10,700$

After 2 years:  $\$10,700 \times 1.07 = \$10,000 \times 1.07 \times 1.07 = \$11,449$

After 3 years:  $\$11,449 \times 1.07 = \$10,000 \times (1.07)^3 = \$12,250.40$

Each year **what you had before** is **multiplied** by 1.07. Thus **compound** interest.

So after 10 years have

$$\$10,000 \times \underbrace{1.07 \times 1.07 \times \cdots \times 1.07}_{10 \text{ times}} = 10,000 \times (1.07)^{10} \approx \boxed{\$20,000}$$

# Compound Interest

At end of each year a bank pays 7% interest into your account.  
Initially have \$10,000 in account. How much after 10 years?

**Think**  $10 \times 7\% = 70\%$  in 10 years, so have **\$17,000** but that is **wrong**.

After 1 year:  $\$10,000 \times 1.07 = \$10,700$

After 2 years:  $\$10,700 \times 1.07 = \$10,000 \times 1.07 \times 1.07 = \$11,449$

After 3 years:  $\$11,449 \times 1.07 = \$10,000 \times (1.07)^3 = \$12,250.40$

Each year **what you had before** is **multiplied** by 1.07. Thus **compound** interest.

So after 10 years have

$$\$10,000 \times \underbrace{1.07 \times 1.07 \times \cdots \times 1.07}_{10 \text{ times}} = 10,000 \times (1.07)^{10} \approx \boxed{\$20,000}$$

**Conclusion:** Money approximately doubles in 10 years!

So in 20 years multiplies by 4, in 30 years by 8,...

# General Compound Interest

If the interest rate is  $r\%$ , then each year money multiplies by

$$m = 1 + \frac{r}{100}.$$

If you start with an initial amount  $A$  of money then after  $t$  years you have

$$A \times m^t = A \times \left(1 + \frac{r}{100}\right)^t$$

- 5.** If you invest \$1000 at 14% interest, how much will you have 5 years later? (Guess!)

$A \approx \$700$      $B \approx \$1400$      $C \approx \$1500$      $D \approx \$1700$      $E \approx \$2000$

# General Compound Interest

If the interest rate is  $r\%$ , then each year money multiplies by

$$m = 1 + \frac{r}{100}.$$

If you start with an initial amount  $A$  of money then after  $t$  years you have

$$A \times m^t = A \times \left(1 + \frac{r}{100}\right)^t$$

- 5.** If you invest \$1000 at 14% interest, how much will you have 5 years later? (Guess!)

$$A \approx \$700 \quad B \approx \$1400 \quad C \approx \$1500 \quad D \approx \$1700 \quad E \approx \$2000 \quad \boxed{E}$$

After 5 years, you have

$$\$1,000 \times \left(1 + \frac{14}{100}\right)^5 = \$1,000 \times (1.14)^5.$$

How much is this?

# General Compound Interest

If the interest rate is  $r\%$ , then each year money multiplies by

$$m = 1 + \frac{r}{100}.$$

If you start with an initial amount  $A$  of money then after  $t$  years you have

$$A \times m^t = A \times \left(1 + \frac{r}{100}\right)^t$$

- 5.** If you invest \$1000 at 14% interest, how much will you have 5 years later? (Guess!)

$A \approx \$700$     $B \approx \$1400$     $C \approx \$1500$     $D \approx \$1700$     $E \approx \$2000$    E

After 5 years, you have

$$\$1,000 \times \left(1 + \frac{14}{100}\right)^5 = \$1,000 \times (1.14)^5.$$

How much is this? **Smart way: 14% in 1 year  $\approx$  7% per year for 2.**

# General Compound Interest

If the interest rate is  $r\%$ , then each year money multiplies by

$$m = 1 + \frac{r}{100}.$$

If you start with an initial amount  $A$  of money then after  $t$  years you have

$$A \times m^t = A \times \left(1 + \frac{r}{100}\right)^t$$

- 6.** If you invest \$1,000 at 14% interest, how many years until you have \$7,000?

$$A = \log(7/1.14) \quad B = \log(7)/\log(1.14) \quad C = \log(1.14)/\log(7)$$

$$D = \log(7) - \log(1.14) \quad E = \text{other}$$



# General Compound Interest

If the interest rate is  $r\%$ , then each year money multiplies by

$$m = 1 + \frac{r}{100}.$$

If you start with an initial amount  $A$  of money then after  $t$  years you have

$$A \times m^t = A \times \left(1 + \frac{r}{100}\right)^t$$

- 6.** If you invest \$1,000 at 14% interest, how many years until you have \$7,000?

$$A = \log(7/1.14) \quad B = \log(7)/\log(1.14) \quad C = \log(1.14)/\log(7)$$

$$D = \log(7) - \log(1.14) \quad E = \text{other} \quad \boxed{B}$$

## §7.9: Population Growth

Assume each generation of bunnies has 3 times as many bunnies as previous one. Initially have 100 bunnies. How many bunnies after  $n$  generations?

$$\begin{array}{lll} A = 100 \times 3n & B = 100 + 3n & C = 100(1 + 3n) \\ D = 100^{3n} & E = 100 \times 3^n & \end{array}$$

Answer: E

Start with 100

After 1 generation have  $100 \times 3$  bunnies

## §7.9: Population Growth

Assume each generation of bunnies has 3 times as many bunnies as previous one. Initially have 100 bunnies. How many bunnies after  $n$  generations?

$$\begin{array}{lll} A = 100 \times 3n & B = 100 + 3n & C = 100(1 + 3n) \\ D = 100^{3n} & E = 100 \times 3^n & \end{array}$$

Answer: E

Start with 100

After 1 generation have  $100 \times 3$  bunnies

After 2 generations have  $100 \times 3 \times 3$  bunnies

## §7.9: Population Growth

Assume each generation of bunnies has 3 times as many bunnies as previous one. Initially have 100 bunnies. How many bunnies after  $n$  generations?

$$\begin{array}{lll} A = 100 \times 3n & B = 100 + 3n & C = 100(1 + 3n) \\ D = 100^{3n} & E = 100 \times 3^n & \end{array}$$

Answer: E

Start with 100

After 1 generation have  $100 \times 3$  bunnies

After 2 generations have  $100 \times 3 \times 3$  bunnies

After 3 generations have  $100 \times 3 \times 3 \times 3$  bunnies

## §7.9: Population Growth

Assume each generation of bunnies has 3 times as many bunnies as previous one. Initially have 100 bunnies. How many bunnies after  $n$  generations?

$$\begin{array}{lll} A = 100 \times 3n & B = 100 + 3n & C = 100(1 + 3n) \\ D = 100^{3n} & E = 100 \times 3^n & \end{array}$$

Answer: E

Start with 100

After 1 generation have  $100 \times 3$  bunnies

After 2 generations have  $100 \times 3 \times 3$  bunnies

After 3 generations have  $100 \times 3 \times 3 \times 3$  bunnies

So...after  $n$  generations have

$$100 \times \underbrace{3 \times 3 \times \cdots \times 3}_{n \text{ times}} = 100 \times 3^n \text{ bunnies.}$$

# More Bunnies

We saw that:

- if we start with 100 bunnies, and
- the bunny population triples every generation,

then we have  $100 \times 3^n$  bunnies after  $n$  generations.

**7.** How many generations until there are  $10^7 = 10,000,000$  bunnies?

$$\begin{array}{lll} A = \log(5/3) & B = 5 - \log(3) & C = 5 / \log(3) \\ D = 5/3 & E = 10^5/3 & \end{array}$$

# More Bunnies

We saw that:

- if we start with 100 bunnies, and
- the bunny population triples every generation,

then we have  $100 \times 3^n$  bunnies after  $n$  generations.

**7.** How many generations until there are  $10^7 = 10,000,000$  bunnies?

$$\begin{array}{lll} A = \log(5/3) & B = 5 - \log(3) & C = 5 / \log(3) \\ D = 5/3 & E = 10^5/3 & \end{array}$$

$$\begin{array}{lll} A \approx 0.22 & B \approx 4.52 & C \approx 10.48 \\ D \approx 1.67 & E \approx 3,333 & \end{array}$$

# More Bunnies

We saw that:

- if we start with 100 bunnies, and
- the bunny population triples every generation,

then we have  $100 \times 3^n$  bunnies after  $n$  generations.

**7.** How many generations until there are  $10^7 = 10,000,000$  bunnies?

$$\begin{array}{lll} A = \log(5/3) & B = 5 - \log(3) & C = 5 / \log(3) \\ D = 5/3 & E = 10^5/3 & \end{array}$$

$$\begin{array}{lll} A \approx 0.22 & B \approx 4.52 & C \approx 10.48 \\ D \approx 1.67 & E \approx 3,333 & \boxed{C} \end{array}$$



# Flu Outbreak

At the start of an outbreak of H1N1 flu in a large herd of cattle, there were 5 infected individuals. The numbers doubles every 3 days. How many days until there are 80 infected cows?

$$A = \log(16)/\log(2) \quad B = \log(16/2) \quad C = 16/\log(2)$$

$$D = 3\log(16)/\log(2) \quad E = \log(48/2)$$

# Flu Outbreak

At the start of an outbreak of H1N1 flu in a large herd of cattle, there were 5 infected individuals. The numbers doubles every 3 days. How many days until there are 80 infected cows?

$$A = \log(16)/\log(2) \quad B = \log(16/2) \quad C = 16/\log(2)$$

$$D = 3 \log(16)/\log(2) \quad E = \log(48/2)$$

At the start of an outbreak of H1N1 flu in a large class of students, there were 5 infected individuals. The numbers doubles every 3 days. How many days until there are 80 infected students?

$$A = \log(16)/\log(2) \quad B = \log(16/2) \quad C = 16/\log(2)$$

$$D = 3 \log(16)/\log(2) \quad E = \log(48/2)$$

# Flu Outbreak

At the start of an outbreak of H1N1 flu in a large herd of cattle, there were 5 infected individuals. The numbers doubles every 3 days. How many days until there are 80 infected cows?

$$A = \log(16)/\log(2) \quad B = \log(16/2) \quad C = 16/\log(2)$$

$$D = 3 \log(16)/\log(2) \quad E = \log(48/2)$$

At the start of an outbreak of H1N1 flu in a large class of students, there were 5 infected individuals. The numbers doubles every 3 days. How many days until there are 80 infected students?

$$A = \log(16)/\log(2) \quad B = \log(16/2) \quad C = 16/\log(2)$$

$$D = 3 \log(16)/\log(2) \quad E = \log(48/2) \quad \boxed{D}$$