

# Welcome Back!

# Differential Calculus

Instructor:

`schley@math.ucsb.edu`

Office Hours:

© 2022 Daryl Cooper, Peter M. Garfield, Ebrahim Ebrahim &  
Nathan Schley

Please do not distribute outside of this course.

A nice thing about derivatives...

$$\begin{aligned}\frac{d}{dx}(a \cdot f(x) + b \cdot g(x)) &= a \frac{d}{dx} f(x) + b \frac{d}{dx} g(x) \\ &= a \cdot f'(x) + b \cdot g'(x)\end{aligned}$$

For example...

A nice thing about derivatives...

$$\begin{aligned}\frac{d}{dx}(a \cdot f(x) + b \cdot g(x)) &= a \frac{d}{dx} f(x) + b \frac{d}{dx} g(x) \\ &= a \cdot f'(x) + b \cdot g'(x)\end{aligned}$$

For example...

$$\begin{aligned}\frac{d}{dx}(3x^2 + 5x) &= 3 \frac{d}{dx} x^2 + 5 \frac{d}{dx} x \\ &= 3(2x) + 5(1) \\ &= 6x + 5\end{aligned}$$

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:** What is the derivative of  $(x^3 + 1)(2x^2 - 3x + 5)$ ?

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:** What is the derivative of  $(x^3 + 1)(2x^2 - 3x + 5)$ ?

**Question:**  $\frac{d}{dx} ((x^3 + 1)(2x^2 - 3x + 5)) = ?$

A =  $10x^4 - 8x^3 + 10x^2 + 12x - 3$     B =  $10x^4 - 12x^3 + 15x^2 + 4x + 5$   
C =  $10x^4 - 12x^3 + 15x^2 + 4x - 3$     D = Other

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:** What is the derivative of  $(x^3 + 1)(2x^2 - 3x + 5)$ ?

**Question:**  $\frac{d}{dx} ((x^3 + 1)(2x^2 - 3x + 5)) = ?$

A =  $10x^4 - 8x^3 + 10x^2 + 12x - 3$     B =  $10x^4 - 12x^3 + 15x^2 + 4x + 5$   
C =  $10x^4 - 12x^3 + 15x^2 + 4x - 3$     D = Other

**Hint:**  $2x^5 - 3x^4 + 5x^3 + 2x^2 - 3x + 6$

# A Warning!



$$\frac{d}{dx} (f(x)g(x)) \neq f'(x) \times g'(x)$$



**Example:** What is the derivative of  $(x^3 + 1)(2x^2 - 3x + 5)$ ?

**Question:**  $\frac{d}{dx} ((x^3 + 1)(2x^2 - 3x + 5)) = ?$

$$\begin{aligned} A &= 10x^4 - 8x^3 + 10x^2 + 12x - 3 & B &= 10x^4 - 12x^3 + 15x^2 + 4x + 5 \\ C &= 10x^4 - 12x^3 + 15x^2 + 4x - 3 & D &= \text{Other} \end{aligned}$$

**Hint:**  $2x^5 - 3x^4 + 5x^3 + 2x^2 - 3x + 6$

**Answer:**



# Differentiating $f(x) = e^{kx}$

$$\frac{d}{dx} (e^{kx}) = ke^{kx}$$

versus

$$\frac{d}{dx} (x^n) = nx^{n-1}$$



These are not polynomials.  $\frac{d}{dx} (e^{kx}) \neq ke^{(k-1)x}$ .



**Question:** Find  $\frac{d}{dx} (4e^{3x} + 5x^3)$

$$A = 12e^{2x} + 15x^2$$

$$B = 12e^{3x} + 15x^3$$

$$C = 4e^{3x} + 15x^2$$

$$D = 12e^{3x} + 15x^2$$

$$E = \text{Other}$$

# Differentiating $f(x) = e^{kx}$

$$\frac{d}{dx} (e^{kx}) = ke^{kx}$$

versus

$$\frac{d}{dx} (x^n) = nx^{n-1}$$




These are not polynomials.  $\frac{d}{dx} (e^{kx}) \neq ke^{(k-1)x}$ .




**Question:** Find  $\frac{d}{dx} (4e^{3x} + 5x^3)$

A =  $12e^{2x} + 15x^2$       B =  $12e^{3x} + 15x^3$       C =  $4e^{3x} + 15x^2$   
D =  $12e^{3x} + 15x^2$       E = Other      D


# Meanings: The First Derivative




# Meanings: The First Derivative




# Meanings: The First Derivative




# Meanings: The First Derivative



# Meanings: The First Derivative



# Meanings: The First Derivative




Point:

$$f'(x) > 0 \iff f(x) \text{ is increasing}$$


$$f'(x) < 0 \iff f(x) \text{ is decreasing}$$




# Meanings: The Second Derivative




# Meanings: The Second Derivative




# Meanings: The Second Derivative




# Meanings: The Second Derivative




# Meanings: The Second Derivative




# Meanings: The Second Derivative



# Meanings: The Second Derivative



# Meanings: The Second Derivative



**Point:**

$$f''(x) > 0 \iff f'(x) \text{ is increasing}$$

$$\iff f(x) \text{ is concave up}$$

$$f''(x) < 0 \iff f'(x) \text{ is decreasing}$$

$$\iff f(x) \text{ is concave down}$$



# Concavity

$$f''(x) > 0 \iff f(x) \text{ is concave up}$$

$$f''(x) < 0 \iff f(x) \text{ is concave down}$$

(1) For which values of  $x$  is  $f(x) = x^3 - 6x^2 + 3x + 2$  concave up?

A when  $x = 0$     B when  $x < 6$     C when  $x > 6$

D when  $x < 2$     E when  $x > 2$

# Concavity

$$f''(x) > 0 \iff f(x) \text{ is concave up}$$

$$f''(x) < 0 \iff f(x) \text{ is concave down}$$

(1) For which values of  $x$  is  $f(x) = x^3 - 6x^2 + 3x + 2$  concave up?

A when  $x = 0$     B when  $x < 6$     C when  $x > 6$

D when  $x < 2$     E when  $x > 2$     ☐ E

# Concavity

$$f''(x) > 0 \iff f(x) \text{ is concave up}$$


$$f''(x) < 0 \iff f(x) \text{ is concave down}$$

(1) For which values of  $x$  is  $f(x) = x^3 - 6x^2 + 3x + 2$  concave up?

A when  $x = 0$     B when  $x < 6$     C when  $x > 6$

D when  $x < 2$     E when  $x > 2$     E

(2) Where is  $f''(x) > 0$ ?



A when  $x < 2$     B when  $x > 2$     C when  $x < 1$

D when  $x > 1$     E when  $-0.7 < x < 1$

# Concavity

$$f''(x) > 0 \iff f(x) \text{ is concave up}$$


$$f''(x) < 0 \iff f(x) \text{ is concave down}$$

(1) For which values of  $x$  is  $f(x) = x^3 - 6x^2 + 3x + 2$  concave up?

A when  $x = 0$     B when  $x < 6$     C when  $x > 6$

D when  $x < 2$     E when  $x > 2$     ☒ E

(2) Where is  $f''(x) > 0$ ?




A when  $x < 2$     B when  $x > 2$     C when  $x < 1$

D when  $x > 1$     E when  $-0.7 < x < 1$     ☒ D

## §8.13: Max/Min problems


Often want to find the biggest, smallest, most, least, maximum, minimum of something.



Here's the graph of  
 $y = f(x) = -x^2 + 4x + 3$

## §8.13: Max/Min problems

Often want to find the biggest, smallest, most, least, maximum, minimum of something.




Here's the graph of  
 $y = f(x) = -x^2 + 4x + 3$

The maximum value or just maximum of the function is 7.

## §8.13: Max/Min problems

Often want to find the biggest, smallest, most, least, maximum, minimum of something.




Here's the graph of  
 $y = f(x) = -x^2 + 4x + 3$

The maximum value or just maximum of the function is 7.

The value of  $x$  which gives the maximum of  $f(x)$  is  $x = 2$

## §8.13: Max/Min problems

Often want to find the biggest, smallest, most, least, maximum, minimum of something.



Here's the graph of  
 $y = f(x) = -x^2 + 4x + 3$

The maximum value or just maximum of the function is 7.


The value of  $x$  which gives the maximum of  $f(x)$  is  $x = 2$

We write  $f(2) = 7$ .



## §8.13: Max/Min problems

Often want to find the biggest, smallest, most, least, maximum, minimum of something.



Here's the graph of  
 $y = f(x) = -x^2 + 4x + 3$

The maximum value or just maximum of the function is 7.

The value of  $x$  which gives the maximum of  $f(x)$  is  $x = 2$

We write  $f(2) = 7$ .

For this example you can see this is the maximum because

$$f(x) = -x^2 + 4x + 3 = -(x - 2)^2 + 7$$

$(x - 2)^2$  is always positive except when  $x = 2$

so the maximum must be at  $x = 2$ . But there is an easier way.

# How To Find A Maximum

- (1) At the highest point, it's not going up or down. So find  $f'(x)$  to look for the flat part.
- (2) Solve  $f'(x) = 0$  for  $x$ . The  $x$  value that gives the max must be one of these! (Usually there is just one.)
- (3) To find the maximum for  $f(x)$ , use the  $x$ -value you just found...because it gives you the maximum!

# How To Find A Maximum

- (1) At the highest point, it's not going up or down. So find  $f'(x)$  to look for the flat part.
- (2) Solve  $f'(x) = 0$  for  $x$ . The  $x$  value that gives the max must be one of these! (Usually there is just one.)
- (3) To find the maximum for  $f(x)$ , use the  $x$ -value you just found...because it gives you the maximum!

- 1.** Use this method to find the maximum of  $f(x) = -x^2 + 8x + 5$ .  
The maximum value is...

$$A = 4 \quad B = 5 \quad C = -2x + 8 \quad D = 21 \quad E = 15$$

# How To Find A Maximum

- (1) At the highest point, it's not going up or down. So find  $f'(x)$  to look for the flat part.
- (2) Solve  $f'(x) = 0$  for  $x$ . The  $x$  value that gives the max must be one of these! (Usually there is just one.)
- (3) To find the maximum for  $f(x)$ , use the  $x$ -value you just found...because it gives you the maximum!

- 1.** Use this method to find the maximum of  $f(x) = -x^2 + 8x + 5$ .  
The maximum value is...

A = 4    B = 5    C =  $-2x + 8$     D = 21    E = 15    D

# How To Find A Maximum

- (1) At the highest point, it's not going up or down. So find  $f'(x)$  to look for the flat part.
- (2) Solve  $f'(x) = 0$  for  $x$ . The  $x$  value that gives the max must be one of these! (Usually there is just one.)
- (3) To find the maximum for  $f(x)$ , use the  $x$ -value you just found...because it gives you the maximum!

1. Use this method to find the maximum of  $f(x) = -x^2 + 8x + 5$ .  
The maximum value is...

A = 4    B = 5    C =  $-2x + 8$     D = 21    E = 15    D

2. Find the value of  $x$  which makes  $f(x) = (2 - x)(x + 6)$  a maximum.

A = 16    B = 1    C =  $-1$     D = 2    E =  $-2$

# How To Find A Maximum

- (1) At the highest point, it's not going up or down. So find  $f'(x)$  to look for the flat part.
- (2) Solve  $f'(x) = 0$  for  $x$ . The  $x$  value that gives the max must be one of these! (Usually there is just one.)
- (3) To find the maximum for  $f(x)$ , use the  $x$ -value you just found...because it gives you the maximum!



1. Use this method to find the maximum of  $f(x) = -x^2 + 8x + 5$ .  
The maximum value is...

A = 4    B = 5    C =  $-2x + 8$     D = 21    E = 15    D



2. Find the value of  $x$  which makes  $f(x) = (2 - x)(x + 6)$  a maximum.

A = 16    B = 1    C =  $-1$     D = 2    E =  $-2$     E

# How To Find A Minimum?



# How To Find A Minimum?



What this technique **actually does** is find both maxima and minima. In Math 34A a problem will have either a maximum **or** a minimum, **but not both**. So the technique will find what you want. In Math 34B you discover how to do problems which have both a maximum and a minimum and find out which is which.



# More Examples

**3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

# More Examples

**3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

Answer: C

# More Examples

**3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

Answer: C

**4.** What is minimum of  $f(x) = x^2 + 16x^{-2}$ ?

$$A = 2 \quad B = 4 \quad C = 6 \quad D = 8 \quad E = 16$$

# More Examples

**3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

Answer:

**4.** What is minimum of  $f(x) = x^2 + 16x^{-2}$ ?

$$A = 2 \quad B = 4 \quad C = 6 \quad D = 8 \quad E = 16$$

Answer:

# More Examples

- 3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

Answer: C

- 4.** What is minimum of  $f(x) = x^2 + 16x^{-2}$ ?

$$A = 2 \quad B = 4 \quad C = 6 \quad D = 8 \quad E = 16$$

Answer: D

- 5.** Find the value of  $x$  which makes  $f(x) = -e^x - e^{-2x}$  a maximum.

$$A = 0 \quad B = \ln(2) \quad C = -\ln(2) \quad D = \ln(2)/3 \quad E = \ln(2)/3$$

# More Examples

**3.** What is the minimum of  $f(x) = (x + 2)(x + 4) + 3$ ?

$$A = 0 \quad B = 1 \quad C = 2 \quad D = 3 \quad E = 4$$

Answer:

**4.** What is minimum of  $f(x) = x^2 + 16x^{-2}$ ?

$$A = 2 \quad B = 4 \quad C = 6 \quad D = 8 \quad E = 16$$

Answer:

**5.** Find the value of  $x$  which makes  $f(x) = -e^x - e^{-2x}$  a maximum.

$$A = 0 \quad B = \ln(2) \quad C = -\ln(2) \quad D = \ln(2)/3 \quad E = \ln(2)/3$$

Answer:

# Word Problem #1

A ball is thrown into the air. After  $t$  seconds the height in meters above the ground of the ball is  $h(t) = 40t - 10t^2$ . How many meters high did the ball go?

$$A = 2 \quad B = 40 - 20t \quad C = 20 \quad D = 40$$

# Word Problem #1

A ball is thrown into the air. After  $t$  seconds the height in meters above the ground of the ball is  $h(t) = 40t - 10t^2$ . How many meters high did the ball go?

$$A = 2$$

$$B = 40 - 20t$$

$$C = 20$$

$$D = 40$$

$$\boxed{D}$$



## Word Problem #2

If an airline sells tickets at a price of  $\$200 + 5x$  each the number of tickets it sells is  $1000 - 20x$ . What price should the tickets be if the airline wants to get the most money?

$$A = 5 \quad B = 25 \quad C = 175 \quad D = 200 \quad E = 225$$

## Word Problem #2

If an airline sells tickets at a price of  $\$200 + 5x$  each the number of tickets it sells is  $1000 - 20x$ . What price should the tickets be if the airline wants to get the most money?

$$A = 5 \quad B = 25 \quad C = 175 \quad D = 200 \quad E = 225 \quad \boxed{E}$$

# Word Problem #3

A fenced garden with an area of  $100 \text{ m}^2$  will be made in the shape of a rectangle. It will be surrounded on all four sides by a fence. What length and width should be used so the least amount of fence is needed?

# Word Problem #3

A fenced garden with an area of  $100 \text{ m}^2$  will be made in the shape of a rectangle. It will be surrounded on all four sides by a fence. What length and width should be used so the least amount of fence is needed?

## Approach:

- (1) Express the total length of fence in terms of only one variable, either  $L$  = length of field, or  $W$  = width of field. This gives a formula for  $P$  = (total length of fence) involving, say,  $W$ .

# Word Problem #3

A fenced garden with an area of  $100 \text{ m}^2$  will be made in the shape of a rectangle. It will be surrounded on all four sides by a fence. What length and width should be used so the least amount of fence is needed?

## Approach:

- (1) Express the total length of fence in terms of only one variable, either  $L$  = length of field, or  $W$  = width of field. This gives a formula for  $P$  = (total length of fence) involving, say,  $W$ .
- (2) Find minimum by solving  $\frac{dP}{dW} = 0$ .

# Word Problem #3

A fenced garden with an area of  $100 \text{ m}^2$  will be made in the shape of a rectangle. It will be surrounded on all four sides by a fence. What length and width should be used so the least amount of fence is needed?

## Approach:

- (1) Express the total length of fence in terms of only one variable, either  $L$  = length of field, or  $W$  = width of field. This gives a formula for  $P$  = (total length of fence) involving, say,  $W$ .
- (2) Find minimum by solving  $\frac{dP}{dW} = 0$ .

Students always find (1) the hardest part.

**You** have been prepared for this by word problems from chapter 3!