

Nonlinear Dynamics & Chaos I

Exercise Set 7 Questions

Question 1

Consider a planar Hamiltonian system

$$\begin{aligned}\dot{x} &= \frac{\partial H(x, y)}{\partial y} + f_1(x, y), \\ \dot{y} &= -\frac{\partial H(x, y)}{\partial x} + f_2(x, y),\end{aligned}$$

where the twice continuously differentiable function $H(x, y)$ is the Hamiltonian associated with the system (say, the energy in classical mechanics, or the stream function in fluid mechanics), and the continuously differentiable $\mathbf{f} = (f_1, f_2)$ is a dissipative term (say, damping in classical mechanics, or compressible terms in fluid mechanics). Assume that $\nabla \cdot \mathbf{f} \neq 0$ for all $(x, y) \in \mathbb{R}^2$. (Linear damping, for instance has this property.) Show that the above system can have no limit cycles.

Question 2 -

Consider a planar dynamical system with the following phase portrait:




Figure 1: Phase portrait of the planar dynamical system

Which of the following statement is true?

- (a) The ω -limit set of γ is empty.
- (b) By the Poincaré-Bendixson theorem, the ω -limit set of γ is composed of the lines $y = 1$ and $y = -1$.
- (c) The Poincaré-Bendixson theorem does not apply to γ .
- (d) None of the above

Question 3 - Accuracy of averaging

Show that on time scales of $\mathcal{O}(1/\varepsilon)$, a solution $\mathbf{x}(t)$ with $\mathbf{x}(0) = \mathbf{x}_0$ of the dynamical system

$$\dot{\mathbf{x}} = \varepsilon \mathbf{f}(\mathbf{x}, t, \varepsilon), \quad \mathbf{x} \in \mathbb{R}^n, \quad (1)$$

(ε is a small parameter and \mathbf{f} is a smooth function that is T -periodic in time) remains $\mathcal{O}(\varepsilon)$ -close to any solution $\mathbf{y}(t)$ with $\mathbf{y}(0) = \mathbf{x}_0 + \mathcal{O}(\varepsilon)$ of the averaged system

$$\dot{\mathbf{y}} = \varepsilon \bar{\mathbf{f}}_0(\mathbf{y}), \quad \mathbf{y} \in \mathbb{R}^n, \quad (2)$$

where

$$\bar{\mathbf{f}}_0(\mathbf{y}) = \frac{1}{T} \int_0^T \mathbf{f}(\mathbf{y}, t, 0) dt.$$

Hint: Subtract (2) from (1) and integrate to obtain an expression for $|\mathbf{x}(t) - \mathbf{y}(t)|$. Estimate $|\mathbf{x}(t) - \mathbf{y}(t)|$ from above using the facts that $\bar{\mathbf{f}}$ is Lipschitz and $|\hat{\mathbf{f}} - \bar{\mathbf{f}}|/\varepsilon$ is uniformly bounded, where $\hat{\mathbf{f}}$ is the right-hand-side of the system into which (1) is transformed by the averaging transformation $\mathbf{x} = \mathbf{y} + \varepsilon \mathbf{w}(\mathbf{y}, t)$. Then use the following generalized Gronwall inequality:

If $u(t), v(t), c(t)$ are non-negative functions, $c(t)$ is differentiable, and

$$v(t) \leq c(t) + \int_0^t u(s)v(s) ds,$$

then

$$v(t) \leq c(0)e^{\int_0^t u(s) ds} + \int_0^t c'(s)e^{\int_s^t u(\tau) d\tau} ds.$$

Question 4 - Unsteady separation in time-periodic fluid flows

Fluid trajectories $\mathbf{x}(t) = (x(t), y(t))$ in a two-dimensional time-periodic flow satisfy the differential equations

$$\begin{aligned} \dot{x} &= u(x, y, t), & u(x, y, t) &= u(x, y, t + T), \\ \dot{y} &= v(x, y, t), & v(x, y, t) &= v(x, y, t + T), \end{aligned} \quad (3)$$

where $T > 0$ is the period, u and v are smooth velocity components satisfying the incompressibility condition $u_x + v_y \equiv 0$. Assume that the fluid is bounded by a wall at $y = 0$, on which the velocity field satisfies the no-slip boundary conditions $u(x, 0, t) = v(x, 0, t) = 0$. As a result, all boundary points are nonhyperbolic fixed points for (3).

We say that a boundary point $\mathbf{p}_0 = (x_0, 0)$ is a separation point for the flow (3) if \mathbf{p}_0 admits an unstable manifold $W^u(\mathbf{p}_0)$. Physically, $W^u(\mathbf{p}_0)$ is a time-dependent curve of fluid particles that shrinks to \mathbf{p}_0 as backward time. In forward time, $W^u(\mathbf{p}_0)$ attracts fluid particles, then ejects them from the vicinity of the wall (see Fig. 1). Show that separation points satisfy the criteria

$$\int_0^T u_y(x_0, 0, t) dt = 0, \quad \int_0^T v_{yy}(x_0, 0, t) dt > 0,$$

i.e., separation takes place where the average of the wall-shear is zero and the average of v_{yy} is positive.

Hint: Use incompressibility and the boundary conditions to show that (3) can be rewritten as

$$\begin{aligned} \dot{x} &= yU(x, y, t), \\ \dot{y} &= y^2V(x, y, t). \end{aligned}$$

To focus on the vicinity of the boundary, introduce the scaled variable $y = \varepsilon\eta$, where $0 \leq \varepsilon \ll 1$. Show that the resulting $(\dot{x}, \dot{\eta})$ equations are slowly varying and find the corresponding averaged equations. For the averaged equations, rescale time on trajectories by letting $\frac{d\tau}{dt} = \eta(t)$ in order to remove the common η factor from the right-hand-side, and look for a hyperbolic fixed point with an unstable manifold off the wall. Use the averaging theorem to relate your results to the original system (3).




Figure 2: Unsteady separation from a no-slip wall