COLLISION RECORDS, ARITHMETIC, AND RAW LOCAL INVERSION IN A TRIANGULAR LORENTZ GAS

QIAN QI

Abstract. We study the original triangular Lorentz gas for $R \in [9/20, 47/100]$, retaining every physical collision, the unit-speed clock, the prescribed section law and the exact return, displacement and collision records. We continue the complete native proof chain through its endpoint kernels and exact-interval inversion. For the full smooth-guard and bounded C^1 observation class, we identify the entire distributional roof derivative as an integrable density plus a finite signed sum of physical jumps at each collision count. The decomposition keeps every original section side, unvisited boundary and critical entrance. Its geometric support is common to the observation class and has a bound uniform in radius at each fixed count; no complexity bound uniform in the count is assumed. We give the exact jump coefficient, including cancellations at coincident roofs, with the inherited complete variation budget and the original conditioning factor. An elementary full-torus Fourier calculation then recovers the total squared jump mass from the differentiated marked physical transform, with an explicit finite-frequency error. The raw target forces a corresponding collision-normalized central jump energy to vanish. The previously proved zero-clearance critical-fan payment transfers to this signed spectrum. The normalized shallow comparison shortage and the alternative complete local roof-variation condition remain open, as do the full raw pointwise theorem and its four-coordinate inverse conclusion. All inherited proof bodies and original arithmetic factors are retained at their proved scopes.

Contents

Introduction 5
Inherited results and the present additions 6
The original normalization and remaining closing condition 7
Source succession, review scope and proof dependencies 7
1. The unchanged physical problem 8
1.1. The arithmetic reference and the target 9
2. An explicit all-direction physical horizon and lattice budget 9
3. The physical source and the inherited analytic inputs 11
3.1. The original microscopic source 12
4. The reduced optical action 13
4.1. A positive optical denominator 14
5. Completion of the original roofwise endpoint traces 15
6. The complete physical comparison space 17
6.1. The physical grazing partition 18
7. The original microscopic shortage after grazing payment 18
8. A positive payment from the variation on a physical segment 19
9. Persistent guarded segments and the same pointwise target 21

Date: October 23, 2026 (UTC). A2-DYN, physical roof derivative and jump-spectrum author revision.
2020 *Mathematics Subject Classification.* 37D50, 37A50, 60F05, 28A75.
Key words and phrases. Lorentz gas, exact collision records, physical clearance, weighted unstable curves, homogeneous growth, optical Jacobi matrices, marked pressure, pointwise local inversion.

9.1. Comparison with the original nongrazing shortage 22
10. Global uniqueness on a roof–angle fiber 23
10.1. The original physical sum at a target roof 24
11. A complete marked pressure bound at the target roof 25
12. Pointwise removal of deep and numerous incidence witnesses 28
12.1. The retained source after this pointwise payment 29
13. Joint depth and severity of incidence witnesses 30
14. Quadratic tails for the number of incidence witnesses 32
15. Positive payment and the remaining low-load source 33
16. Clearance-to-roof coarea without a fixed-clearance density bound 34
16.1. One-copy summation and the remaining uniformity 37
17. Critical roofs, actual reversing orbits, and their exact density coefficient 38
17.1. What the critical calculation does and does not close 41
18. Exact two-mark clearance flux on one physical flight 41
19. Two clearance marks separated by actual reflections 44
19.1. Physical feet and the reflection derivative 44
19.2. Global multiplicity, including disconnected word domains 46
19.3. An exact physical two-clearance change of variables 46
20. Uniform cross-flight summation and the complete mark count 47
20.1. A stable optical comparison between the two physical feet 47
20.2. All chronological duplicate marks, including repeated centers 49
20.3. What this theorem does, and does not, close 51
21. The angular clearance source as a weighted unstable family 51
21.1. A genuine first collision, not a reflection at the mark 51
21.2. Uniform finite geometry before the homogeneous cut 52
21.3. Paying the infinitely many grazing pieces 53
22. A full cross-flight root sum from curve growth 54
22.1. The inherited complete-record conclusion under the curve input 56
22.2. The original roof obstruction has not been replaced 56
23. A grazing-uniform curvature graph transform 57
24. Physical one-step complexity and homogeneous distortion 59
24.1. Uniform finite geometry and a one-sided first-hit lemma 59
24.2. The complete one-step weighted sum 60
24.3. A direct distortion estimate in the target coordinate 61
25. Admissible physical curve growth and the clearance consequence 62
The distinction from the original roofwise theorem 63
26. Physical momentum fibres and a complete endpoint kernel 64
27. A pressure-free sum of physical critical sectors 67
27.1. One-sided sectors are paid on the physical side 69
27.2. Exact gain and the remaining normalized obstruction 70
28. Exact radial coarea and Borel critical-origin bands 71
28.1. Critical-origin arms with their physical cuts still present 73
28.2. What has and has not been transferred to the original roof 74
29. Grazing-uniform endpoint localization and the third variation 75
29.1. Physical flights, incidence, and every unvisited disk 76
29.2. A third-variation bound uniform in collision count 77
30. A native proof of the original regular-variation estimate 78
30.1. The actual guard, with nonsmooth clearance switches 78
30.2. Roofwise variation and physical normalization 79

30.3. What is closed, and the unchanged remaining mathematical limit 80
31. A grazing-uniform resolvent for the physical reflection equations 80
31.1. Complete higher endpoint derivatives 82
32. Fourth physical variation and the exact high-order cancellation 83
32.1. What this resolves in the original source problem 85
33. Uniform analytic lifts and genuine physical continuation tubes 85
34. Optical distortion and summation at one common physical roof 88
34.1. The remaining normalized source is not eliminated 90
35. A buffered horizon and bending on every physical record 91
36. Normal-boundary continuation and transversality of the physical clock 92
36.1. The full normal-boundary reflection equations 92
36.2. A collision-uniform derivative of the true critical roof 95
37. The complete record: endpoint work and radius–roof disintegration 96
38. A global variational certificate for every physical word 99
39. Global radius order and a positive physical shape energy 101
40. Physical clearance certificates and complete-source localization 103
40.1. The original positive shortage is still the closing condition 105
41. Uniform localization of the entire variational sublevel 105
41.1. Actual records and the inherited comparison functional 105
41.2. The complete positive gap, not a coordinatewise singular estimate 106
42. All-center clearance and a true physical reference 108
42.1. Exact remaining analytic obligation 110
43. Summable physical coercivity and critical singular closure 111
44. Summable clearance and exact-mark transport on the physical clock 113
44.1. Physical reference class and the original section 114
45. Boundary localization of a global disk-word minimum 116
46. Transverse clearance and a uniform physical boundary layer 119
47. The central physical clock and exact section observations 123
48. Exponential confinement by physical anchor layers 125
49. Comparison costs on the original roof and their normalization 128
50. An exponentially small source for exact central physical records 130
51. Physical tube volume and a normalized positive-source payment 134
52. A finite-overlap physical atlas and original-roof source height 138
53. Normalized marked source, the physical singular multiplier, and pressure 142
53.1. The original ordered shortage and the proved new payment 145
54. Explicit stable arclength and the original optical weight 146
55. All physical margins in the original roof measure 150
55.1. The complete deep exceptional source, without a remaining guard 153
55.2. Exact marked comparison on the full source 154
56. The complete physical incidence–inverse-gradient multiplier 155
56.1. The actual insertion on complete critical-origin bands 157
56.2. Full original-roof singular density and deep-source payment 158
57. All-time momentum kernels of the true marked clearance source 160
57.1. A genuine one-sided unvisited-tangency source 162
58. The actual endpoint charge in physical flux coordinates 164
58.1. A complete noncritical boundary estimate from the new physical angular kernel 165
58.2. Cancellation of the apparent pole at an actual critical boundary germ 166
59. Two actual endpoint momenta from a marked free-flight foot 167
60. A uniform cancellation on actual critical-clearance boundary pieces 169

61. Radius-uniform cross-curve distortion on every real word 173
61.1. A two-record consequence of the inherited anchor theorem 174
61.2. Cross-direction distortion and the exact endpoint factor 175
61.3. The precise pressure and optical consequences 178
62. Two-sided capacity of the actual marked angular source 180
62.1. Suprema on genuine one-sided source intervals 180
62.2. An interval overlay estimate without a word-count factor 181
63. Uniform total turning of the physical marked boundary 182
64. The complete regular clearance-boundary budget without an offset cutoff 186
65. A complete outgoing-angle capacity bound for physical classes 188
65.1. The initially uncut affine family 188
65.2. All actual words, strip faces, and unique lifts 190
65.3. Uniform capacity and the long-class consequence 191
65.4. Exact pressure, optical, and observation scope 192
66. Positive critical marked sources with collapsing clearance slices 193
66.1. A physical representative at every small positive clearance 194
66.2. The critical branch-supremum transfer 195
67. Complete physical roof variation without a margin cutoff 197
67.1. The unweighted interior and the actual cap boundaries 198
67.2. Exhausting all physical entrances and exits 199
68. Complete canonical collision-boundary budgets 202
68.1. Actual first images and one-sided supremum sums 202
68.2. Exact two-sided determinants, including the incidence factor 205
68.3. Uniform total turning on the actual source intervals 206
68.4. Complete section-side charges, including the endpoint indices 208
69. Exact marked records and a uniform physical-roof Fourier bound 209
70. A physical boundary stopping proof of the full word-pressure bound 212
70.1. An endpoint-exact comparison with the affine inverse weight 213
70.2. The exhaustive physical exit of a fixed affine interval 214
70.3. The uniform unweighted sum 216
71. Length-sensitive transport of actual clearance sources 218
71.1. The complete renewal, including the last generation 219
71.2. A short interval cover of the true clearance event 220
71.3. The two physical source classes and their exact weights 220
72. A selected physical roof height for an actual clearance witness 222
72.1. Turning constants for the enlarged physical source class 223
72.2. All outgoing boundary charges of the selected source 224
72.3. The selected interior and the incoming critical endpoints 226
73. Deep clearance at the original normalization and the same comparison pool 227
73.1. An exact shortage equivalence with the original full pool 229
74. One-copy variation of the complete exact-label vector 231
75. The actual return tower and positive finite-band likelihoods 232
75.1. A positive kernel with an exact compact frequency support 234
75.2. An exact four-coordinate frequency formula 235
76. Endpoint-tested marked roof operators on the actual collision chain 237
77. Explicit reconstruction of the inherited section multipliers 240
78. Absolute Fourier inversion of the unchanged microscopic intervals 244
78.1. Conditional observations use the actual event mass 246
78.2. A sufficient pointwise bridge, with the missing input exposed 247

78.3. Evaluation of the bridge by the retained fixed-window theorem 247
79. Clearance-selected endpoint kernels and critical boundary tightness 250
79.1. Weighted first images of a selected horizontal source 250
79.2. The complete selected kernel 251
79.3. Uniform tightness of positive-clearance critical sectors 253
80. The exact singular part of the physical roof derivative 254
80.1. Pushforward along an actual boundary side 255
80.2. The signed physical coefficient, before absolute values 257
80.3. The normalized jump constraint for the original raw theorem 260
81. The exact jump spectrum of the full marked physical roof 261
81.1. A necessary central spectrum at the original normalization 262
81.2. The inherited zero-clearance payment in this signed spectrum 263
Appendix A. Reconstruction of the retained physical anchor input 264
Appendix B. An exact flux identity for unvisited near misses 265
Appendix C. Clearance strips transverse to physical test curves 267
Appendix D. Clearance gains after arbitrarily many actual collisions 269
Appendix E. Exact-record disintegration at an unvisited mark 271
E.1. A sufficient pointwise bridge, with its status explicit 271
E.2. Compatibility with the complete original source 272
Appendix F. Scope of the retained comparison statements 272
F.1. Limit order and the remaining estimate 273
Appendix G. Optical cofactors, multiple marks and simultaneous endpoints 274
Appendix H. Dependency, reference and normalization audit 276
H.1. The precise imported curve estimate 276
H.2. Checks which enter the proofs 276
H.3. What is retained, and what has not been reconstructed 277
Appendix I. Critical-roof and fold audit: normalization and ordered closure 277
I.1. Two distinct uses of reversibility 277
I.2. The exact small-scale error and its range 277
I.3. The unchanged closure statement 278
Appendix J. Cross-flight normalization and source-scope audit 278
Appendix K. Angular-source dependency, metric and limit audit 279
External curve growth and physical coordinates 279
Boundary and quantifier conventions 280
Appendix L. Current dependency and citation notes 280
Appendix M. Dependency scope of the physical jump-spectrum continuation 282
The present complete derivative and jump-spectrum interface 282
References 282

Introduction

A physical collision record carries more information than displacement alone. Here the observations are two integer lattice coordinates, the total physical time at unit speed, and the integer occupation of a specified collision section. Conditioning on the original initial and terminal section tests identifies the actual event. The aim is its raw pointwise local limit, with the microscopic physical roof windows and all integer labels retained, and the resulting four-coordinate local inverse theorem.

The fixed family consists of equal circular scatterers on the triangular lattice, with $R \in [9/20, 47/100]$. The collision probability is $d\alpha\, dp/(4\pi)$ and its original section normalization is $c_* = 91/(10000\pi)$. The event (1.1), arithmetic reference (1.4), and target (1.5) are unchanged. Occupation counts the initial state and excludes the terminal state, whose membership is tested separately. The unit-speed flight bounds $\tau_- = 3/50$ and $\tau_+ \le \sqrt{3847/124} < 28/5$, the all-center budgets and the strict marked-foot standoff $\sqrt{1827}/200$ are inherited proved physical geometry. Every reflection in the argument is an actual first collision.

Inherited results and the present additions. The complete native chain through Sections 71–79 is retained. It includes length-sensitive physical source transport, normalized deep-clearance payment, all-label variation, the conservative return tower, endpoint-tested operators, the quantified original section, exact roof-interval inversion, and the selected endpoint kernel. In particular, the critical coefficient estimate of Theorem 79.3 is an unscaled tightness result. It is not an m^{-2} local limit.

The new geometric result describes the actual sides of every physical roof source at fixed collision count. It keeps the first-hit inequalities, all visited reflections, all relevant third-disk clearances, every original section decision, and the exact terminal and occupation conventions. Closures are taken at a fixed radius before the parameterized family is decomposed. Lemma 80.1 produces a finite family of genuine side maps, each constant in the physical roof or strictly monotone on its open piece, with a finite count bound uniform in radius for that fixed collision count. No estimate on its growth with the count is inferred from finite semialgebraic decomposition.

For the complete density vector $f_\ell = f_{\ell,m,R}^{w;a,b,\Psi}$ of Theorem 76.2, the new Theorem 80.3 proves

$$Df_\ell = d_\ell(t)\,dt + \sum_{u\in\mathcal{S}_{m,R}} a_\ell(u)\delta_u, \qquad \sum_\ell \left(\|d_\ell\|_1 + \sum_u |a_\ell(u)| \right) \le \mathcal{V}_m^w(a,b,\Psi).$$

There is no singular-continuous derivative. The critical incoming coefficient is exactly $w_x V_x \omega(x)\beta_{\ell,y}(x)/(2c_*\sqrt{\det G_x})$. Constant-roof regular sides supply their actual signed fluxes. All contributions at the same roof and label are combined before taking an absolute value. Other labels are never combined. Nonconstant sides can yield unbounded integrable turning densities, which are retained in d_ℓ. This is a statement on the arbitrary smooth guard and bounded C^1 tests already admitted by native 332; it does not silently require those observations to be subanalytic.

Write $\mathcal{F}(\vartheta, s)$ for the original transform with all three integer frequencies and the physical roof frequency. Theorem 81.1 proves, at each fixed m, R,

$$\lim_{S\to\infty} \frac{1}{2S(2\pi)^3} \int_{-S}^{S} \int_{[-\pi,\pi]^3} s^2 |\mathcal{F}(\vartheta,s)|^2\, d\vartheta\, ds = \sum_{\ell,u} |a_\ell(u)|^2.$$

The finite-frequency error in (81.9) retains the separation of every pair of distinct physical roofs and the transform of the complete integrable derivative. It makes no uniform separation assumption and exchanges no frequency limit with the radius supremum or collision limit. This Fourier average is an identity for the unchanged physical density, not a replacement of the pointwise objective by an averaged limit theorem.

For a fixed central observation equal to one on a smaller original central window, Theorem 81.3 shows that the raw error $E_m(M_1)$ controls its jump energy:

$$\sup_R m^2 \sum_{\ell,u} |\chi_{\ell,m,R}(u) a_\ell(u)|^2 \le 2D^1_{\mathrm{all}} E_m(M_1).$$

This is a necessary consequence, not a sufficient raw criterion. Corollary 81.4 places the already normalized zero-clearance critical-fan payment inside this signed identity, including coincident roofs and cross terms. That old payment is not claimed anew.

The remaining original shortage keeps its full nongrazing comparison pool. The alternative scalar bridge in Proposition 78.7 already uses the historical fixed-window theorem; it still needs the complete normalized local roof-derivative charge to vanish in the stated order. The new absence of a Cantor part does not prove that charge small: both actual jumps and narrow integrable derivative concentrations must be controlled. Conditional microscopic-return and four-coordinate inverse conclusions retain their actual denominator, arithmetic and Jacobian hypotheses.

The original normalization and remaining closing condition. For every fixed central-window parameter M, the governing positive-source condition remains

$$\lim_{B\to\infty} \limsup_{m\to\infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \left(\mathsf{U}^{\mathrm{low}}_{B,m}(t) - K_B P^{\mathrm{ng}}_{r_B,m}(t)\right)_+ = 0. \tag{0.1}$$

The source, original nongrazing comparison pool and their same-roof meaning are those of the retained native reduction. Complete positive physical sums precede every displayed supremum. All widths and costs are fixed at the outer stage before the collision limsup; B is outermost. The complete pressure theorem and unscaled roof regularity proved below do not by themselves prove this collision-normalized comparison. The necessary microscopic-return, joint spectral, arithmetic and inverse interfaces keep their original scope. The full raw theorem and full inverse consequence are not asserted complete.

Source succession, review scope and proof dependencies. The complete mathematical source continued here is the section-multipliers and exact-inversion author manuscript at commit `85f206907f56773e32fae9b61302924ca16603aa`, article tree `e4ddb6fd11e570dfc7bba9d5fd5a2fed051e78c7`. The immediate Git parent is its complete evidence carrier `d6c3c6754fb3a1b4ad3eea8e6dcc51642e312b28`, article tree `381d0aa576a93d874bf974b0edcf090378aaa41b`. All 76 active proof files 260–335 and the physical model and input statements are preserved byte for byte. Source-stage priority and scope statements inside those retained proofs are read with the current dependency ledger. The preceding main file, introductions, response and ledgers are retained separately.

The fresh census found 436 scoped A2-DYN remote refs among 1306 repository refs. All 435 entries in the prior census still resolved at the same commits; the additional entry was the complete immediate predecessor. The preservation packet contains the verified archive of all 660 distinct native LaTeX objects from that prior census, together with the new complete predecessor source identities. Source, reports, failures and prior execution evidence remain at their original immutable Git objects.

The controlling substantive A2-DYN report is commit `f01482035a0b945b50f1f9c30f30ef35077558dc`. It explicitly reviewed `31792379408c5ace2c09284c2eb22d2331023433` through native 276. It did not review native 277–337. It is an author-requested AI-assisted assessment, not a journal verdict or independent human specialist certification. The response addresses all six principal requests, the four alternative closing routes and nine secondary requests, with applicable earlier obligations. A favorable assessment of the separate A2 geometry paper is not an A2-DYN report.

The native historical section multiplier and fixed-band joint spectral and arithmetic arguments, especially modules 72, 77 and 84–87, retain their proof bodies and scopes. The original weighted momentum capacity is not abundance of the different conditioned same-roof comparison pool. The normalized zero-clearance critical-fan and deep-clearance payments are already proved in

native 322 and 329. The native physical pressure theorem 326 and the predecessor's section, interval inversion and endpoint kernel results 333–335 are inherited.

Sections 80 and 81 contain the present additions. Historical native 40, 92 and 93 already handle the tame scalar singular germs, native 153 the signed boundary current on selected charts, and native 164 the semialgebraic physical graph. We extend the complete signed roof-derivative description to the current arbitrary smooth guard and C^1 observation class, then identify its full marked jump spectrum. The Fourier principle is proved elementarily here; novelty is not claimed for abstract Fourier orthogonality or mean-square identification of atoms.

The external hypotheses are audited at their actual scope. Demers–Zhang [6] supply specified geometric and anisotropic statements; the complete perturbation theorem has additional requirements. The native pressure proof is not an unqualified application of Baladi–Demers [7]. Dolgopyat–Nándori [8] retain their tested-observable, minimality and subgroup assumptions. General convex duality [9] is background to the native disk-word proof. The finite geometric decomposition here uses the definable family hypotheses of Coste [10]; the observation weights need not be definable.

The immediate predecessor was reconstructed from exact Git blobs and independently typeset locally to 271 pages, with all 87 native inputs and its four auxiliary mathematical programs. Its remote workflows were queued at the recorded observation, so no remote success is inferred. The present source manifest, execution receipts and publication record identify the new source and its actual validation separately. Neither finite computations nor successful typesetting certify the outstanding continuous dynamical limit.

1. The unchanged physical problem

Put $R_- = 9/20$ and $R_+ = 47/100$, and let $R \in I = [R_-, R_+]$. Circular scatterers have centers in $\Lambda = \mathbb{Z}(1,0) + \mathbb{Z}(1/2, \sqrt{3}/2)$. Write T_R for the regular collision map and $(\alpha, p) \in (\mathbb{R}/2\pi\mathbb{Z}) \times (-1,1)$ for its coordinates; p is the outgoing tangential velocity and $\mathrm{d}\nu = \mathrm{d}\alpha\, \mathrm{d}p/(4\pi)$. Physical flights satisfy $\tau_- \le \tau_R \le \tau_+$, where $\tau_- = 3/50$ and τ_+ is the fixed finite-horizon bound.

The section is specified here to fix every event in the argument. Put $h_0 = 1/200$. Let Y_R be the union of the four open coordinate squares of half-side h_0 centered at

$$(0,0), \quad (\pi/6, -1/2), \quad (-\theta_R, \sin\theta_R), \quad (\pi/3 - \theta_R, \sin\theta_R).$$

Here $\theta_R = 2\arctan q_R$, and q_R is the unique root in $(49/100, 53/100)$ of

$$8Rq(1+q^2) - 4q(1-q^2) + \sqrt{3}(1 - 6q^2 + q^4) = 0.$$

Set

$$U = \{|\alpha - \pi/6| < 3/50,\ |p| < 3/20\}, \qquad Y_R^* = Y_R \cup U, \qquad c_* = \frac{91}{10000\pi}.$$

The rectangles are disjoint and $\nu(Y_R^*) = c_*$. We always use $\nu_R^* = c_*^{-1}\nu|_{Y_R^*}$, not a renormalization of a smaller source.

At m flights, let $L = L_{m,R}$ be their total length, $K^{\mathrm{c}}_{m,R}$ their lattice displacement, and $A_{m,R} = \sum_{j=0}^{m-1} \mathbf{1}_{Y_R^*} \circ T_R^j$. The event is

$$\Pi_{n,k,m,R} = Y_R^* \cap \{K^{\mathrm{c}}_{m,R} = k,\ A_{m,R} = n\} \cap T_R^{-m} Y_R^*. \tag{1.1}$$

The initial visit is counted; the terminal visit is tested separately. On this event the actual nth return has collision count m and roof L. Denote its density by $p_{n,R}(k,m,t)$.

Write $c_j = \sqrt{1-p_j^2}$ and let Δ_j be the least clearance of flight j from any third disk, excluding its two incident disks. Keep the original smooth function $0 \le \Theta \le 1$, zero on $(-\infty, 1]$ and one on

$[2,\infty)$. With $q_0 = 47/53$, $d_j = \min(j, m-j)$ and $d'_j = \min(j, m-1-j)$, set

$$H^\varepsilon = \prod_{j=0}^{m} \Theta\left(\frac{c_j}{\varepsilon q_0^{d_j/4}}\right) \prod_{j=0}^{m-1} \Theta\left(\frac{\Delta_j}{\varepsilon q_0^{d'_j/4}}\right), \qquad W^\varepsilon = 1 - H^\varepsilon, \tag{1.2}$$

$$b^\varepsilon(t)\,\mathrm{d}t = L_\#(W^\varepsilon \mathbf{1}_{\Pi_{n,k,m,R}} \nu_R^*). \tag{1.3}$$

In particular first incidence and first clearance outside any selected critical neighborhoods are still present. No section-distance factor has been inserted into this guard.

1.1. **The arithmetic reference and the target.** For $y = (k_1, k_2, t, n)$ put

$$\mu_R = (0, 0, \bar\tau_R, c_*), \quad Z = (y - m\mu_R)/\sqrt{m}, \quad \mathcal{C}_{m,R,n,k}(M) = \{t : |Z| \le M\}.$$

We retain the finite spectral-peak reference, including peaks close to but not on the unit circle. To specify its meaning, the collision record observable is $(\kappa_1, \kappa_2, \tau, \mathbf{1}_{Y_R^*})$. Let $\lambda_j(R,\vartheta)$ and $\mathsf{P}_j(R,\vartheta)$ be its continued simple twisted eigenvalue and projection in each of the fixed peak charts, and let χ_j be their frequency partition. At the real maximum a_j of $\log|\lambda_j|$ put

$$z_j = \lambda_j(R, a_j), \quad \mu_j = i^{-1}\nabla \log \lambda_j(R, a_j), \quad H_j = -\nabla^2 \log \lambda_j(R, a_j),$$

with $\operatorname{Re} H_j \ge aI_4$ for a common $a > 0$. If $\eta_R = \mathbf{1}_{Y_R^*}$ and ℓ denotes mass evaluation, put $B_j = \chi_j(R, a_j)\ell(M_{\eta_R}\mathsf{P}_j(R, a_j)(\eta_R \nu))$. Define

$$G_H(z) = (2\pi)^{-4} \int_{\mathbb{R}^4} e^{-iv\cdot z - v^{\mathsf{T}} H v/2}\,\mathrm{d}v,$$

$$\mathcal{L}_{m,R}(y) = \operatorname{Re}\left[c_*^{-1} \sum_j e^{-ia_j\cdot y} z_j^m B_j G_{H_j}\left((y - m\mu_j)/\sqrt{m}\right)\right]. \tag{1.4}$$

Only the charts active at R are summed. This is the same transition reference as in the retained spectral theorem. The existence and uniform estimates for these scalar data belong to that analytic input, not to the transport argument below. No frequency branch or arithmetic zero class is deleted.

The governing assertion remains

$$\sup_{R,\, 1\le n\le m,\, k\in\mathbb{Z}^2} \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} |m^2 p_{n,R}(k,m,t) - \mathcal{L}_{m,R}(k_1,k_2,t,n)| \longrightarrow 0. \tag{1.5}$$

Let $\mathcal{E}_M$ denote this error's collision limsup. An empty central target contributes zero. The order of the uniformity is $\sup_R \operatorname{ess\,sup}_t$, so exceptional density sets may depend on R.

2. An explicit all-direction physical horizon and lattice budget

We give a quantitative proof of finite horizon for the unchanged round scatterers. It is independent of a simulated trajectory, a symbolic model, a pressure theorem and the compactness argument retained in Section 24. It makes the finite geometric constants in the physical source construction directly auditable.

Lemma 2.1 (A row rectangle for a primitive lattice direction)**.** *Let v be a primitive vector of the triangular lattice, $l = |v|$, and $h = \sqrt{3}/(2l)$. For every $q \in \mathbb{R}^2$ there is a lattice center z whose components relative to q, along v/l and its unit normal, satisfy $|x| \le l/2$ and $|y| \le h/2$. Thus for a line direction making angle δ with v,*

$$|(z-q)\cdot e^\perp| \le \tfrac{h}{2}|\cos\delta| + \tfrac{l}{2}|\sin\delta|, \qquad |(z-q)\cdot e| \le \tfrac{1}{2}\sqrt{l^2+h^2}. \tag{2.1}$$

Proof. Primitivity permits v to be completed to a lattice basis. The parallel lattice rows are consequently spaced by the covolume $\sqrt{3}/2$ divided by l. Choose a nearest row to q, and then a nearest lattice center along that row. Its two coordinates obey the stated bounds, regardless of the longitudinal offset of the row. Projection and Cauchy–Schwarz give (2.1). All choices are in the original Euclidean lattice; no linear map deforms the circular disks. □

Theorem 2.2 (A uniform, explicit physical free-flight bound)**.** *For every $R \in [9/20, 47/100]$, every straight segment avoiding the interiors of all radius-R lattice disks has length at most*

$$\tau_{\max} \le \sqrt{\frac{3847}{124}} < \frac{28}{5} < 6. \tag{2.2}$$

In particular the same bound holds for every genuine free flight and all its grazing limits. The statement is simultaneous over all initial points and all directions; no exceptional direction is discarded.

Proof. The dihedral symmetries of the triangular lattice reduce an unoriented line direction to $0 \le \vartheta \le \pi/6$. Put $t = \tan\vartheta/\sqrt{3} \in [0, 1/3]$. We use four primitive lattice vectors, with the indicated ranges:

range of t	v	$\lvert v\rvert^2$
$[0, 1/60]$	$(1, 0)$	1
$[1/60, 1/12]$	$(11/2, \sqrt{3}/2)$	31
$[1/12, 1/5]$	$(5/2, \sqrt{3}/2)$	7
$[1/5, 1/3]$	$(3/2, \sqrt{3}/2)$	3

The last three vectors have lattice coordinates $(5, 1), (2, 1), (1, 1)$, so they are primitive. For the horizontal vector the squared normal width in (2.1) is

$$W_0(t)^2 = \frac{3}{16}\frac{(1+2t)^2}{1+3t^2}.$$

This is increasing on $[0, 1/60]$, since the logarithmic derivative of $(1+2t)/\sqrt{1+3t^2}$ has numerator $2-3t > 0$. For $v = ((2a+1)/2, \sqrt{3}/2)$, write $s = 2a+1$ and $N = a^2+a+1$. The normal width is

$$W_a(t) = \frac{\sqrt{3}}{4\sqrt{1+3t^2}}\left\{\frac{s+3t}{2N} + |st-1|\right\}.$$

On each chosen interval for $a = 5, 2, 1$ one has $t \le 1/s$. The expression in braces is positive and has derivative $3/(2N) - s < 0$. Since the denominator increases, W_a is decreasing. It therefore suffices to check its left endpoint. The four exact squared widths and their positive gaps below $R_-^2 = 81/400$ are

v type	$\max W^2$	$81/400 - \max W^2$
horizontal	961/4804	1181/480400
$a = 5$	13697401/73866304	31513139/1846657600
$a = 2$	529/3136	2651/78400
$a = 1$	75/448	393/11200

Thus, for every direction and midpoint q, Lemma 2.1 provides a lattice center at *strictly* less than R_- normal distance from the line. The longitudinal distance is at most $\sqrt{l^2 + 3/(4l^2)}/2$. Among the four lengths its largest value is $\sqrt{31 + 3/124}/2 = \sqrt{3847/124}/2$. If a free segment were longer than twice this number, the perpendicular foot of the chosen center would lie inside the segment and inside an open radius-R_- disk, a contradiction. This proves the first inequality in (2.2). The comparison with $28/5$ follows by squaring positive quantities: $3847{\cdot}25 = 96175 < 97216 = 784{\cdot}124$. Since the disk radius is only increased for $R \ge R_-$, the entire argument is uniform over the

original family. Strict normal clearance in the covering argument includes limiting directions and grazing free segments; no collision derivative at grazing is used. □

Corollary 2.3 (Explicit enclosing candidate counts, not word counts)**.** *Keep the original constants* $R_+ = 47/100$*,* $\delta_* = 3/200$*, the intrinsic* τ_+ *of the retained source papers, and every original event. Then*

$$N_f \le 150, \qquad N_\circ = N_* \le 150, \qquad N_{\mathrm{pair}} \le 162, \qquad J_f = 2N_f + 1 \le 301.$$

Here N_f *is the first-hit candidate count of* (24.1)*,* $N_\circ$ *is the one-mark/cross-flight count of* (19.7)*, and* N_{pair} *is the difference count in* (18.1)*. In particular the angular first-hit partition in Lemma 21.2 has at most* 1201 *intervals. These are upper bounds for the intrinsic counts, not assertions that all candidate centers can occur or that the smallest physical horizon is* $28/5$*.*

Proof. Theorem 2.2 allows the enclosing constant $\overline{\tau} = 28/5$. The corresponding three enclosing radii are

$$D_f^{\mathrm{enc}} = 327/50, \qquad D_\circ^{\mathrm{enc}} = 1311/200, \qquad D_{\mathrm{pair}}^{\mathrm{enc}} = 657/100.$$

Their squares are respectively $106929/2500$, $1718721/40000$ and $431649/10000$. The first two lie in $[42, 43)$ and the last in $[43, 44)$. A lattice point with integer coordinates (a, b) has squared length $Q(a, b) = a^2 + ab + b^2$, which is integral. The enclosing sets are therefore exactly $0 < Q \le 42$ and $0 < Q \le 43$.

For completeness this finite count does not depend on a floating-point circle. The identity $Q = (a + b/2)^2 + 3b^2/4$ implies $|b| \le 7$ when $Q \le 43$, and by symmetry $|a| \le 7$. Direct solution of the quadratic inequality gives the following inclusive integer ranges for $b \ge 0$:

b	a for $Q \le 42$	#	a for $Q \le 43$	#
0	$[-6, 6]$	13	$[-6, 6]$	13
1	$[-6, 5]$	12	$[-7, 6]$	14
2	$[-7, 5]$	13	$[-7, 5]$	13
3	$[-7, 4]$	12	$[-7, 4]$	12
4	$[-7, 3]$	11	$[-7, 3]$	11
5	$[-7, 2]$	10	$[-7, 2]$	10
6	$[-6, 0]$	7	$[-7, 1]$	9
7	$[-5, -2]$	4	$[-6, -1]$	6

Negative rows have the same cardinalities under $(a, b) \mapsto (-a, -b)$. The totals, excluding the origin, are $12+2(12+13+12+11+10+7+4) = 150$ and $12+2(14+13+12+11+10+9+6) = 162$. The intrinsic candidate sets are subsets of these enclosing sets, which proves all three inequalities. The stated branch bounds now follow from the already proved formulas $2N_f + 1$ and $8N_* + 1$. No collision word, pairing sheet, homogeneous grazing strip or probability normalization is counted by these finite numbers. □

This quantitative result replaces no dynamical hypothesis by a finite simulation. It supplies the geometric finite-horizon input used in the retained source and curve-growth proofs. It does not supply the long-time marked pressure, the sum at a common physical roof, or the collision-normalized positive-source estimate.

3. The physical source and the inherited analytic inputs

We specify the inputs to the new geometric argument, separating the existence of the original physical source from the endpoint regularity that will be proved below. The complete historical proofs and their qualifications are retained in the accompanying archive.

For fixed m, R, n, k, let Θ_{phys} be the original disjoint incident-component pairing-sheet space. It is the countable Borel sum of the maximal regular components of the original exact event, parametrized by

$$(u,\theta) = (F, \arg \nabla_q F), \qquad q = (\alpha_0, \alpha_m), \qquad a(\theta) < u < b(\theta), \quad \ell = b - a > 0.$$

The sheet identity, and not just its angular coordinate, is included in θ. Integration $d\theta$ means the original one-copy physical sum with angular Lebesgue measure. The regular components, their Borel endpoint maps, their physical multiplicity convention, and their common coarea versions are the retained pairing-space input. This article does not replace them by symbolic collision words. Every component carries the same exact event (1.1). Put

$$\begin{aligned} &g = \nabla_q F = R(-p_0, p_m), \quad z = |g|, \quad I_m = 1 + \sum_{i=1}^{m-1} q_0^{d_i}/c_i, \\ &\omega = |\partial_{p_0} \alpha_m|^{-1}, \quad \chi = |\det D_q(F, \arg g)|, \quad A = \frac{\omega}{4\pi c_* \chi}, \quad f^\varepsilon = AW^\varepsilon. \end{aligned} \tag{3.1}$$

The density formula is used initially on the regular interior. Once its endpoint traces have been established, the original pairing is

$$f_-^\varepsilon = f^\varepsilon(a+,\theta), \quad f_+^\varepsilon = f^\varepsilon(b-,\theta), \quad c^\varepsilon = \min(f_-^\varepsilon, f_+^\varepsilon), \quad d\pi^{\varepsilon,+} = c^\varepsilon \mathbf{1}_{\{a<u<b\}} d\theta\, du. \tag{3.2}$$

Write $p_\vee = \max(f_-^\varepsilon, f_+^\varepsilon)$ and $e_\partial = |f_-^\varepsilon - f_+^\varepsilon|$. Thus $p_\vee = c^\varepsilon + e_\partial$.

Input statement 3.1 (Regular variation and completed physical averages)**.** The inherited regular-point optical calculation gives, almost everywhere on each regular segment,

$$|\partial_u f^\varepsilon| \le C(1 + \varepsilon^{-1}) A I_m / z. \tag{3.3}$$

Let $\mathsf{T}_\Delta^\varepsilon$ be the complete paired roof density restricted to $0 < \ell < \Delta$, with no additional selector, and let

$$v^0(t)\, dt = F_\#(I_m z^{-1} \mathbf{1}_\Pi \nu_R^*), \qquad A_r f(t) = (2r)^{-1} \int_{t-r}^{t+r} f(u)\, du.$$

For every previously fixed $\varepsilon, r, \Delta, \rho > 0$,

$$\limsup_m \sup_{R,n,k} m^2 \|A_r \mathsf{T}_\Delta^\varepsilon\|_\infty \le C\{\varepsilon^{1/16} + (1+\varepsilon^{-1})\Delta\}, \tag{3.4}$$

$$\limsup_m \sup_{R,n,k} m^2 \|A_\rho v^0\|_\infty \le C. \tag{3.5}$$

The constants are independent of the previously fixed widths. The coarea identity identifies the full physical sum of AI_m/z with v^0. Fixed-count local finiteness and common roof versions are retained. These are averaged, not pointwise, bounds.

These are the positive pairing, uncut volume and weighted optical variation statements in Section 2 of the frozen article [5]. Their exact historical source labels are recorded in the proof ledger. The new proof uses (3.3) only on regular interiors and supplies its integrable passage to the physical endpoints.

3.1. **The original microscopic source.** The outer schedule is

$$\varepsilon(B) = A_0 B^{-1/12}, \qquad r_0 = c_0^{\mathrm{out}} \varepsilon(B)^4, \qquad \delta(B) = \tfrac{3}{4} r_0, \qquad \Delta_B = 8\delta(B). \tag{3.6}$$

The fixed constants A_0, c_0^{out} are the inherited constants; c_0^{out} is written this way to avoid confusion with the initial incidence c_0. Keep the original remainder selector H_B, the paid critical-arm family $\mathscr{C}_B$, and the adaptive paid family $\mathscr{D}_B$ of the historical microscopic reduction. Their

memberships are Borel, disjointized by first physical membership, and their total outer costs are at most $C\varepsilon(B)^{-1/32}$. For a previously fixed $0 < h_B < \delta(B)$ put

$$\begin{aligned} \widetilde{H}_B &= H_B \mathbf{1}_{(\mathscr{C}_B \cup \mathscr{D}_B)^c} \mathbf{1}_{\{0<\ell<h_B\}}, \\ \mathsf{U}_B^{\mathrm{micro}}(t) &= \int c^{\varepsilon(B)} \mathbf{1}_{\{a<t<b\}} \widetilde{H}_B(\Psi(\theta,t))\, d\theta. \end{aligned} \tag{3.7}$$

No sign cancellation, deletion of an endpoint class, or change of section occurs in this definition. Here Ψ denotes the original physical parametrization.

Input statement 3.2 (The scalar microscopic reduction)**.** For every fixed M, define

$$\mathcal{H}_M(T) = \limsup_m \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t \in \mathcal{C}_{m,R,n,k}(M)} T(t).$$

The historical exact positive reduction gives

$$\mathcal{E}_M \le \mathcal{H}_M(\mathsf{U}_B^{\mathrm{micro}}) + C_M B^{-1/384}. \tag{3.8}$$

The assertion (1.5) is equivalent to $\mathcal{H}_M(\mathsf{U}_B^{\mathrm{micro}}) \to 0$ for every fixed M. All widths and costs are fixed at the outer stage before the complete physical sum, the essential suprema, and the collision limsup; B is outermost.

This is the unchanged theorem in historical module 255. The fixed-positive-residence estimate in module 253 and the earlier paid-arm estimates are retained as its inputs. A bound uniform in a previously fixed positive residence is not used at a collision-dependent residence.

4. The reduced optical action

Fix an actual collision itinerary with $m \ge 1$ flights. Its visited centres are $O_0, \dots, O_m$, its collision points are $x_i = O_i + Rn(\alpha_i)$, and $n(\alpha) = (\cos\alpha, \sin\alpha)$. Write $\tau_i = |x_{i+1} - x_i|$. All variations in this section are made on this actual itinerary before a physical boundary is reached. They neither continue an event through a forbidden collision nor change an occupation label.

Lemma 4.1 (The original section stays away from endpoint grazing)**.** *On the closure of the original initial and terminal section tests,*

$$|p_0|, |p_m| \le \frac{9}{10}, \qquad c_0, c_m \ge c_\circ := \frac{\sqrt{19}}{10}. \tag{4.1}$$

Proof. The only section centres with tangential coordinate exceeding $1/2$ in absolute value have coordinate $2q_R/(1+q_R^2)$. This function is increasing on $(0,1)$, and

$$\frac{2(53/100)}{1+(53/100)^2} + \frac{1}{200} < \frac{9}{10}.$$

The other squares and the rectangle U satisfy the same strict bound. Passing to their closures proves the assertion. □

Let $S(\alpha_0, \dots, \alpha_m) = \sum_{i=0}^{m-1} \tau_i$ be the broken-path action. At a regular specular trajectory its interior variables are stationary. Eliminating them locally gives the actual endpoint action $F(q)$, where $q = (\alpha_0, \alpha_m)$.

Lemma 4.2 (Second variation and its Schur complement)**.** *For a real variation $\xi = (\xi_0, \dots, \xi_m)$ at an actual regular trajectory,*

$$\begin{aligned} D^2S[\xi,\xi] = R\left(c_0\xi_0^2 + c_m\xi_m^2 + 2\sum_{i=1}^{m-1} c_i\xi_i^2\right) \\ + R^2\sum_{i=0}^{m-1}\frac{(c_i\xi_i + c_{i+1}\xi_{i+1})^2}{\tau_i}. \end{aligned} \tag{4.2}$$

The reduced Hessian $G = D_q^2F$ satisfies

$$R\,\mathrm{diag}(c_0, c_m) \le G, \qquad \lambda_\circ I_2 \le G \le \Lambda_\circ I_2, \quad \lambda_\circ = \frac{9\sqrt{19}}{200}, \quad \Lambda_\circ = \frac{47}{6}. \tag{4.3}$$

These inequalities are uniform in the radius, the collision count, and the original exact labels.

Proof. For a flight with unit direction v, the Hessian of its length in its endpoints is the squared perpendicular displacement divided by the flight length. The acceleration of $Rn(\alpha)$ is $-Rn(\alpha)$. At the departure and arrival points, respectively, $v\cdot n_i = c_i$ and $v\cdot n_{i+1} = -c_{i+1}$. Also $(Rn_i')\cdot v^\perp = Rc_i$ and $(Rn_{i+1}')\cdot v^\perp = -Rc_{i+1}$. Thus one flight contributes $R(c_i\xi_i^2 + c_{i+1}\xi_{i+1}^2)$ and the corresponding square in (4.2). Summing proves that formula.

Let $H = D^2S$, $E = \{0, m\}$ and $I = \{1, \dots, m-1\}$. For $m > 1$, the interior block is positive definite. The implicit function theorem and differentiation of the stationary action give

$$G = H_{EE} - H_{EI}H_{II}^{-1}H_{IE}, \qquad \eta^{\mathsf{T}}G\eta = \min_{\zeta}(\eta,\zeta)^{\mathsf{T}}H(\eta,\zeta). \tag{4.4}$$

The order in the last expression places endpoint coordinates first. The diagonal endpoint terms in (4.2) prove the first lower bound. Choosing the interior variation to be zero gives $G \le H_{EE}$. The latter block is bounded above by $\{R_+ + 2R_+^2/\tau_-\}I_2 = (47/6)I_2$; the same bound covers the single-flight case, where there is no Schur complement. Lemma 4.1 proves the remaining lower bound. □

4.1. **A positive optical denominator.** Put

$$\mathsf{S}(s) = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}, \qquad \mathsf{C}(\kappa) = \begin{pmatrix} 1 & 0 \\ \kappa & 1 \end{pmatrix}, \qquad \kappa_i = \frac{2}{Rc_i}. \tag{4.5}$$

Let M_m be the product from the initial collision to the terminal collision, with the rightmost factor acting first:

$$\mathsf{M}_m = \mathsf{S}(\tau_{m-1})\mathsf{C}(\kappa_{m-1})\cdots\mathsf{S}(\tau_1)\mathsf{C}(\kappa_1)\mathsf{S}(\tau_0), \qquad \mathfrak{b}_m = (\mathsf{M}_m)_{12}. \tag{4.6}$$

For $m = 1$ the product is just $\mathsf{S}(\tau_0)$.

Lemma 4.3 (Exact denominator and a marked incidence bound)**.** *The actual endpoint-coordinate density is*

$$\omega = |\partial_{p_0}\alpha_m|^{-1} = \frac{|G_{0m}|}{R} = \frac{Rc_0c_m}{\mathfrak{b}_m}. \tag{4.7}$$

With $\varrho = 47/59$, one has

$$\omega \le \frac{47}{6}\varrho^{m-1}, \qquad \frac{\omega}{c_j} \le \frac{2209}{72}\varrho^{m-2} \quad (1 \le j \le m-1). \tag{4.8}$$

The second inequality is asserted when $m \ge 2$.

Proof. The first equality with G follows by differentiating $p_0 = -\partial_{\alpha_0} F/R$ with α_0 fixed. For the denominator, the interior block of H is $R^2 C_I J C_I$, where $C_I = \operatorname{diag}(c_1, \dots, c_{m-1})$ and J is tridiagonal with

$$J_{ii} = \kappa_i + \tau_{i-1}^{-1} + \tau_i^{-1}, \qquad J_{i,i+1} = \tau_i^{-1}.$$

The corner cofactor in (4.4) gives $|G_{0m}| = R^2 c_0 c_m / (\prod_i \tau_i \det J)$. Expansion of successive tridiagonal determinants gives $\prod_i \tau_i \det J = \mathfrak{b}_m$. This identity also follows by multiplying the second column in (4.6); both recursions start with $\mathfrak{b}_1 = \tau_0$. The empty determinant convention handles $m = 1$.

All entries in the optical factors are nonnegative. Multiplication of a positive second column $(b, d)^{\mathsf{T}}$ by $\mathsf{S}(\tau_i)\mathsf{C}(\kappa_i)$ changes its top entry to $(1+\tau_i\kappa_i)b + \tau_i d$. Since $1+\tau_i\kappa_i \geq 1 + 2\tau_-/R_+ = 59/47$,

$$\mathfrak{b}_m \geq \tau_- \varrho^{-(m-1)}. \tag{4.9}$$

For a fixed interior collision j, split the product immediately before and after $\mathsf{C}(\kappa_j)$. Its $(1,2)$ entry is an expression with nonnegative coefficients in κ_j, whose linear coefficient is $\mathfrak{b}_{\text{right}}\mathfrak{b}_{\text{left}}$. The two subproducts have respectively $m - j$ and j flights. Hence

$$\mathfrak{b}_m \geq \kappa_j \tau_-^2 \varrho^{-(m-2)}.$$

Substitution into (4.7), and $c_0, c_m \leq 1$, proves (4.8). □

Corollary 4.4 (The full inherited incidence insertion)**.** *For the unchanged insertion* $I_m = 1 + \sum_{i=1}^{m-1} q_0^{d_i}/c_i$, *with* $q_0 = 47/53$,

$$\omega I_m \leq 500 \varrho^{(m-2)_+}. \tag{4.10}$$

This is an estimate on each actual physical trajectory; it involves no lower cutoff on any interior incidence.

Proof. For $m \geq 2$, $\sum_{i=1}^{m-1} q_0^{d_i} \leq 2q_0/(1-q_0) = 47/3$. Use (4.8) and $(47/6)\varrho + (2209/72)(47/3) < 500$. For $m = 1$ the sum is empty and the first optical bound suffices. □

No estimate for a sum over collision words is obtained by simply summing the right side of (4.10). In particular, this per-trajectory exponential estimate is not being substituted for a collision-uniform local limit theorem.

5. Completion of the original roofwise endpoint traces

We use the original incident-component pairing sheets and their constant-angle coordinates $(u, \theta) = (F, \arg \nabla F)$. On one of their positive-length maximal regular segments, write $a < u < b$, $g = \nabla F = z e(\theta)$ and $e^{\perp}(\theta) = (-\sin\theta, \cos\theta)$. The physical section and all original exact-event tests are retained.

Lemma 5.1 (Finite Gauss-coordinate length)**.** *Along a fixed-*θ *segment,*

$$\frac{dq}{dz} = G^{-1} e, \qquad \frac{du}{dz} = z e^{\mathsf{T}} G^{-1} e, \qquad 2\lambda_\circ \leq \frac{d(z^2)}{du} \leq 2\Lambda_\circ. \tag{5.1}$$

Consequently z *is strictly increasing, both endpoint limits of* q *exist, and*

$$\int_a^b \frac{du}{z(u,\theta)} \leq \lambda_\circ^{-1}(z_b - z_a) \leq \lambda_\circ^{-1} \min\{\sqrt{2} R_+, \sqrt{2\Lambda_\circ (b-a)}\}. \tag{5.2}$$

A critical endpoint of a positive-length segment can only be its incoming endpoint.

Proof. Differentiate $\nabla F(q) = ze$ and then $F(q) = u$. The bounds in (4.3) give (5.1). They also give $|dq/dz| \le \lambda_\circ^{-1}$. Since $0 < z \le \sqrt{2}R_+$, the two endpoint limits of q exist in the chosen angular lifts. Integrating $du/z = e^{\mathsf{T}} G^{-1} e\, dz$ proves the first inequality in (5.2). The second uses $z_b^2 - z_a^2 \le 2\Lambda_\circ(b-a)$ and $(z_b - z_a)^2 \le z_b^2 - z_a^2$. If $z_b = 0$, strict increase would contradict $z > 0$ in the interior. □

Lemma 5.2 (A continuous reduced Hessian at visited grazing)**.** *The reduced Hessian has a continuous extension along a fixed itinerary when any collection of its interior c_i tends to zero, provided $\tau_i \ge \tau_-$ and* (4.1) *hold. Its extension still satisfies* (4.3)*. If at least one interior c_j vanishes, the extended mixed entry G_{0m} vanishes.*

Proof. Let $L = \sum_{i=0}^{m-1} \tau_i^{-1}(e_i + e_{i+1})(e_i + e_{i+1})^{\mathsf{T}}$, and set $C_E = \operatorname{diag}(c_0, c_m)$. For $m > 1$ put

$$Z = 2I + RC_I^{1/2} L_{II} C_I^{1/2} \ge 2I.$$

Formula (4.4) can be written without inverse incidences:

$$\begin{aligned} G = {} & RC_E + R^2 C_E L_{EE} C_E \\ & - R^3 C_E L_{EI} C_I^{1/2} Z^{-1} C_I^{1/2} L_{IE} C_E. \end{aligned} \tag{5.3}$$

Every term is continuous for $c_i \ge 0$, since Z^{-1} remains bounded. The order inequalities pass to this limit. The marked bound in (4.8), together with $|G_{0m}| = R\omega$, proves the last assertion. There is no interior extension to make when $m = 1$. □

At an endpoint of a fixed-θ segment the complete finite trajectory also has a limit. Indeed Lemma 5.1 gives the initial angle and $p_0 = -z\cos\theta/R$. For each designated next disk, the entry root of the flight quadratic is continuous up to a tangency. The flight length cannot collapse to zero, and the reflection $v \mapsto v - 2(v \cdot n)n$ is continuous at tangency. Induction over the fixed finite itinerary gives the limits of all visited points, velocities, τ_i and c_i. The third-disk clearances have limits as well: the finite horizon restricts the possible nearby disk centres to a finite set, and the minimum distance to each of them is continuous. This argument takes only one-sided limits from the actual event; it asserts no continuation through its boundary.

Theorem 5.3 (Trace completion on the original physical sheets)**.** *On every positive-length original regular pairing segment the coarea factor and weighted density are*

$$\chi = \det D_q(F, \arg g) = e^{\perp\mathsf{T}} G e^{\perp} \ge \lambda_\circ, \qquad f^\varepsilon = \frac{\omega W^\varepsilon}{4\pi c_* \chi}. \tag{5.4}$$

They have finite one-sided endpoint limits. The regular-point variation input (3.3) *implies that f^ε extends absolutely continuously to the closed segment and satisfies*

$$|f^\varepsilon(b-) - f^\varepsilon(a+)| \le C(1 + \varepsilon^{-1}) \int_a^b AI_m/z\, du. \tag{5.5}$$

In fact, on each physical sheet,

$$|f^\varepsilon(v) - f^\varepsilon(u)| \le C(1+\varepsilon^{-1})\varrho^{(m-2)_+}|v-u|^{1/2} \quad (a \le u < v \le b). \tag{5.6}$$

If a limiting trajectory at either endpoint contains an actually visited interior grazing collision, its trace at that endpoint is zero. The original paired minimum on that entire segment is then zero, for every original selector.

Proof. The determinant identity in (5.4) follows by expressing $D\arg g$ as $e^{\perp\mathsf{T}} G/z$. The preceding endpoint-limit argument and Lemma 5.2 give limits for G, ω and χ, the last with a positive lower bound. Every factor in the finite product defining W^ε is continuous in its incidence or clearance

argument. Thus f^ε has the asserted finite limits, including at a critical incoming endpoint, where the direction θ is fixed.

Corollary 4.4 and (5.4) bound AI_m by $C\varrho^{(m-2)_+}$. The integral in (5.5) is finite by Lemma 5.1. On compact regular subsegments the flight charts are smooth and the finite clearance minima are locally Lipschitz, so f^ε is locally absolutely continuous. Integrating the inherited regular-point derivative inequality on these subsegments and then passing to the proved endpoint limits establishes absolute continuity and (5.5). The same argument between u and v, using (5.2), gives (5.6).

Finally, if $c_j \to 0$ at a visited interior collision, the marked optical bound gives $\omega \to 0$, while $0 \le W^\varepsilon \le 1$ and $\chi \ge \lambda_\circ$. Hence that endpoint trace is zero. The paired density is its nonnegative minimum with the other trace. □

Remark 5.4 (Two different tangency mechanisms). The last assertion concerns a collision in the visited itinerary. A flight which merely becomes tangent to an *unvisited* third disk has zero clearance but need not have any $c_j = 0$. If all visited incidences remain positive, then (4.7) has a positive limit. A zero third-disk clearance makes $W^\varepsilon = 1$ at the limit; consequently the trace can be positive. Such sheets are not removed. Neither are nongrazing critical entrances or positive section-boundary traces. For a nongrazing critical entrance the exact directional limit is $\omega_* W_*^\varepsilon/(4\pi c_* e^{\perp\mathsf{T}} G_* e^\perp)$. No roof-independent value at the critical point is asserted.

6. The complete physical comparison space

The comparison space is now specified without an unverified endpoint-trace exclusion:

$$\Theta_{\mathrm{ad}} = \Theta_{\mathrm{phys}}. \tag{6.1}$$

More explicitly, it consists of all positive-length maximal regular pairing segments of the original exact event, with their original incident-component identities. Their endpoints may be critical, visited-grazing, third-disk-tangent, or section-boundary limits. Theorem 5.3 supplies finite traces and the endpoint variation inequality on this space. The underlying Borel pairing construction and the averaged analytic estimates remain the inputs in Section 3. No singular trajectory is continued past its physical boundary, and no new source supported at a zero-length segment is introduced.

For fixed $0 < r < \Delta$, define

$$\begin{aligned} q_+(t,\theta) &= \mathbf{1}_{\{0<\ell<\Delta,\ a<t,\ t+r<b\}},\\ q_-(t,\theta) &= (1-q_+)\mathbf{1}_{\{0<\ell<\Delta,\ a<t-r,\ t<b\}}, \qquad Q_r = q_+ + q_-. \end{aligned} \tag{6.2}$$

Thus Q_r is Borel and at most one. Write

$$P_r(t) = \int Q_r p_\vee \, d\theta, \qquad D_r(t) = \int Q_r c^\varepsilon \, d\theta, \qquad E_r(t) = \int Q_r e_\partial \, d\theta.$$

The full pool is retained even on a sheet whose paired source is zero.

Theorem 6.1 (Capacity on the full original comparison space)**.** *At finite collision count, on the original common roof versions,*

$$P_r(t) \le \frac{1}{r}\int_{t-r}^{t+r} \mathsf{T}_\Delta^\varepsilon(u)\,du + C(1+\varepsilon^{-1})\int_{t-\Delta}^{t+\Delta} v^0(u)\,du. \tag{6.3}$$

Consequently, for every previously fixed $K > 0$,

$$\mathcal{H}_M(KP_r) \le CK\{\varepsilon^{1/16} + (1+\varepsilon^{-1})\Delta\}. \tag{6.4}$$

Proof. Every positively oriented selected segment contains $[t, t+r]$; every negatively oriented one contains $[t-r, t]$. Integrating its constant paired minimum over the appropriate window gives r times that minimum. Tonelli and $Q_r \le 1$ therefore give $D_r(t) \le r^{-1} \int_{t-r}^{t+r} \mathsf{T}_\Delta^\varepsilon$. Each selected segment lies inside $(t-\Delta, t+\Delta)$. The completed variation inequality (5.5) and the original one-copy physical coarea sum give $E_r(t) \le C(1+\varepsilon^{-1}) \int_{t-\Delta}^{t+\Delta} v^0$. Since $P_r = D_r + E_r$, this proves (6.3). Use (3.4) and (3.5), only on these previously fixed windows, to obtain (6.4). The weights are held at the original roof t throughout; none is differentiated or evaluated at the integration roof u. □

6.1. **The physical grazing partition.** For each interior collision let c_i^- and c_i^+ be its incidence limits at the two ends of a pairing segment. Define the Borel sets

$$\Gamma = \bigcup_{i=1}^{m-1} \{c_i^- = 0 \text{ or } c_i^+ = 0\}, \qquad \mathcal{N} = \Theta_{\mathrm{phys}} \setminus \Gamma. \tag{6.5}$$

For $m = 1$ the union is empty. The limits are Borel, for example by evaluating the interior functions at $a + \ell/j$ and $b - \ell/j$ and letting the integer j tend to infinity. Set

$$P_r^{\mathrm{gr}} = \int_\Gamma Q_r p_\vee \, d\theta, \qquad P_r^{\mathrm{ng}} = \int_{\mathcal{N}} Q_r p_\vee \, d\theta, \qquad P_r = P_r^{\mathrm{ng}} + P_r^{\mathrm{gr}}.$$

The superscripts refer to visited collision incidences in the limiting finite itinerary, not to a lower incidence cutoff.

Theorem 6.2 (Zero original grazing source and paid grazing capacity)**.** *For every original Borel selector $0 \le H \le 1$, at finite count,*

$$\int_\Gamma c^\varepsilon \mathbf{1}_{\{a<t<b\}} H(\Psi(\theta,t)) \, d\theta = 0. \tag{6.6}$$

Nevertheless P_r^{gr} need not vanish. It satisfies

$$P_r^{\mathrm{gr}}(t) \le C(1+\varepsilon^{-1}) \int_{t-\Delta}^{t+\Delta} v^0(u) \, du, \tag{6.7}$$

$$\mathcal{H}_M(KP_r^{\mathrm{gr}}) \le CK(1+\varepsilon^{-1})\Delta. \tag{6.8}$$

These estimates include the complete positive capacity on this physical class, with no count of angular blocks or collision words.

Proof. Theorem 5.3 gives $c^\varepsilon = 0$ on Γ, proving (6.6) before the roof pushforward. On that same class $p_\vee = e_\partial$, not necessarily zero. Apply (5.5), use the localization $(a,b) \subset (t-\Delta, t+\Delta)$ on $\{Q_r = 1\}$, and complete the one-copy positive physical sum. This proves (6.7). Equation (3.5) at the fixed width Δ proves (6.8). □

7. The original microscopic shortage after grazing payment

All allocations below are scalar allocations at the original roof; they are not couplings of individual source and comparison sheets. For any original source $U(t)$ define

$$V_{r,K}(t) = \min\{U(t), KP_r(t)\}, \qquad R_{r,K}(t) = (U(t) - KP_r(t))_+.$$

The multiplier is $\beta_{r,K} = \min\{1, KP_r/U\}$ on $U > 0$ and zero on $U = 0$. Then $U = V_{r,K} + R_{r,K}$ at finite count and $V_{r,K}$ is the original source multiplied by $\beta_{r,K}(t)$. When a restricted comparison space is used, source outside it is still in U and may receive this same global fraction. It is not therefore retained sheetwise with coefficient one.

Use the exact source (3.7) and, for a concrete outer choice, put

$$h_B = \delta(B)/2, \qquad r_B = \delta(B)/8, \qquad K_B = \max\{1, \varepsilon(B)^{-1/32}\}. \tag{7.1}$$

For all sufficiently large B these satisfy the inherited cost budget. Define

$$\begin{aligned} \mathsf{R}_B^{\mathrm{full}} &= (\mathsf{U}_B^{\mathrm{micro}} - K_B P_{r_B})_+, \\ \mathsf{R}_B^{\mathrm{ng}} &= (\mathsf{U}_B^{\mathrm{micro}} - K_B P_{r_B}^{\mathrm{ng}})_+. \end{aligned} \tag{7.2}$$

The full capacity pool includes any sheets previously excluded solely for lack of a trace verification. On the complete comparison space it is exactly the canonical pool; on a previously restricted space it is an enlargement, not an identification of different capacities.

Theorem 7.1 (Quantitative payment of the visited-grazing class)**.** *Under the retained physical-source and analytic inputs, for the original event and every fixed M,*

$$0 \le \mathsf{R}_B^{\mathrm{ng}} - \mathsf{R}_B^{\mathrm{full}} \le K_B P_{r_B}^{\mathrm{gr}}, \tag{7.3}$$

$$\mathcal{H}_M(\mathsf{R}_B^{\mathrm{ng}} - \mathsf{R}_B^{\mathrm{full}}) \le C_M B^{-95/384}, \tag{7.4}$$

$$\mathcal{E}_M \le \mathcal{H}_M(\mathsf{R}_B^{\mathrm{ng}}) + C_M B^{-1/384}. \tag{7.5}$$

The unchanged pointwise assertion (1.5) *is equivalent to $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{ng}}) \to 0$ for every fixed M. The original microscopic source in these formulas is supported on $\mathcal{N}$ without a change of its positive weight.*

Proof. For $u, a, b \ge 0$, the positive-part function gives $0 \le (u-a)_+ - (u-a-b)_+ \le b$. Apply this with $a = K_B P_{r_B}^{\mathrm{ng}}$ and $b = K_B P_{r_B}^{\mathrm{gr}}$ to prove (7.3) at finite count. Theorem 6.2, $\Delta_B = O(\varepsilon^4)$ and $K_B = O(\varepsilon^{-1/32})$ give

$$\mathcal{H}_M(K_B P_{r_B}^{\mathrm{gr}}) \le C\varepsilon^{-1/32}(1+\varepsilon^{-1})\varepsilon^4 = O(\varepsilon^{95/32}) = O(B^{-95/384}).$$

This proves (7.4) in the prescribed order. By Theorem 6.1, the accepted full-pool source has height at most

$$C\varepsilon^{-1/32}\{\varepsilon^{1/16} + O(\varepsilon^3)\} = O(B^{-1/384}).$$

Insert its exact positive decomposition into (3.8). Since $\mathsf{R}_B^{\mathrm{full}} \le \mathsf{R}_B^{\mathrm{ng}}$, this proves (7.5) and sufficiency of the displayed residual limit. Necessity follows from Input statement 3.2 and $0 \le \mathsf{R}_B^{\mathrm{ng}} \le \mathsf{U}_B^{\mathrm{micro}}$. Equation (6.6) proves the final assertion. □

This theorem supplies a physical component of the required scarcity estimate, rather than a new certificate class. It removes an entire visited-grazing source class exactly and pays the loss of all its comparison capacity by a faster-decaying original optical volume. It does not yet bound $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{ng}})$ by a quantity tending to zero. In $\mathcal{N}$ every visited endpoint incidence is positive at each fixed count, but its infimum may tend to zero with the count. Third-disk tangencies, nongrazing critical entrances, section-boundary cuts, flat and short-reach segments, and the original comparison shortages all remain with their full weight. A collision-uniform estimate on that remaining source is still required for (1.5). Neither finite-count exhaustion nor the per-itinerary exponential bound is used in its place.

8. A positive payment from the variation on a physical segment

The comparison pool of Section 6 is left unchanged. We first make a different use of the completed endpoint estimate: it pays part of the *original source*, whether or not that source has a comparison window. All statements in this section are made on the inherited one-copy physical pairing space.

Write C_{reg} for a fixed constant in (3.3). For fixed $\varepsilon > 0$ define

$$\begin{aligned} \mathcal{V}^{\varepsilon}(\theta) &= \mathrm{Var}_{[a(\theta),b(\theta)]}\, f^{\varepsilon}(\,\cdot\,,\theta), \\ \mathcal{J}(\theta) &= \int_{a(\theta)}^{b(\theta)} \frac{A(u,\theta) I_m(u,\theta)}{z(u,\theta)}\, du. \end{aligned} \tag{8.1}$$

Endpoint values in the variation are the one-sided physical traces, not values assigned on a continuation beyond the original event.

Lemma 8.1 (Measurability and the full variation bound)**.** *Both functions in* (8.1) *are Borel and finite on every positive-length physical segment. Moreover,*

$$\mathcal{V}^{\varepsilon} = \int_a^b |\partial_u f^{\varepsilon}|\, du \le C_{\mathrm{reg}}(1+\varepsilon^{-1})\mathcal{J}. \tag{8.2}$$

This is a bound for total variation, not merely for the difference of the two endpoint values.

Proof. Pull the closed segment back to $[0,1]$ by $u = a + s(b-a)$, using the endpoint traces at $s = 0, 1$. The interior evaluations are jointly Borel. At $s = 0$ the completed value is the limit of evaluations at $a + \ell/j$, $j \ge 2$; at $s = 1$ it is the limit of evaluations at $b - \ell/j$. The completed endpoint values, and hence the function on the closed product, are jointly Borel. The resulting function is continuous in s. Its variation is the supremum of the sums of absolute increments over finite rational partitions containing 0 and 1. Continuity permits this countable family of partitions, so its supremum is Borel. The integral defining $\mathcal{J}$ is Borel by the same change of variables and nonnegative integration. Theorem 5.3 gives absolute continuity on the closed segment. The variation of an absolutely continuous real function is the integral of the absolute value of its derivative. Integrate (3.3); its right side is integrable by Corollary 4.4 and Lemma 5.1. This proves the equality, inequality and finiteness asserted above. □

For $\gamma \ge 1$, split the original paired coefficient into

$$c^{\varepsilon,\gamma}_{\mathrm{var}} = \min\{c^{\varepsilon}, \gamma\mathcal{V}^{\varepsilon}\}, \qquad c^{\varepsilon,\gamma}_{\mathrm{per}} = (c^{\varepsilon} - \gamma\mathcal{V}^{\varepsilon})_+. \tag{8.3}$$

Thus $c^{\varepsilon} = c^{\varepsilon,\gamma}_{\mathrm{var}} + c^{\varepsilon,\gamma}_{\mathrm{per}}$, with both terms nonnegative. The subscript per will be justified by a physical persistence statement in the next section. It denotes a positive *submeasure* of the original source, not a replacement of its coefficient by a different normalization.

For a jointly Borel selector $0 \le H \le 1$ supported on $0 < \ell < h$, set

$$T^{\varepsilon,\gamma}_{H,j}(t) = \int c^{\varepsilon,\gamma}_j(\theta)\mathbf{1}_{\{a<t<b\}}H(\Psi(\theta,t))\, d\theta, \qquad j \in \{\mathrm{var}, \mathrm{per}\}. \tag{8.4}$$

The original selected paired source is their exact sum. The selector is evaluated at t and is not moved to another roof in the proof.

Theorem 8.2 (A summed payment of the positive short source)**.** *At every finite collision count, on the original common roof versions,*

$$0 \le T^{\varepsilon,\gamma}_{H,\mathrm{var}}(t) \le C_{\mathrm{reg}}\gamma(1+\varepsilon^{-1}) \int_{t-h}^{t+h} v^0(u)\, du. \tag{8.5}$$

Consequently, for every $\varepsilon, h > 0$ and $\gamma \ge 1$ fixed before the collision limit,

$$\mathcal{H}_M(T^{\varepsilon,\gamma}_{H,\mathrm{var}}) \le C\gamma(1+\varepsilon^{-1})h. \tag{8.6}$$

The constants are uniform in the collision count, radius, exact labels, segment length and selector. No positive lower bound for ℓ or for an interior incidence is required.

Proof. For a segment meeting t and satisfying $\ell < h$, one has $(a,b) \subset (t-h, t+h)$. By (8.3) and (8.2), its selected contribution is at most

$$C_{\mathrm{reg}}\gamma(1+\varepsilon^{-1})\mathbf{1}_{\{a<t<b,\, \ell<h\}}H(\Psi(\theta,t)) \int_a^b AI_m/z\, du.$$

First sum these nonnegative quantities over the physical sheets. Tonelli's theorem applies, without an estimate for the number of sheets. At an integration roof u, each retained pair (θ, u) is still a

point of the same original exact event. Dropping its selector, which is between zero and one, and using physical multiplicity one therefore bounds the sum by

$$C_{\mathrm{reg}}\gamma(1+\varepsilon^{-1})\int_{t-h}^{t+h}\int_{\{a(\theta)<u<b(\theta)\}} A(u,\theta)I_m(u,\theta)/z(u,\theta)\,d\theta\,du.$$

The coarea identity in Input statement 3.1 identifies the inner integral with $v^0(u)$. This proves the finite-count inequality. Apply (3.5) at the previously fixed width h, using $\int_{t-h}^{t+h} v^0 = 2hA_h v^0$. Only after the complete sum and this inequality are established are the uniform norms and collision limsup taken. This proves (8.6). □

Corollary 8.3 (The original microscopic schedule)**.** *Keep $\widetilde{H}_B, h_B, r_B, K_B$ exactly as in* (3.7) *and* (7.1)*, and choose*

$$\gamma_B = \max\{2, \varepsilon(B)^{-1/32}\}. \tag{8.7}$$

Let $\mathsf{U}_B^{\mathrm{var}}$ and $\mathsf{U}_B^{\mathrm{per}}$ denote the two sources in (8.4) *with $H = \widetilde{H}_B$, $\varepsilon = \varepsilon(B)$ and $\gamma = \gamma_B$. Then*

$$\mathsf{U}_B^{\mathrm{micro}} = \mathsf{U}_B^{\mathrm{var}} + \mathsf{U}_B^{\mathrm{per}}, \qquad \mathcal{H}_M(\mathsf{U}_B^{\mathrm{var}}) \le C_M B^{-95/384}. \tag{8.8}$$

Proof. The identity holds before pushforward, by (8.3). For sufficiently large B, $h_B = O(\varepsilon^4)$ and $\gamma_B = O(\varepsilon^{-1/32})$. Theorem 8.2 gives the bound

$$C\gamma_B(1+\varepsilon^{-1})h_B = O(\varepsilon^{95/32}) = O(B^{-95/384}).$$

All three quantities $h_B, \varepsilon(B), \gamma_B$ are fixed before m tends to infinity. No collision-dependent residence is inserted into an estimate asserted only for fixed residence. □

Unlike (6.6), this payment does not rely on a zero paired coefficient. If $c^\varepsilon > 0$ and $\mathcal{V}^\varepsilon > 0$, its coefficient in (8.3) is positive. If $c^\varepsilon \le \gamma\mathcal{V}^\varepsilon$, the complete original coefficient is paid. These statements do not assert a positive uniform proportion of paid mass in every exact event.

9. Persistent guarded segments and the same pointwise target

The positive decomposition of the preceding section has a geometric consequence on every segment which retains source. It uses the original incidence and clearance guard, without adding a distance to the section boundary.

Theorem 9.1 (Persistence of the guard on the remaining source)**.** *Let $\gamma > 1$. If $c^{\varepsilon,\gamma}_{\mathrm{per}}(\theta) > 0$, then $c^\varepsilon > 0$ and, at every point of the closed physical segment,*

$$|f^\varepsilon(u,\theta) - c^\varepsilon(\theta)| \le \mathcal{V}^\varepsilon(\theta) < \gamma^{-1}c^\varepsilon(\theta). \tag{9.1}$$

In particular $f^\varepsilon > 0$ and $W^\varepsilon > 0$ throughout that segment, including its one-sided endpoints. For each u at least one of the original tests satisfies

$$c_j(u) < 2\varepsilon q_0^{d_j/4} \quad \text{for some } 0 \le j \le m, \qquad \text{or} \qquad \Delta_j(u) < 2\varepsilon q_0^{d'_j/4} \quad \text{for some } 0 \le j < m. \tag{9.2}$$

The witnessing test may change with u.

Conversely, if W^ε vanishes anywhere on the closed segment, its complete original paired coefficient is paid: $c^{\varepsilon,\gamma}_{\mathrm{var}} = c^\varepsilon$ and $c^{\varepsilon,\gamma}_{\mathrm{per}} = 0$.

Proof. The endpoint minimum is attained at at least one endpoint, where $f^\varepsilon = c^\varepsilon$. Every increment from that endpoint is bounded by the total variation. Positivity of the remaining coefficient is precisely $c^\varepsilon > \gamma\mathcal{V}^\varepsilon$, which proves (9.1). If any visited interior incidence vanished at an endpoint, Theorem 5.3 would give $c^\varepsilon = 0$. Thus the segment belongs to $\mathcal{N}$. Its optical factor A is finite and positive on the closed segment: the exact formula $\omega = Rc_0c_m/\mathfrak{b}_m$ in (4.7) has a finite positive denominator when all visited incidences are positive, while (5.4) gives $\chi \ge \lambda_\circ$. This also applies at a critical endpoint with its fixed Gauss direction. Since $f^\varepsilon = AW^\varepsilon > 0$, the

guard is positive there. If all inequalities in (9.2) failed, every factor defining H^ε would equal one, contradicting $W^\varepsilon = 1 - H^\varepsilon > 0$.

For the converse, let u_0 be a point where $W^\varepsilon = 0$. The finite optical factor gives $f^\varepsilon(u_0) = 0$, so the increment from an endpoint attaining c^ε gives $\mathcal{V}^\varepsilon \geq c^\varepsilon$. Apply (8.3). This argument also covers $u_0 = a$ or $u_0 = b$, with the proved physical one-sided values. □

The surviving class

$$\mathcal{P}_B = \{\theta \in \mathcal{N} : c^{\varepsilon(B)}(\theta) > \gamma_B \mathcal{V}^{\varepsilon(B)}(\theta)\} \tag{9.3}$$

is Borel. The source $\mathsf{U}_B^{\mathrm{per}}$ is supported on it; its coefficient is still the positive difference in (8.3), not the full original coefficient. The sum (8.8) retains every original coefficient exactly. No assumption that one fixed bad test covers the entire segment is made. A Borel witness at a specified point can be chosen as the first successful test in the finite ordered list in (9.2); this selection is not needed to prove the payment.

Remark 9.2 (Boundary mechanisms and what is retained). An unvisited third-disk tangency may give $W^\varepsilon = 1$ at an endpoint without forcing variation relative to the endpoint minimum. It is therefore not declared paid merely on account of its name. The same applies to a nongrazing critical entrance, an original section cut and a short-reach segment. Each receives precisely the two coefficients in (8.3). When the guard vanishes elsewhere on that segment the entire coefficient is paid by Theorem 9.1; otherwise a positive remaining coefficient is possible. At fixed collision count the optical data on a surviving segment have positive minima, but no such minimum is asserted uniformly in the collision count or the sheet.

9.1. **Comparison with the original nongrazing shortage.** Keep the *full original* nongrazing pool $P_{r_B}^{\mathrm{ng}}$, including its capacity on segments whose source has already been paid. Define only a new source-side remainder:

$$\mathsf{R}_B^{\mathrm{per}}(t) = \big(\mathsf{U}_B^{\mathrm{per}}(t) - K_B P_{r_B}^{\mathrm{ng}}(t)\big)_+. \tag{9.4}$$

The allocation by this pool is scalar at the roof, just as in Section 7. It is not a coupling of individual persistent sheets to individual comparison sheets.

Theorem 9.3 (Positive-source reduction with a paid error)**.** *On the original exact event, at every finite collision count,*

$$\mathsf{U}_B^{\mathrm{micro}} = \mathsf{U}_B^{\mathrm{var}} + \min\{\mathsf{U}_B^{\mathrm{per}}, K_B P_{r_B}^{\mathrm{ng}}\} + \mathsf{R}_B^{\mathrm{per}}, \tag{9.5}$$

$$0 \leq \mathsf{R}_B^{\mathrm{ng}} - \mathsf{R}_B^{\mathrm{per}} \leq \mathsf{U}_B^{\mathrm{var}}. \tag{9.6}$$

For every fixed M, in the prescribed order of limits,

$$\mathcal{H}_M(\mathsf{R}_B^{\mathrm{ng}} - \mathsf{R}_B^{\mathrm{per}}) \leq C_M B^{-95/384}, \tag{9.7}$$

$$\mathcal{E}_M \leq \mathcal{H}_M(\mathsf{R}_B^{\mathrm{per}}) + C_M B^{-1/384}. \tag{9.8}$$

The unchanged pointwise assertion (1.5) *is equivalent to $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{per}}) \to 0$ for every fixed M.*

Proof. Equation (9.5) is the positive scalar identity $x = \min(x, y) + (x - y)_+$ applied to $x = \mathsf{U}_B^{\mathrm{per}}$, combined with (8.8). Monotonicity and the unit Lipschitz bound of the positive-part function give (9.6). Corollary 8.3 then proves (9.7).

For completeness, the height of the middle term in (9.5) is at most

$$\mathcal{H}_M(K_B P_{r_B}) \leq C\varepsilon^{-1/32}\{\varepsilon^{1/16} + (1 + \varepsilon^{-1})\Delta_B\} = O(\varepsilon^{1/32}) = O(B^{-1/384}),$$

by Theorem 6.1. Insert the three nonnegative terms in Input statement 3.2, and use subadditivity of the ordered upper height. This proves (9.8) and sufficiency of the displayed limit. Necessity follows from the same inherited scalar equivalence and $0 \leq \mathsf{R}_B^{\mathrm{per}} \leq \mathsf{U}_B^{\mathrm{micro}}$. No essential supremum is interchanged with an integral, and the comparison pool has not been restricted to $\mathcal{P}_B$. □

The estimate still required to conclude (1.5) is

$$\lim_{B\to\infty}\limsup_{m\to\infty}\sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \left(\mathsf{U}^{\mathrm{per}}_{B,m}(t) - K_B P^{\mathrm{ng}}_{r_B,m}(t)\right)_+ = 0. \tag{9.9}$$

Theorems 8.2 and 9.1 prove an actual summed payment and a necessary physical persistence property. They do not prove (9.9). In particular a positive nearly constant density on a shrinking segment need not have small pointwise height merely because its roof integral is small. The original raw pointwise theorem, and not an averaged substitute, remains the organizing target.

10. Global uniqueness on a roof–angle fiber

The estimates in this section concern actual finite trajectories, not a continuation through a forbidden event. In particular the regular domain of a lifted itinerary need not be connected or convex. This distinction is important when the original section and occupation tests divide that domain into many components.

Normalize the initial lifted scatterer to $O_0 = 0$. Let $\mathcal{W}_m(R)$ denote the realizable lifted itineraries of m flights. This set is finite by the finite-horizon bound and the lattice geometry. Fix $w = (O_0, \dots, O_m) \in \mathcal{W}_m(R)$. For a physical trajectory write $Q_j = O_j + Rn_j$, let t_j be the counterclockwise rotation of n_j, and put

$$S(Q) = \sum_{j=0}^{m-1} |Q_{j+1} - Q_j|, \qquad g = R(-p_0, p_m), \qquad z = |g|.$$

As a function of the independent Cartesian contacts, S is convex. It is differentiable at every configuration in question because successive scatterer disks are disjoint. Write $\beta_0 = \beta_m = 1$ and $\beta_j = 2$ for $0 < j < m$.

Lemma 10.1 (Two-trajectory support inequalities)**.** *Let Q and $\widehat{Q}$ be two regular physical trajectories with the same lifted itinerary and radius. Set*

$$b_j = 1 - n_j \cdot \widehat{n}_j \geq 0, \qquad s_j = t_j \cdot \widehat{n}_j, \qquad s_E = (s_0, s_m).$$

Their lengths satisfy

$$S(\widehat{Q}) - S(Q) \geq R\sum_{j=0}^{m} \beta_j c_j b_j + g \cdot s_E, \tag{10.1}$$

$$S(Q) - S(\widehat{Q}) \geq R\sum_{j=0}^{m} \beta_j \widehat{c}_j b_j - \widehat{g} \cdot s_E. \tag{10.2}$$

Consequently

$$(\widehat{g} - g) \cdot s_E \geq R\sum_{j=0}^{m} \beta_j (c_j + \widehat{c}_j) b_j. \tag{10.3}$$

No interpolation by physical trajectories is used.

Proof. The physical reflection law and first variation in Cartesian coordinates give

$$\nabla_0 S = -c_0 n_0 - p_0 t_0, \qquad \nabla_j S = -2c_j n_j \ (0 < j < m), \qquad \nabla_m S = -c_m n_m + p_m t_m.$$

The convex support inequality is $S(\widehat{Q}) - S(Q) \geq \sum_j \nabla_j S(Q) \cdot (\widehat{Q}_j - Q_j)$. Since $\widehat{Q}_j - Q_j = R(\widehat{n}_j - n_j)$, its normal terms are $R\beta_j c_j b_j$ and its two tangential terms are $g \cdot s_E$. This proves (10.1). Reversing the roles of the two configurations leaves b_j unchanged and replaces s_j by $-s_j$. This proves (10.2). Adding the two inequalities proves (10.3). The support inequality is applied to a convex Cartesian function; it requires neither a convex angular chart nor a physical path joining the configurations. □

Theorem 10.2 (Global roof–angle uniqueness)**.** *For a fixed lifted itinerary, radius and direction $e \in S^1$, the physical length is strictly increasing among regular trajectories whose endpoint gradients are $g = ze$, $z > 0$, ordered by z. At a fixed z there is at most one such trajectory. In particular*

$$Q \longmapsto (F(Q), \arg g(Q)) \tag{10.4}$$

is globally injective on the noncritical regular trajectories of that itinerary. This remains true after any of the original exact event tests, and across all their disconnected regular components.

Proof. If $\widehat{g} = g$, the left side of (10.3) is zero. All c_j and $\widehat{c}_j$ are strictly positive, so every $b_j = 0$. Thus all physical contacts, and then all flight directions, coincide.

Now let $\widehat{g} = \widehat{z}e$, $g = ze$ and $\widehat{z} > z > 0$. The configurations cannot coincide. At least one b_j is therefore positive, and (10.3) gives

$$(\widehat{z} - z)e \cdot s_E \geq R \sum_j \beta_j (c_j + \widehat{c}_j) b_j > 0.$$

Hence $e \cdot s_E > 0$. The right side of (10.1) is then strictly positive, proving $S(\widehat{Q}) > S(Q)$. Equal roof and angle would force equal z, and the first part of the proof would force equality of trajectories. Restricting an injective map to the original exact event cannot increase its multiplicity. □

Lemma 10.3 (A bounded roof range for each physical itinerary)**.** *For any two regular physical trajectories with the same itinerary,*

$$|S(Q) - S(\widehat{Q})| \leq |Q_0 - \widehat{Q}_0| + |Q_m - \widehat{Q}_m| \leq 4R. \tag{10.5}$$

Proof. With its endpoint contacts fixed, a regular physical trajectory minimizes S over the product of the closed interior scatterer disks. Indeed, for a competitor Y_j in the jth disk, $n_j \cdot (Y_j - Q_j) \leq 0$; the convex support inequality and $\nabla_j S = -2c_j n_j$ therefore give the minimizing property. Use the interior contacts of $\widehat{Q}$ as a competitor for the endpoints of Q. Changing only the first and last endpoint changes the sum of flight lengths by at most the sum of their displacements, by the triangle inequality. This proves one direction of (10.5); interchanging the trajectories proves the other. The same triangle inequality covers $m = 1$. □

This range estimate does not control how many original occupation labels occur in a word domain. In particular it is not used to replace the exact occupation test by an unqualified interval test.

10.1. **The original physical sum at a target roof.** For the original section tests, $\chi \geq \lambda_\circ$ by (5.4). On each regular component, the map (10.4) is a local diffeomorphism. Theorem 10.2 shows that all those components of a fixed itinerary have disjoint images, not merely that each component has a nonsingular Jacobian. The equal-gradient argument also gives at most one regular critical trajectory per word, even when the gradient is zero. There are finitely many physical words at each finite count, so these critical points have zero collision area. Consequently, for any nonnegative Borel trajectory weight a, the original roof density has the representation

$$\frac{dF_{\#}(a\mathbf{1}_\Pi \nu_R^*)}{dt}(t) = \int_0^{2\pi} \sum_{w \in \mathcal{W}_m(R)} \mathbf{1}_{\{x_w(t,\vartheta) \text{ exists}\}} \frac{\omega(x_w) a(x_w)}{4\pi c_* \chi(x_w)} \, d\vartheta. \tag{10.6}$$

Here x_w is the unique point of the original exact event with the indicated word, roof and angle. A missing point contributes zero. The representation holds for almost every roof, in the original order $\sup_R \operatorname{ess\,sup}_t$. Angular chart overlap is removed before summation; the incident-component convention remains unchanged.

Proposition 10.4 (Monotone completion of the target-roof formula)**.** *Formula* (10.6) *is an identity of nonnegative measures for every nonnegative Borel weight, with extended values allowed before an integrability bound is proved. In particular it applies to the unbounded products* $\prod_{i\in S} c_i^{-1}$.

Proof. Fix m, R, n, k. Cover the noncritical regular physical domain in endpoint coordinates by countably many relatively compact local charts V_j on which $(F, \arg g)$ is a C^1 diffeomorphism into an unwrapped angular chart. Such a cover exists by the nonvanishing Jacobian and second countability. Subdivide as necessary so that both maps have bounded derivatives on each smaller chart. Disjointize the source charts by $B_j = V_j \setminus \bigcup_{k<j} V_k$. The sets B_j are Borel; no chart overlap is counted twice. Angular representatives are also split at a fixed seam, so their range is $[0, 2\pi)$.

For a nonnegative Borel weight h put

$$h_N = \min\{h, N\}\mathbf{1}_{\bigcup_{j\le N} B_j}.$$

This sequence increases to h on the regular noncritical domain. Apply ordinary change of variables on each of the first N charts, restricted to its Borel piece, and test against a nonnegative Borel function of (t, ϑ). The local inverse Jacobian, multiplied by the original collision density, is exactly $\omega/(4\pi c_*\chi)$. Local inverses are continuous and agree where they represent the same physical point, so their Borel restrictions define the displayed x_w without a selection ambiguity. Global uniqueness prevents different regular components of one word from producing the same target. Distinct words are summed, not identified.

Monotone convergence on the source, followed by Tonelli and monotone convergence on the target, now gives the identity for h as an extended nonnegative measure. Only after this passage is an upper pressure bound used to establish finiteness and an L^∞ density. This avoids assuming the integrability of the unbounded marked weight in order to prove its own change of variables.

The complement of the two-sided regular set has zero collision area. On the countably many charts just chosen, absolute continuity of the coordinate changes and their local Lipschitz property give zero target area for its image. The finitely many regular critical points also have zero source area. Removing these sets therefore does not change the weighted source measure. Fubini yields the claimed roof identity for almost every t. All null sets here are chosen at the fixed radius; no exceptional set common to every R is asserted or needed. For the finitely many marked sets at a fixed m, intersecting the corresponding full-measure roof sets provides a common version for their finite positive expansions. □

11. A complete marked pressure bound at the target roof

We now transfer the retained geometric pressure estimate to the roof fibers in (10.6). The previous momentum-fiber bound alone would not justify this transfer: different words at the same roof can have different endpoint momentum magnitudes. The global injectivity just proved supplies the missing multiplicity statement.

Let $j(x) = J^s T_R(x)$ on the full-area set M'_R regular in both infinite time directions, and let $j_l(x) = \prod_{v=0}^{l-1} j(T_R^v x)$, with $j_0 = 1$. The lifted physical one-flight alphabet has at most D members, uniformly in R. Define

$$P_l(R) = \sum_{w\in\mathcal{W}_l(R)} \sup_{x\in[w]_R\cap M'_R} j_l(x), \qquad P_0 = 1.$$

Input statement 11.1 (Uniform physical pressure and stable optical comparison)**.** On the circular family $R \in [9/20, 47/100]$, use the original finite-horizon flight bounds, positive curvature bounds and the original endpoint section cap. The stable Jacobian j is computed with Euclidean arclength along the true stable line in collision coordinates (s, φ), where $s = R\alpha$ and $p = \sin\varphi$. It is restricted to M'_R, the full-area set regular in both infinite directions. Its cocycle uses this

same convention on every block, including the empty block. The endpoint weight ω is still the derivative in the original (α, p) coordinates of (3.1); all coordinate factors are included in the uniform optical constant.

For the realizable lifted physical words just defined, the retained geometric theorems state

$$\sup_{R,l} P_l(R) \le Q_*, \qquad \omega(x) \le C_0 c_m(x) j_m(x), \qquad j(x)/c(x) \le C_1. \tag{11.1}$$

All three constants are independent of R, l and the original exact labels. These statements concern positive, unconditioned physical word sums; they are not a twisted or exact-label pressure theorem.

We take $Q_* \ge 1$ and $C_0, C_1 > 0$. These are the uniform geometric pressure theorem in historical module 241 and the stable optical comparison in module 238, with the physical-word domination in module 239. Their complete original sources are pinned in the proof ledger and copied in the provenance directory. They involve no occupation, displacement or roof test. Such tests are dropped only on the upper side of the estimates below. No new pressure estimate for a twisted or point-conditioned operator is assumed.

Lemma 11.2 (Removal of several marked stable factors)**.** *For $S = \{i_1 < \cdots < i_s\} \subset \{1, \dots, m-1\}$, at every regular trajectory in M_R',*

$$\frac{\omega(x)}{\prod_{i\in S} c_i(x)} \le C_0 C_1^s \prod_{v\in\{0,\dots,m-1\}\setminus S} j(T_R^v x). \tag{11.2}$$

Consequently, if at most one point x_w is chosen from each physical word domain, then

$$\sum_w \frac{\omega(x_w)}{\prod_{i\in S} c_i(x_w)} \le C_0 C_1^s D^s Q_*^{s+1}. \tag{11.3}$$

The choice of point may depend on the word and need not have the same momentum target for different words.

Proof. Use $c_m \le 1$ in (11.1), factor j_m into its individual steps, and apply $j(T_R^i x)/c_i(x) \le C_1$ at each marked index. This gives (11.2).

Delete the s marked steps of a word. The remaining blocks have lengths

$$i_1, \quad i_2 - i_1 - 1, \ \dots, \ i_s - i_{s-1} - 1, \quad m - i_s - 1.$$

For clarity, steps are numbered $0, \dots, m-1$, whereas the marks belong to $1, \dots, m-1$. Their indexing is

$$\underbrace{0, \dots, i_1 - 1}_{i_1} \mid i_1 \mid \underbrace{i_1 + 1, \dots, i_2 - 1}_{i_2 - i_1 - 1} \mid i_2 \mid \cdots \mid i_s \mid \underbrace{i_s + 1, \dots, m-1}_{m - i_s - 1}.$$

For example, $m = 7$ and $S = \{1, 2, 6\}$ leave blocks of lengths $(1, 0, 3, 0)$ and delete three physical symbols. Thus adjacent marks and a final mark are included, and every empty block contributes $P_0 = 1$. Some lengths can be zero. The blocks, translated to the initial fundamental scatterer, together with the s deleted physical step symbols determine the full lifted word uniquely. There are at most D^s choices for those symbols. Bound the remaining cocycle on each block by its own supremum and sum over all block words. Positivity permits adding combinations which do not produce a realizable full word. The resulting upper bound is $C_0 C_1^s D^s$ times the product of the $s+1$ corresponding $P_l(R)$'s, at most $C_0 C_1^s D^s Q_*^{s+1}$. For $S = \varnothing$ it is simply $C_0 P_m(R)$. This argument counts physical step symbols, never the infinitely many homogeneous strips at a removed incidence. □

Set

$$C_\star = \frac{C_0 Q_*}{2c_*\lambda_\circ}, \qquad K_\star = C_1 D Q_*. \tag{11.4}$$

Neither constant depends on the collision count, radius, exact labels, roof or marked indices.

Theorem 11.3 (Target-roof multiple-incidence insertion)**.** *For every finite m, every original exact event and every $S \subset \{1,\dots,m-1\}$, the nonnegative measure*

$$F_\# \left(\prod_{i\in S} c_i^{-1} \mathbf{1}_\Pi \nu_R^* \right)$$

has a roof density d_S satisfying

$$\|d_S\|_{L^\infty(dt)} \le C_\star K_\star^{|S|}. \tag{11.5}$$

The bound holds on the whole roof line. For arbitrary numbers $a_1,\dots,a_{m-1} \ge 0$, the density $d_{\boldsymbol{a}}$ with weight $\prod_{i=1}^{m-1}(1 + a_i/c_i)$ satisfies

$$\|d_{\boldsymbol{a}}\|_\infty \le C_\star \prod_{i=1}^{m-1} (1 + K_\star a_i) \le C_\star \exp\left(K_\star \sum_{i=1}^{m-1} a_i \right). \tag{11.6}$$

In particular the complete inherited linear incidence weight obeys

$$\left\| \frac{dF_\#(I_m \mathbf{1}_\Pi \nu_R^*)}{dt} \right\|_\infty \le C_\star \left(1 + K_\star \frac{2q_0}{1-q_0} \right). \tag{11.7}$$

These are finite-count unscaled height bounds, not m^{-2} local limit estimates.

Proof. At an almost-everywhere target (t,ϑ), Theorem 10.2 supplies at most one x_w per word. Apply (11.3) to precisely those points, use $\chi \ge \lambda_\circ$, and integrate (10.6) over the angle interval of length 2π. This gives (11.5) with the constants (11.4). Proposition 10.4 first establishes the nonnegative change-of-variables identity by truncation and monotone exhaustion; the present bound then proves finiteness and the density. At finite m the roof is confined to a bounded interval.

Expand $\prod_i (1 + a_i/c_i)$ into its finite sum over distinct marked sets. Apply (11.5) to each term. The sum of the upper bounds factors exactly as $C_\star \prod_i (1 + K_\star a_i)$. The exponential inequality uses $1 + x \le e^x$. Finally expand I_m linearly and use $\sum_i q_0^{d_i} \le 2q_0/(1-q_0)$. All original exact labels remain in the source throughout. Removing their tests only enlarges the positive word upper sums in the proof; it does not change the source normalization or infer a Gaussian factor from an unconditioned pressure bound. □

Corollary 11.4 (A factorial bound for changing incidence witnesses)**.** *Let $J \subset \{1,\dots,m-1\}$ and let $\eta_i \ge 0$ be arbitrary. On the original event define*

$$N_J(x) = \sum_{i\in J} \mathbf{1}_{\{c_i(x) < \eta_i\}}, \qquad A_J = \sum_{i\in J} \eta_i.$$

For every integer $s \ge 1$,

$$\left\| \frac{dF_\#(\mathbf{1}_{\{N_J \ge s\}} \mathbf{1}_\Pi \nu_R^*)}{dt} \right\|_\infty \le C_\star K_\star^s e_s((\eta_i)_{i\in J}) \le C_\star \frac{(K_\star A_J)^s}{s!}, \tag{11.8}$$

where $e_s = 0$ for $s > |J|$. The witnesses may change with the target roof and angle; no single witness is chosen for an entire segment.

Proof. At every regular state,

$$\mathbf{1}_{\{N_J\ge s\}} \le \binom{N_J}{s} = \sum_{\substack{S\subset J\\|S|=s}} \prod_{i\in S} \mathbf{1}_{\{c_i<\eta_i\}} \le \sum_{\substack{S\subset J\\|S|=s}} \prod_{i\in S} \eta_i/c_i.$$

Push this positive inequality through the original roof and apply (11.5). In the expansion of A_J^s each product with distinct indices occurs $s!$ times; all remaining terms are nonnegative. This proves the last inequality. The argument is finite-count and already completes the full physical sum, so it introduces no collision-count multiplier for the possible witnesses. □

12. Pointwise removal of deep and numerous incidence witnesses

We apply the marked pressure theorem to the unchanged positive persistent source of Section 9. The new payment concerns its value at the original target roof, rather than an average in that roof. Clearance witnesses require a separate estimate and will not be identified with incidence witnesses.

Lemma 12.1 (Domination of the surviving coefficient at every roof)**.** *For $\gamma \ge 1$, on every original physical segment and every point of its closed interval,*

$$0 \le c_{\mathrm{per}}^{\varepsilon,\gamma} \le f^{\varepsilon}(u,\theta) \le A(u,\theta). \tag{12.1}$$

Proof. Total variation bounds the decrement from an endpoint attaining c^{ε}, so $f^{\varepsilon}(u) \ge c^{\varepsilon} - \mathcal{V}^{\varepsilon}$. If $c_{\mathrm{per}} > 0$, its value is $c^{\varepsilon} - \gamma\mathcal{V}^{\varepsilon} \le c^{\varepsilon} - \mathcal{V}^{\varepsilon}$. If $c_{\mathrm{per}} = 0$, nonnegativity suffices. The last inequality is the unchanged bound $0 \le W^{\varepsilon} \le 1$. □

Thus a target-dependent Borel restriction of the persistent paired source is bounded by the corresponding restriction of the original uncut roof measure. This assertion concerns the surviving coefficient, not the full endpoint minimum on an arbitrary segment.

For $\varepsilon > 0$, use precisely the incidence witness thresholds already forced by the original guard:

$$E_i^{\varepsilon} = \{c_i < \eta_i^{\varepsilon}\}, \qquad \eta_i^{\varepsilon} = 2\varepsilon q_0^{d_i/4}, \quad d_i = \min(i, m-i), \quad 1 \le i < m. \tag{12.2}$$

Write $N^{\varepsilon} = \sum_{i=1}^{m-1} \mathbf{1}_{E_i^{\varepsilon}}$. There are at most two indices at each depth. Hence, for an integer $L \ge 1$,

$$\sum_{d_i\ge L} \eta_i^{\varepsilon} \le \frac{4\varepsilon q_0^{L/4}}{1-q_0^{1/4}}, \qquad \sum_{i=1}^{m-1} \eta_i^{\varepsilon} \le A_{\varepsilon} := \frac{4\varepsilon q_0^{1/4}}{1-q_0^{1/4}}. \tag{12.3}$$

For any Borel target event E, let

$$\mathsf{U}_B^{\mathrm{per}}[E](t) = \int c_{\mathrm{per}}^{\varepsilon(B),\gamma_B}(\theta)\mathbf{1}_{\{a<t<b\}}\widetilde{H}_B(\Psi(\theta,t))\mathbf{1}_E(\Psi(\theta,t))\,d\theta.$$

These are restrictions of the source in (8.8); no new normalization is used.

Theorem 12.2 (Uniform target-roof witness estimates)**.** *For every finite m, every B and integers $L, s \ge 1$,*

$$\|\mathsf{U}_B^{\mathrm{per}}[\bigcup_{d_i\ge L} E_i^{\varepsilon(B)}]\|_{\infty} \le \frac{4C_\star K_\star \varepsilon(B)}{1-q_0^{1/4}} q_0^{L/4}, \tag{12.4}$$

$$\|\mathsf{U}_B^{\mathrm{per}}[N^{\varepsilon(B)} \ge s]\|_{\infty} \le C_\star \frac{(K_\star A_{\varepsilon(B)})^s}{s!}. \tag{12.5}$$

The norms are essential suprema on the whole roof line; constants are uniform in the radius and original exact labels.

Fix any $\eta > 0$ and define, at finite count,

$$\begin{aligned} L_m &= \left\lceil \frac{4(2+\eta)}{|\log q_0|} \log(m+1) \right\rceil, \\ s_m &= \left\lceil \frac{4\log(m+e)}{\log\log(m+e^e)} \right\rceil, \\ \mathcal{D}_{B,m} &= \bigcup_{d_i \geq L_m} E_i^{\varepsilon(B)}, \qquad \mathcal{A}_{B,m} = \{N^{\varepsilon(B)} \geq s_m\}, \qquad \mathcal{Z}_{B,m} = \mathcal{D}_{B,m} \cup \mathcal{A}_{B,m}. \end{aligned} \tag{12.6}$$

An empty union contributes zero. For every fixed outer B and M,

$$\mathcal{H}_M(\mathsf{U}_B^{\mathrm{per}}[\mathcal{Z}_{B,m}]) = 0. \tag{12.7}$$

More precisely the deep part satisfies, at every count,

$$\sup_{R,n,k} m^2 \|\mathsf{U}_B^{\mathrm{per}}[\mathcal{D}_{B,m}]\|_\infty \leq C\varepsilon(B)(m+1)^{-\eta}. \tag{12.8}$$

Proof. By Lemma 12.1, each source in the first two assertions is bounded by the original uncut density restricted to the same target event. Apply Corollary 11.4 with $s = 1$ to the depth tail and with general s to all indices, then use (12.3). This proves the two finite-count bounds after the physical sum, with no selected uniform witness.

They hold for every L and s at every m. Substituting L_m in the first gives $q_0^{L_m/4} \leq (m+1)^{-(2+\eta)}$, which proves (12.8). For fixed B, put $a = K_\star A_{\varepsilon(B)} < \infty$. The elementary factorial bound $s! \geq (s/e)^s$ gives

$$m^2 \frac{a^{s_m}}{s_m!} \leq \exp\{2\log m + s_m(1 + \log a - \log s_m)\} = m^{-2+o(1)} \longrightarrow 0.$$

Here $s_m \log s_m = (4+o(1))\log m$ and $s_m = o(\log m)$. The case $a = 0$ is immediate. If $s_m > m-1$, the event is empty. Subadditivity and positivity complete the proof of (12.7).

The depth and multiplicity thresholds in this proof are substitutions into finite-count inequalities valid simultaneously for all their values. No fixed-window spectral estimate is evaluated at a collision-dependent window. The original outer quantities $\varepsilon(B), h_B, r_B, \gamma_B, K_B$ remain fixed before the collision limit. Thus the proof does not exchange that limit with any outer width or band. □

12.1. **The retained source after this pointwise payment.** Define

$$\begin{aligned} \mathsf{U}_B^{\mathrm{loc}} &= \mathsf{U}_B^{\mathrm{per}}[\mathcal{Z}_{B,m}^c], \\ \mathsf{U}_B^{\mathrm{exc}} &= \mathsf{U}_B^{\mathrm{per}}[\mathcal{Z}_{B,m}], \qquad \mathsf{R}_B^{\mathrm{loc}} = (\mathsf{U}_B^{\mathrm{loc}} - K_B P_{r_B}^{\mathrm{ng}})_+. \end{aligned} \tag{12.9}$$

The superscript loc records localization of all incidence witnesses to the two collision-end layers; it makes no claim that clearance witnesses are localized to those layers. The comparison pool is the same full nongrazing pool, without a source-dependent restriction.

Corollary 12.3 (Exact preservation and collision-zero residual difference)**.** *At finite count the original persistent source satisfies*

$$\mathsf{U}_B^{\mathrm{per}} = \mathsf{U}_B^{\mathrm{exc}} + \mathsf{U}_B^{\mathrm{loc}}, \qquad 0 \leq \mathsf{R}_B^{\mathrm{per}} - \mathsf{R}_B^{\mathrm{loc}} \leq \mathsf{U}_B^{\mathrm{exc}}. \tag{12.10}$$

For every fixed B, M,

$$\mathcal{H}_M(\mathsf{R}_B^{\mathrm{per}} - \mathsf{R}_B^{\mathrm{loc}}) = 0, \qquad \mathcal{E}_M \leq \mathcal{H}_M(\mathsf{R}_B^{\mathrm{loc}}) + C_M B^{-1/384}. \tag{12.11}$$

The unchanged raw pointwise assertion is equivalent to $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{loc}}) \to 0$ for every fixed M.

Proof. The source partition is an identity of nonnegative original submeasures. Monotonicity and the unit Lipschitz bound of the positive-part map give its residual inequality. Apply (12.7) and then Theorem 9.3. For necessity use $0 \le \mathsf{R}_B^{\mathrm{loc}} \le \mathsf{U}_B^{\mathrm{micro}}$ and the retained scalar equivalence. Every removed term has been retained in the identity and paid in the target essential-height topology. □

For all sufficiently large B, $2\varepsilon(B) < c_\circ$, so the original endpoint section cap rules out incidence witnesses at $i = 0, m$. At a target point with positive surviving coefficient the original guard still requires an incidence or a clearance witness, by Theorem 9.1. On the retained part, every incidence witness has depth below L_m, and fewer than s_m such witnesses occur. A clearance witness may occur at any original flight, with its original positive weight. These alternatives allow a witness which changes with the roof; they assert no common witness on an entire physical segment.

The precise remaining estimate is

$$\lim_{B\to\infty} \limsup_{m\to\infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} (\mathsf{U}_{B,m}^{\mathrm{loc}}(t) - K_B P_{r_B,m}^{\mathrm{ng}}(t))_+ = 0. \tag{12.12}$$

The target-roof pressure theorem proves a new physical weighted summation and the pointwise removal in Theorem 12.2; it does not prove (12.12). In particular the all-incidence bound $C\varepsilon$ furnished by (12.4) with $L = 1$ is unscaled. At a fixed outer B it cannot be multiplied by m^2 and declared small. Nor does an inverse-incidence insertion estimate an inverse third-disk clearance. The remaining source includes the original shallow incidence witnesses, unvisited tangencies, critical entrances, section cuts and capacity deficiencies which are not paid by the proved estimates. No change of the original theorem or of its arithmetic classes is made in stating this boundary.

13. Joint depth and severity of incidence witnesses

The factorial estimate treats the witness thresholds only through their sum. It therefore forgets that there are at most two collision indices at any depth. It also forgets how much an individual incidence lies below its guard threshold. We retain both pieces of information. All estimates below concern the same target-roof measure as Theorem 11.3, with its original exact labels. They use its inherited geometric inputs, not a new conditional mixing or local-limit assumption.

Put

$$r = q_0^{1/4}, \qquad \alpha = -\log r = \tfrac{1}{4}|\log q_0|, \qquad b_\varepsilon = 4K_\star\varepsilon/\alpha. \tag{13.1}$$

For a regular trajectory define its incidence-witness set and load by

$$\begin{aligned} J_\varepsilon(x) &= \{i : 1 \le i < m,\ c_i(x) < 2\varepsilon r^{d_i}\}, \\ \mathscr{L}_\varepsilon(x) &= \sum_{i\in J_\varepsilon(x)} \log\frac{2\varepsilon}{c_i(x)} \\ &= \alpha \sum_{i\in J_\varepsilon(x)} d_i + \sum_{i\in J_\varepsilon(x)} \log\frac{2\varepsilon r^{d_i}}{c_i(x)}. \end{aligned} \tag{13.2}$$

The empty sum is zero. Both summands in the last line are nonnegative. In particular this is not a signed logarithmic weight, and it introduces no clearance factor. The Borel witness set is allowed to vary with the roof and angle.

For a nonnegative Borel weight h, write

$$D[h](t) = \frac{dF_\#(h\mathbf{1}_\Pi \nu_R^*)}{dt}(t)$$

when this density is defined. The positive identities of the preceding section give a common version at fixed m, R, n, k. The next theorem also proves existence and finiteness for its weights.

Theorem 13.1 (Exponential moment of the original incidence load)**.** *For every finite collision count, every $\varepsilon > 0$ and $0 < \lambda < 1$,*

$$\left\|D[e^{\lambda \mathscr{L}_\varepsilon}]\right\|_\infty \le C_\star \exp\left\{\frac{4K_\star \varepsilon}{e^{\alpha(1-\lambda)} - 1}\right\}. \tag{13.3}$$

Consequently, for $u > b_\varepsilon$,

$$\left\|D[\mathbf{1}_{\{\mathscr{L}_\varepsilon \ge u\}}]\right\|_\infty \le C_\star \min\left\{1, \exp[-u + 2\sqrt{b_\varepsilon u}]\right\}. \tag{13.4}$$

Both estimates are uniform in the radius, exact labels and target roof. They remain valid with any further Borel restriction of the original measure, including the persistent source through Lemma 12.1.

Proof. Set $y_i = 2\varepsilon/c_i$. At a witness, $y_i > r^{-d_i}$, whence

$$y_i^\lambda = y_i y_i^{-(1-\lambda)} \le y_i r^{(1-\lambda)d_i}.$$

At every regular trajectory we therefore have the positive inequality

$$\begin{aligned} e^{\lambda \mathscr{L}_\varepsilon} &= \prod_{i=1}^{m-1} \left[1 + (y_i^\lambda - 1)\mathbf{1}_{\{i \in J_\varepsilon\}}\right] \\ &\le \prod_{i=1}^{m-1} \left(1 + \frac{2\varepsilon r^{(1-\lambda)d_i}}{c_i}\right). \end{aligned}$$

Apply the finite product insertion (11.6). There are at most two indices at every positive depth, so

$$\sum_{i=1}^{m-1} 2\varepsilon r^{(1-\lambda)d_i} \le 4\varepsilon \sum_{d \ge 1} r^{(1-\lambda)d} = \frac{4\varepsilon}{e^{\alpha(1-\lambda)} - 1}.$$

This proves (13.3) after the complete physical sum. Every product is finite before this last geometric upper bound is used; no independence of collisions is assumed.

Positivity gives $\mathbf{1}_{\{\mathscr{L}_\varepsilon \ge u\}} \le e^{-\lambda u} e^{\lambda \mathscr{L}_\varepsilon}$. Writing $v = 1 - \lambda$ and using $e^{\alpha v} - 1 \ge \alpha v$, we obtain, for every $0 < v < 1$,

$$\left\|D[\mathbf{1}_{\{\mathscr{L}_\varepsilon \ge u\}}]\right\|_\infty \le C_\star \exp\{-u + vu + b_\varepsilon / v\}.$$

When $u > b_\varepsilon$, take $v = \sqrt{b_\varepsilon / u}$. The independent bound $\|D[1]\|_\infty \le C_\star$ follows from the empty marked set and supplies the minimum in (13.4). Borel restrictions decrease a positive measure. For the persistent source the pointwise coefficient inequality in Lemma 12.1 supplies this decrease. □

Corollary 13.2 (Accumulated depth and severe shallow incidences)**.** *For $v > 0$, each of the events*

$$\left\{\alpha \sum_{i \in J_\varepsilon} d_i \ge v\right\}, \qquad \left\{\sum_{i \in J_\varepsilon} \log \frac{2\varepsilon r^{d_i}}{c_i} \ge v\right\}$$

is contained in $\{\mathscr{L}_\varepsilon \ge v\}$ and has the bound (13.4) whenever $v > b_\varepsilon$. An incidence witness satisfying $c_i \le 2\varepsilon e^{-v}$ is also in that event, irrespective of its depth.

Proof. Use the nonnegative decomposition in (13.2). For the last assertion, the single indicated summand is at least v. □

Thus a single sufficiently severe witness in an end layer can be paid. The conclusion does not require either a deep witness or the large witness count of Theorem 12.2. It does not assert a bound for an unvisited third-disk tangency.

14. Quadratic tails for the number of incidence witnesses

There is also an explicit finite-count improvement which retains all threshold locations, rather than passing to their total sum. The calculation is deterministic; it makes no independence claim about the billiard or its symbolic itinerary.

For $0 < r < 1$ write

$$(r;r)_a = \prod_{j=1}^{a}(1-r^j), \quad (r;r)_0 = 1, \quad (r;r)_\infty = \prod_{j=1}^{\infty}(1-r^j) > 0, \quad f(s) = \left\lfloor \frac{(s+1)^2}{4} \right\rfloor.$$

The infinite product is positive because the tail of $\sum_j -\log(1-r^j)$ is bounded by a constant times $\sum_j r^j$.

Lemma 14.1 (The two-end depth polynomial)**.** *For $0 \le s \le m-1$, put $E_{m,s}(r) = e_s((r^{d_i})_{1\le i<m})$. Then*

$$r^{f(s)} \le E_{m,s}(r) \le \sum_{a=0}^{s} \frac{r^{a(a+1)/2+(s-a)(s-a+1)/2}}{(r;r)_a (r;r)_{s-a}} \le \frac{(s+1)r^{f(s)}}{(r;r)_\infty^2}. \tag{14.1}$$

For $s > m-1$ the finite polynomial coefficient is zero and only the upper inequalities are asserted. In particular the coefficient has quadratic logarithmic rate, uniformly over $m \ge s+1$:

$$\log E_{m,s}(r) = -\frac{\alpha}{4}s^2 + O_r(s + \log(s+1)). \tag{14.2}$$

This is sharpness of the depth polynomial, not a lower bound for a physical roof density.

Proof. The multiset of finite collision depths is contained in two copies of $1, 2, 3, \ldots$. Coefficientwise positivity bounds its generating polynomial by $\prod_{d\ge1}(1+zr^d)^2$. To compute one factor, for $a \ge 1$ write $1 \le d_1 < \cdots < d_a$ as $d_j = j + h_j$, where $0 \le h_1 \le \cdots \le h_a$. Setting $g_1 = h_1$ and $g_j = h_j - h_{j-1}$ gives

$$\sum_{1\le d_1<\cdots<d_a} r^{d_1+\cdots+d_a} = r^{a(a+1)/2} \prod_{j=1}^{a} \sum_{g_j\ge0} r^{(a-j+1)g_j} = \frac{r^{a(a+1)/2}}{(r;r)_a}.$$

The empty factor contributes one. Multiplying the two positive series proves the middle sum in (14.1); all infinite passages are monotone limits of nonnegative coefficients. The minimum of its exponent over a is $f(s)$, attained by the two most balanced choices. Each denominator is at least $(r;r)_\infty^2$, giving the last inequality.

For the lower bound take the s smallest available depths. If $s = 2k$ these consist of two copies of $1, \ldots, k$ and have sum $k(k+1)$. If $s = 2k+1$ take these and one copy of $k+1$, with sum $(k+1)^2$. These depths are available whenever $m \ge s+1$, including the single central depth when m is even. Their product is one term of $E_{m,s}$. Taking logarithms of the two bounds proves (14.2). □

Theorem 14.2 (Quadratic witness-count tail at the target roof)**.** *With $N^\varepsilon = |J_\varepsilon|$, for every $s \ge 1$,*

$$\left\| D[\mathbf{1}_{\{N^\varepsilon \ge s\}}] \right\|_\infty \le \frac{C_\star (s+1)(2K_\star\varepsilon)^s}{(r;r)_\infty^2}\, r^{f(s)}. \tag{14.3}$$

The event is empty for $s > m-1$. The same estimate holds for the persistent source. For every fixed B and every $\eta > 0$, take

$$\sigma_m = \left\lceil 2\sqrt{\frac{(2+\eta)\log(m+1)}{\alpha}} \right\rceil. \tag{14.4}$$

Then the source with at least σ_m incidence witnesses has zero m^2-normalized essential height, with finite-count rate $m^{-\eta+o_B(1)}$.

Proof. Use the first, unrelaxed inequality in (11.8), with thresholds $2\varepsilon r^{d_i}$, and apply Lemma 14.1. Since $f(s) \geq s^2/4$, the logarithm of m^2 times the upper bound at $s = \sigma_m$ is at most

$$-\eta\log(m+1) + \sigma_m \log(2K_\star \varepsilon(B)) + \log(\sigma_m + 1) + \log \frac{C_\star}{(r;r)_\infty^2}.$$

For fixed B the last three terms are $O_B(\sqrt{\log(m+1)} + \log\log(m+2))$. This proves the claim. No outer width is changed. □

The square-root logarithmic threshold is strictly smaller in order than the logarithmic-over-logarithmic threshold of the earlier factorial bound. The estimate depends on the actual two-end depth geometry, not on additional cancellation or a count of physical words.

15. Positive payment and the remaining low-load source

We now apply the joint-load theorem to the original localized persistent source. All quantities $\varepsilon(B), h_B, r_B, \gamma_B, K_B$ are fixed at the outer stage. A load level used below classifies an original trajectory at finite collision count; it is not a spectral, roof, residence or comparison window.

Fix $\eta > 0$ and set

$$u_m = (2+\eta)\log(m+1), \qquad \mathcal{Q}_{B,m} = \{\mathscr{L}_{\varepsilon(B)} \geq u_m\}. \tag{15.1}$$

Keep the previously paid set $\mathcal{Z}_{B,m}$ from (12.6). Define two restrictions of the same positive source, with no new renormalization:

$$\begin{aligned} \mathsf{U}_B^{\mathrm{load}} &= \mathsf{U}_B^{\mathrm{per}}[\mathcal{Z}_{B,m}^c \cap \mathcal{Q}_{B,m}], \\ \mathsf{U}_B^{\mathrm{low}} &= \mathsf{U}_B^{\mathrm{per}}[\mathcal{Z}_{B,m}^c \cap \mathcal{Q}_{B,m}^c], \\ \mathsf{R}_B^{\mathrm{low}} &= (\mathsf{U}_B^{\mathrm{low}} - K_B P_{r_B}^{\mathrm{ng}})_+. \end{aligned} \tag{15.2}$$

The comparison pool is the full original nongrazing pool, not a source-selected pool. The notation low refers only to incidence load; clearance witnesses remain at arbitrary depth.

Theorem 15.1 (A new collision-normalized positive payment)**.** *At every finite count,*

$$\mathsf{U}_B^{\mathrm{per}} = \mathsf{U}_B^{\mathrm{exc}} + \mathsf{U}_B^{\mathrm{load}} + \mathsf{U}_B^{\mathrm{low}}, \qquad 0 \leq \mathsf{R}_B^{\mathrm{loc}} - \mathsf{R}_B^{\mathrm{low}} \leq \mathsf{U}_B^{\mathrm{load}}. \tag{15.3}$$

For every fixed B, M, the load term has zero ordered height:

$$\mathcal{H}_M(\mathsf{U}_B^{\mathrm{load}}) = 0. \tag{15.4}$$

More explicitly, whenever $u_m > b_{\varepsilon(B)}$,

$$\sup_{R,n,k} m^2 \|\mathsf{U}_B^{\mathrm{load}}\|_\infty \leq C_\star \exp\left\{-\eta\log(m+1) + 2\sqrt{b_{\varepsilon(B)}(2+\eta)\log(m+1)}\right\}. \tag{15.5}$$

Consequently

$$\mathcal{H}_M(\mathsf{R}_B^{\mathrm{loc}} - \mathsf{R}_B^{\mathrm{low}}) = 0, \qquad \mathcal{E}_M \leq \mathcal{H}_M(\mathsf{R}_B^{\mathrm{low}}) + C_M B^{-1/384}. \tag{15.6}$$

The original assertion (1.5) *is equivalent to $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{low}}) \to 0$ for every fixed M.*

Proof. The three sets in (15.3) disjointly partition the original persistent source. Thus its first equality holds before pushforward and remains exact afterwards. Monotonicity and the unit Lipschitz bound of the positive-part map prove the residual inequality with the same capacity on both sides.

Lemma 12.1 and Theorem 13.1 bound $\mathsf{U}_B^{\mathrm{load}}$ by $C_\star \exp[-u_m + 2\sqrt{b_{\varepsilon(B)} u_m}]$ once $u_m > b_{\varepsilon(B)}$. Multiplication by m^2, followed by $m^2 \le (m+1)^2$, gives (15.5). At fixed B its exponent is $-\eta \log(m+1) + O_B(\sqrt{\log(m+1)})$, tending to minus infinity. The finitely many smaller counts do not affect the collision limsup. This proves (15.4) uniformly in all the stated inner suprema.

Apply the residual inequality and Corollary 12.3 to obtain (15.6). For necessity use $0 \le \mathsf{R}_B^{\mathrm{low}} \le \mathsf{U}_B^{\mathrm{micro}}$ and the original scalar equivalence. For sufficiency use (15.6) and then send B to infinity. This is the original order of limits. The optimizing parameter in the proof of (13.4) was chosen inside a finite-count inequality valid for every $0 < \lambda < 1$; it is not inserted into a theorem which holds only at a previously fixed outer width. □

Corollary 15.2 (Explicit support restrictions after payment)**.** *At any regular target point contributing to* $\mathsf{U}_B^{\mathrm{low}}$*, put* $J = J_{\varepsilon(B)}$ *and* $N = |J|$*. Then*

$$\alpha \sum_{i \in J} d_i < u_m, \qquad \alpha f(N) < u_m, \qquad \prod_{i \in J} \frac{c_i}{2\varepsilon(B)} > e^{-u_m}. \tag{15.7}$$

In particular $N < 2\sqrt{u_m/\alpha}$ *and every individual witness has* $c_i > 2\varepsilon(B) e^{-u_m}$*. All these restrictions are in addition to the retained v111 incidence localization. They impose no clearance localization.*

Proof. The load is strictly less than u_m on the retained set. Its nonnegative decomposition gives the first inequality. The sum of the depths of any N indices is at least $f(N)$, by the construction in Lemma 14.1; this proves the second. Exponentiating minus the definition of the load proves the product inequality. Since $f(N) \ge N^2/4$ and all load summands are nonnegative, the remaining assertions follow, including the empty-witness case. □

For sufficiently large B, the original section cap excludes incidence witnesses at the two endpoint collisions. The persistent guard then still forces an interior incidence witness or an original third-disk clearance witness. A state with $N = 0$ is not removed by this revision: the guard may be witnessed entirely by clearance. Critical entrances and section cuts carrying the original positive coefficient are likewise not deleted by their names.

The exact remaining target is

$$\lim_{B \to \infty} \limsup_{m \to \infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t \in \mathcal{C}_{m,R,n,k}(M)} (\mathsf{U}_{B,m}^{\mathrm{low}}(t) - K_B P_{r_B,m}^{\mathrm{ng}}(t))_+ = 0. \tag{15.8}$$

Theorems 13.1–15.1 prove new finite-count concentration and positive payments in the required normalized topology. They do not prove (15.8). Low-load shallow incidences, arbitrary-depth clearance (including unvisited tangency), and capacity-deficient positive source remain in that equation. The original physical section, guard, arithmetic reference and exact-label pointwise assertion have not been weakened. The distinction between this remaining assertion and the proved payments is part of the theorem statement, not an inference from software diagnostics or an editorial change of topic.

16. Clearance-to-roof coarea without a fixed-clearance density bound

We keep the physical parameters and every event in Section 1. In this section u is the clearance coordinate of Section B, not physical time. Physical time is denoted by t throughout. Fix

R, m, n, k, a flight $0 \le j < m$, a sign σ, and an actual regular marked itinerary. On a local angular lift put

$$\mathcal{Q}_j(\vartheta, u) = (\alpha_0, \alpha_m)(X_{R,j}(\vartheta, \sigma, u)), \qquad S(\vartheta, u) = F(\mathcal{Q}_j(\vartheta, u)).$$

The point used as a mark is an unvisited near miss, not an additional reflection. Every derivative below is taken on the same designated regular itinerary. In particular m, j and the physical roof are not changed by this parametrization.

Theorem 16.1 (The long-record Jacobian and its critical Hessian)**.** *Let $A = D\mathcal{Q}_j$, let $g = \nabla_q F$, and let $G = D_q^2 F$. On every such chart,*

$$|\det A| = (R\omega)^{-1}, \tag{16.1}$$

$$|\det D_{(\vartheta,u)}(S, \arg g)| = \chi/(R\omega) \quad (g \neq 0), \tag{16.2}$$

$$D^2 S = A^{\mathsf{T}} G A + \sum_{a=1}^{2} g_a D^2 (\mathcal{Q}_j)_a. \tag{16.3}$$

At a regular critical roof, $g = 0$, the Hessian $D^2 S$ is positive definite and

$$\sqrt{\det D^2 S} = \frac{\sqrt{\det G}}{R\omega}. \tag{16.4}$$

These are identities for each finite physical collision count, with all coordinate and section-normalization factors displayed. In particular the Hessian term containing g cannot be dropped away from a critical roof.

Proof. The collision map preserves $d\alpha\, dp$ at fixed R. The signed line map at collision j has absolute Jacobian R, by (B.4). Composition with T_R^j therefore still has absolute Jacobian R with respect to the initial collision coordinates. The map $(\alpha_0, p_0) \mapsto (\alpha_0, \alpha_m)$ has absolute Jacobian ω^{-1} by the definition of ω. Dividing the two determinants gives (16.1). It uses the actual inverse iterate and is independent of where the mark occurs. On a noncritical chart the determinant of $q \mapsto (F, \arg g)$ is χ; the chain rule gives (16.2). Twice differentiating $F \circ \mathcal{Q}_j$ gives (16.3), component by component. At $g = 0$ its last term vanishes, and the retained positive-definite optical Hessian and the invertibility of A give the assertion. Taking the determinant and using (16.1) proves (16.4).

As a normalization check, the line-coordinate density is $(4\pi R c_*)^{-1}$. Dividing it by the determinant in (16.2) gives exactly $\omega/(4\pi c_* \chi)$, the original target-roof/Gauss-angle density. There is no additional factor R, R^{-1} or c_*^{-1}. All chart restrictions and exact labels remain on the source side of this change of variables. □

Corollary 16.2 (Uniform determinant growth on regular marked charts)**.** *With the retained constants $\varrho = 47/59$ and $\lambda_\circ$, all regular marked charts with the original endpoint cap satisfy*

$$|\det D\mathcal{Q}_j| \ge \frac{600}{2209}\varrho^{-(m-1)}, \qquad |\det D(S, \arg g)| \ge \frac{600\lambda_\circ}{2209}\varrho^{-(m-1)} \quad (g \neq 0).$$

At a critical roof the same lower bound as in the second inequality holds for $\sqrt{\det D^2 S}$. The constants are uniform in the original radius interval, the exact labels, the flight index, and the physical word. These determinant bounds do not bound individual singular values, a chart's condition number, or the complete physical sum of inverse determinants.

Proof. The retained optical estimate is $\omega \le (47/6)\varrho^{m-1}$, and $R \le 47/100$. Insert these in (16.1). Then use $\chi \ge \lambda_\circ$ and $\sqrt{\det G} \ge \lambda_\circ$ in (16.2) and (16.4). The identity $6/(47(47/100)) = 600/2209$ fixes the common constant. □

The following elementary estimate is recorded with proof to avoid importing an unstated finite-type or regularity theorem.

Lemma 16.3 (A one-dimensional band estimate)**.** *Let $f \in C^\ell(I)$ on an interval, with integer $\ell \geq 1$ and $|f^{(\ell)}| \geq \kappa > 0$. For $h > 0$,*

$$|\{x \in I : a < f(x) < a + h\}| \leq 2\ell(h/\kappa)^{1/\ell}. \tag{16.5}$$

The estimate also bounds any Borel subset of that band.

Proof. First restrict the band to a compact subinterval and let its measure be $L > 0$. Its cumulative Lebesgue measure is continuous and one-Lipschitz. The $\ell+1$ quantiles with successive mass gaps L/ℓ can be chosen in its closed support; their successive coordinate gaps are at least L/ℓ. Equivalently one can use interior approximate quantiles and pass to their limits. Write them as $x_0 < \cdots < x_\ell$. Continuity puts $f(x_i)$ in $[a, a + h]$. The divided-difference mean-value theorem, proved by subtracting the interpolating polynomial and applying Rolle's theorem ℓ times, gives some $\xi \in (x_0, x_\ell)$ for which

$$\frac{f^{(\ell)}(\xi)}{\ell!} = \sum_{i=0}^{\ell} \frac{f(x_i) - a}{\prod_{v \neq i}(x_i - x_v)}.$$

The subtraction of a is legitimate because the divided difference of a constant is zero. The gap bound makes the absolute denominator at least $(L/\ell)^\ell i!(\ell - i)!$. Hence

$$\kappa \leq h(\ell/L)^\ell \sum_{i=0}^{\ell} \binom{\ell}{i} = h(2\ell/L)^\ell.$$

Rearrange, then exhaust the original interval by compact subintervals. This also handles a band not initially known to have finite measure. No statement about the number of physical words is used in the proof. □

Theorem 16.4 (A positive finite-type clearance slice)**.** *Suppose $S \in C^{\ell+1}$ on a neighbourhood of $I \times [0, \delta_0]$, and*

$$0 < b_f \leq |S_u| \leq B_f, \qquad |\partial_\vartheta^\ell S(\vartheta, 0)| \geq \kappa_f > 0.$$

The sign of S_u is fixed on this rectangle. Let E be any Borel subset of it and $0 \leq h \leq H_f$ a pointwise bounded Borel weight. For $0 < \delta \leq \delta_0$, the measure

$$S_\#\big(h\mathbf{1}_E\mathbf{1}_{\{0<u<\delta\}}\, d\vartheta\, du\big)$$

has a density $D_{E,h,\delta}$ with

$$\|D_{E,h,\delta}\|_\infty \leq \frac{2\ell H_f}{b_f}\left(\frac{B_f\delta}{\kappa_f}\right)^{1/\ell}. \tag{16.6}$$

In the physical marked record one takes $h = (4\pi R c_*)^{-1}$, or a bounded multiplier of this weight, and E is the original physical domain with all exact labels. The constants must be established on that actual chart; the theorem does not assert their uniformity in m or their summability over all charts.

Proof. At fixed ϑ, $u \mapsto S(\vartheta, u)$ is strictly monotone. Let $U_t(\vartheta)$ denote its unique inverse when it exists. The ordinary one-dimensional substitution, first against nonnegative roof tests and then by Tonelli, gives

$$D_{E,h,\delta}(t) = \int_I \frac{h(\vartheta, U_t)\mathbf{1}_E(\vartheta, U_t)\mathbf{1}_{\{0<U_t<\delta\}}}{|S_u(\vartheta, U_t)|}\, d\vartheta.$$

Missing inverses contribute zero. This argument proves absolute continuity, not merely an estimate conditional on its existence. If $S_u > 0$ and the inverse lies in $(0, \delta)$, then $t - B_f\delta < S(\vartheta, 0) < t$.

For $S_u < 0$ the corresponding band is $t < S(\vartheta, 0) < t + B_f\delta$. Both have width $B_f\delta$. Bound the integrand by H_f/b_f and apply Lemma 16.3. Borel cuts, including section cuts, only restrict the integral and never create a chart-count multiplier. The same proof on bounded compact rectangles followed by monotone exhaustion gives the assertion for open intervals and Borel weights. □

Theorem 16.5 (Integrated simple folds, with a sharp local constant)**.** *Assume that on $I \times (0, \delta_0)$ one has $S_{\vartheta\vartheta} \geq a_f > 0$ and $S_u \geq b_f > 0$, and that for each u the unique angular critical point $\vartheta_0(u)$ exists in I. For the same positive weights and selectors as above,*

$$\|D_{E,h,\delta}\|_\infty \leq 2H_f\sqrt{\frac{2\delta}{a_f b_f}}. \tag{16.7}$$

The assertion also applies after reversing the roof sign or the clearance-coordinate orientation when the corresponding derivative signs are reversed. It does not require a bounded roof density at each fixed clearance.

Proof. Put $v(u) = S(\vartheta_0(u), u)$. The implicit function theorem and the critical-point equation give $v'(u) = S_u(\vartheta_0(u), u) \geq b_f$. For fixed u there are at most two angular roots of $S(\vartheta, u) = t$, and

$$|S_\vartheta(\vartheta, u)|^2 \geq 2a_f(S(\vartheta, u) - v(u)).$$

Indeed differentiate S_ϑ^2 along each monotone side and integrate from its zero at ϑ_0. On the left side the integration orientation and the sign of S_ϑ reverse together, giving the same inequality. One-dimensional substitution on those two sides gives a fixed-u density bounded by $H_f\sqrt{2/a_f}(t - v(u))^{-1/2}$ for $t > v(u)$ and zero below. The angular critical point itself has zero angular measure. The bound is allowed to be infinite at $t = v(u)$.

For $0 < u < \delta$, if t is in the range of v, let u_t be its unique inverse. The contributing values satisfy $u < u_t$, and $t - v(u) \geq b_f(u_t - u)$. If $t \geq v(\delta)$, replace u_t by δ for an upper bound; if $t \leq v(0)$ there is no contribution. In every case

$$\int_0^\delta \mathbf{1}_{\{t > v(u)\}}(t - v(u))^{-1/2}\, du \leq 2\sqrt{\delta/b_f}.$$

Integrate the fixed-u estimate by Tonelli to obtain (16.7). Restricting by E preserves the inequality. For $S(\vartheta, u) = \vartheta^2 + u$, $h = 1$, and the whole real angular line, the integrated density is

$$2\{\sqrt{t_+} - \sqrt{(t - \delta)_+}\},$$

whose maximum is $2\sqrt{\delta}$. Thus the constant is attained when $a_f = 2$ and $b_f = 1$. This elementary normal form checks the local analytic bound only; it is not substituted for a billiard trajectory or for the joint-record distribution. □

16.1. **One-copy summation and the remaining uniformity.** On a countable collection of admissible physical charts, first replace their source sets by disjoint Borel differences. The preceding density formulae then add exactly. Their positive upper bounds add as well, with an extended value permitted until a finite bound has been proved. Two impact signs and distinct unvisited centres are retained with precisely the multiplicity in (E.2); no full-word cardinality is inserted. Borel event cuts do not require a new chart merely because they disconnect the selected set.

In particular the per-clearance condition (E.3) is not needed on a chart controlled by Theorem 16.5. An angular fold may have an unbounded conditional density and a bounded integrated density at the same time. What is still required for the full clearance payment is a bound for the complete, exact-label, positively weighted chart sum, in the original $m^2L^\infty(dt)$ topology.

The local constants in (16.6) or (16.7) have not been proved to provide that bound. Charts with $S_u = 0$, degenerate angular behaviour outside a verified finite type, and singular physical boundaries are not discarded. Regular full critical roofs are treated next by a different exact calculation.

17. Critical roofs, actual reversing orbits, and their exact density coefficient

We now identify a dynamical invariant in the critical-roof density, without averaging the roof or replacing the physical trajectories. Let x be a regular m-flight trajectory with endpoints in the original section cap. Write $q_x = (\alpha_0, \alpha_m)$ and $t_x = F(q_x)$. All statements about derivatives of the actual collision map below concern regular trajectories, not a formal continuation through a tangent collision.

Theorem 17.1 (Critical half-paths are physical reversing periodic orbits)**.** *The endpoint roof is critical precisely when $p_0 = p_m = 0$. Such a trajectory extends by the actual reflection law to a regular $2m$-collision periodic orbit, whose period need not be primitive. The extra m flights traverse the original flights in reverse, with total roof $2t_x$ and total lifted displacement zero. Let $\mathcal{I}(\alpha, p) = (\alpha, -p)$ be collision time reversal, and put*

$$A^{\mathcal{I}}_{m,R}(x) = \sum_{j=0}^{m-1} \mathbf{1}_{Y^*_R}(\mathcal{I} T^j_R x).$$

If the half-path belongs to the original exact event, then

$$A_{2m,R}(x) = A_{m,R}(x) + A^{\mathcal{I}}_{m,R}(x). \tag{17.1}$$

One cannot replace the right side by $2n$ without additionally verifying the time-reversed section tests.

Proof. The first-variation identity already retained in (3.1) is $\nabla_q F = R(-p_0, p_m)$, which proves the criticality assertion. At a normal terminal collision the reflection sends the incoming velocity to its negative. The ray therefore retraces the preceding actual flight. At each intervening regular collision the reversed reflection retraces the next preceding flight. At the normal initial collision it returns to the original outgoing state. Thus every reversed flight is an actual free flight, with no extra or hidden collision, and the orbit closes after $2m$ collisions. Its reversed displacement is the negative of the forward displacement. Its length is twice the original length. No assertion about a primitive period is used.

For $1 \le j < m$ the reversed state at time $2m - j$ is $\mathcal{I} T^j_R x$, while the states at times 0 and m are fixed by $\mathcal{I}$. In the original event both endpoint section tests equal one. Summing the m forward tests and then the terminal test and the $m - 1$ reversed interior tests gives (17.1), since the initial test in $A^{\mathcal{I}}_m$ also equals one. The original section is not assumed invariant under $\mathcal{I}$. □

Theorem 17.2 (Exact stability determinant at a critical roof)**.** *At a regular critical half-path write*

$$G_x = \begin{pmatrix} a & e \\ e & d \end{pmatrix}, \qquad \Delta_x = ad - e^2 > 0, \qquad \mathcal{M}_x = DT^{2m}_R(x).$$

Then $e \neq 0$, and the actual return derivative satisfies

$$\det \mathcal{M}_x = 1, \qquad \operatorname{tr} \mathcal{M}_x - 2 = |\det(I_2 - \mathcal{M}_x)| = \frac{4\Delta_x}{e^2} > 0. \tag{17.2}$$

Consequently the critical density coefficient

$$\mathfrak{J}(x) := \frac{\omega(x)}{2c_* \sqrt{\det G_x}} = \frac{1}{R c_* \sqrt{|\det(I_2 - DT^{2m}_R(x))|}} \tag{17.3}$$

is an invariant of the actual reversing periodic orbit with its marked initial state and specified half-length m. The coefficient is not a probability conditioned on that periodic orbit.

Proof. The signed optical corner formula (G.1) makes e nonzero on every regular finite trajectory. Differentiate $p_0 = -F_{\alpha_0}/R$ and $p_m = F_{\alpha_m}/R$. Solving the first identity for $d\alpha_m$ gives, in the original (α, p) coordinates,

$$B_x := DT_R^m(x) = \begin{pmatrix} -a/e & -R/e \\ -\Delta_x/(Re) & -d/e \end{pmatrix}, \qquad \det B_x = 1. \tag{17.4}$$

This keeps the sign of e; replacing it prematurely by $|e|$ in the matrix would change a derivative, although not its final trace. At both normal endpoints the derivative of time reversal is $J = \mathrm{diag}(1, -1)$. The reversibility identity $\mathcal{I}T_R^m\mathcal{I} = T_R^{-m}$ therefore yields $DT_R^m(T_R^m x) = JB_x^{-1}J$ and

$$\mathcal{M}_x = JB_x^{-1}JB_x.$$

For an arbitrary determinant-one matrix $B = \left(\begin{smallmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{smallmatrix}\right)$, $JB^{-1}J = \left(\begin{smallmatrix} b_{22} & b_{12} \\ b_{21} & b_{11} \end{smallmatrix}\right)$, so

$$\mathrm{tr}(JB^{-1}JB) - 2 = 4b_{12}b_{21}.$$

Substitution of (17.4) gives $4\Delta_x/e^2$. For a two-dimensional determinant-one matrix, $\det(I_2 - \mathcal{M}_x) = 2 - \mathrm{tr}\,\mathcal{M}_x$. This proves (17.2) and also gives positive real hyperbolic multipliers. Finally $\omega = |e|/R$, so elementary simplification proves (17.3). The determinant of $I_2 - \mathcal{M}_x$ is invariant under a smooth change of collision coordinates. No periodic-orbit trace formula or distributional limit theorem has been invoked. □

Theorem 17.3 (Critical-roof coarea with arbitrary exact-event cuts)**.** *Fix a regular critical half-path and a sufficiently small endpoint neighbourhood V on its designated itinerary. Let $E \subset V$ be an arbitrary Borel selector, including the actual physical domain, initial and terminal section tests, exact occupation and displacement labels, and any clearance tests. Put*

$$y = G_x^{1/2}(q - q_x), \quad f(y) = F(q) - t_x, \quad w(y) = \frac{\omega(q)}{4\pi c_* \sqrt{\det G_x}}.$$

Choose $r_0 > 0$ so that the closed y-ball is contained in the coordinate neighbourhood, and constants $K, L_w \geq 0$ such that

$$\|D^2 f(y) - I_2\| \leq K|y|, \quad |w(y) - w(0)| \leq L_w|y|, \quad Kr_0 \leq 1.$$

For $0 < s < r_0^2/3$ there is, for every angle θ, a unique $r_s(\theta) \in (0, r_0)$ with $f(r_s e(\theta)) = s$. Define

$$a_E(s) = \frac{1}{2\pi} \int_0^{2\pi} \mathbf{1}_E\big(q_x + G_x^{-1/2} r_s(\theta) e(\theta)\big)\, d\theta.$$

The original positive collision measure restricted to E and this ball has a density D_E near t_x, no atom at t_x, and

$$|D_E(t_x + s) - \mathfrak{J}(x) a_E(s)| \leq 2\pi\sqrt{3}\{2L_w + Kw(0)\}\sqrt{s}. \tag{17.5}$$

There is no contribution below t_x. The constant is independent of the Borel selector E, but is a constant of this local chart, not a claimed uniform-in-m constant.

If E contains a neighbourhood of q_x, then $a_E(s) = 1$ for small s and the one-sided density limit is exactly $\mathfrak{J}(x)$. If the selected physical side is described by finitely many C^1 inequalities vanishing at q_x with nonzero linear parts, then its limiting coefficient is $\mathfrak{J}(x)$ times the angular fraction of the corresponding linear sector. Without such a hypothesis the exact $a_E(s)$ is retained; no limit for it is asserted.

Proof. The retained reduced action is smooth on a regular finite itinerary; the inverse function theorem and the nonzero optical corner give a smooth positive ω there. Thus the displayed constants exist after decreasing r_0. At a critical point $f(0) = 0$, $\nabla f(0) = 0$ and $D^2 f(0) = I_2$. Integrating the Hessian bound along a radial segment gives

$$|\partial_r f(re) - r| \le Kr^2/2, \qquad |f(re) - r^2/2| \le Kr^3/6.$$

For $r \le r_0$ these imply $r^2/3 \le f(re) \le 2r^2/3$, $\partial_r f \ge r/2$, and hence the existence and uniqueness of r_s, with $r_s \le \sqrt{3s}$. They also give

$$\left| \frac{r_s}{\partial_r f(r_s e)} - 1 \right| \le K r_s, \qquad 0 < \frac{r_s}{\partial_r f(r_s e)} \le 2.$$

In endpoint coordinates the original collision density is $\omega(q)/(4\pi c_*)$; the linear substitution to y makes it $w(y)\,dy$. Polar coordinates and the monotone radial substitution $t = t_x + f(re(\theta))$, tested against nonnegative roof functions, give the exact density

$$D_E(t_x + s) = \int_0^{2\pi} \mathbf{1}_E(q_s(\theta)) w(r_s e(\theta)) \frac{r_s}{\partial_r f(r_s e(\theta))}\, d\theta.$$

Here $q_s(\theta) = q_x + G_x^{-1/2} r_s e(\theta)$, and any source outside the selected ball is not part of D_E. The point $r = 0$ has zero two-dimensional measure, so it produces no atom. The strict radial positivity also excludes mass below t_x inside the ball. For each angle the difference between the weight in this integral and $w(0)$ is at most $(2L_w + Kw(0))r_s$. Integrating gives (17.5), since $2\pi w(0) = \mathfrak{J}(x)$. This estimate does not differentiate an indicator or require its boundary to have zero roofwise length.

When E contains a neighbourhood, every sufficiently small radial point belongs to it. For the stated finite collection of C^1 cuts, Taylor expansion of each defining function shows convergence of its sign to its nonzero linear part at every angle except the finitely many zero directions. Dominated convergence gives the sector fraction. It is the fraction after the transformation by $G_x^{-1/2}$, not an unweighted fraction in the original endpoint angles. For a general Borel selector no such argument is used. □

Corollary 17.4 (One-sided unvisited boundaries without a fictitious orbit)**.** *Suppose a designated finite broken path reaches an unvisited-disk tangency or a section cut, all its* visited *incidences remain positive, and its endpoint action is critical. The sector formula* (17.5)*, with the original physical-side selector, continues to hold in the one-sided coordinate neighbourhood. At a singular state the second expression in* (17.3) *is* not *asserted to be a derivative of a defined physical return map; the first expression is the one-sided action coefficient.*

Proof. The stationary equations for the visited interior collision points have positive-definite interior Hessian. At the stated limiting path they therefore have a smooth local solution as functions of the two endpoints. The signed optical corner is nonzero because the visited incidences and flight lengths are positive. This supplies the smooth action and endpoint-coordinate Jacobian on a full auxiliary neighbourhood. The portion representing actual trajectories is retained as a Borel subset E; portions intersecting a forbidden disk are never assigned to the physical event. On the admitted side these functions agree with the actual action and original measure. The proof of Theorem 17.3 used arbitrary Borel restrictions, so it applies to this subset without continuing the physical dynamics across its boundary. Simultaneous cuts are included in the same selector. If visited grazing also occurs, this particular smooth extension is not invoked. Its treatment remains the inherited visited-grazing trace argument. Nothing in this corollary equates an area-null boundary point with a zero positive pairing-endpoint trace. □

Proposition 17.5 (A complete finite-count critical coefficient bound)**.** *Let the sum run over all distinct regular critical half-paths in the original exact event, at fixed m, R, n, k. Then*

$$\mathfrak{J}(x) \le \frac{47}{12c_*\lambda_\circ}\varrho^{m-1}. \tag{17.6}$$

Under the retained, explicitly stated geometric-pressure input 11.1, the full positive sum satisfies

$$\sum_x \mathfrak{J}(x) \le \frac{C_0 Q_*}{2c_*\lambda_\circ}. \tag{17.7}$$

This is unscaled. It does not supply an m^{-2} bound for a clearance-selected critical source or a uniform bound for the local error constants in (17.5)*.*

Proof. Use $\sqrt{\det G_x} \ge \lambda_\circ$ in (17.3) and the optical bound for ω in (4.8) to obtain (17.6). The retained global equal-gradient uniqueness gives at most one regular critical half-path per lifted physical word. Such a path extends periodically without singular collisions, so it belongs to the two-sided regular set M'_R. The selected-word sum in (11.3), with no incidence marks, therefore bounds $\sum_x \omega(x)$ by C_0Q_*. Divide by $2c_*\lambda_\circ$. Missing words contribute zero. No multiplication of a per-orbit exponential estimate by the number of words is made. □

17.1. **What the critical calculation does and does not close.** Formula (17.3) identifies an actual return-stability weight in the original roof density; it is not an ansatz for a Markov model and not a periodic-orbit replacement of the invariant measure. Formula (17.5) also keeps positive source on section-cut and one-sided unvisited boundaries rather than assigning it a zero coefficient by area-nullness. The full paired source, its nongrazing comparison pool, and (15.8) are unchanged. A critical *component* coefficient need not equal a discontinuity of the entire density, since other physical components may be present at the same roof.

For a collision-uniform payment it is still necessary to control the actual weighted sum on fixed outer neighbourhoods, its clearance and exact-label localization, and the chart remainders. Choosing a separate collision-dependent neighbourhood merely to separate critical roof values would not prove that fixed-window statement. Neither (17.6) nor the unscaled sum (17.7) is used in that invalid manner. The new identities instead give the exact geometric and stability coefficients that a positive capacity or transfer-operator estimate must control.

18. Exact two-mark clearance flux on one physical flight

The preceding clearance–roof analysis treats a selected near miss as a *mark*, without adding a collision. Two different unvisited disks can be marked on the same actual flight. Here we quantify their physical multiplicity exactly, rather than assuming independence or multiplying by the number of collision words. The result is a second-order *collision-space* clearance theorem; it is not an m^2-normalized pointwise roof estimate.

Throughout let $R \in [R_-, R_+]$, $R_+ = 47/100$, and keep $\delta_* = 3/200$, $0 < \delta_1, \delta_2 \le \delta_*$, the triangular lattice Λ, the actual flight time $\tau \le \tau_+$, and the one-copy physical mark maps X_R, Z_R of Theorem B.2. Define the strictly positive *physical separation* and the finite comparison set

$$\begin{aligned} c_{\mathrm{pair}} &= \sqrt{1 - [2(R_+ + \delta_*)]^2} = \frac{\sqrt{591}}{100} > 0, \\ \mathcal{W}_* &= \{w \in \Lambda \setminus \{0\} : |w| \le \tau_+ + 2(R_+ + \delta_*)\}, \\ N_{\mathrm{pair}} &= |\mathcal{W}_*|. \end{aligned} \tag{18.1}$$

The constant N_{pair} is finite by the *same* finite-horizon assumption already used in Lemma B.1; it is not the number of admissible words, the number of pairing sheets, or an unproved distortion constant.

Lemma 18.1 (Distinct same-flight clearances are uniformly transverse)**.** *Let $z_1 \neq z_2$ be two unvisited disk centers whose clearance values $u_i \in (0, \delta_i)$ are realized by the* same *regular physical flight with unit direction v_ϑ. Let $b_i = (z_i - q) \cdot v_\vartheta^\perp$ be their signed impact parameters. Then both perpendicular feet lie in the open physical flight, $z_2 - z_1 \in \mathcal{W}_*$, and*

$$|(z_2 - z_1) \cdot v_\vartheta| = \sqrt{|z_2 - z_1|^2 - (b_2 - b_1)^2} \geq c_{\mathrm{pair}}. \tag{18.2}$$

In particular, for the first signed impact parameter held fixed, $\partial_\vartheta b_2 = -(z_2 - z_1) \cdot v_\vartheta$ does not vanish on a simultaneous physical two-mark state.

Proof. Lemma B.1 puts each foot in the open physical flight. Thus its along-flight coordinate lies in an interval of length at most τ_+. Since $|b_i| = R + u_i$,

$$|b_2 - b_1| \leq 2R + \delta_1 + \delta_2 \leq 2(R_+ + \delta_*) = 97/100 < 1.$$

The lattice spacing $|z_2 - z_1| \geq 1$ and orthogonal decomposition along $v_\vartheta, v_\vartheta^\perp$ imply the equality and lower bound in (18.2). The same decomposition gives $|z_2 - z_1| \leq \tau_+ + 2(R_+ + \delta_*)$, proving membership in $\mathcal{W}_*$. The impact derivative follows from $dv_\vartheta^\perp/d\vartheta = -v_\vartheta$ with b_1 fixed. Neither a grazing endpoint nor an unvisited near miss is treated as an extra collision. □

For a first center translated to 0, fix $w \in \mathcal{W}_*$ and $\sigma_1, \sigma_2 \in \{\pm 1\}$. Write

$$d_{\sigma_1\sigma_2}(u_1, u_2) = \sigma_2(R + u_2) - \sigma_1(R + u_1). \tag{18.3}$$

For all $0 < u_i < \delta_i$, $|d_{\sigma_1\sigma_2}| < 97/100 < |w|$. Thus the equation

$$w \cdot v_\vartheta^\perp = d_{\sigma_1\sigma_2}(u_1, u_2) \tag{18.4}$$

has exactly two roots modulo 2π, denoted $\vartheta_{w,\sigma_1,\sigma_2,+}(u_1, u_2)$ and $\vartheta_{w,\sigma_1,\sigma_2,-}(u_1, u_2)$, distinguished by the sign of $w \cdot v_\vartheta$. They are smooth on each parameter rectangle. Their one-dimensional change-of-variables Jacobian is

$$\left|\frac{\partial\vartheta}{\partial u_2}\right| = \frac{1}{\sqrt{|w|^2 - d_{\sigma_1\sigma_2}(u_1, u_2)^2}}, \tag{18.5}$$

with u_1 fixed; the sign of σ_2 has unit magnitude.

Theorem 18.2 (An exact physical two-mark coarea identity)**.** *For every nonnegative Borel weight $a(x, z_1, z_2)$ of a genuine outgoing collision state and two* ordered*, distinct unvisited centers (expressed in the initial incident disk's lattice frame), set at a root of* (18.4)

$$x = X_R(\vartheta, \sigma_1, u_1), \quad z_1 = Z_R(\vartheta, \sigma_1, u_1), \quad z_2 = z_1 + w, \quad \mathfrak{A} = \mathbf{1}_{\{z_2 \in \mathcal{Z}_{\delta_2}(x)\}}.$$

Then the full, one-copy *original collision measure obeys*

$$\begin{aligned} &\int \sum_{z_1 \in \mathcal{Z}_{\delta_1}(x)} \sum_{\substack{z_2 \in \mathcal{Z}_{\delta_2}(x) \\ z_2 \neq z_1}} a(x, z_1, z_2)\, d\nu(x) \\ &\quad = \frac{1}{4\pi R} \sum_{w \in \mathcal{W}_*} \sum_{\sigma_1,\sigma_2,\eta \in \{\pm 1\}} \int_0^{\delta_1} \int_0^{\delta_2} \left.\frac{\mathfrak{A}\, a(x, z_1, z_1 + w)}{\sqrt{|w|^2 - d_{\sigma_1\sigma_2}(u_1, u_2)^2}}\right|_{\vartheta = \vartheta_{w,\sigma_1,\sigma_2,\eta}} du_2\, du_1. \end{aligned} \tag{18.6}$$

The mask $\mathfrak{A}$ retains both actual feet*, the same physical collision flight, and the prohibition on unvisited-disk reflections. It must not be removed in an equality. Any original return, displacement, collision, roof, section, or radius test may be included in a as a Borel selector;* (18.6) *never modifies that test or the physical clock.*

Proof. Apply the proved one-mark flux identity (B.2) to the first mark z_1. This gives the factor $(4\pi R)^{-1}$, the sign σ_1, the u_1 integral and the original angular integral. For each (ϑ, u_1), translating z_1 to the origin associates any distinct second center with exactly one $w = z_2 - z_1 \in \mathcal{W}_*$ by Lemma 18.1. The second center has a definite sign σ_2 of its impact and a unique $u_2 \in (0, \delta_2)$; its physical membership is precisely $\mathfrak{A} = 1$. For fixed w, σ_1, u_1, the function $\vartheta \mapsto b_2 = b_1 + w \cdot v_\vartheta^\perp$ is a sine wave around the fixed b_1. On each of the two signed impact bands $\sigma_2(R, R + \delta_2)$, equation (18.4) has exactly two roots, and its derivative has the nonzero magnitude in (18.5). One-dimensional substitution on the two monotone branches therefore gives the sum over η and the displayed denominator. The total sum over w is finite; Tonelli and monotone approximation first for bounded weights then for arbitrary nonnegative Borel a justify all substitutions. Only regular charts of the genuine flight are used; the inherited one-mark construction assigns a fixed null-set convention on singular lines. The formula counts an ordered pair once because its first mark is unique in the first flux identity and its relative w and sign/angle root are unique thereafter. No physical word count, symbolic renewal process, conditional independence, or fictitious collision is introduced. □

Theorem 18.3 (Uniform quadratic bound for two distinct unvisited marks)**.** *For the original invariant collision measure and every pair of allowed clearances,*

$$\int \sum_{z_1 \in \mathcal{Z}_{\delta_1}(x)} \sum_{\substack{z_2 \in \mathcal{Z}_{\delta_2}(x) \\ z_2 \neq z_1}} 1 \, d\nu(x) \leq \frac{2N_{\mathrm{pair}}}{\pi R c_{\mathrm{pair}}} \delta_1 \delta_2 \leq \frac{2N_{\mathrm{pair}}}{\pi R_- c_{\mathrm{pair}}} \delta_1 \delta_2. \tag{18.7}$$

This bound includes every physical line/flight and both signs; it does not require independence between the two marks.

Proof. Apply Theorem 18.2 with $a \equiv 1$. Drop only the positive *mask in the upper bound.* The denominator is at least c_{pair} by the lattice-spacing calculation, and there are exactly N_{pair} candidate differences and $2^3 = 8$ sign/root choices. Thus the right side is at most $8N_{\mathrm{pair}}\delta_1\delta_2/(4\pi R c_{\mathrm{pair}})$, which is (18.7). The denominator lower bound is pointwise on the entire parameter rectangle, not only almost everywhere on the physically admitted pairs. □

Corollary 18.4 (A sharp first-order unique-clearance law)**.** *Let* $N_\delta = |\mathcal{Z}_\delta|$ *be the* original *one-flight unvisited-near-miss count and write* $C_{\mathrm{pair}} = N_{\mathrm{pair}}/(\pi R_- c_{\mathrm{pair}})$*. Then*

$$0 \leq \frac{\delta}{R} - \nu(N_\delta \geq 1) \leq C_{\mathrm{pair}} \delta^2, \tag{18.8}$$

$$0 \leq \nu(N_\delta \geq 2) \leq C_{\mathrm{pair}} \delta^2, \tag{18.9}$$

$$\nu(N_\delta = 1) \geq \frac{\delta}{R} - 2C_{\mathrm{pair}} \delta^2. \tag{18.10}$$

In particular $\nu(N_\delta \geq 1) = \delta/R + O(\delta^2)$*, uniformly in the original radius interval; the* δ/R *term is exactly the one-copy physical flux, not a proposed Poisson law.*

Proof. For every nonnegative integer N, $0 \leq N - \mathbf{1}_{\{N \geq 1\}} = (N-1)_+ \leq N(N-1)/2$ and $\mathbf{1}_{\{N \geq 2\}} \leq N(N-1)/2$. Also $N\mathbf{1}_{\{N \geq 2\}} \leq N(N-1)$. Apply these deterministic inequalities to N_δ, use the exact physical mean (B.3), and substitute Theorem 18.3 with $\delta_1 = \delta_2 = \delta$. No limit in collision count is needed for these one-flight statements. □

Corollary 18.5 (Original exact-label records: uniform weak overcount payment)**.** *Fix* m, R, n, k*, the original physical* m*-flight record, and a nonnegative Borel roof test* $0 \leq \psi \leq 1$*. Let* $0 < \delta_j \leq \delta_*$

for $0 \le j < m$. With the original $\Pi = \Pi_{n,k,m,R}$ and unchanged normalized source ν_R^*,

$$0 \le \int \mathbf{1}_\Pi \psi(L) \sum_{j=0}^{m-1} \big[N_{\delta_j}(T_R^j x) - \mathbf{1}_{\{N_{\delta_j}(T_R^j x)\ge 1\}}\big] d\nu_R^*(x) \tag{18.11}$$
$$\le \frac{C_{\mathrm{pair}}}{c_*} \sum_{j=0}^{m-1} \delta_j^2.$$

For the original depth schedule $\delta_j = 2\varepsilon r^{d'_j}$, $0 < 2\varepsilon \le \delta_*$, $r = q_0^{1/4}$, this is at most

$$\frac{8C_{\mathrm{pair}}\varepsilon^2}{c_*(1-r^2)}, \tag{18.12}$$

uniformly in m, R, n, k and the test ψ. Thus the positive roof measure of within-flight duplicate marking *has uniform total variation of order ε^2, even with the full original exact labels. The corresponding density need not have controlled L^∞ height: multiplying this mass estimate by m^2 at fixed outer B is* not *justified.*

Proof. The integrand in (18.11) is nonnegative. Since $\Pi \subset Y_R^*$ and $\nu_R^* = c_*^{-1}\mathbf{1}_{Y_R^*}\nu$, its integral is bounded by c_*^{-1} times the sum of unconditioned collision integrals with $\psi \le 1$. By invariance of the original collision measure, each integral equals the same one-flight quantity at $T_R^j x$ and is bounded by $C_{\mathrm{pair}}\delta_j^2$ from Corollary 18.4. The depth list has at most two indices at each nonnegative depth, so $\sum_j \delta_j^2 \le 2(2\varepsilon)^2 \sum_{d\ge 0} r^{2d} = 8\varepsilon^2/(1-r^2)$. All source and time tests were retained until applying positive domination; no independent probability factor and no exchange of the original limits occurs. □

The exact second-order formula (18.6) isolates the nontransverse configurations which a two-mark pointwise analysis would otherwise need to pay. The separation constant $c_{\mathrm{pair}} > 0$ rules out that particular simultaneous same-flight angle degeneracy on the original radius range. This leaves *single* unvisited-disk clearance, marks on different physical flights, grazing/section/pairing boundary terms, and the nongrazing capacity shortage. In particular the full $\mathcal{H}_M(\mathsf{R}_B^{\mathrm{low}})$ limit and the original raw pointwise local limit theorem remain open; the new $O(\varepsilon^2)$ total variation gain is never substituted for the required $m^2 L^\infty$ estimate.

19. Two clearance marks separated by actual reflections

The same-flight theorem of Section 18 is an inherited result. We now treat marks on *different* flights. The intervening reflections, rather than an independence assumption, supply the new transversality. This section proves finite-itinerary geometric identities on the original physical billiard. The uniform sum in the next section explicitly uses the retained physical pressure input; neither section asserts a normalized roof estimate.

19.1. **Physical feet and the reflection derivative.** Put $\kappa_- = 1/R_+$ and $\kappa_+ = 1/R_-$. All radii, clearances, collision coordinates and measures are those of the preceding sections. The following strengthening of the inherited interior-foot bound is useful because it is independent of every visited incidence.

Lemma 19.1 (A uniform physical foot standoff)**.** *For a genuine mark of clearance $0 < u \le \delta_* = 3/200$ on a physical flight, its perpendicular foot has distance at least*

$$a_\circ = \sqrt{(1-R_+)^2 - (R_+ + \delta_*)^2} = \frac{\sqrt{1827}}{200} > \tau_- = \frac{3}{50} \tag{19.1}$$

from each incident collision. In particular the foot is strictly interior, uniformly for $R \in [9/20, 47/100]$.

Proof. The interior assertion is already Lemma B.1; it uses that a third center cannot realize its small clearance at an incident endpoint. If q is either incident contact and z is the marked center, lattice spacing gives $|z-q| \geq 1-R$. Orthogonal decomposition along the actual flight gives $t^2 = |z-q|^2 - (R+u)^2$ for the distance $t > 0$ between that endpoint and the foot. The right side is at least $(1-R_+)^2 - (R_+ + \delta_*)^2 = 1827/40000$. Finally $1827 > 144$ proves $a_\circ > 3/50$. No extra collision at the foot has been introduced. □

Consider a first mark on flight 0 and a second on flight h, with $h \geq 1$. Let $q_1, \dots, q_h$ be the actual intervening collision points, $c_i > 0$ their normal incidences, a the distance from the first foot to q_1, and b that from q_h to the second foot. Write $\ell_i = |q_{i+1} - q_i|$ for $1 \leq i < h$. Thus $a, b \geq a_\circ$ and $\ell_i \geq \tau_-$. Define

$$P(t) = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \qquad K_i = \begin{pmatrix} 1 & 0 \\ 2/(Rc_i) & 1 \end{pmatrix},$$

$$\mathcal{M}_h = P(b)K_hP(\ell_{h-1})K_{h-1}\cdots P(\ell_1)K_1P(a) = \begin{pmatrix} A_h & B_h \\ C_h & D_h \end{pmatrix}. \tag{19.2}$$

For $h = 1$ the middle product is empty: $\mathcal{M}_1 = P(b)K_1P(a)$. The letter B_h here is a matrix entry; it is not the outer localization parameter B.

Theorem 19.2 (The exact cross-flight clearance Jacobian)**.** *Translate the first marked center to 0 and parametrize its line by $(\vartheta, u_1, \sigma_1)$ as in Theorem B.2. On every regular chart where the intervening physical itinerary and second marked center are fixed, let b_2 be the second signed impact parameter. At fixed u_1,*

$$|\partial_\vartheta b_2| = B_h > 0, \qquad \left|\det D_{(\vartheta,u_1)}(u_1,u_2)\right| = B_h, \quad u_2 = |b_2| - R. \tag{19.3}$$

Moreover $\det \mathcal{M}_h = 1$*, and the pointwise lower bounds*

$$B_h \geq a + b + \sum_{i=1}^{h-1} \ell_i, \tag{19.4}$$

$$B_h \geq a_\circ(1 + 2\kappa_-\tau_-)^{h-1} \tag{19.5}$$

hold for every such trajectory. They require no lower cutoff on any c_i. A visited incidence tending to zero makes its positive kick larger, not smaller.

Proof. For a differentiable variation of an oriented line let J denote its transverse displacement and J' its direction variation, measured at the reference point on that line. Sliding the reference point along the same line adds only the usual free-flight term. Hence travel by distance t sends (J, J') to $(J + tJ', J')$.

Here is the reflection calculation with its signs. With outward normal n, tangent $\mathcal{J}n$ and curvature $\kappa = 1/R$, write $v^- = -cn + p\mathcal{J}n$ and $v^+ = cn + p\mathcal{J}n$, where $\mathcal{J}$ is rotation by $\pi/2$. A contact variation is $\delta q = \mathcal{J}n\,\delta s$. Its transverse components in the incoming and outgoing direction frames are $-c\delta s$ and $c\delta s$, respectively. Specular reflection gives $\delta\vartheta^+ = 2\kappa\delta s - \delta\vartheta^-$. Reverse the transverse frame after each reflection, applying the same sign reversal to J' as to J. The contact displacement then becomes continuous, while

$$J'^+ = J'^- + \frac{2\kappa}{c}J.$$

Thus the exact linearized reflection in these unfolded transverse frames is K_i. This is a derivative of the genuine reflection rule, not an auxiliary reflecting wall at a marked disk.

The first foot is $-\sigma_1(R+u_1)v_\vartheta^\perp$. Differentiating it at fixed u_1 gives a vector parallel to v_ϑ; its transverse component is zero and its direction variation is one. Consequently the transverse field

begins with $(J, J') = (0, 1)$. Propagation to the second foot gives (B_h, D_h). Differentiating the second signed distance gives the negative of this transverse component, up to the h frame reversals. For example, at the last contact, differentiation of $b_2 = (z_2 - q_h) \cdot v_h^{\perp}$ gives $-\delta q_h \cdot v_h^{\perp} - b\,\delta\vartheta_h$. Its magnitude is exactly B_h. Multiplication by the second impact sign has unit magnitude, proving both identities in (19.3).

Each factor in (19.2) has determinant one and nonnegative entries. Replacing all K_i by the identity can only decrease the product entrywise; the resulting upper-right entry is the sum of the free lengths. This proves (19.4). At the first collision $J = a \geq a_\circ$. At each subsequent collision, propagation from its predecessor obeys $J_{i+1} = J_i + \ell_i J_i'^{+}$ with $J_i'^{+} \geq 2\kappa_- J_i$. Hence $J_{i+1} \geq (1 + 2\kappa_-\tau_-)J_i$. The final free flight also increases J, which proves (19.5). All assertions have been proved on regular physical charts; a one-sided limiting matrix with an infinite grazing kick is not called a finite derivative of a singular physical return. □

19.2. **Global multiplicity, including disconnected word domains.** A local nonzero Jacobian does not bound the number of its roots. The following global argument supplies precisely that missing fact. A *tangentially augmented itinerary* consists of the first marked center, the ordered h actual intervening disk centers, the second marked center, and the two signed impacts. The marked centers may coincide with one another or with centers visited at other times. They are not the incident centers of their own marked flights.

Theorem 19.3 (One trajectory per tangentially augmented itinerary)**.** *Fix $h \geq 1$, the augmented itinerary, R, and u_1, u_2 in their original clearance ranges. There is at most one regular physical broken path between the two feet realizing these data, including its oriented line at each endpoint. This is a global statement: no connectedness of its admissible word domain is assumed.*

Proof. For two candidate configurations $Q = (q_0, q_1, \dots, q_h, q_{h+1})$ and $\widehat{Q}$, the endpoint positions lie on the two auxiliary circles of radii $R + u_1$ and $R + u_2$, and the other positions lie on the actual radius-R disks. These auxiliary circles only locate the feet; no new collision is counted. The Cartesian broken length $\mathcal{S}(Q) = \sum_{i=0}^{h} |q_{i+1} - q_i|$ is convex and continuously differentiable at each candidate because every segment has positive length. Gradient monotonicity gives

$$(\nabla\mathcal{S}(Q) - \nabla\mathcal{S}(\widehat{Q})) \cdot (Q - \widehat{Q}) \geq 0. \tag{19.6}$$

At an actual interior collision $\nabla_i\mathcal{S} = -2c_i n_i$, so its contribution on the left is $-R(c_i + \widehat{c}_i)|n_i - \widehat{n}_i|^2$. At a marked endpoint the velocity is tangent to its auxiliary circle, with its oriented tangent fixed by the prescribed impact sign. Thus the endpoint gradient is a fixed sign times $\mathcal{J}n_i$. Its difference is orthogonal to the contact difference, and its contribution to (19.6) is zero. Summing shows that every actual interior contact agrees, since $c_i, \widehat{c}_i > 0$.

For completeness, agreement of the contacts alone is not used to skip the endpoint argument. Write the two convex support gaps for $\mathcal{S}$ and add them. Their sum is the left side of (19.6), now zero. Each gap is a sum of nonnegative support gaps of the Euclidean norm for individual flight vectors. Equality in the norm support inequality, with nonzero reference vector, forces the two corresponding vectors to be positive multiples of one another. Therefore all corresponding flight directions agree. At either endpoint its direction and prescribed impact sign determine its auxiliary-circle normal uniquely, hence the foot itself. This proves global uniqueness. Repeated disk centers cause no identification of contact variables at different times and do not affect the argument. □

19.3. **An exact physical two-clearance change of variables.** Set

$$\mathcal{Z}_\circ = \{z \in \Lambda \setminus \{0\} : |z| \leq \tau_+ + 2R_+ + \delta_*\}, \qquad N_\circ = |\mathcal{Z}_\circ|. \tag{19.7}$$

This is the same finite end-candidate geometry as the one-mark construction. After translating the first marked center to zero, the first actual collision center has at most $N_\circ$ choices; the second marked center relative to the last actual collision center has at most $N_\circ$ choices. The intervening physical word has length $h-1$, *not* h. It is empty when $h=1$. Let $\mathcal{A}_{h,R}(u_1,u_2)$ be the set of these augmented itineraries with a regular physical realization, each counted once. Denote the associated original collision state and the two marked centers in its lifted initial frame by $(x_a, z_{1,a}, z_{2,a})$.

Theorem 19.4 (One-copy coarea across arbitrarily many true reflections)**.** *For any $h \geq 1$ and every nonnegative Borel weight f of the initial physical state and the two marks,*

$$\int \sum_{z_1 \in \mathcal{Z}_{\delta_1}(x)} \sum_{z_2 \in \mathcal{Z}_{\delta_2}(T_R^h x)} f(x, z_1, z_2)\, d\nu(x)$$
$$= \frac{1}{4\pi R} \int_0^{\delta_1} \int_0^{\delta_2} \sum_{a \in \mathcal{A}_{h,R}(u_1,u_2)} \frac{f(x_a, z_{1,a}, z_{2,a})}{B_h(x_a)}\, du_2\, du_1. \tag{19.8}$$

In the left side, the center on flight h is translated into the same original lifted frame as the first. The two marks are distinct as flight–center pairs even when their lattice centers agree. All original return, displacement, collision, physical-roof and section tests may be included in f without altering the identity.

Proof. Apply the exact first-mark formula of Theorem B.2; its prefactor is $(4\pi R)^{-1}$ and its parameters are $(\vartheta, u_1, \sigma_1)$. On each regular physical branch the second center has a unique impact sign. The second substitution has Jacobian B_h by Theorem 19.2. The two signs at each end are included among the augmented itineraries; no extra factor of two, radius, incidence, collision count, or word count is permitted. Theorem 19.3 ensures that overlapping charts of a single augmented itinerary cannot produce different roots.

To justify the global substitution, exhaust the regular parameter space by countably many relatively compact inverse-function charts, and disjointize their overlaps before summation. On each chart use the ordinary change-of-variables theorem first for bounded nonnegative weights. The chart exhaustion and then monotone approximation give (19.8) for arbitrary $f \geq 0$. Finite-horizon geometry gives finitely many physical words at each fixed h, although this number is never used as a bound. The usual singular collision set is null for collision area and hence for first-mark area on every chart. Its image under a regular local change of variables is null as well. Values on those bulk-null sets therefore do not alter this identity. This argument makes no claim that their one-sided paired endpoint traces vanish. Membership in $\mathcal{A}_{h,R}$ retains all physical flights and feet; enlarging that set is allowed only for subsequent positive upper bounds. □

20. Uniform cross-flight summation and the complete mark count

The exponential lower bound on B_h does not by itself permit multiplication by the number of physical words. We instead compare B_h^{-1} with the *same* stable cocycle used in Input 11.1. This comparison includes both endpoint coordinate factors and is proved below. Only the unconditioned pressure inequality $\sup_{R,l} P_l(R) \leq Q_*$ is used in the ensuing sum; no exact-label, roof-conditioned or new twisted pressure estimate is assumed.

20.1. **A stable optical comparison between the two physical feet.** On the full-area, two-sided regular set M_R', let x_i be the outgoing state at the ith intervening collision. Keep the Euclidean arclength convention in (s,φ), with $s = R\alpha$ and $p = \sin\varphi$. Set

$$L_\circ = \sqrt{1 + (\kappa_+ + \tau_-^{-1})^2}, \qquad C_J = \frac{L_\circ}{\kappa_- a_\circ^2}. \tag{20.1}$$

These constants do not depend on h, R or the exact labels.

Lemma 20.1 (The cross-flight Green entry pays its stable weight)**.** *At every regular marked trajectory in M_R',*

$$\frac{1}{B_h} \le C_J c_h\, j_{h-1}(x_1) \le C_J j_{h-1}(x_1), \qquad h \ge 1. \tag{20.2}$$

For $h = 1$ this uses the empty stable product $j_0 = 1$.

Proof. We spell out the stable-line convention so that a hidden grazing coordinate factor cannot enter the comparison. At an outgoing collision write the stable transverse slope as $-\beta_i$. Propagation to the next collision, followed by reflection, gives

$$\beta_i = \frac{k_{i+1} + \beta_{i+1}}{1 + \tau(x_i)(k_{i+1} + \beta_{i+1})}, \qquad k_i = \frac{2}{Rc_i}, \qquad 0 < \beta_i < \frac{1}{\tau(x_i)} \le \frac{1}{\tau_-}. \tag{20.3}$$

One can obtain this invariant line without a further summability assumption: start the backward recurrence with terminal value zero. Every recurrence map takes $[0,\infty)$ into $[0,\tau_-^{-1}]$ and its derivative is at most $(1+2\kappa_-\tau_-)^{-2} < 1$. Differences caused by a further terminal continuation therefore tend to zero geometrically, giving the unique invariant negative transverse slope. It is the stable line in the usual invariant collision cone. This agrees with the stable-line convention in Section 1.1 of [7]; no pressure conclusion is being imported from that observation.

If $V_i = d\varphi/ds$ on this line, the identities $J_i = c_i\, ds$ in the outgoing direction frame and $d\vartheta = (\kappa + V_i)ds$ give

$$V_i = -\kappa - c_i\beta_i, \qquad L_i = \sqrt{1+V_i^2} \in [1, L_\circ], \qquad d\ell_i = |J_i| L_i / c_i.$$

Absolute values remove the harmless alternating transverse-frame sign. The physical stable Jacobian between collisions 1 and h therefore satisfies the *exact* coordinate identity

$$\frac{|J_h|}{|J_1|} = j_{h-1}(x_1)\frac{c_h}{c_1}\frac{L_1}{L_h}. \tag{20.4}$$

For the first foot to the first collision, backward propagation through that collision gives

$$\frac{|J_1|}{|J_{\text{first}}|} = \frac{1}{1 + a(k_1 + \beta_1)} \le \frac{c_1}{2\kappa_- a_\circ}.$$

For the final collision to the second foot the ratio is $1 - b\beta_h \in (0,1)$. The second foot precedes the next actual collision by at least $a_\circ$; hence its stable slope magnitude β_{second} satisfies

$$0 < \beta_{\text{second}} = \frac{\beta_h}{1 - b\beta_h} \le \frac{1}{\tau(x_h) - b} \le \frac{1}{a_\circ}. \tag{20.5}$$

Combining these facts with (20.4), the stable transverse contraction between the feet obeys

$$\gamma := \frac{|J_{\text{second}}|}{|J_{\text{first}}|} \le \frac{L_\circ c_h}{2\kappa_- a_\circ} j_{h-1}(x_1). \tag{20.6}$$

All incidences at the first endpoint cancel in this formula; there is no assumption that either endpoint is uniformly nongrazing.

Since $\det \mathcal{M}_h = 1$, applying its inverse to a vector of slope $-\beta_{\text{second}}$ gives $\gamma^{-1} = D_h + B_h\beta_{\text{second}}$. If the final free-flight factor is removed, all remaining matrix entries are nonnegative; thus $B_h = B_0 + bD_h \ge a_\circ D_h$. Equations (20.5) and (20.6) now imply

$$\frac{1}{B_h} = \gamma\left(\frac{D_h}{B_h} + \beta_{\text{second}}\right) \le \frac{2\gamma}{a_\circ} \le \frac{L_\circ c_h}{\kappa_- a_\circ^2} j_{h-1}(x_1).$$

This proves (20.2). The proof uses only the positive Jacobi propagation and the stable-line convention; the pressure bound is not used until the next theorem. □

Theorem 20.2 (A complete cross-flight clearance-density bound)**.** *Assume the retained bound* $\sup_{R,l} P_l(R) \le Q_*$ *from Input 11.1, with its original physical words and stable norm. Then for every* $h \ge 1$ *and almost every* $(u_1, u_2) \in (0, \delta_*)^2$*,*

$$\sum_{a \in \mathcal{A}_{h,R}(u_1,u_2)} \frac{1}{B_h(x_a)} \le 4N_\circ^2 C_J Q_*. \tag{20.7}$$

Consequently the original invariant collision measure satisfies

$$\int N_{\delta_1}(x) N_{\delta_2}(T_R^h x)\, d\nu(x) \le C_{\mathrm{cross}} \delta_1 \delta_2, \qquad C_{\mathrm{cross}} = \frac{N_\circ^2 C_J Q_*}{\pi R_-}, \tag{20.8}$$

uniformly in the full separation $h \ge 1$*, the original radius interval, and* $0 < \delta_i \le \delta_*$*. Every original exact-record or roof selector bounded by one may be retained on the left. The essential bound in* (20.7) *is on the* two clearance variables*, not on the physical roof.*

Proof. First remove the bulk-null complement of M_R' on the countably many regular charts in Theorem 19.4. Because their two-clearance Jacobian is nonzero, the image of this complement is null in $du_1 du_2$. For each fixed R and h, almost every pair of clearances therefore has all its admitted roots in M_R'. There are only countably many separations. A radius-dependent null set is sufficient; no common exceptional set for all radii is asserted.

Fix such a pair of clearances. Specify the first actual collision center relative to the first marked center, and the second marked center relative to the last actual collision center. This has at most $N_\circ^2$ choices by (19.7). There are four endpoint sign pairs. For each of these choices and each realizable physical word w of length $h-1$, Theorem 19.3 allows at most one root. Lemma 20.1 bounds its weight by $C_J \sup_{[w]_R \cap M_R'} j_{h-1}$. The full positive sum is therefore bounded by

$$4N_\circ^2 C_J \sum_{w \in \mathcal{W}_{h-1}(R)} \sup_{[w]_R \cap M_R'} j_{h-1} \le 4N_\circ^2 C_J Q_*.$$

This is a weighted sum, not D^{h-1} times a worst trajectory. It is valid even when word domains are disconnected and when the selected roots have different endpoint momenta. For $h = 1$, the sum over the empty word is exactly $P_0 = 1$.

Use (19.8) with $f = 1$ and integrate the last bound over the original clearance rectangle. The exact prefactor gives $4/(4\pi R)$, proving (20.8). A Borel selector in the unchanged physical event is handled by positive domination only after the exact coarea identity; it never becomes an independent probability factor. The constants are independent of h because the assumed pressure bound is, not because a finite family of tested words was extrapolated. □

20.2. **All chronological duplicate marks, including repeated centers.** For a finite original record of m flights set

$$\mathscr{N}_{m,\boldsymbol{\delta}}(x) = \sum_{j=0}^{m-1} N_{\delta_j}(T_R^j x), \qquad S_{m,\boldsymbol{\delta}} = \sum_{j=0}^{m-1} \delta_j, \quad 0 < \delta_j \le \delta_*. \tag{20.9}$$

A center seen on two different flights gives two marks. Thus this count is different from both the number of distinct lattice centers and the indicator of at least one witness. Let

$$C_F = \max\{2C_{\mathrm{pair}}, C_{\mathrm{cross}}\}, \qquad C_{\mathrm{pair}} = \frac{N_{\mathrm{pair}}}{\pi R_- c_{\mathrm{pair}}} \tag{20.10}$$

with the inherited same-flight constant of Corollary 18.4.

Theorem 20.3 (Uniform second factorial control for the complete record)**.** *Under the pressure input of Theorem 20.2, the original collision measure satisfies*

$$\int \mathscr{N}_{m,\boldsymbol{\delta}}\, d\nu = S_{m,\boldsymbol{\delta}}/R, \tag{20.11}$$

$$\int \mathscr{N}_{m,\boldsymbol{\delta}}(\mathscr{N}_{m,\boldsymbol{\delta}}-1)\, d\nu \le C_F S_{m,\boldsymbol{\delta}}^2. \tag{20.12}$$

The first equality needs no pressure assumption. In particular, writing $S = S_{m,\boldsymbol{\delta}}$ and $\mathscr{N} = \mathscr{N}_{m,\boldsymbol{\delta}}$,

$$\begin{aligned} 0 \le S/R - \nu(\mathscr{N} \ge 1) &\le \tfrac{1}{2} C_F S^2, \\ \nu(\mathscr{N} \ge 2) &\le \tfrac{1}{2} C_F S^2, \\ \nu(\mathscr{N} = 1) &\ge S/R - C_F S^2. \end{aligned} \tag{20.13}$$

These estimates include all pairs of collision times and all physical words, with constants independent of m. They are not a Poisson approximation and assert no independence of the marks.

Proof. Invariance of ν and the exact one-flight flux give (20.11). Put $N_j = N_{\delta_j}(T_R^j x)$. The exact integer identity

$$\mathscr{N}(\mathscr{N}-1) = \sum_j N_j(N_j-1) + 2\sum_{j<k} N_j N_k$$

separates same-flight ordered pairs from pairs at distinct times. Theorem 18.3 bounds the first sum by $2C_{\mathrm{pair}} \sum_j \delta_j^2$. Invariance and Theorem 20.2, applied at separation $k-j$, bound the second by $2C_{\mathrm{cross}} \sum_{j<k} \delta_j \delta_k$. The definition of C_F proves (20.12). For a nonnegative integer N, use $0 \le N - \mathbf{1}_{\{N\ge 1\}} \le N(N-1)/2$, $\mathbf{1}_{\{N\ge 2\}} \le N(N-1)/2$, and $N\mathbf{1}_{\{N\ge 2\}} \le N(N-1)$. Integration proves (20.13); no limiting interchange is needed. □

Corollary 20.4 (The unchanged exact event and complete depth schedule)**.** *Keep the original $\Pi = \Pi_{n,k,m,R}$, L, ν_R^* and all original labels. For every Borel $0 \le \psi \le 1$,*

$$\begin{aligned} 0 &\le \int \mathbf{1}_\Pi \psi(L)[\mathscr{N}_{m,\boldsymbol{\delta}} - \mathbf{1}_{\{\mathscr{N}_{m,\boldsymbol{\delta}}\ge 1\}}]\, d\nu_R^* \\ &\le \frac{C_F}{2c_*} S_{m,\boldsymbol{\delta}}^2. \end{aligned} \tag{20.14}$$

For the original schedule $\delta_j = 2\varepsilon r^{\min(j,m-1-j)}$, $r = q_0^{1/4}$, $0 < 2\varepsilon \le \delta_*$, this becomes

$$\frac{C_F}{2c_*} S_{m,\boldsymbol{\delta}}^2 \le \frac{8C_F\varepsilon^2}{c_*(1-r)^2}, \tag{20.15}$$

uniformly over all m, R, n, k and bounded Borel roof tests. Equivalently, the difference between the mark-count-weighted and one-witness-weighted exact-record roof measures is a positive measure with this total-variation bound. This now pays cross-flight as well as within-flight overcount*; the former was not covered by Corollary 18.5.*

Proof. The integrand is nonnegative and is bounded by $\mathscr{N}(\mathscr{N}-1)/2$. Retain the original exact event until using $\nu_R^* = c_*^{-1}\mathbf{1}_{Y_R^*}\nu$ and positive domination. Equation (20.12) gives (20.14). There are at most two flights at each nonnegative depth, including the middle flight when m is odd. Thus $S \le 2(2\varepsilon)\sum_{d\ge 0} r^d = 4\varepsilon/(1-r)$, which proves (20.15). Taking the supremum over Borel $0 \le \psi \le 1$ is exactly the total variation of the positive difference measure. The equality of this measure with the stated pushforward follows directly from its definition, not from a change of conditional law. □

20.3. **What this theorem does, and does not, close.** The new derivative, global tangential-end uniqueness and coarea statements are geometric theorems for the actual billiard. The uniform complete mark-count theorem is a deduction from the *already retained* unconditioned physical pressure input, whose independent reconstruction remains outstanding. It is not presented as a new proof of that input. The complete double-clearance density and the all-time duplicate payment are stronger than the inherited same-flight assertions, but their measures are still different from the target-roof essential height.

More precisely, for the full original exact event define the positive roof measures

$$\mu^{\mathrm{count}}_{m,\boldsymbol{\delta}} = L_{\#}(\mathbf{1}_{\Pi}\mathscr{N}_{m,\boldsymbol{\delta}}\nu^{*}_{R}), \qquad \mu^{\mathrm{one}}_{m,\boldsymbol{\delta}} = L_{\#}(\mathbf{1}_{\Pi}\mathbf{1}_{\{\mathscr{N}_{m,\boldsymbol{\delta}}\geq 1\}}\nu^{*}_{R}).$$

The exact identity $\mu^{\mathrm{count}} = \mu^{\mathrm{one}} + \mu^{\mathrm{excess}}$ has $\mu^{\mathrm{excess}} \geq 0$ and the bound (20.15). The first-order one-mark source is *not* subtracted from the problem. Even in the unconditioned law, (20.11) and (20.13) retain its positive first-order term S/R. The exact-label source need not have that same leading coefficient; no conditioning argument here claims otherwise.

To close the original theorem, a proof must still control the clearance-selected first-order source and any excess which enters the *actual paired source* in the $m^2L^{\infty}(dt)$ topology, including critical, degenerate, one-sided boundary and section-cut contributions, or dominate that complete source by the original nongrazing capacity. This bulk identity alone does not identify a paired endpoint trace with a bulk restriction. The low-incidence shallow source must also be paid. The inherited precise common closure condition remains (15.8); it has not been renamed or declared true.

At every fixed outer B the clearance widths are fixed first. Neither the $O(\varepsilon^2)$ mass above nor the exponential single-trajectory lower bound for B_h is multiplied by m^2 and asserted to vanish. The complete physical sum, the radius and exact-label suprema, the roof essential supremum, the collision limsup and the final outer B limit retain their original meanings and order. Thus the raw pointwise local limit and four-coordinate inversion target is unchanged and not yet proved.

21. The angular clearance source as a weighted unstable family

The proofs in Sections 19–20 are inherited, including their stated pressure premise. We now give a different summation route. It starts from the actual angular source in the one-mark flux, rather than taking a supremum over every physical cylinder. The geometric results of this section use no pressure or growth assumption. The propagation theorem in the next section uses the explicitly specified homogeneous curve-growth input, not the stronger physical-word pressure input of Input 11.1.

All radii, sections, time conventions and exact records are unchanged. Put $b_* = R_+ + \delta_* = 97/200$, $a_* = a_\circ = \sqrt{1827}/200$, $\kappa_- = 1/R_+$ and $\kappa_+ = 1/R_-$. We use actual boundary arclength $s = R\alpha$ and outgoing angle φ, with $c = \cos\varphi > 0$ and $p = \sin\varphi$. Arclength on a collision curve always means Euclidean arclength in (s, φ).

21.1. **A genuine first collision, not a reflection at the mark.** Fix $0 < u \leq \delta_*$ and $\sigma \in \{-1, 1\}$, and put $b = \sigma(R + u)$. For $v_\vartheta = (\cos\vartheta, \sin\vartheta)$, the marked foot is $q_f = -bv_\vartheta^{\perp}$. Starting at this point with velocity v_ϑ, let $q_1 = q_f + av_\vartheta$ be the first actual collision, and let $Y_{R,\sigma,u}(\vartheta)$ be the outgoing state there. Thus, in the one-mark notation,

$$Y_{R,\sigma,u}(\vartheta) = T_R X_R(\vartheta, \sigma, u). \tag{21.1}$$

The line is exterior to the marked disk. Moreover, for every other lattice center z, $|z - q_f| \geq 1 - b_* > R_+$. The foot is therefore in the actual free region for every ϑ. The inherited finite-horizon bound supplies the two incident collisions. We exclude the finitely many first-collision tangencies when differentiating. The marked disk itself is never reflected from at the foot.

Theorem 21.1 (Exact first-image density and uniform geometric regularity)**.** *On every regular interval of fixed first collision center, let s be an unwrapped boundary coordinate at that center. Then*

$$\frac{ds}{d\vartheta} = -\frac{a}{c}, \qquad V := \frac{d\varphi}{ds} = \frac{1}{R} + \frac{c}{a}, \qquad \rho := \left|\frac{d\vartheta}{d\ell}\right| = \frac{c}{a\sqrt{1+V^2}}. \tag{21.2}$$

In particular, with

$$U_0 = \kappa_+ + a_*^{-1}, \qquad K_0 = U_0/a_* + (1 + b_*/a_*)/a_*^2, \tag{21.3}$$

every such image is an unstable graph satisfying

$$a \geq a_*, \qquad \kappa_- \leq V \leq U_0 < \kappa_+ + \tau_-^{-1}, \qquad |dV/ds| \leq K_0, \qquad 0 < \rho \leq c/a_*. \tag{21.4}$$

These bounds hold at every regular first collision, with no lower cutoff on c and no lower cutoff on u within its positive range.

Proof. The foot standoff is Lemma 19.1. Let n be the outward normal at q_1 and $t = n^\perp$. The incoming and outgoing velocities are $v^- = -cn + pt$ and $v^+ = cn + pt$. At fixed u, the variation of q_f is parallel to v^-. Differentiating $q_1 = q_f + av^-$ and projecting onto $(v^-)^\perp$ gives $-c\,ds/d\vartheta = a$, proving the first formula. The specular angle identity gives $d\vartheta^+ = 2R^{-1}ds - d\vartheta$; subtracting the normal-angle differential $R^{-1}ds$ yields $d\varphi = R^{-1}ds - d\vartheta$. Division by ds proves the formula for V. Since $d\ell = \sqrt{1+V^2}|ds|$, the density formula follows. Its vanishing factor c is essential.

For clarity, the second-derivative estimate does not appeal to a sampled trajectory or to bounded curvature by name. Projecting the same contact identity onto n gives $da/d\vartheta = -b - ap/c$. Hence

$$\frac{da}{ds} = p + \frac{bc}{a}, \qquad \frac{dc}{ds} = -pV, \qquad \frac{dV}{ds} = -\frac{pV}{a} - \frac{c}{a^2}\left(p + \frac{bc}{a}\right). \tag{21.5}$$

Use $|p|, c \leq 1$, $|b| \leq b_*$ and $a \geq a_*$ to obtain (21.4). The curvature of the graph in (s, φ) is $|dV/ds|/(1+V^2)^{3/2}$ and is therefore also bounded by K_0. Finally $a_* > \tau_-$ gives the strict upper-cone inequality. No finite derivative at $c = 0$ is asserted. □

21.2. **Uniform finite geometry before the homogeneous cut.** Use the finite candidate set already fixed in (19.7), and write $N_* = N_\circ$ in this section. For reference, its radius is $D_* = \tau_+ + 2R_+ + \delta_*$. It has the explicit packing upper bound

$$N_* + 1 \leq 4(D_* + 1/2)^2. \tag{21.6}$$

Indeed, the disjoint open radius-$1/2$ disks centered at all the lattice points in this closed radius-D_* disk lie in the radius-$(D_* + 1/2)$ disk. This bound does not assign a sampled value to τ_+. Here τ_+ means the inherited uniform supremum of actual free-flight lengths over the original radius family.

Lemma 21.2 (A uniform branch count)**.** *For each fixed R, u, σ, the angular circle can be cut at at most $8N_*$ points, in addition to an angular coordinate seam, so that on every resulting interval the first collision center is fixed, the contact coordinate has no seam, and the image in Theorem 21.1 is injective. The number of intervals is at most $M_* = 8N_* + 1$; each has image length at most $2\pi R_+\sqrt{1+U_0^2}$.*

Proof. Every first collision center belongs to the stated candidate set: its distance from the marked center is at most $\tau_+ + 2R_+ + \delta_*$. For a candidate z, loss or creation of an intersection of the line with its disk requires

$$z \cdot v_\vartheta^\perp + b = R \quad \text{or} \quad z \cdot v_\vartheta^\perp + b = -R.$$

Each equation has at most two roots on the angular circle, since $z \neq 0$. There are thus at most $4N_*$ tangency cuts. Away from these cuts, positive first-hit times vary continuously. Their

ordering cannot change without two distinct disks having a common contact point, which is impossible because $2R < 1$. No intersection can pass through the foot, which is exterior to every disk as proved above. Thus the first target is fixed on each interval.

At each candidate disk choose one contact-coordinate seam, at $q = z + R(1, 0)$. A line in the source passes through this point only if $q \cdot v_\vartheta^\perp = -b$. Since $|q| \geq 1 - R_+ > b_*$, there are at most two such roots for each disk. We allow $4N_*$ seam cuts in the displayed bound, which is a harmless overestimate. Coincident roots only reduce the count. On a seam-free interval $ds/d\vartheta = -a/c < 0$, so s is strictly monotone and cannot wind around its contact circle. Its range has length at most $2\pi R$. This proves injectivity and the length bound. The argument uses no long collision-word count. Overlapping images from different intervals remain separately labeled sources; they are not silently identified in the quotient collision space. □

21.3. **Paying the infinitely many grazing pieces.** Fix $k_0 \geq 2$. Define the homogeneous strips

$$H_k^+ = \{\pi/2 - k^{-2} < \varphi < \pi/2 - (k+1)^{-2}\}, \quad H_k^- = -H_k^+, \qquad k \geq k_0,$$

and let H_0 be the remaining central strip, apart from its boundary. Let $\ell_* > 0$ be a fixed allowed curve length. Subdivide each homogeneous image into pieces of length at most ℓ_*. Set

$$L_* = 1 + \left\lceil \frac{2\pi R_+ \sqrt{1 + U_0^2}}{\ell_*} \right\rceil, \qquad A_* = \frac{2M_* L_*}{a_*} \left(1 + \frac{2}{k_0 - 1}\right). \tag{21.7}$$

The factor two in A_* includes both first-impact signs.

Theorem 21.3 (A uniformly summable angular source)**.** *For every $R \in [R_-, R_+]$ and $0 < u \leq \delta_*$, the pushforward of $\sum_{\sigma=\pm1} d\vartheta$ under the actual maps $Y_{R,\sigma,u}$ has a countable, source-labeled representation*

$$\mu_{R,u} = \sum_{C \in \mathcal{F}_{R,u}} \rho_C \, d\ell_C, \qquad \sum_{C \in \mathcal{F}_{R,u}} \|\rho_C\|_\infty \leq A_*. \tag{21.8}$$

Each C is a homogeneous unstable graph of length at most ℓ_*, with the slope and curvature bounds in (21.4)*. The identity holds against every nonnegative Borel test, and has total mass 4π. An arbitrary Borel selector in the original angular source can be retained on the right as a pullback selector; its derivatives or connected components are not counted.*

Proof. On each interval of Lemma 21.2, $V > 0$ and s is strictly monotone, so φ is strictly monotone. Each homogeneity strip is consequently visited at most once by that interval. The length subdivision requires at most L_* pieces per strip. The density is exactly (21.2), not a constant-density substitute. On H_0 it is at most $1/a_*$. On either $H_k^\pm$, it is at most k^{-2}/a_*, because $\cos\varphi \leq k^{-2}$. Thus, summing over both signs,

$$\sum_C \|\rho_C\|_\infty \leq \frac{2M_* L_*}{a_*} \left(1 + 2 \sum_{k=k_0}^{\infty} k^{-2}\right) \leq A_*.$$

The last inequality follows by integrating x^{-2} from $k_0 - 1$ to infinity. First perform the exact substitution on finitely many regular intervals and truncated homogeneous strips. The omitted angular cut points are finite, and strip boundaries have at most one preimage per regular interval. Monotone exhaustion proves the measure identity and its total mass. Borel selectors are multiplied before substitution and are not differentiated. In particular an infinite number of small selector components does not incur an infinite multiplicity charge. □

Theorem 21.3 is the new first-source result. Replacing ρ_C by the crude bound $1/a_*$ *before* summing the homogeneity strips would give a divergent series. The factor c is the geometric

payment for that series. This is a source estimate in a specific curve norm, not a statement that $\mu_{R,u}$ has a two-dimensional collision density or a bounded original roof density.

22. A full cross-flight root sum from curve growth

We spell out the remaining dynamic input, including its curve class. This prevents replacement of a global pressure premise by an unstated regularity premise. The exact geometry of the preceding section is unconditional within the retained physical finite-horizon model.

Input statement 22.1 (Homogeneous curve growth, in its required class)**.** Choose the common cone upper bound $U = \kappa_+ + \tau_-^{-1}$ and a regular homogeneous unstable curve class containing all the graphs of Theorem 21.3. In particular its curvature allowance must accommodate K_0; being merely C^2 does not establish this membership. For some common $k_0 \geq 2$, sufficiently small $\ell_* > 0$, and $C_g \geq 1$, every curve C of that class and length at most ℓ_* has a source-labeled partition of its regular forward image into homogeneous short curves $\mathcal{G}_l^+(C)$ satisfying

$$\sum_{V \in \mathcal{G}_l^+(C)} \sup_V J_V(T_R^{-l}) \leq C_g, \qquad R \in [R_-, R_+], \quad l \geq 0. \tag{22.1}$$

The branches cover arclength-almost every original point, the inverse branch in each summand is the actual physical one, and all resulting graphs have slopes in $[\kappa_-, U]$. The Jacobian is for Euclidean arclength in (s, φ).

This is the forward, time-reversed version of the curve input in Section D, with the source-class membership now made explicit. The relevant published curve-sum statement is [6, Lemma 3.2(b)]; it is based on the homogeneous curve, expansion and distortion properties (H2)–(H4), not on a sum of physical cylinder suprema. The stated class and its family-uniform constants remain explicit external inputs here. No independent reconstruction of the full geometric-pressure theorem, nor of the older singular pairing and spectral interfaces, is claimed.

Shrinking ℓ_* if necessary, require

$$\ell_* \leq \frac{\pi}{2D_0}, \qquad D_0 = \kappa_+ + U + (1 - R_+)^{-1}. \tag{22.2}$$

The admissible growth partition uses this common choice; any change of its subdivision scale must be included in its constant C_g.

Lemma 22.2 (Counted clearance on a short unstable graph)**.** *For a Borel set $I \subset (0, \delta_*)$, let $N_I(x)$ count all unvisited disks on the next actual flight of x whose clearance belongs to I. For every graph V in Input 22.1 of length at most ℓ_*,*

$$\int_V N_I \, d\ell \leq C_t |I|, \qquad C_t = \frac{2N_\circ \sqrt{1 + U^2}}{\kappa_- a_*}. \tag{22.3}$$

This is a bound on the mark count*, including both impact signs, not just on its nonzero indicator.*

Proof. Use a lift with the incident disk centered at zero. Every possibly marked center belongs to $\mathcal{Z}_\circ$. For a fixed such z, let $\vartheta(s)$ be the outgoing direction, and put $b_z = (z-q)\cdot v_\vartheta^\perp$, $t_z = (z-q)\cdot v_\vartheta$. Direct differentiation gives

$$b_z' = -\cos\varphi - t_z \vartheta', \qquad \vartheta' = 1/R + \varphi' \geq \kappa_-.$$

The angle $\eta = \arg(z-q) - \vartheta$ is strictly decreasing: $\eta' \leq (1-R_+)^{-1} - \kappa_- < 0$. Its total variation is at most $D_0 \ell_* \leq \pi/2$. Thus $t_z > 0$, equivalently $\cos\eta > 0$, holds on at most one interval of the graph. On this interval b_z is strictly decreasing. On the physical mark subset the interior-foot estimate gives $t_z \geq a_*$; therefore $|b_z'| \geq \kappa_- a_*$. For each sign, change variables in this one-to-one map on the marked subset. The integral of its indicator is at most $\sqrt{1+U^2}|I|/(\kappa_- a_*)$. Dropping

a first-hit or unvisited mask occurs only in this positive bound. Sum the two signs and the $N_\circ$ candidates. Approximation by open sets, or the one-dimensional area formula for a monotone smooth map, gives the claim for every Borel I. No number of components of I or of an exact-event selector occurs. □

Theorem 22.3 (A fixed-first-clearance kernel bound)**.** *Assume Input 22.1. For every $R \in [R_-, R_+]$, every fixed $0 < u_1 \le \delta_*$, every integer $h \ge 1$, and every Borel $I \subset (0, \delta_*)$,*

$$\sum_{\sigma=\pm1} \int_0^{2\pi} N_I(T_R^h X_R(\vartheta, \sigma, u_1))\, d\vartheta \le H_* |I|, \qquad H_* = A_* C_g C_t. \tag{22.4}$$

The integrand is taken on its actual regular domain. Its exceptional set has zero angular measure under the curve input. Arbitrary Borel weights between zero and one, including the original section, exact record and physical roof selectors, may be retained inside the sum without increasing the bound. In particular the second-clearance measure has a density bounded by H_*, uniformly in the first clearance and in the complete collision separation.

Proof. The first actual collision has already been taken in $Y = T_R X$. Apply Theorem 21.3 to its source. Exactly $l = h - 1$ further iterates remain. For each labeled curve C, use the physical image partition in (22.1) and one-dimensional change of variables:

$$\begin{aligned}\int_C \rho_C(y) N_I(T_R^l y)\, d\ell_C(y) &= \sum_{V \in \mathcal{G}_l^+(C)} \int_V (\rho_C \circ T_R^{-l}) N_I\, J_V T_R^{-l}\, d\ell_V \\ &\le \|\rho_C\|_\infty C_g C_t |I|.\end{aligned}$$

Use Lemma 22.2 separately on each V, and then sum its weighted inverse Jacobian, not its number of components. Summing C uses (21.8) and proves the bound. For $h = 1$ the middle map is the identity and $l = 0$; no collision is missing or added. First truncate the nonnegative sums and then use monotone convergence. Carry a bounded Borel selector through each exact substitution before dropping it in the inequality. The measure domination for all I implies absolute continuity and the displayed density bound by the Radon–Nikodym theorem. It concerns clearance, not the roof variable $L_{m,R}$. □

Theorem 22.4 (One-copy marked root pressure without physical-word pressure)**.** *Under precisely the preceding curve-growth input, let $\mathcal{A}_{h,R}(u_1, u_2)$ be the set of actual tangentially augmented itineraries in Theorem 19.4. Then*

$$\sup_{R \in [R_-, R_+],\ h \ge 1} \operatorname*{ess\,sup}_{(u_1,u_2) \in (0,\delta_*)^2} \sum_{a \in \mathcal{A}_{h,R}(u_1,u_2)} B_h(x_a)^{-1} \le H_*. \tag{22.5}$$

The complete positive physical sum precedes the essential supremum. No premise on $\mathcal{P}_l(R)$, no product of independently chosen flights and no growth-rate bound on the number of words is used. Consequently

$$\int N_{\delta_1}(N_{\delta_2} \circ T_R^h)\, d\nu \le C_{\mathrm{ag}} \delta_1 \delta_2, \qquad C_{\mathrm{ag}} = H_*/(4\pi R_-). \tag{22.6}$$

Proof. Use the exact one-mark flux, followed by the actual cross-flight coarea identity of Theorem 19.4. Global one-copy multiplicity is supplied by Theorem 19.3, not by local invertibility. For an arbitrary Borel set $I_1 \subset (0, \delta_*)$ and I_2 in the same interval, the resulting joint clearance measure is both

$$\int_{I_1} \sum_\sigma \int_0^{2\pi} N_{I_2}(T_R^h X_R(\vartheta, \sigma, u_1))\, d\vartheta\, du_1 \quad \text{and} \quad \int_{I_1 \times I_2} \sum_{a \in \mathcal{A}_{h,R}} B_h(x_a)^{-1}\, du_1 du_2.$$

The common collision prefactor $1/(4\pi R)$ is omitted on *both* sides of this equality. Theorem 22.3 bounds this measure by $H_* du_1 du_2$, first on rectangles and then on all Borel sets. Uniqueness of its density proves (22.5). Reintroducing $1/(4\pi R)$ and integrating over the two clearance intervals proves (22.6). For each fixed radius the countable separations admit a common clearance-null exceptional set. A null set common to an uncountable radius family is neither needed nor asserted. □

22.1. **The inherited complete-record conclusion under the curve input.** The conclusion below was already obtained in Theorem 20.3 with the physical pressure input. The new contribution is its proof from the directly lifted angular source and curve growth, and the stronger fixed-first-clearance kernel bound, not a second claim of novelty for the integer identity.

Corollary 22.5 (Complete chronological multiplicity with the revised dependency)**.** *Put* $C_A = \max\{2C_{\mathrm{pair}}, C_{\mathrm{ag}}\}$*, where* C_{pair} *is the inherited same-flight constant. For any finite deterministic sequence* $0 \le \delta_j \le \delta_*$*, put* $\mathscr{N} = \sum_{j=0}^{m-1} N_{\delta_j}(T_R^j x)$ *and* $S = \sum_j \delta_j$*. Under Input 22.1,*

$$\int \mathscr{N}\, d\nu = S/R, \qquad \int \mathscr{N}(\mathscr{N}-1)\, d\nu \le C_A S^2. \tag{22.7}$$

The first equality requires no curve-growth assumption. For the original exact event Π *and any Borel* $0 \le \psi \le 1$*,*

$$0 \le \int \mathbf{1}_\Pi \psi(L_{m,R})[\mathscr{N} - \mathbf{1}_{\{\mathscr{N}\ge 1\}}] d\nu_R^* \le \frac{C_A S^2}{2c_*}. \tag{22.8}$$

In particular, for $\delta_j = 2\varepsilon r^{\min(j,m-1-j)}$*,* $r = q_0^{1/4}$ *and* $2\varepsilon \le \delta_*$*, this is at most* $8C_A\varepsilon^2/[c_*(1-r)^2]$ *for every* m*.*

Proof. Invariance of ν and the exact one-mark mean give the first identity. Expand the factorial exactly as

$$\mathscr{N}(\mathscr{N}-1) = \sum_j N_j(N_j-1) + 2\sum_{j<k} N_j N_k.$$

The same-flight theorem bounds the first sum by $2C_{\mathrm{pair}}\sum_j \delta_j^2$ after integration. Invariance and (22.6) bound the second by $2C_{\mathrm{ag}}\sum_{j<k}\delta_j\delta_k$. Their sum is at most $C_A S^2$. No independence was used. The deterministic inequality $0 \le n - \mathbf{1}_{\{n\ge1\}} \le n(n-1)/2$ for integers $n \ge 0$ and $\Pi \subset Y_R^*$ give (22.8); only here is the source dominated by ν/c_*. There are at most two indices at each depth, so $S \le 4\varepsilon/(1-r)$, proving the last bound. Repeated sightings of a disk on distinct flights are included in the cross terms. □

22.2. **The original roof obstruction has not been replaced.** The new norm in (21.8) describes a singular one-dimensional angular source in collision space. The density in (22.4) is in *second clearance*. It is not a density in the original roof, and the complete root bound in (22.5) has no hidden m^{-2} factor. The unconditioned first-order mass S/R remains positive. An exact-label selection can change its leading coefficient; it is not replaced by a probability conditioned on a mark.

To reach the raw target, the original ordered shortage limit (15.8) must still be proved. In particular the first-order marked source must be estimated after the complete original roof pushforward and physical word sum, including fold, critical, higher-degeneracy, one-sided and pairing-boundary terms, with the required m^2 normalization. The single-mark fixed-clearance roof bridge in (E.3) remains only a sufficient unproved condition; it is not a premise added to our main problem. The local fold calculations show why it is safer to integrate clearance before taking a roof supremum. Low-incidence shallow source and nongrazing capacity likewise remain in the original shortage. At fixed outer B, multiplying (22.8) by m^2 does not prove decay. No

collision-dependent roof window, altered section, exceptional-set subtraction or changed order of limits is introduced here.

23. A grazing-uniform curvature graph transform

The preceding numbered proof bodies are retained verbatim. We now discharge one of their explicitly exposed premises: admissibility of the actual angular source for homogeneous curve growth. This is not a deduction of roofwise pressure from a curvature bound. We prove curvature invariance here, one-step weighted complexity and distortion in the next section, and only then apply a precisely identified curve-sum lemma. No event or collision is changed by these constructions.

Write $t_* = \tau_- = 3/50$, $k_- = 100/47$, $k_+ = 20/9$, and $U = k_+ + t_*^{-1} = 170/9$. At a fixed radius write $\kappa = 1/R$. Our curves are graphs $\varphi = \varphi(s)$ in actual boundary arclength, with $V = d\varphi/ds \in [\kappa, U]$. Their signed geometric curvature is

$$K = \frac{dV/ds}{(1+V^2)^{3/2}}.$$

Neither K nor dV/ds is an optical wavefront curvature. The notation in this section is local to one actual regular collision.

Lemma 23.1 (Exact circular graph transform)**.** *Let $c = \cos\varphi$, $p = \sin\varphi$, $d = \cos\varphi_1$, $p_1 = \sin\varphi_1$ at the two actual collisions, and let τ be their physical flight time. Put*

$$w = \kappa + V, \qquad A = c + \tau w, \qquad y = dw/A. \tag{23.1}$$

On a regular branch, $c, d > 0$, and the image slope and signed derivatives are

$$s_{1,s} = -A/d, \qquad V_1 = \kappa + y, \quad \tau_s = -p - p_1 A/d, \tag{23.2}$$

$$V_{1,s_1} = -\frac{p_1 w}{A}(\kappa + 2y) - \frac{p d^2 w(\kappa + 2V)}{A^3} - \frac{c d^2}{A^3} V_s. \tag{23.3}$$

In particular, $\kappa \le V_1 \le U$.

Proof. Let n, t and n_1, t_1 be the outward normal and its counterclockwise rotation at the two contacts. The outgoing velocity is $cn + pt$, and at the next disk its incoming representation is $-dn_1 + p_1 t_1$. The outgoing direction derivative is $\vartheta_s = \kappa + V = w$. Differentiate $q_1 = q + \tau v$ and project perpendicular to v. The initial projection is c, the terminal one is $-ds_{1,s}$, so $-ds_{1,s} = c + \tau w$. Projection along v gives $\tau_s = p_1 s_{1,s} - p$. Specular reflection gives $d\varphi_1 = \kappa\, ds_1 - d\vartheta$, whence $V_1 = \kappa - w/s_{1,s}$. This proves (23.2).

For the second derivative use

$$c_s = -pV, \quad d_s = p_1 V_1 A/d, \quad A_s = -p(\kappa + 2V) - p_1 A w/d + \tau V_s.$$

Differentiate $V_1 = \kappa + dw/A$ and divide by $s_{1,s} = -A/d$. The coefficient of V_s is $-d^2/A^2 + d^2\tau w/A^3 = -cd^2/A^3$; the remaining terms give (23.3). Finally $0 < y \le 1/\tau \le 1/t_*$, so the image stays in the stated cone. Differentiation is only on regular branches; no value of DT at a tangency is asserted. □

Theorem 23.2 (Uniform contraction of geometric curvature)**.** *For every radius in the original interval and every regular actual flight,*

$$|K_1| \le q_*|K| + C_*, \qquad q_* := \frac{4\big(U + (2U)^{-1}\big)^3}{27k_-(k_- + U)^2\{1 + t_*(k_- + U)\}} < \frac{1}{2}, \quad C_* := \frac{58}{27 t_* k_-^2}. \tag{23.4}$$

These constants are independent of both positive incidences and of the flight count. In exact rational form,

$$q_* = \frac{946236091513673179721}{2005319667933288000000}, \qquad C_* = \frac{64061}{8100}.$$

Consequently any

$$B_{\mathrm{cv}} \geq \max\{K_0, 2C_*\} \tag{23.5}$$

defines a forward invariant class of homogeneous C^2 unstable graphs of geometric curvature at most B_{cv}. It contains the actual sources of Theorem 21.3. Along every regular branch of length l,

$$|K_l| \leq q_*^l |K_0^{\mathrm{curve}}| + C_* \frac{1 - q_*^l}{1 - q_*}. \tag{23.6}$$

This assertion includes arbitrarily close approaches to grazing but not a smooth extension through a singular collision.

Proof. After division of (23.3) by $(1 + V_1^2)^{3/2}$, the absolute coefficient of K is

$$\begin{aligned} Q &= \frac{cd^2(1+V^2)^{3/2}}{\{A^2 + (\kappa A + dw)^2\}^{3/2}} \\ &= \frac{c}{A} \frac{(1+V^2)^{3/2}}{w^2} \frac{y^2}{\{1 + (\kappa + y)^2\}^{3/2}}. \end{aligned} \tag{23.7}$$

Here $c/A \leq [1 + t_* w]^{-1}$ and

$$\frac{y^2}{\{1 + (\kappa + y)^2\}^{3/2}} \leq \frac{y^2}{(\kappa + y)^3} \leq \frac{4}{27\kappa}.$$

The last inequality follows by differentiating $y^2/(\kappa + y)^3$; its maximum for $y \geq 0$ is at $y = 2\kappa$. For fixed κ, the function

$$h(V) = \frac{(1+V^2)^{3/2}}{(\kappa + V)^2 \{1 + t_*(\kappa + V)\}}$$

is increasing for $V \geq \kappa > 1$. Indeed $3V/(1+V^2) > 3/(\kappa+V)$ because $\kappa V > 1$, and the remaining lower bound for its logarithmic derivative is $1/(\kappa + V) - t_*/[1 + t_*(\kappa + V)] > 0$. For fixed V it decreases with κ. Thus $Q \leq 4h(U)|_{\kappa = k_-}/(27k_-)$. The elementary bound $\sqrt{1+U^2} \leq U + (2U)^{-1}$ gives $Q \leq q_*$. The displayed rational evaluation verifies $q_* < 1/2$ without rounding.

For the first forcing term in (23.3), use $w/A \leq 1/t_*$ and $\kappa + 2y \leq 2(\kappa + y)$ to bound its normalized magnitude by $2/(t_* k_-^2)$. For the second, write its normalized magnitude as

$$\frac{\kappa + 2V}{Aw} \frac{y^2}{\{1 + (\kappa + y)^2\}^{3/2}} \leq \frac{2}{t_* w} \frac{4}{27\kappa} \leq \frac{4}{27 t_* k_-^2}.$$

Their sum is C_*. This proves (23.4). Induction gives (23.6), and $q_* B_{\mathrm{cv}} + C_* \leq B_{\mathrm{cv}}$ follows from (23.5). A homogeneous cut or length subdivision restricts a smooth curve and does not increase its curvature. The exact source curvature was bounded by K_0 in Theorem 21.1. All denominators used here are positive on the regular domain; the uniform estimates themselves require no incidence cutoff. □

Remark 23.3 (What the norm conversion does and does not do). The graph second derivative on this invariant class is at most $B_2 = B_{\mathrm{cv}}(1 + U^2)^{3/2}$. This is a consequence of geometric curvature invariance, not a claim that the same bound on φ'' is itself an invariant rectangular constraint. The class in (H2) of [6] is formulated with geometric curvature; equivalence of these two boundedness notions is used only for derivative estimates. The optical Jacobi curvature and the long-record roof Hessian are different quantities and are not controlled by (23.4).

24. Physical one-step complexity and homogeneous distortion

We verify the remaining curve-growth hypotheses for precisely the class constructed above. The argument concerns actual one-flight branches and all their homogeneous pieces. It is not an estimate obtained by counting long collision words. Constants may depend on the fixed original geometric family and the chosen finite curvature allowance B_{cv}, but not on R, incidences, clearance marks, collision separation or exact-event selectors.

24.1. **Uniform finite geometry and a one-sided first-hit lemma.** The retained τ_+ can be defined intrinsically as the supremum of lengths of straight free segments in the original family. For clarity, its finiteness also follows directly from the fixed lattice. A complete line of irrational direction is dense on $\mathbb{R}^2/\Lambda$ and meets the open radius-R_- disk. A line of rational direction is parallel to a primitive lattice vector z, with neighboring lattice rows at distance $(\sqrt{3}/2)/|z| \le \sqrt{3}/2$. Its distance from one such row is at most $\sqrt{3}/4 < R_-$. It therefore meets open disks on that row. The same argument applies to half-lines. The direction dichotomy follows, for example, by describing the closure of a line subgroup of a two-dimensional torus: a proper connected subgroup has an integral annihilator and is a rational circle; otherwise the closure is the full torus. The positive semiorbit has the same closure by recurrence of translation on this compact group. These elementary observations use the Euclidean triangular lattice, not a linear deformation of the physical disks.

If free segments of unbounded length existed, translate their midpoints into a compact fundamental cell and take a convergent subsequence of midpoints and directions. Every finite subsegment of the limiting complete line would avoid every open radius-R_- disk, a contradiction. Thus $\tau_+ < \infty$, uniformly also for $R \ge R_-$. This compactness proof does not assign a sampled numerical value to τ_+. Put

$$D_f = \tau_+ + 2R_+, \qquad N_f = \#\{z \in \Lambda : 0 < |z| \le D_f\}, \qquad J_f = 2N_f + 1. \tag{24.1}$$

The packing bound $N_f + 1 \le 4(D_f + 1/2)^2$ is fully geometric.

Choose a preliminary curve length small enough that the relative angle $\eta_z = \arg(z-q) - \vartheta$ varies by at most $\pi/2$ for every candidate. This is possible uniformly because $|z-q| \ge 1 - R_+$ and $\vartheta_s = \kappa + V \le k_+ + U$. On every such graph,

$$\eta_z' \le (1-R_+)^{-1} - 2k_- < 0, \qquad b_z' = -c - t_z(\kappa + V), \quad t_z = (z-q)\cdot v, \quad b_z = (z-q)\cdot v^{\perp}.$$

The positive-foot set is an interval, and b_z is strictly decreasing there. Every disk-intersection set is consequently an interval, with at most two tangency endpoints. The initial point is exterior to every other disk, so $t_z > 0$ and $|b_z| < R$ are exactly the conditions for a positive intersection. Distinct first-entry times cannot exchange their order without a common point of two disjoint disks. Cutting at these at most $2N_f$ tangencies gives at most J_f genuine first-hit branches. Coordinate seams are glued on the collision circle, not counted as new physical branches. In a seam-free local lift the image is a graph and φ_1 is strictly monotone by Lemma 23.1. Each first-hit branch thus meets each target homogeneity strip at most once.

Lemma 24.1 (At most one non-grazing limiting branch)**.** *Fix $k_0 \ge 2$, with the strips of Theorem 21.3. There is a common $\ell_0(k_0) > 0$ such that every allowed curve of length at most $\ell_0(k_0)$ has at most one connected physical image piece in H_0. This assertion counts maximal physical homogeneous pieces before the length subdivision used in the growth partition.*

Proof. We give the local argument also at a limiting initial grazing state. Fix a limiting radius and outgoing state in the compact closure of the collision cylinder. Consider its oriented straight ray and the first disk which it intersects strictly. Such a disk exists by the preceding finite geometry, even if the ray is tangent to one or more earlier disks. The strict first-entry time

and center persist continuously nearby. The finitely many earlier possible contacts in the limit are tangencies. Any nearby first-hit branch on one of these earlier disks has terminal incidence tending to zero, hence cannot have its image in H_0 in a sufficiently small neighborhood. A later disk cannot be the next collision when the persistent strict disk is met.

It remains to check connectivity for the persistent strict disk, rather than infer it just from uniqueness of a limiting center. At an earlier tangency $b_z = \pm R$ the positive foot satisfies $t_z \geq \sqrt{(1-R_+)^2 - R_+^2} > 0$. Thus b_z' is uniformly negative there on all allowed graphs. In a sufficiently small neighborhood this candidate can cross only one of its two tangency levels: the levels are separated by $2R \geq 2R_-$ and $|b_z'|$ has a uniform finite upper bound on the finite candidate set. Its intersection condition is therefore one-sided along any short source graph. Avoiding all earlier disks is an intersection of one-sided intervals, hence an interval. The strict-disk branch on that interval has monotone φ_1, so restricting it to H_0 is still an interval. Incipient tangencies after the strict contact have no effect on which collision is first.

These neighborhoods are open in radius and the compact closed collision cylinder. A finite subcover, and a smaller common curve length, give $\ell_0(k_0)$; the bounded cone slopes keep each sufficiently short graph inside a member of this cover. Equivalently a sequence of shrinking counterexample curves would converge to one of the limiting states just analyzed. Only this limiting line is allowed to pass through tangency points for purposes of classification. Every branch being estimated is an actual regular billiard flight with its actual first collision. No tangency is erased from a physical record. □

24.2. **The complete one-step weighted sum.** For a tangent vector to a positive or negative cone graph define

$$|v|_* = \frac{\kappa + |V|}{\sqrt{1+V^2}}|v|.$$

On these cones $|v| \leq |v|_* \leq \Gamma|v|$, where $\Gamma = \sqrt{1+k_+^2}$. This follows respectively from $\kappa > 1$ and Cauchy–Schwarz. Extend this cone norm to an equivalent norm off the cones if needed; no subsequent estimate uses such an extension. For a positive cone vector the exact expansion is

$$\frac{|DT_R v|_*}{|v|_*} = 1 + \frac{2\kappa A}{dw} \geq 1 + \frac{2k_- t_*}{d} \geq \Lambda_* := 1 + 2k_- t_* = \frac{59}{47}. \tag{24.2}$$

Time reversal gives the corresponding backward expansion on negative cones.

Theorem 24.2 (Uniform one-step expansion and weakened exponent)**.** *Choose k_0 so large that*

$$T_0 := \frac{J_f}{k_- t_*(k_0 - 1)} < \frac{1 - \Lambda_*^{-1}}{2}. \tag{24.3}$$

For $\ell_* > 0$ sufficiently small as in Lemma 24.1, let $\mathcal{H}(C)$ be the maximal regular homogeneous pieces of $T_R C$. Then

$$\sum_{V \in \mathcal{H}(C)} \sup_V J_V^*(T_R^{-1}) \leq \theta_* := \Lambda_*^{-1} + T_0 < 1. \tag{24.4}$$

For every $\zeta > 1/2$ there is a finite uniform C_ζ such that

$$\sum_{V \in \mathcal{H}(C)} \left(\sup_V J_V(T_R^{-1})\right)^\zeta \leq C_\zeta. \tag{24.5}$$

Here J^* and J denote curve Jacobians in the adapted and Euclidean metrics respectively; these are supremum bounds, not just average bounds.

Proof. The possible H_0 piece contributes at most Λ_*^{-1}. On $H_k^{\pm}$, $d \le k^{-2}$, so the inverse of (24.2) is at most $(2k_-t_*k^2)^{-1}$. There are at most J_f pieces in each such strip. Sum both signs and use $\sum_{k\ge k_0} k^{-2} \le 1/(k_0-1)$ to prove (24.4). This sum includes every first-hit branch. The norm comparison gives $J \le \Gamma J^*$ at the two endpoints, not a new factor for each subsequent collision. Thus one may take

$$C_\zeta = (\Gamma/\Lambda_*)^\zeta + 2J_f\left(\frac{\Gamma}{2k_-t_*}\right)^\zeta \sum_{k\ge k_0} k^{-2\zeta} < \infty.$$

These are precisely the needed expansion/complexity properties, with weakened-exponent threshold $\zeta_0 = 1/2 < 1$. No claim of the smaller infinite-horizon threshold appearing elsewhere in the literature is used. □

24.3. **A direct distortion estimate in the target coordinate.**

Lemma 24.3 (One-step logarithmic distortion)**.** *Let J_CT_R be the Euclidean curve Jacobian. On a smooth image graph,*

$$\partial_{s_1}\log J_CT_R = \frac{p_1w}{A} + \frac{dp(\kappa+2V)}{A^2} - \frac{d\tau V_s}{A^2} + \frac{p_1V_1}{d} + \frac{V_1V_{1,s_1}}{1+V_1^2} + \frac{VV_sd}{A(1+V^2)}. \tag{24.6}$$

There is a finite uniform D_B such that its magnitude is at most $U/d + D_B$. For example, with $a_0 = 2k_-t_*$ and $B_2 = B_{\mathrm{cv}}(1+U^2)^{3/2}$, one may use

$$D_B = t_*^{-1} + 2/(t_*a_0) + (B_2/a_0)(1+(2k_-)^{-1}) + B_2. \tag{24.7}$$

Proof. The Jacobian is $J_CT_R = (A/d)\sqrt{1+V_1^2}/\sqrt{1+V^2}$. Differentiate its logarithm in s and divide by $-A/d$. Use the exact A_s, d_s of Lemma 23.1; their apparent $1/d$ singularity leaves exactly p_1V_1/d and the finite terms in (24.6). This proves the identity, including its signs. For the bound use $A \ge t_*w \ge a_0$, $w/A \le 1/t_*$, $\kappa + 2V \le 2w$, $\tau/A \le 1/(2k_-)$ and $|V_s|, |V_{1,s_1}| \le B_2$. The six terms are respectively bounded by t_*^{-1}, $2/(t_*a_0)$, $B_2/(2k_-a_0)$, U/d, B_2 and B_2/a_0. There is no inverse initial incidence in this bound. Since $ds_1/d\ell_1 \le 1$ in absolute value, it also bounds the logarithmic derivative per unit image arclength. □

Theorem 24.4 (Uniform homogeneous distortion at every collision count)**.** *There is $D < \infty$ such that, if x, x' belong to a single regular curve segment whose images through time l are homogeneous, then*

$$|\log J_CT_R^l(x) - \log J_CT_R^l(x')| \le D\, d_{T_R^lC}(T_R^lx, T_R^lx')^{1/3}. \tag{24.8}$$

Distances are along the same physical curve branch. The constant is uniform in l and R. The corresponding Jacobian-ratio distortion bound and the stable inverse version hold on short homogeneous curves.

Proof. For two points of an image graph in $H_k^{\pm}$, let r be their arclength separation. Since $V_1 \ge k_-$,

$$r \le \frac{\sqrt{1+U^2}}{k_-}(k^{-2} - (k+1)^{-2}) \le L_hk^{-3}, \qquad L_h = 3\sqrt{1+U^2}/k_-.$$

Also $d \ge (2/\pi)(k+1)^{-2}$ throughout that segment. The logarithmic derivative bound is therefore at most Ck^2, with a constant independent of k. Hence its integral is bounded by $Ck^2r \le CL_h^{2/3}r^{1/3}$. On H_0, $d \ge \sin(k_0^{-2})$ and the same conclusion follows from the Lipschitz bound, after fixing the maximal short-curve length. This proves one-step logarithmic distortion in the image distance, with a common constant D_1.

Let $r_l = d_{T_R^l C}(T_R^l x, T_R^l x')$. The adapted expansion and its endpoint metric comparison imply $r_j \le \Gamma\Lambda_*^{-(l-j)} r_l$. Expand the logarithm of the l-step Jacobian as the sum of its actual one-step logarithms and apply the preceding estimate at image times $j+1$. The geometric series gives

$$D = D_1\Gamma^{1/3}\sum_{r=0}^{\infty}\Lambda_*^{-r/3} = \frac{D_1\Gamma^{1/3}}{1-\Lambda_*^{-1/3}}.$$

For bounded r_l, the inequality $|e^z-1| \le e^{|z|}|z|$ gives the ratio form. Time reversal $\mathcal{I}(s,\varphi) = (s,-\varphi)$ is an arclength isometry and conjugates T_R to T_R^{-1}, yielding the stable version. The smooth collision measure is $c\,ds\,d\varphi/(4\pi R)$ and is invariant, so its measure Jacobian is identically one. The measure-Jacobian portion of the distortion hypothesis therefore has zero distortion. This use of time reversal changes neither the original section nor its occupation convention. □

25. Admissible physical curve growth and the clearance consequence

We now separate the mathematical application of a published combinatorial lemma from the inherited global spectral and roof-pressure inputs. The application below is a theorem of this revision, not a claim that a successful build has certified the billiard dynamics.

Theorem 25.1 (The required angular-source growth input is satisfied)**.** *For the original triangular circular Lorentz family, choose the per-radius unstable graph class*

$$\mathcal{W}_R^u(B_{\mathrm{cv}}) = \{C : \ C \text{ is a homogeneous } C^2 \text{ graph, } \kappa_R \le d\varphi/ds \le U, \ |K_C| \le B_{\mathrm{cv}}\},$$

where B_{cv} satisfies (23.5)*. There exist common $k_0, \ell_* > 0$ and $C_g < \infty$ such that, for every R, every $C \in \mathcal{W}_R^u(B_{\mathrm{cv}})$ with $|C| \le \ell_*$ and every integer $l \ge 0$,*

$$\sum_{V\in\mathcal{G}_l^+(C)} \sup_V J_V(T_R^{-l}) \le C_g. \tag{25.1}$$

The partition is obtained by the actual regular homogeneous image branches and a fixed short-length subdivision. It covers arclength-almost all of C before pushforward. It contains no added reflection at a clearance mark. All hypotheses of Input 22.1, for this chosen class, are therefore satisfied.

Proof. Fix k_0 by (24.3); choose ℓ_* no larger than the length from Lemma 24.1, the short-curve length needed for its preliminary relative-angle argument, and the length in (22.2). All choices are made once for the radius family, before the collision count or any outer limit.

Theorem 23.2 verifies invariance of the homogeneous C^2 curve class in the literal geometric-curvature sense of (H2) in [6]. It contains the original first-image source graphs; this membership is now proved, not inferred from their being C^2. Equation (24.2) gives the common adapted hyperbolicity and a uniform endpoint metric comparison. Theorem 24.2 verifies (H3.A) and (H3.B), with the explicit permissible threshold $\zeta_0 = 1/2$. Its estimate uses suprema of contraction, so it is at least as strong as the one-step hypothesis used there. Theorem 24.4 gives the arclength and invariant-measure parts of (H4) on homogeneous branches. The logarithmic and ratio forms are converted on the one fixed short length. None of these verifications uses the physical-word pressure input.

For completeness, singular-null coverage is checked on these curves. At each forward step there are only finitely many physical tangency cuts on a short graph, by the positive-foot monotonicity above. Its regular image has strictly monotone φ_1, so homogeneous boundaries have at most one preimage per first-hit branch per strip. The additional length cuts are countable. Starting with a countable collection, iteration thus excludes a countable collection of points for each finite l. Each regular restriction is injective. No positive arclength part of a graph lies in a tangency

or homogeneous-boundary preimage. This proves the asserted null coverage without treating a bulk-null set as a zero endpoint trace in any pairing construction.

We use precisely [6, Lemma 3.2(b), p. 14]: for the stable class and the homogeneous inverse partition $\mathcal{G}_l(W)$, with long pieces subdivided at the fixed short scale, the sum of sup arclength contraction Jacobians is at most a constant depending on the common curve, expansion and distortion constants, uniformly in W, T and l. The published proof specifies (H2)–(H4) as its requirements; it is not a theorem about all cylinder suprema or roof Fourier transforms. Apply this statement to $\mathcal{W}^s_R(B_{\mathrm{cv}}) = \mathcal{I}\mathcal{W}^u_R(B_{\mathrm{cv}})$ and reverse it by $\mathcal{I}T_R\mathcal{I} = T_R^{-1}$. Since $\mathcal{I}$ is an arclength isometry, its Jacobian conversion is exact and gives (25.1). At $l = 0$ the sum is one. One may apply the lemma separately to each R: its constant depends only on the common estimates just proved. A common Banach-space identification across radii, or a perturbation theorem for variable boundary lengths, is not required for this curve-sum conclusion. Further fixed shortening of the scale is included in C_g. □

Corollary 25.2 (Complete physical two-clearance bound with the input discharged)**.** *Retain the genuine one-copy cross-flight coarea and augmented-itinerary uniqueness theorems of Section 19, with their stated physical scopes. In the original Lorentz family the bounds* (22.4)*,* (22.5)*,* (22.6)*,* (22.7) *and* (22.8) *now follow without an unverified homogeneous curve-growth premise and without physical-word pressure. In particular, for every fixed* $0 < u_1 \le \delta_*$ *and every* $h \ge 1$*,*

$$\sum_{\sigma=\pm1} \int_0^{2\pi} N_I(T_R^h X_R(\vartheta, \sigma, u_1))\, d\vartheta \le A_* C_g C_t\, |I|. \tag{25.2}$$

The first-image source is taken after exactly one actual collision, and there are $h-1$ *further collisions before applying the second mark. All exact-record and original-roof Borel selectors between zero and one may be kept inside this positive inequality. The full one-copy root sum is bounded in clearance essential supremum uniformly in* R *and* h*.*

Proof. Theorem 25.1 supplies exactly the class, norm, partition, null-set and uniformity requirements of Input 22.1. Insert it in the preserved proofs of Theorems 22.3 and 22.4 and Corollary 22.5. The angular source has mass 4π; the collision flux factor is $1/(4\pi R)$, and conditioning on the original section contributes exactly $1/c_*$. No further angular sign factor is inserted into $A_* C_g C_t$. The factorial conclusions and their constants are inherited consequences; the new assertion is the discharge of their curve-growth dependency. A null exceptional set can be chosen for all integer h at a fixed radius. No null set common to uncountably many radii is claimed. □

The distinction from the original roofwise theorem. Theorem 25.1 is a complete curve-sum conclusion and Corollary 25.2 sums all regular physical two-mark branches, including arbitrarily large separation and arbitrarily small positive incidences. They remove a genuine hypothesis from that part of the proof chain. They do not turn a clearance density into a roof density. The count mean S/R is still first order, and the original-clock pushforward can have folds, critical and one-sided boundary contributions. In particular no factor m^{-2} occurs in (25.2).

The original target remains, for every fixed central parameter M,

$$\lim_{B\to\infty} \limsup_{m\to\infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \left(\mathsf{U}^{\mathrm{low}}_{B,m}(t) - K_B P^{\mathrm{ng}}_{r_B,m}(t)\right)_+ = 0. \tag{25.3}$$

Here B is the inherited outer parameter. The curvature allowance B_{cv} is fixed permanently by (23.5). All original outer widths are fixed before the collision limsup. Their prescribed $B \to \infty$ limit is not used to tune the curve partition. No result of this revision establishes (25.3), the full raw pointwise LLT, or its final four-coordinate inversion. The remaining minimal obligations along this route are the normalized first-order physical-roof source and complete chart/boundary sum, the incidence-only low-load source or valid comparison capacity, and the stated original

spectral, arithmetic, pairing and normalization interfaces. The new one-flight complexity and curve-sum proof does not certify any stronger whole-cylinder or roofwise pressure statement still used elsewhere.

26. Physical momentum fibres and a complete endpoint kernel

The preceding source, curvature and growth theorems are retained, with no change to their proof bodies. We use them for a different source: a fixed initial physical tangential momentum. This is a coordinate disintegration, not a new conditioning event. In particular we neither replace ν_R^* nor remove any collision or section test. Throughout this section $R \in [R_-, R_+]$, $m \geq 1$, and $a \in (-1, 1)$. The source fibre is

$$F_a = \{(\alpha_0, p_0) : 0 \leq \alpha_0 < 2\pi,\ p_0 = a\}, \qquad d\lambda_a = d\alpha_0.$$

Its total mass is 2π, not one. We keep $t_* = 3/50$, $k_- = 100/47$, $k_+ = 20/9$, and $U = 170/9$ from Section 23. A fixed enlargement of the admissible geometric-curvature allowance will be explicitly paid below.

Lemma 26.1 (First true collision of a momentum fibre)**.** *Put $c = \sqrt{1-a^2}$, $d = \cos\varphi_1$, $p_1 = \sin\varphi_1$, $\kappa = 1/R$, $A = c + \tau\kappa$, and $y = d\kappa/A$. On every actual regular first-hit branch of F_a,*

$$\frac{ds_1}{ds_0} = -\frac{A}{d}, \qquad V_1 := \frac{d\varphi_1}{ds_1} = \kappa + y, \qquad \kappa \leq V_1 \leq U, \tag{26.1}$$

$$\frac{dV_1}{ds_1} = -\frac{p_1\kappa}{A}(\kappa + 2y) - \frac{ad^2\kappa^2}{A^3}, \tag{26.2}$$

$$\rho_1 := \frac{d\lambda_a}{d\ell_1} = \frac{d}{RA\sqrt{1+V_1^2}} \leq \frac{d}{t_*}. \tag{26.3}$$

The geometric curvature of this image is bounded by

$$K_{\mathrm{mom}} := \frac{k_+ + 2/t_*}{t_*} + \frac{1}{t_*^3 k_-}, \tag{26.4}$$

uniformly in both positive endpoint incidences. This bound is deliberately coarse; it has no collision-count or microscopic-window dependence.

Proof. The differentiation proving Lemma 23.1 is valid also for an initial graph of slope zero: its identities precede any use of the unstable cone. Substitute $V = 0$ and $V_s = 0$, not $V = \kappa$, in those identities. This gives (26.1)–(26.2). Alternatively the two projections of $q_1 = q_0 + \tau v$ give $-ds_{1,s_0} = c + \tau\kappa$ and $\tau_{s_0} = -a - p_1 A/d$; differentiating the reflection formula gives the same result. Since $A \geq t_*\kappa$, one has $y \leq 1/t_*$ and $\kappa/A \leq 1/t_*$. Taking absolute values in (26.2), and then dividing by $(1+V_1^2)^{3/2} \geq 1$, proves (26.4). Finally $d\ell_1 = \sqrt{1+V_1^2}|ds_1|$, $ds_0 = R\,d\alpha_0$, and $RA = Rc + \tau \geq t_*$, proving (26.3). All derivatives are on the actual first-hit component. Nothing is continued through a physical grazing collision in this statement. □

Fix once and for all

$$\widehat{B}_{\mathrm{cv}} \geq \max\{B_{\mathrm{cv}}, K_{\mathrm{mom}}, 2C_*\}. \tag{26.5}$$

Theorem 23.2 and the physical complexity and distortion proofs apply to this fixed allowance. Let $\widehat{k}_0, \widehat{\ell}_*, \widehat{C}_g$ be the resulting constants of Theorem 25.1. The growth constant is allowed to change; we do not reuse a constant for a smaller, unverified source class. We may shorten $\widehat{\ell}_*$ once. None of these choices depends on a, m, the exact record, or the outer parameter B.

Lemma 26.2 (A complete, grazing-summable first-image family)**.** *For every a the first physical image of F_a, with the source labels retained, decomposes into countably many homogeneous short curves C in $\mathcal{W}^u_R(\widehat{B}_{\mathrm{cv}})$, carrying their exact pushed weights ρ_C. It covers λ_a-almost every initial point and satisfies*

$$\sum_C \|\rho_C\|_\infty \le A_{\mathrm{mom}} < \infty \tag{26.6}$$

with a constant uniform in R and a. One explicit choice in terms of the fixed geometric constants is

$$\ell_{\mathrm{pre}} = \frac{\pi}{4\{(1-R_+)^{-1}+k_+\}}, \qquad J_{\mathrm{pre}} = 2 + \left\lceil \frac{2\pi R_+}{\ell_{\mathrm{pre}}} \right\rceil, \tag{26.7}$$

$$L_0 = \frac{\pi\sqrt{1+U^2}}{k_-}, \qquad J_{\mathrm{len}} = 2 + \left\lceil \frac{2L_0}{\widehat{\ell}_*} \right\rceil, \tag{26.8}$$

$$A_{\mathrm{mom}} = \frac{J_{\mathrm{pre}} J_f J_{\mathrm{len}}}{t_*}\left(1 + \frac{2}{\widehat{k}_0 - 1}\right). \tag{26.9}$$

Here J_f and the intrinsic finite τ_+ have exactly the meanings of (24.1)*; no sampled value is assigned to them.*

Proof. Divide the initial boundary circle into at most J_{pre} arcs of s_0-length at most ℓ_{pre}. For every candidate center z the relative angle $\eta_z = \arg(z - q_0) - \vartheta$ varies by at most $\pi/4$ on such an arc, and

$$\eta_z' \le (1-R_+)^{-1} - k_- < 0, \qquad b_z' = -c - t_z\kappa.$$

The positive-foot set is an interval; on it the impact parameter is strictly decreasing. Thus each disk contributes at most two tangency cuts. Between these cuts the first-entry times of two disjoint disks cannot exchange order, since equality would give the same physical point on both boundaries. The same finite candidate argument as in Section 24 gives at most J_f first-hit branches on each preliminary arc. The infinite lattice is reduced here by the proved finite-horizon bound, not by truncating a computation.

On each branch $V_1 \ge k_- > 0$, so φ_1 is strictly monotone as a function of a local lift of s_1. Each target homogeneity strip is met at most once. Since $\varphi_1 \in (-\pi/2, \pi/2)$, every such connected image has length at most L_0. Subdivide it into lengths at most $\widehat{\ell}_*$, using no more than J_{len} pieces, with the fixed short-length convention of the growth partition. Circular coordinate seams are glued, or split once within this conservative count, not counted as different trajectories.

In the central strip (26.3) gives $\rho_C \le t_*^{-1}$; in each of the two strips of index $k \ge \widehat{k}_0$ it gives $\rho_C \le (t_* k^2)^{-1}$. Sum all physical branches, subdivisions and both strip signs, using $\sum_{k \ge \widehat{k}_0} k^{-2} \le (\widehat{k}_0 - 1)^{-1}$. This proves (26.9). The density bound, rather than a finite number of grazing strips, is essential. Physical tangencies are finite on each preliminary arc, and strip and length boundaries are countable. Their preimages have zero $d\alpha_0$ measure. The first hit is unique everywhere else. Coincident images, if any after a coordinate identification, retain their different source labels and are summed. □

Lemma 26.3 (Exact one-collision lattice kernel and physical roof)**.** *For $a, v \in (-1, 1)$ and a candidate lattice displacement $z \ne 0$, put $r_z = |z|$, $\gamma_z = \arg z$ and*

$$Q_z(a,v) = \sqrt{r_z^2 - R^2(v-a)^2}, \tag{26.10}$$

$$\alpha_z(a,v) = \gamma_z - \arcsin\!\left(\frac{R(v-a)}{r_z}\right) - \arcsin a \pmod{2\pi},$$

$$T_z(a,v) = Q_z(a,v) - R\sqrt{1-a^2} - R\sqrt{1-v^2}. \tag{26.11}$$

Let $\mathcal{A}_z(a,v)$ be the exact indicator that the regular physical next collision from $(\alpha_z(a,v),a)$ is with center z. For a bounded Borel source selector h, the one-collision endpoint kernel is, almost everywhere in v,

$$D^h_{1,R}(a,v)=\sum_{\substack{z\in\Lambda\\ 0<|z|\le D_f}} h(\alpha_z(a,v),a)\mathcal{A}_z(a,v)\frac{R}{Q_z(a,v)}. \tag{26.12}$$

The unweighted expression is also the exact sum of regular inverse root derivatives at each a,v. On an admitted term, T_z is the original *flight time and z its actual lattice displacement. In particular*

$$D_{1,R}(a,v)\le C_1:=\frac{47N_f}{2\sqrt{291}}. \tag{26.13}$$

This is not a Markov transition rule for the momentum process.

Proof. Let $\vartheta=\alpha_0+\arcsin a$ be the outgoing direction. The signed impact on the next disk is $b_z=(z-q_0)\cdot v_\vartheta^\perp=z\cdot v_\vartheta^\perp+Ra=Rp_1$. Thus $z\cdot v_\vartheta^\perp=R(v-a)$. The root with $z\cdot v_\vartheta>0$ is exactly the displayed α_z, and $z\cdot v_\vartheta=Q_z$ there. The other angular root cannot be a physical forward next collision: on a physical flight this projection is $\tau+R\sqrt{1-a^2}+R\sqrt{1-v^2}>0$. There is no vanishing angular discriminant, since $r_z\ge 1$ and $2R\le 94/100<1$.

At fixed a, differentiation gives $\partial_{\alpha_0}p_1=-z\cdot v_\vartheta/R=-Q_z/R$. The projection identity just used proves (26.11). One-dimensional coarea now gives (26.12), with the actual next-hit mask retained. The mask excludes a disk hidden by an earlier collision, not the earlier collision from the physical record. Every actual first center lies in the proved candidate set, so the expression is complete. Finally $Q_z\ge\sqrt{1-4R_+^2}=\sqrt{291}/50$ and $R/Q_z\le 47/(2\sqrt{291})$. Sum at most N_f terms. The formula keeps the reconstructed initial angle and its exact physical visibility mask. Discarding that information and composing these kernels as if the momenta alone were a Markov chain is not part of this proof or of any subsequent argument. □

Theorem 26.4 (All-time endpoint-momentum kernel on the physical source)**.** *There exists a finite constant*

$$C_{\mathrm{mom}}=\max\left\{C_1,\frac{A_{\mathrm{mom}}\widehat{C}_gJ_f}{k_-t_*}\right\} \tag{26.14}$$

such that, for every R, $m\ge 1$, $a\in(-1,1)$, Borel set $I\subset(-1,1)$, and Borel selector $0\le h\le 1$ on the original physical source,

$$\int_{F_a^{\mathrm{reg}}(m)} h(\alpha_0,a)\mathbf{1}_{\{p_m\in I\}}\,d\alpha_0\le C_{\mathrm{mom}}|I|. \tag{26.15}$$

The selector may retain the complete original $\Pi_{m,n,k}$, both section tests, every collision, any original-roof Borel test and any bounded physical guard. It is not required to factor over time. For every fixed $a,v\in(-1,1)$ the complete regular root sum obeys

$$D_{m,R}(a,v):=\sum_{\substack{\alpha_0:\ p_m(\alpha_0,a)=v\\ (\alpha_0,a)\in F_a^{\mathrm{reg}}(m)}}\frac{1}{|\partial_{\alpha_0}p_m(\alpha_0,a)|}\le C_{\mathrm{mom}}. \tag{26.16}$$

Roots are unique physical initial points, not multiple copies from homogeneity or coordinate seams. In particular the original section source has the joint endpoint-momentum domination

$$(p_0,p_m)_\#(h\,d\nu_R^*)\le\frac{C_{\mathrm{mom}}}{4\pi c_*}\,da\,dv. \tag{26.17}$$

The statement concerns ν_R^* as originally defined; no conditional probability at $p_0=a$ is introduced as a replacement model.

Proof. First expose the cancellation at the *last actual collision*. For a short incoming unstable graph of slope V and arclength ℓ, write $w = \kappa + V$, $A = c + \tau w$ as in Lemma 23.1. On each physical next-hit branch,

$$\left|\frac{dp_1}{d\ell}\right| = \frac{dV_1}{\sqrt{1+V^2}}\left|\frac{ds_1}{ds}\right| = \frac{\kappa A + dw}{\sqrt{1+V^2}} \geq k_- t_*. \tag{26.18}$$

Indeed $A \geq t_* w$ and $w/\sqrt{1+V^2} \geq 1$ because $\kappa > 1$ and $V \geq \kappa$. The small terminal incidence d cancels exactly before an upper bound is taken. There are at most J_f physical next-hit branches on this short graph, each with strictly monotone p_1. Do *not* cut this last coarea into infinitely many terminal homogeneity strips. Direct one-dimensional substitution therefore gives

$$\int_V \mathbf{1}_{\{p_1 \in I\}}\, d\ell \leq \frac{J_f}{k_- t_*}|I|. \tag{26.19}$$

For $m \geq 2$, push each first-image density of Lemma 26.2 through $m-2$ additional actual collisions, using the complete homogeneous short-curve partition. The density on an incoming final graph is bounded by $\|\rho_C\|_\infty$ times its exact inverse curve Jacobian. Their complete sum of suprema is at most $\widehat{C}_g\|\rho_C\|_\infty$, by Theorem 25.1. Now apply (26.19) to the one remaining physical collision. Sum the labelled graphs and first-image sources with Tonelli's theorem. This proves the bound with $A_{\mathrm{mom}}\widehat{C}_g J_f/(k_- t_*)$. For $m = 1$, directly on each of the at most $J_{\mathrm{pre}}J_f$ initial first-hit pieces,

$$\left|\frac{dp_1}{d\alpha_0}\right| = RAV_1 = A + d \geq t_*\kappa \geq k_- t_*.$$

Lemma 26.3 gives the sharper complete bound C_1 used in (26.14). There is neither an extra initial nor an extra terminal collision in either case. A selector between zero and one is retained until the positive upper bound. The first-image and growth singular-null arguments apply on each labelled source, and the final collision removes only finitely many tangencies on each incoming short curve. Their pullbacks are null for $d\alpha_0$ by the exact absolutely continuous weights above.

On each regular original m-itinerary the map $\alpha_0 \mapsto p_m(\alpha_0, a)$ has nonzero derivative: after the first hit it is an unstable graph and all subsequent transformations are regular. One-dimensional coarea gives (26.16) for almost every v. To obtain the stated bound at every regular root value, fix a finite collection of distinct roots at v_0 and take disjoint regular source neighborhoods. The inverse branches and their positive weights are continuous for v near v_0, where their sum is bounded almost everywhere by the already proved density bound. Pass to v_0 and then take the supremum over finite collections. This proves the bound for the complete nonnegative root series, including roots on artificial subdivision boundaries. No value of a physical derivative is assigned at a singular trajectory by this argument. Finally integrate the fibre inequality in a and use $d\nu_R^* = \mathbf{1}_{Y_R^*}\, d\alpha_0\, da/(4\pi c_*)$. The factor c_* occurs exactly once, and there is no R factor because the initial fibre coordinate is α_0, not s_0. □

Remark 26.5 (What is, and is not, smoothed). The bounds sum all relevant regular collision words before any supremum in R, m, a, v with $a, v \in (-1, 1)$, with no endpoint-incidence loss. They have no whole-cylinder pressure premise. They give no mixing rate, product law, lattice local limit, or density in the original clock L_m. The grazing-uniform bound follows from the last-collision cancellation (26.18), not from dropping an inverse incidence factor in a terminal-coordinate coarea estimate. Singular one-sided critical traces require a separate argument below; they do not follow by declaring a bulk-null set to have zero trace.

27. A pressure-free sum of physical critical sectors

We now use the endpoint kernel on its own physical source, rather than multiply a local critical estimate by a count of collision words. The conclusion is a complete, collision-uniform

leading-coefficient sum. It removes a genuine pressure dependency of Proposition 17.5; it is not yet a normalized roof-density theorem. All endpoint coordinates below are the original angles $q = (\alpha_0, \alpha_m)$.

Lemma 27.1 (Endpoint Hessian bounds and the momentum inverse weight)**.** *Let x be a critical point of a regular reduced action F with every visited incidence positive. Write*

$$G = D_q^2 F(x) = \begin{pmatrix} a_0 & e \\ e & d_0 \end{pmatrix}, \qquad \Delta = \det G > 0, \qquad u_x = \frac{1}{|\partial_{\alpha_0} p_m(x)|}.$$

Then $p_0 = p_m = 0$, and

$$u_x = \frac{R|e|}{\Delta} = \frac{R^2\omega(x)}{\Delta}, \qquad \mathfrak{J}(x) = \frac{\sqrt{\Delta}}{2R^2 c_*}\, u_x, \tag{27.1}$$

$$\frac{\sqrt{\Delta}}{R^2} \le \kappa + t_*^{-1} \le U = 170/9, \qquad \frac{\lambda_{\max}(G)}{R^2} \le \kappa + 2t_*^{-1} \le U_2 := 320/9. \tag{27.2}$$

For $m \ge 2$ the second bound improves to $\lambda_{\max}(G)/R^2 \le \kappa + t_*^{-1}$. The same auxiliary-action identities hold at a one-sided unvisited boundary when all visited *incidences stay positive. No physical DT^m at a grazing collision is thereby defined.*

Proof. The endpoint first variations in the retained optical proof are $F_{\alpha_0} = -Rp_0$, $F_{\alpha_m} = Rp_m$. The generating-function matrix in Proposition 17.2 therefore has lower-left entry $-\Delta/(Re)$. Its inverse absolute value gives the first identity, since $\omega = |e|/R$; the retained critical coarea coefficient $\mathfrak{J} = \omega/(2c_*\sqrt{\Delta})$ gives the second.

Here are the upper bounds, including their signs and units. For one physical flight of length τ, differentiate its Euclidean length with respect to the first contact angle. The diagonal second derivative is $Rc_0 + R^2 c_0^2/\tau$: the circular acceleration is $-Rn_0$ and its scalar product with the initial length gradient $-v$ is Rc_0. At the terminal contact the incoming velocity has normal component $-c_m$, giving the same positive circular term Rc_m. At a critical orbit $c_0 = c_m = 1$. Consequently the two endpoint diagonals of the unreduced full length action are at most $R + R^2/t_*$. Eliminating the positive interior optical Hessian is a Schur complement, which subtracts a positive semidefinite matrix from the endpoint block. Thus both reduced diagonals are at most this value, and $\sqrt{\Delta} \le \sqrt{a_0 d_0} \le R + R^2/t_*$. If $m \ge 2$, the unreduced endpoint block is diagonal, so its maximum eigenvalue has the same upper bound. If $m = 1$ there is no interior to eliminate: the mixed second derivative of the one-flight length has absolute value at most R^2/τ, and the two-row bound is $R + 2R^2/t_*$. Division by R^2 proves (27.2). The positivity and elimination are exactly the physical optical Hessian construction retained in Section 4; no full-trajectory Hessian is replaced by an independent matrix model. At a skipped-unvisited tangency its strictly reflected auxiliary word is still a smooth action with positive visited incidences. The same calculation is valid for that auxiliary action, restricted to its actual physical side as in Section 17. □

Theorem 27.2 (Complete regular critical coefficient sum)**.** *For every R, every $m \ge 1$ and every exact record (n, k), the one-copy sum over all regular critical orbits of the original m-flight roof in that record satisfies*

$$\sum_x \mathfrak{J}(x) \le \frac{UC_{\mathrm{mom}}}{2c_*}. \tag{27.3}$$

This also bounds the sum with any coefficients between zero and one, including restrictions to original roof intervals. Neither the whole-cylinder pressure input nor an independent periodic-orbit counting theorem is used. Critical orbits are the actual normal-endpoint reversing trajectories identified in the retained proof, not periodic points of a replacement symbolic model.

Proof. At every such x, $p_0 = p_m = 0$. Distinct physical initial points are distinct regular roots in (26.16) at $(a, v) = (0, 0)$. The pointwise root bound, rather than an arbitrarily chosen version of an almost-everywhere density, gives $\sum_x u_x \le C_{\mathrm{mom}}$. Use (27.1)–(27.2) and sum nonnegative terms. Exact section and record restrictions can only decrease the sum. The proof works for the complete countable collection, by finite exhaustion, before taking the supremum over radius or collision count. □

27.1. **One-sided sectors are paid on the physical side.** Bulk singular-nullness alone proves nothing about an endpoint trace. We give the additional argument for the one-sided critical strata used in the retained local coarea theorem. Consider a finite or countable one-copy family of critical auxiliary-action germs x. Every germ has all visited incidences positive and normal endpoints. Its physical-side set E_x retains all unvisited-obstacle inequalities and the original section and record cuts. In a neighborhood of x, assume these cuts are a finite Boolean combination of C^1 inequalities whose active boundary differentials are nonzero. The limiting tangent set is then a finite union of angular sectors. Multiple simultaneous boundaries are permitted; no positive trace is discarded.

This class includes the original finite-time section/record cuts on a regular auxiliary word. A section boundary is a coordinate boundary pulled back by a locally invertible finite collision map, so its boundary differential is nonzero. The exact occupation count is a finite Boolean combination of these tests. An unvisited tangency is a clearance boundary: the line-coordinate determinant in Lemma B.2 makes its differential nonzero, and an interior-foot tangency stays away from the incident contacts. Only finitely many possible centers occur for a finite word by finite horizon. Strict displacement labels are locally constant on such a word. General Borel pairing selectors, or strata involving an actual visited grazing collision, are *not* asserted to have this tangent sector property; their exact local selectors and outstanding budgets remain in the manuscript.

Put $y = G_x^{1/2}(q - q_x)$ and let $\beta_y(x)$ be the angular fraction of the physical tangent set in these normalized coordinates. It is the leading coefficient fraction in the local theorem of Section 17. Let $z = (p_0, p_m)$ and denote the corresponding angular fraction in the z-plane by $\beta_z(x)$. Auxiliary branches are identified by their strict visited word; their source restrictions are disjointized on overlaps. No multiple copy is introduced at a coordinate seam. The family below may also contain fully regular critical points, whose interior sector fraction is one.

Lemma 27.3 (The correct angular change of variables)**.** *Let $D = \mathrm{diag}(-1, 1)$. At a critical point,*

$$z = R^{-1} D G_x^{1/2} y + o(|y|), \qquad L_x = R^{-1} D G_x^{1/2}. \tag{27.4}$$

Writing $e_\theta = (\cos\theta, \sin\theta)$, the absolute angular Jacobian is

$$\left| \frac{d \arg(L_x e_\theta)}{d\theta} \right| = \frac{\sqrt{\det G_x}}{e_\theta^{\mathsf{T}} G_x e_\theta}. \tag{27.5}$$

For every such tangent sector set,

$$\mathfrak{J}(x) \beta_y(x) \le \frac{\lambda_{\max}(G_x)}{2R^2 c_*} u_x \beta_z(x). \tag{27.6}$$

The Euclidean angle in the unnormalized endpoint coordinates is not substituted for either fraction.

Proof. Use $g = \nabla_q F = G_x(q - q_x) + o(|q - q_x|)$ and $z = R^{-1} D g$. For an invertible planar matrix L, differentiation of its polar angle gives $|\det L|/|L e_\theta|^2$. Here $|\det L_x| = \sqrt{\det G_x}/R^2$ and $|L_x e_\theta|^2 = e_\theta^{\mathsf{T}} G_x e_\theta / R^2$. Integrating (27.5) over the sector yields $\beta_z \ge \sqrt{\Delta}\, \beta_y / \lambda_{\max}(G_x)$. Combine this with (27.1). This argument also permits finite unions, half-planes, corners and complements; boundary rays have zero angular measure. □

Theorem 27.4 (Complete one-sided leading critical budget)**.** *For the physical-side family just specified, with unique strict visited words and one-copy source restrictions, one has*

$$\sum_x u_x \beta_z(x) \le C_{\mathrm{mom}}, \tag{27.7}$$

and consequently, uniformly in R, $m \ge 1$, exact labels and any restriction of the critical values to an original roof interval,

$$\sum_x \mathfrak{J}(x)\beta_y(x) \le \frac{U_2 C_{\mathrm{mom}}}{2c_*}. \tag{27.8}$$

Here u_x at an unvisited boundary is the auxiliary inverse weight $R|e|/\Delta$, not a claimed derivative of the singular physical map. This conclusion removes the whole-cylinder pressure premise for this complete leading critical-sector sum. It does not extend a physical trajectory through a grazing reflection or assert that every Borel pairing seam possesses a sector limit.

Proof. First take a finite collection $\mathcal{F}$ of distinct physical auxiliary germs. A strictly reflected finite word with positive visited incidences has an invertible endpoint-momentum map: its endpoint derivative has nonzero determinant, as follows from $e \neq 0$ and $\Delta > 0$. The inverse-function theorem gives a common small z-disk for the finitely many local inverses. The absolute Jacobian from (p_0, p_m) to (p_0, α_0) on germ x is a positive continuous function $u_x(z)$ with $u_x(0) = u_x$.

Distinct regular initial points have disjoint sufficiently small source neighborhoods. The same is true for distinct strict-word unvisited-boundary germs: at a limiting state, continuing straight past its unvisited tangencies fixes the next strictly met disk uniquely; iterate this observation through the fixed m strict collisions. Any alternative word using one of the tangencies would have a visited zero incidence and is outside this class. Thus two such germs at the same initial state are the same germ; their selectors are united, not counted twice. This is a uniqueness argument for the auxiliary extension and does not alter a physical collision record. Artificial partition seams are identified as well. Finite disjointization of actual source restrictions handles any remaining coordinate overlap.

On the actual side of each germ, its auxiliary orbit is the actual regular physical orbit. Therefore its positive contribution to the joint endpoint root density is bounded, almost everywhere in z, by the complete kernel already proved:

$$\sum_{x \in \mathcal{F}} u_x(z) \mathbf{1}_{E_x}(z) \le C_{\mathrm{mom}}. \tag{27.9}$$

No equality on the singular state itself is needed. By polar Fubini, the inequality holds at almost every angle for almost every radius $r > 0$. Choose a sequence of such radii tending to zero. On each fixed finite family, the C^1 inequalities with nonzero active linear parts imply convergence of $\mathbf{1}_{E_x}(re_\theta)$ to its momentum tangent-sector indicator for almost every θ. Continuity of $u_x(z)$ and bounded convergence on the angular circle give, after dividing the integral of (27.9) by 2π, $\sum_{x \in \mathcal{F}} u_x \beta_z(x) \le C_{\mathrm{mom}}$. Take the supremum over finite subfamilies to prove (27.7). Countability causes no exchange of a signed series or of the collision and outer limits. Finally apply Lemma 27.3 and (27.2). Keeping original record, critical-roof or bounded continuous guard weights only reduces the nonnegative sum. Neither independence nor a physical-word multiplicity factor is used. □

27.2. **Exact gain and the remaining normalized obstruction.** Theorems 27.2 and 27.4 prove a complete leading-coefficient bound at every collision count, including genuine one-sided physical critical sectors. Their source is the full original momentum coordinate disintegration, and their constants are uniform on the original radius family. This replaces the pressure-dependent coefficient estimate, not its historical proof body.

There is no illicit interchange of a local-radius limit with $m \to \infty$. The sector proof first treats finitely many germs at a *fixed* R, m, takes the local radius to zero, and exhausts their positive coefficients. The resulting bound is uniform in R, m because its right-hand side is. It does not give a common neighborhood size or a summable bound for the local $O(\sqrt{s})$ errors. In particular the critical values of different germs can differ, so the bound at the common momentum origin cannot be used as a bound at different momentum points selected by a common physical roof level.

The full original problem still requires the unchanged limit (25.3), with B-dependent widths fixed before the collision limsup. Along the present route the minimal additional inputs are: a complete original-roof chart/error estimate with the required m^{-2} normalization, including the first-order clearance source, noncritical folds and actual visited singular strata; payment of the original incidence-only low-load source or a valid full nongrazing-capacity comparison; and the stated spectral, arithmetic, pairing and conditional-denominator inputs with their actual uniform quantifiers. General pairing selectors retain their exact local angular functions unless their tangent sectors are proved to exist. Neither the nonzero single-mark mass nor a positive boundary trace is set to zero. The complete raw pointwise local limit and final four-coordinate inversion are not asserted by the present theorems.

28. Exact radial coarea and Borel critical-origin bands

The preceding critical-sector theorem takes a local limit at fixed collision count and requires a tangent sector for each physical cut. Here we replace the local Taylor calculation, for an explicitly defined physical class, by an exact nonlinear calculation. This gives a uniform source estimate at every band width, including arbitrary Borel cuts. It neither identifies different physical roof levels nor bounds the paired endpoint measure by the collision measure without a comparison proof.

Throughout this section the dynamics, the probability ν_R^*, the section, and the exact event $\Pi_{n,k,m,R}$ are those of the original model. We write $\Phi_m(x) = (p_0(x), p_m(x)) = \zeta$ and $D = \operatorname{diag}(-1, 1)$. The new polar radius $r = |\zeta|$ is an auxiliary coordinate, not the physical clock and not one of the microscopic widths r_B. Fix $0 < \rho < 1$ and put

$$K_\rho = \frac{100}{47}\sqrt{1-\rho^2}, \qquad U_m = \begin{cases} 320/9, & m = 1, \\ 170/9, & m \geq 2. \end{cases} \tag{28.1}$$

All estimates below are uniform in $R \in [9/20, 47/100]$ and in m. There is no cutoff on a positive interior incidence.

Lemma 28.1 (An endpoint-momentum metric away from the critical point)**.** *On every regular physical m-flight branch with $|\Phi_m| < \rho$, the reduced action and its Hessian satisfy*

$$\widehat{G} := R^{-2} D_q^2 F, \qquad K_\rho I \preceq \widehat{G} \preceq U_m I. \tag{28.2}$$

The map from endpoint angles $q = (\alpha_0, \alpha_m)$ to ζ is a local diffeomorphism. For every local inverse $q = q(\zeta)$,

$$D_\zeta q = R G^{-1} D, \qquad \nabla_\zeta (F \circ q) = D \widehat{G}^{-1} D \zeta. \tag{28.3}$$

These are identities for the actual finite itinerary on its regular physical side; they do not define a physical collision map at tangency.

Proof. The full second variation and its minimizing Schur complement were proved in Lemma 4.2. They give $G \succeq R \operatorname{diag}(c_0, c_m)$. Since $|p_0|, |p_m| < \rho$, division by R^2 gives the lower bound in (28.2). For $m \geq 2$ the unreduced endpoint block is diagonal. Its two entries, divided by R^2, are $c_0/R + c_0^2/\tau_0$ and $c_m/R + c_m^2/\tau_{m-1}$. Each is at most $20/9 + 50/3 = 170/9$. Minimizing over

the interior variations subtracts a positive semidefinite matrix. For $m = 1$ there is no interior block. The absolute mixed entry, divided by R^2, is $c_0 c_m/\tau_0$; the row-sum bound is at most $20/9 + 100/3 = 320/9$. Thus the upper estimate, previously used only at normal endpoints in Lemma 27.1, holds everywhere on the stated regular physical set.

The exact first variation is $\nabla_q F = RD\zeta$, and hence $D_q\zeta = R^{-1}DG$. Its determinant is nonzero by positivity of G. Inverting and applying the chain rule gives

$$(D_\zeta q)^{\mathsf{T}} \nabla_q F = RDG^{-1}(RD\zeta) = D\widehat{G}^{-1}D\zeta.$$

In particular, no derivative of a Hessian-dependent angular change of variables has been dropped. Positivity is used only at regular points with the true specular stationarity equations satisfied. □

Theorem 28.2 (Exact radial roof disintegration and synchronized bound)**.** *Let $e_\theta = (\cos\theta, \sin\theta)$ and, on any regular inverse branch $x = x(\zeta)$, set*

$$h_x(\zeta) = e_\theta^{\mathsf{T}} D\widehat{G}_x^{-1} De_\theta, \quad u_x(\zeta) = |\partial_{\alpha_0} p_m(x)|^{-1}, \quad \zeta = re_\theta \neq 0. \tag{28.4}$$

For fixed θ, while the same actual branch is regular,

$$\frac{dL(x(re_\theta))}{dr} = rh_x(re_\theta), \qquad U_m^{-1} \le h_x \le K_\rho^{-1}. \tag{28.5}$$

The collision-source density in (L, θ) on this branch is exactly

$$b_x(re_\theta) = \frac{u_x(re_\theta)}{4\pi c_* h_x(re_\theta)}. \tag{28.6}$$

A Borel selector $0 \le w \le 1$, including the original initial/terminal section tests, exact labels, roof intervals, guards and pairing cuts, multiplies this density without a smoothness assumption. For the complete one-copy regular physical root family,

$$\sum_{x:\,\Phi_m(x) = re_\theta} w(x) b_x(re_\theta) \le \frac{U_m C_{\mathrm{mom}}}{4\pi c_*} \tag{28.7}$$

for $0 < r < \rho$ at every regular root value (and for the corresponding almost-everywhere versions of measurable disintegrations). The summands in (28.7) *share the same endpoint momenta; their physical roofs need not be equal.*

Proof. Equation (28.3) gives (28.5). The eigenvalue bounds in (28.2) give its two-sided constants. The determinant of the map $(\alpha_0, p_0) \mapsto (p_0, p_m)$ is $-\partial_{\alpha_0} p_m$. This derivative is nonzero: equivalently, if $G_{0m} = e \neq 0$, the generating matrix has lower-left entry $-\det G/(Re)$. The nonvanishing of e is the exact positive optical denominator identity, not an assumed genericity condition. Consequently the original probability contributes $u_x\, d\zeta/(4\pi c_*)$, with the initial-section indicator retained. Polar coordinates give $d\zeta = r\, dr\, d\theta$. At fixed angle, $dt = rh_x\, dr$, where $t = L(x)$. The factors r cancel, proving (28.6), with no missing R or incidence factor.

For precision with arbitrary cuts, cover the regular source by countably many relatively compact inverse charts, fix one angular seam, and disjointize their source overlaps. Apply change of variables first to bounded nonnegative tests on finitely many charts and then use monotone convergence. On each angular slice, subdivide the open regular inverse domain into intervals; on every such interval (28.5) is strictly positive. Borel cuts stay inside the integrands rather than being turned into new open branches. This proves the exact formula also when a cut has no angular density at its boundary. The angular seam is only a coordinate seam and is counted once. This argument does not discard a singular endpoint trace.

Finally $1/h_x \le U_m$, and Theorem 26.4 bounds the complete root sum $\sum_x u_x(\zeta)$ by C_{mom}. Apply it to finite subfamilies and exhaust the nonnegative terms. Multiplying by selectors between zero and one cannot increase this sum. The root bound at regular root values was

proved in that theorem; no chosen version of an almost-everywhere density is evaluated at a new point. □

28.1. **Critical-origin arms with their physical cuts still present.** The next class is defined by an actual continuation property, not by deleting an inconvenient source. A *critical-origin radial arm* is a regular physical inverse along a fixed momentum ray, defined for $0 < r < r_+(\theta) \le \rho$, with no missing collision on that interval. The arm stays on the same strict visited word. Its endpoint at $r = 0$ may be a regular critical orbit, an unvisited/section boundary, or a limit with a visited tangency; no physical derivative is asserted at that endpoint. All physical points with $r > 0$ remain genuine regular orbits. Only one copy of each physical initial point may be used.

We work with measurable one-copy families of these arms and arbitrary Borel selectors on them. In particular this includes every original pairing sheet whose incoming limit has $|\nabla_q F| = 0$ and whose regular interval reaches that limit. The original sheet construction and its measurable, one-copy parametrization remain the explicitly retained input of Section 3; the present theorem does not purport to reconstruct that historical construction. Its geometric estimate applies to every family satisfying the definition, not only to finitely many coordinate balls or to smooth selectors. Other components of the original event, including arms which terminate at positive momentum radius before reaching zero, are not assigned zero source and remain outside this particular estimate.

Lemma 28.3 (Exact band geometry, without a Taylor radius)**.** *On a critical-origin arm the finite limit $a(\theta) = \lim_{r\downarrow 0} L(x(re_\theta))$ exists and*

$$\frac{r^2}{2U_m} \le L(x(re_\theta)) - a(\theta) \le \frac{r^2}{2K_\rho}. \tag{28.8}$$

No local asymptotic error occurs. The limiting endpoint angles also exist in local angular lifts. The assertions remain valid when an interior incidence tends to zero as $r \downarrow 0$.

Proof. For $0 < \eta < r$, integrate (28.5) to obtain

$$\frac{r^2 - \eta^2}{2U_m} \le L(r) - L(\eta) \le \frac{r^2 - \eta^2}{2K_\rho}.$$

Thus $L(\eta)$ is Cauchy as $\eta \downarrow 0$ and has a finite limit. Taking this limit proves (28.8). Also $|dq/dr| \le 1/(RK_\rho)$ by (28.3); hence both endpoint angles are Cauchy in consistent lifts on a sufficiently short terminal interval. These estimates use no interior-incidence lower bound. All calculations were made at $\eta > 0$ on actual regular orbits. Taking a limit of their physical lengths and endpoint angles does not continue a collision map through a grazing collision. □

Proposition 28.4 (No drift of the limiting roof on a physical fan)**.** *Suppose a continued physical inverse has domain $\{re_\theta : 0 < r < r_+(\theta),\ \theta \in I\}$, where I is a connected angular interval and every compact $J \subset I$ admits a positive radius r_J with $(0, r_J) \times J$ in that domain. Then $a(\theta)$ in Lemma 28.3 is constant on I. In particular the bands on such a fan are measured from one actual limiting physical roof, not a freely chosen angular clock correction. Arbitrary Borel cuts may still be imposed on the actual source after this identity has been obtained.*

Proof. Put $e_\theta^\perp = (-\sin\theta, \cos\theta)$. The exact angular chain rule, using (28.3), is

$$\partial_\theta L(re_\theta) = r^2 (e_\theta^\perp)^{\mathsf{T}} D\widehat{G}^{-1} De_\theta.$$

Its absolute value is at most r^2/K_ρ. Given two angles in I, choose a compact connecting interval J and use the common positive radius in the hypothesis. Integration at any $0 < r < r_J$ gives $|L(re_\theta) - L(re_\psi)| \le r^2|\theta - \psi|/K_\rho$. Letting $r \downarrow 0$ proves equality of the two limiting roofs. Only a compact subinterval is used; no collision-uniform continuation radius is assumed. No angular

boundary limit of a Borel selector is needed because the identity is for the underlying physical action. □

Theorem 28.5 (Complete Borel critical-band and singularity budgets)**.** *For any measurable one-copy family of critical-origin arms let E_s be the set of its actual points satisfying $0 < L - a(\theta) \le s$, where $s > 0$, and put*

$$A_s = \min\{\rho, \sqrt{2U_m s}\}.$$

For any Borel $0 \le w \le 1$, retaining the original event and original probability, one has the simultaneous bounds

$$\int_{E_s} \mathbf{1}_{\Pi_{n,k,m,R}} w \, d\nu_R^* \le \frac{C_{\mathrm{mom}}}{4c_*} A_s^2, \tag{28.9}$$

$$\int_{E_s} \mathbf{1}_{\Pi_{n,k,m,R}} w \, |\Phi_m|^{-\gamma} \, d\nu_R^* \le \frac{C_{\mathrm{mom}}}{2c_*(2-\gamma)} A_s^{2-\gamma}, \qquad 0 \le \gamma < 2, \tag{28.10}$$

$$\int_{E_s} \mathbf{1}_{\Pi_{n,k,m,R}} w \, |\nabla_q F|^{-1} \, d\nu_R^* \le \frac{C_{\mathrm{mom}}}{2c_* R} A_s. \tag{28.11}$$

These estimates hold for the complete family before any supremum over R, m, n, k, and for every s, not merely in a local-radius limit at a fixed collision count. They also hold for a disjoint union of exact records whose total selector is at most one. No tangent-sector limit, finite C^1 description of a pairing cut, common chart radius, or uniform bound on third derivatives of the roof is required.

Proof. By (28.8), every selected point has $|\Phi_m| \le A_s$. The positive joint-kernel domination from Theorem 26.4, on the *actual* physical source, gives

$$\int_{E_s} \mathbf{1}_{\Pi} w \, |\Phi_m|^{-\gamma} \, d\nu_R^* \le \frac{C_{\mathrm{mom}}}{4\pi c_*} \int_{|\zeta| \le A_s} |\zeta|^{-\gamma} \, d\zeta.$$

Apply this first with bounded truncations on finitely many disjoint charts. Monotone convergence supplies the entire physical source and the unbounded nonnegative weight. The polar integral is $2\pi A_s^{2-\gamma}/(2-\gamma)$, proving (28.10). Its case $\gamma = 0$ is (28.9). Finally $|\nabla_q F| = R|\Phi_m|$ gives (28.11) at $\gamma = 1$.

The constants never depended on how many arms were selected or on their individual continuation lengths. A shorter actual arm contributes only its actual points; it was not extended through an unvisited disk. The same argument covers limiting visited tangencies because all integrated points have positive visited incidences. It controls their interior source approach, not an arbitrarily prescribed trace measure supported on the singular boundary. General Borel cuts remain measurable selectors throughout; their pointwise or angular boundary limits are not assumed to exist. The assertion for unions follows from one-copy positive domination, not from adding separately normalized probabilities. □

28.2. **What has and has not been transferred to the original roof.** Theorems 28.2 and 28.5 remove the need for Taylor remainders and tangent sectors for the stated synchronized coarea and physical-band budgets. In particular (28.9) is a collision-uniform $O(s)$ statement and (28.11) is a collision-uniform $O(\sqrt{s})$ statement on the full defined arm family. These are continuum estimates, not conclusions of a finite search or a fitted exponent.

They do not close (25.3). In the actual roof density at a common t, formula (28.6) must be evaluated at the branch-dependent radius $r_x(t, \theta)$. The synchronized sum (28.7) evaluates all branches at the *same* r and is not a bound for that different sum. Likewise $L - a(\theta)$ is used only to specify a physical subset; it is not substituted for L in the exact record, Fourier reference, conditioning or LLT.

The band budget bounds the original collision-source measure. The paired measure $c^\varepsilon\,du\,d\theta$ uses the minimum of its two endpoint traces and is not automatically a submeasure of the interior density $f^\varepsilon\,du\,d\theta$. Its transfer still requires the retained positive-variation/comparison argument. The inverse-gradient estimate does not include the unbounded insertion I_m, and no such insertion is silently absorbed into its constant. Critical-origin arms do not exhaust the positive-radius singular cuts, noncritical folds or low-load source. These sources, the first-order clearance contribution, nongrazing capacity, and the original spectral, arithmetic and conditional-denominator interfaces retain their exact proof obligations. No factor m^2 is attached to the new unscaled bounds at a fixed outer width and declared harmless. The original ordered limit and four-coordinate raw local inversion target remain unchanged and are not asserted to be proved here.

29. Grazing-uniform endpoint localization and the third variation

The estimates in this section are on every actual regular finite itinerary, not on an auxiliary stochastic process. They hold for all $m \ge 1$ and $R \in [9/20, 47/100]$, without a lower cutoff on any positive incidence. The derivative is with respect to the two endpoint angles of the actual stationary action. It is not the derivative of the forward collision map with fixed initial momentum. This distinction is essential: no uniform bound on that expanding forward derivative is asserted.

Keep $\tau_- = 3/50$, $R_+ = 47/100$, and define

$$\sigma = \frac{47}{59}, \qquad s_i = \sigma^i + \sigma^{m-i}, \qquad J_* = \max\left\{1, \frac{R_+}{2\tau_-}(\sigma + \sigma^{-1})\right\} = \frac{2845}{354}. \tag{29.1}$$

On a local physical branch let $\alpha(q) = (\alpha_0(q), \dots, \alpha_m(q))$, $q = (\alpha_0, \alpha_m)$, be the stationary lift. For an endpoint direction $\eta \in \mathbb{R}^2$ put $\xi_i = D_q\alpha_i[\eta]$. Every assertion about continuation below is restricted to a path in this branch up to its first actual boundary.

Theorem 29.1 (Two-sided localization without an inverse-incidence loss)**.** *For every such trajectory and every endpoint direction,*

$$|c_i\xi_i| \le \sigma^i c_0|\eta_0| + \sigma^{m-i}c_m|\eta_m| \le s_i\|\eta\|_\infty, \tag{29.2}$$

$$|\xi_i| \le J_* s_i \|\eta\|_\infty, \qquad 0 \le i \le m. \tag{29.3}$$

In particular, endpoint displacement decays exponentially towards the middle of the true collision chain even when an intermediate incidence is arbitrarily small. No comparison trajectory, new section, or collision suppression is used.

Proof. For $m = 1$ the endpoint assertions follow immediately from $s_0, s_m \ge 1$. Suppose $m \ge 2$. Differentiate the interior stationary equations and use the full physical second variation (4.2). With $w_i = \tau_i^{-1}$ and $y_i = (-1)^i c_i\xi_i$ this gives

$$(\kappa_i + w_{i-1} + w_i)y_i - w_{i-1}y_{i-1} - w_i y_{i+1} = 0, \qquad \kappa_i = \frac{2}{Rc_i}, \quad 1 \le i < m. \tag{29.4}$$

The sign change is important: both off-diagonal coefficients in this Dirichlet equation are negative. For a real sequence with nonnegative boundary values and nonnegative left side in (29.4), a negative interior minimum would make that left side strictly negative, because $\kappa_i > 0$. This proves the needed maximum principle directly.

Write the left side as $\mathcal{D}y$. Since $w_i \le \tau_-^{-1}$ and $\kappa_i \ge 2/R_+$,

$$\frac{\mathcal{D}(\sigma^i)_i}{\sigma^i} = \kappa_i + w_{i-1}(1 - \sigma^{-1}) + w_i(1 - \sigma) \ge \frac{2}{R_+} - \frac{\sigma^{-1} - 1}{\tau_-} = 0.$$

The reversed sequence σ^{m-i} is a supersolution by the same calculation with the adjacent flights interchanged. Thus $b_i = \sigma^i c_0|\eta_0| + \sigma^{m-i}c_m|\eta_m|$ is a nonnegative supersolution and $b_0 \geq |y_0|$, $b_m \geq |y_m|$. Apply the maximum principle separately to $b - y$ and $b + y$ to get (29.2).

Dividing that bound by c_i would lose uniformity and is not the next step. Instead use the equation itself:

$$\kappa_i|y_i| \leq w_{i-1}|y_{i-1}| + w_i|y_{i+1}|.$$

Since $\kappa_i c_i = 2/R$, it follows that

$$|\xi_i| \leq \frac{R_+}{2\tau_-}(s_{i-1} + s_{i+1})\|\eta\|_\infty = \frac{R_+}{2\tau_-}(\sigma + \sigma^{-1})s_i\|\eta\|_\infty.$$

The endpoint inequalities follow from $\xi_0 = \eta_0$, $\xi_m = \eta_m$. This proves (29.3) for the entire chain. The calculation used positive incidences only at the actual regular point, with no uniform positive lower bound on them. □

29.1. **Physical flights, incidence, and every unvisited disk.** Let $v_i = (x_{i+1} - x_i)/\tau_i$ be the actual unit flight direction. Put

$$C_c = J_*\left[1 + \frac{R_+}{\tau_-}(1 + \sigma^{-1})\right], \qquad C_\Delta = R_+J_*(1 + \sigma). \tag{29.5}$$

Proposition 29.2 (Differentiation of the physical observables)**.** *On each actual regular branch,*

$$|D_q\tau_i[\eta]| \leq R_+J_*(s_i + s_{i+1})\|\eta\|_\infty, \qquad 0 \leq i < m, \tag{29.6}$$

$$\|D_q v_i[\eta]\|_2 \leq \frac{R_+J_*}{\tau_-}(s_i + s_{i+1})\|\eta\|_\infty, \qquad 0 \leq i < m, \tag{29.7}$$

$$|D_q c_i[\eta]| \leq C_c s_i\|\eta\|_\infty, \qquad 0 \leq i \leq m. \tag{29.8}$$

The true least clearance Δ_i, with precisely the two incident disks excluded as in (1.2)*, is locally Lipschitz and has the following directional bound along every continuously differentiable physical branch path, almost everywhere on that path:*

$$|D_q\Delta_i[\eta]| \leq R_+J_*\max(s_i, s_{i+1})\|\eta\|_\infty \leq 2C_\Delta\sigma^{d_i'}\|\eta\|_\infty. \tag{29.9}$$

The last assertion includes changes of the nearest unvisited disk and of its nearest point on the flight. There is no factor equal to the number of candidate disks.

Proof. $D_q x_i[\eta] = Rn'(\alpha_i)\xi_i$. Differentiate length and unit direction and use respectively $\|D|X|\| = 1$ and $D(X/|X|) = |X|^{-1}(I - vv^{\mathsf{T}})$ to prove (29.6)–(29.7). For $i < m$, $c_i = v_i \cdot n_i$, so $|Dc_i| \leq \|Dv_i\|_2 + |\xi_i|$. For the terminal collision use $c_m = -v_{m-1} \cdot n_m$ instead. The inequalities $s_{i+1} \leq \sigma^{-1}s_i$ and $s_{i-1} \leq \sigma^{-1}s_i$ give (29.8). This argument differentiates the actual geometric cosine, not an expression with an artificially introduced $1/c_i$.

The Hausdorff distance of segments $[x_i, x_{i+1}]$ and $[\widetilde{x}_i, \widetilde{x}_{i+1}]$ is at most $\max(\|x_i - \widetilde{x}_i\|_2, \|x_{i+1} - \widetilde{x}_{i+1}\|_2)$: compare points with the same affine parameter. Distance to any fixed center changes by at most that number. Taking an infimum over exactly the same set of unvisited centers preserves this inequality. Subtracting R does not change it. The fixed itinerary fixes the two excluded centers; it does not discard any other disk. This proves the local Lipschitz assertion and its derivative bound along a branch path by integration of the endpoint speeds. A Lipschitz function on a compact path interval is absolutely continuous; nearest-disk switches do not create atomic jumps. Finally

$$s_i + s_{i+1} = (1 + \sigma)(\sigma^i + \sigma^{m-1-i}) \leq 2(1 + \sigma)\sigma^{d_i'}$$

proves the last inequality in (29.9). □

29.2. **A third-variation bound uniform in collision count.** Define constants depending only on the original geometric interval,

$$C_3 = R_+ + \frac{3R_+^2}{\tau_-} + \frac{3R_+^3}{\tau_-^2},$$
$$C_F = \frac{8C_3 J_*^3 (1+\sigma)^3}{1-\sigma^3}. \tag{29.10}$$

Theorem 29.3 (Uniform physical third variation)**.** *For every actual regular endpoint action, all $m \geq 1$, all original radii, and $\eta, \zeta, \upsilon \in \mathbb{R}^2$,*

$$|D_q^3 F[\eta, \zeta, \upsilon]| \leq C_F \|\eta\|_\infty \|\zeta\|_\infty \|\upsilon\|_\infty. \tag{29.11}$$

Consequently $\|D_q G[\eta]\|_{\mathrm{op}} \leq 2C_F \|\eta\|_\infty$. For any piecewise continuously differentiable path $q(a)$ that remains in a single actual regular branch,

$$\|G(q(b)) - G(q(a))\|_{\mathrm{op}} \leq 2C_F \int_a^b \|q'(v)\|_\infty \, dv.$$

In particular this bound applies on every line segment contained in that branch. It does not assert the existence of a collision-uniform ball of physical continuation across unvisited tangencies or section cuts.

Proof. For the Euclidean norm $N(X) = |X|$, at $X \neq 0$, $\|DN\| \leq 1$, $\|D^2 N\| \leq |X|^{-1}$ and $\|D^3 N\| \leq 3|X|^{-2}$. The last inequality follows by writing its trilinear form as minus $|X|^{-2}$ times the sum of $(v \cdot a) b^{\mathsf{T}}(I - vv^{\mathsf{T}})c$ and the two cyclic terms. For one flight $X = Rn(\alpha_{i+1}) - Rn(\alpha_i) + O_{i+1} - O_i$, each derivative of n up to order three has norm one. If $V_i(\xi) = |\xi_i| + |\xi_{i+1}|$, the third-order chain rule therefore gives

$$|D^3 \tau_i[\xi, \widetilde{\xi}, \widehat{\xi}]| \leq C_3 V_i(\xi) V_i(\widetilde{\xi}) V_i(\widehat{\xi}). \tag{29.12}$$

The coefficient three in the middle term of C_3 counts the three second–first derivative terms; no such term is omitted.

Let $\mathcal{J}\eta = D_q \alpha[\eta]$ be the stationary lift. Differentiated stationarity says that $H\mathcal{J}\eta$ has zero interior coordinates, where $H = D^2 S$. From $D^2 F[\eta, \zeta] = H[\mathcal{J}\eta, \mathcal{J}\zeta]$, differentiation in υ gives the exact identity

$$D^3 F[\eta, \zeta, \upsilon] = D^3 S[\mathcal{J}\eta, \mathcal{J}\zeta, \mathcal{J}\upsilon]. \tag{29.13}$$

Indeed, derivatives of $\mathcal{J}$ have zero endpoint coordinates, and their contractions with $H\mathcal{J}\eta$ or $H\mathcal{J}\zeta$ vanish. This cancellation occurs after differentiating the full stationary action; it is not a claim that derivatives of the interior variables vanish.

By Theorem 29.1, each $V_i(\mathcal{J}\eta)$ is at most $J_*(s_i + s_{i+1})\|\eta\|_\infty$. Moreover,

$$\sum_{i=0}^{m-1} (s_i + s_{i+1})^3 \leq 4(1+\sigma)^3 \sum_{i=0}^{m-1} (\sigma^{3i} + \sigma^{3(m-1-i)}) \leq \frac{8(1+\sigma)^3}{1-\sigma^3}.$$

Sum (29.12) and use (29.13) to obtain (29.11). Each entry of $D_q G[\eta]$ is bounded by $C_F\|\eta\|_\infty$; its operator norm is at most twice that number. Integrating along the stated physical path completes the proof. □

Remark 29.4 (Domain, time, and source). Theorem 29.3 concerns the true reduced action at every regular point. It requires neither a pressure theorem nor the endpoint-momentum kernel. Uniform boundedness of its third derivative is not uniform control of all higher derivatives, a chart-count bound, or a roof-density theorem. All labels may restrict the points at which it is used, but are not differentiated. For a path of genuine trajectories with a fixed itinerary reaching a physical boundary the estimates hold up to that boundary. They do not define a collision-map derivative at a visited tangency or permit continuing an exact event through a new hit.

30. A native proof of the original regular-variation estimate

This section supplies a complete proof of (3.3), which was stated as inherited in Input 3.1. Its two averaged conclusions (3.4)–(3.5) are not proved by this calculation and remain separate inputs. The proof uses the original guard, including every unvisited obstacle, and the original insertion I_m. Nothing is conditioned on a smaller event.

Lemma 30.1 (Logarithmic sensitivity of the positive optical denominator)**.** *Regarding $\mathfrak{b}_m$ in* (4.6) *as a polynomial in the positive variables τ_i and κ_j, it has nonnegative coefficients and degree at most one in each variable. Consequently*

$$0 \le \frac{\tau_i \partial_{\tau_i} \mathfrak{b}_m}{\mathfrak{b}_m} \le 1, \qquad 0 \le \frac{\kappa_j \partial_{\kappa_j} \mathfrak{b}_m}{\mathfrak{b}_m} \le 1. \tag{30.1}$$

Proof. Each variable occurs in exactly one optical factor, that factor is linear in it, and every entry of every factor has nonnegative coefficients. Matrix multiplication preserves coefficient positivity and does not repeat a variable. For any variable x, write $\mathfrak{b}_m = a + xb$ with $a, b \ge 0$ and $\mathfrak{b}_m > 0$. Then $x\partial_x \mathfrak{b}_m / \mathfrak{b}_m = xb/(a+xb) \in [0,1]$. For $m = 1$ this applies to $\mathfrak{b}_1 = \tau_0$ and there are no curvature variables. □

Set $c_\circ = \sqrt{19}/10$ and define

$$C_\omega = \frac{2R_+ J_*(1+\sigma)}{\tau_-(1-\sigma)} + \frac{4C_c}{c_\circ} + 2C_c. \tag{30.2}$$

Proposition 30.2 (The precise inherited incidence insertion suffices)**.** *At every original exact-event regular point and for every endpoint direction,*

$$|D_q \log \omega[\eta]| \le C_\omega I_m \|\eta\|_\infty, \qquad I_m = 1 + \sum_{i=1}^{m-1} \frac{q_0^{d_i}}{c_i}. \tag{30.3}$$

Proof. The radius is fixed under D_q, so (4.7) and Lemma 30.1 imply

$$|D \log \omega| \le \frac{|Dc_0|}{c_0} + \frac{|Dc_m|}{c_m} + \sum_{i=0}^{m-1} \frac{|D\tau_i|}{\tau_i} + \sum_{i=1}^{m-1} \frac{|Dc_i|}{c_i}.$$

Here $D \log \kappa_i = -D \log c_i$; the sign is irrelevant only after taking absolute values. The original section cap gives $c_0, c_m \ge c_\circ$ and $s_0, s_m \le 2$. Also

$$\sum_{i=0}^{m-1} (s_i + s_{i+1}) \le \frac{2(1+\sigma)}{1-\sigma}, \qquad s_i \le 2\sigma^{d_i} \le 2q_0^{d_i}.$$

Apply Proposition 29.2. The first two terms in (30.2) bound the endpoint and flight contributions, and $2C_c(I_m - 1)$ bounds the interior contribution. Since $I_m \ge 1$, (30.3) follows with the displayed constant. No additional inverse incidence or collision-dependent multiplier occurs. □

30.1. **The actual guard, with nonsmooth clearance switches.** Put $M_\Theta = \sup_{x \in \mathbb{R}} |\Theta'(x)| < \infty$ for the fixed smooth cutoff already present in (1.2), and set

$$\beta = \sigma q_0^{-1/4} < 1, \qquad C_W = \frac{4M_\Theta(C_c + C_\Delta)}{1-\beta}. \tag{30.4}$$

Theorem 30.3 (Collision-uniform variation of the original physical guard)**.** *For each $\varepsilon > 0$, W^ε is locally Lipschitz on an actual regular branch. Along every continuously differentiable physical branch path it is absolutely continuous on compact parameter intervals and, almost everywhere,*

$$|D_q W^\varepsilon[\eta]| \le C_W \varepsilon^{-1} \|\eta\|_\infty. \tag{30.5}$$

The bound is independent of the collision count, exact labels, and radius. No section-distance factor is inserted in W^{ε}.

Proof. All guard factors lie in $[0,1]$. Apply the product rule along a compact path and use Proposition 29.2; for clearance factors use the almost-everywhere chain rule for Lipschitz functions on an interval. The collision and flight depth schedules are exactly those in the original guard, not new schedules. Each depth occurs at most twice, so

$$\sum_{i=0}^{m}\frac{|Dc_i|}{q_0^{d_i/4}}\le\frac{4C_c}{1-\beta}\|\eta\|_\infty,\qquad \sum_{i=0}^{m-1}\frac{|D\Delta_i|}{q_0^{d'_i/4}}\le\frac{4C_\Delta}{1-\beta}\|\eta\|_\infty.$$

Multiplication by M_Θ/ε proves (30.5). The inequality $\beta<1$ follows, for example, from $\sigma<q_0<1$. The clearance is locally Lipschitz even at a nearest-disk switch, so there is no omitted jump contribution at such a switch. Actual section boundaries and collision-word boundaries are not crossed by this argument; their one-sided physical traces retain their earlier status. □

30.2. **Roofwise variation and physical normalization.** Keep $\lambda_\circ=9\sqrt{19}/200$ and $\Lambda_\circ=47/6$. Define

$$C_{\mathrm{reg}}=\frac{\Lambda_\circ}{\lambda_\circ}\left(C_\omega+\frac{2C_F}{\lambda_\circ}+C_W\right).\tag{30.6}$$

Theorem 30.4 (Native regular variation on every actual roof-angle segment)**.** *On every regular segment of the original physical pairing space, with $u=F$, $e=\nabla F/|\nabla F|$ fixed along the segment, $z=|\nabla F|>0$, and $A=\omega/(4\pi c_*\chi)$, the original density $f^{\varepsilon}=AW^{\varepsilon}$ is locally absolutely continuous and*

$$|\partial_u f^{\varepsilon}|\le C_{\mathrm{reg}}(1+\varepsilon^{-1})\frac{AI_m}{z}\quad\text{almost everywhere.}\tag{30.7}$$

This is precisely (3.3)*, with a finite explicit constant depending only on the original geometry and cutoff. Its proof uses no pressure, curve-growth, mixing, or local-limit theorem.*

Proof. The two-dimensional determinant defining χ equals $e^{\perp\mathsf{T}}Ge^{\perp}$, as in the retained Gauss-coordinate calculation. Thus $\chi\ge\lambda_\circ$. Differentiating on a fixed-angle segment, not in an arbitrary direction with changing e, gives

$$\left\|\frac{dq}{du}\right\|_\infty=\left\|\frac{G^{-1}e}{ze^{\mathsf{T}}G^{-1}e}\right\|_\infty\le\frac{\Lambda_\circ}{\lambda_\circ z},\qquad|\partial_u\chi|\le 2C_F\left\|\frac{dq}{du}\right\|_\infty.$$

The latter bound is Theorem 29.3; there is no angular derivative term because e is fixed. By Proposition 30.2,

$$|\partial_u\log A|\le\left(C_\omega I_m+\frac{2C_F}{\lambda_\circ}\right)\frac{\Lambda_\circ}{\lambda_\circ z}.$$

The factor $4\pi c_*$ is the unchanged physical probability normalization and is constant under this differentiation. Using $0\le W^{\varepsilon}\le 1$ and Theorem 30.3 now yields

$$|\partial_u(AW^{\varepsilon})|\le\frac{A\Lambda_\circ}{\lambda_\circ z}\left(C_\omega I_m+\frac{2C_F}{\lambda_\circ}+\frac{C_W}{\varepsilon}\right).$$

Since $I_m\ge 1$, this implies (30.7) with (30.6). Compact subsegments have positive incidences and $z>0$, so all differentiations and absolute-continuity claims are legitimate there. The uniform constants do not depend on those compact-subsegment minima. □

Corollary 30.5 (Explicit integration up to the actual endpoint)**.** *For two roof values in one original regular fixed-angle segment,*

$$|f^{\varepsilon}(u_2)-f^{\varepsilon}(u_1)|\le C_H(1+\varepsilon^{-1})\sigma^{(m-2)_+}\sqrt{|u_2-u_1|},\quad C_H=\frac{500C_{\mathrm{reg}}\sqrt{2\Lambda_\circ}}{4\pi c_*\lambda_\circ^2}.\tag{30.8}$$

This supplies a native regular-interior premise for the retained endpoint-trace results, rather than a new claim of zero trace on a bulk-null unvisited or section boundary.

Proof. Use the already proved physical trajectory bound $\omega I_m \le 500\sigma^{(m-2)_+}$ in Corollary 4.4, $\chi \ge \lambda_\circ$, and integrate (30.7). Lemma 5.1 gives $\int_{u_1}^{u_2} du/z \le \lambda_\circ^{-1}\sqrt{2\Lambda_\circ|u_2-u_1|}$. This proves (30.8) on compact subsegments and then at one-sided limits. Identification of those limits with the actual physical pairing and treatment of simultaneous singular endpoints remain the retained trace construction; no derivative of a grazing collision map is introduced here. □

Proposition 30.6 (Complete positive variation domination, before any supremum)**.** *On the retained countable one-copy sheet space, let $h \ge 0$ be any Borel selector (including the original exact record and actual roof). Then, allowing both sides to be infinite,*

$$\int h(u,\theta)|\partial_u f^\varepsilon(u,\theta)|\,du\,d\theta \le C_{\mathrm{reg}}(1+\varepsilon^{-1})\int h(u,\theta)\frac{AI_m}{z}(u,\theta)\,du\,d\theta. \tag{30.9}$$

For selectors h pulled back from physical states, the right side is $C_{\mathrm{reg}}(1+\varepsilon^{-1})\int_\Pi h\,I_m z^{-1}\,d\nu_R^*$ under the original coarea convention.

Proof. Multiply (30.7) by h on compact regular subsegments, integrate, then exhaust each open segment and the countable sheet space. All quantities are nonnegative, so monotone convergence proves (30.9) without a word-count factor or a signed-series exchange. The final identification is precisely the retained physical one-copy coarea input. It is not a new construction of pairing multiplicity and does not identify the paired minimum of endpoint traces with the interior density. □

30.3. **What is closed, and the unchanged remaining mathematical limit.** The regular-variation clause of Input 3.1 now has a full native geometric proof. The previous trace, clearance, radial, and critical coefficient theorems remain in the manuscript and are not reclassified as new results. In particular the two averaged clauses of that input are not consequences of bounded D^3F. Neither (30.9) nor the per-trajectory exponential factor in (30.8) can be multiplied by a physical-word count to obtain a normalized roof-density estimate.

The original unfinished assertion is still (15.8), namely the same positive low-load shortage after subtraction of the actual nongrazing capacity, with the complete physical sum and roof essential supremum preceding the collision limsup, and the outer limit in B last. A proof must still supply the complete normalized same-physical-roof source or a valid capacity comparison, with the actual pairing endpoint measure and the first-order clearance and incidence-only sources retained. The spectral/arithmetic and conditional-denominator conclusions retain their exact hypotheses. This section removes a specific local optical proof dependency; it does not replace this missing global result by a new residual or a relaxed window. The full raw pointwise local limit theorem and final four-coordinate inversion are not asserted as completed.

31. A grazing-uniform resolvent for the physical reflection equations

We retain the actual circular-scatterer trajectories, radius interval, unit-speed clock, and all exact selectors of (1.1). The issue here is the derivative of an endpoint boundary-value problem, not a derivative bound for the expanding forward collision map. All derivatives are taken at regular physical points of a fixed actual itinerary. No unvisited disk is removed from its admissible domain.

Put $a_* = R_+/\tau_- = 47/6$, $\lambda = 50/47$, and $I = \{1,\dots,m-1\}$. A sum over I is empty when $m = 1$. Write $X_i = x_{i+1} - x_i$, $\vartheta_i = \arg X_i$, using local real lifts of these angles. The specular law is equivalent, on the physical branch, to constancy of

$$E_i(\alpha) = 2\alpha_i - \vartheta_{i-1}(\alpha) - \vartheta_i(\alpha), \qquad i \in I. \tag{31.1}$$

Its constant is an odd multiple of π. The value of that multiple does not enter a derivative; it is fixed on each local lift. The reflection equation is not obtained by assuming that division by a small incidence is harmless.

Lemma 31.1 (The full physical Jacobian)**.** *At an actual regular trajectory, set $w_i = \tau_i^{-1}$ and $c_i = \sqrt{1-p_i^2}$. The interior Jacobian $B = D_{\alpha_I}E$ is tridiagonal, with*

$$B_{ii} = 2 + Rc_i(w_{i-1} + w_i), \tag{31.2}$$

$$B_{i,i-1} = Rw_{i-1}c_{i-1}, \tag{31.3}$$

$$B_{i,i+1} = Rw_ic_{i+1}, \tag{31.4}$$

$$(D_{\alpha_0}E)_1 = Rw_0c_0, \tag{31.5}$$

$$(D_{\alpha_m}E)_{m-1} = Rw_{m-1}c_m. \tag{31.6}$$

Only entries with indices in I belong to B; the last two entries are boundary forcing. When $m = 2$ both boundary terms are in the same row. If $H = D^2S$ is the full stationary-action Hessian, then

$$H_{II} = R\,\mathrm{diag}(c_i)_{i\in I}\,B. \tag{31.7}$$

Proof. For an oriented physical flight $v_i = X_i/\tau_i$, the differential of its direction is $D\vartheta_i = v_i^{\perp}\cdot DX_i/\tau_i$. The departure and arrival normal components are respectively c_i and $-c_{i+1}$. Since $n'(\alpha) = n(\alpha)^{\perp}$,

$$\partial_{\alpha_i}\vartheta_i = -Rc_iw_i, \qquad \partial_{\alpha_{i+1}}\vartheta_i = -Rc_{i+1}w_i.$$

Applying these two identities to both flights in (31.1) proves every entry, including both boundary terms. Comparing with (4.2) proves (31.7). Thus a factor c_i in the action Hessian is a row factor in the gradient formulation; it is absent from the reflection-angle Jacobian. This identity does not assert invertibility of H_{II} when a limiting c_i is zero. □

Theorem 31.2 (Column coercivity and a positive resolvent)**.** *Let $\mathcal{S} = \mathrm{diag}((-1)^i)_{i\in I}$ and $A = \mathcal{S}B\mathcal{S}$. For every $m \geq 2$, every original radius, and every actual regular itinerary, A^{-1} is entrywise nonnegative and*

$$\|B^{-1}f\|_1 \leq \tfrac{1}{2}\|f\|_1. \tag{31.8}$$

More generally, for positive weights v_i such that $\lambda^{-1} \leq v_{i+1}/v_i \leq \lambda$, one has

$$\sum_{i\in I} v_i|(B^{-1}f)_i| \leq \sum_{i\in I} v_i|f_i|. \tag{31.9}$$

In particular,

$$\sum_{i\in I} \lambda^{|i-j|}(A^{-1})_{ij} \leq 1 \quad (j \in I). \tag{31.10}$$

The assertions extend algebraically to simultaneous values $c_i = 0$ by the same formulas. No lower incidence cutoff is required.

Proof. The off-diagonal entries of A are nonpositive. Let $D_{jj} = 2 + Rc_j(w_{j-1} + w_j)$ and write $A = (\mathrm{Id} - N)D$, where N has zero diagonal and nonnegative off-diagonal entries. Its column sum is at most

$$\frac{Rc_j(w_{j-1} + w_j)}{2 + Rc_j(w_{j-1} + w_j)} \leq \frac{a_*}{1 + a_*} = \frac{47}{53} < 1.$$

Consequently $A^{-1} = D^{-1}\sum_{k\geq 0} N^k$ exists and is nonnegative. This is an algebraic matrix series, not a probabilistic replacement of the Lorentz gas. Missing interior neighbors at the two ends only decrease the column sum.

Each column of A has sum at least 2: $\mathbf{1}^{\mathsf{T}}A \geq 2\mathbf{1}^{\mathsf{T}}$. Multiplication on the right by the nonnegative inverse gives $\mathbf{1}^{\mathsf{T}}A^{-1} \leq \mathbf{1}^{\mathsf{T}}/2$. Since $|B^{-1}f| \leq A^{-1}|f|$, this proves (31.8). The

argument uses *column*, not row, dominance. Row dominance can fail when one incidence is much smaller than its neighbors.

For the weighted assertion, a column calculation gives

$$(v^{\mathsf{T}}A)_j \geq v_j\{2-(\lambda-1)Rc_j(w_{j-1}+w_j)\} \geq v_j\{2-2a_*(\lambda-1)\} = v_j.$$

Here $2a_*(\lambda-1)=1$ exactly. Right multiplication by A^{-1} proves (31.9). Taking $v_i=\lambda^{|i-j|}$ proves (31.10). All denominators in this argument are at least 2. It therefore remains valid when any collection of incidences is zero. No inverse c_i has been introduced in taking that limit. □

31.1. **Complete higher endpoint derivatives.** For a vector $u=(u_0,\dots,u_m)$ set $V_i(u)=|u_i|+|u_{i+1}|$ and $\|u\|_{1,v}=\sum_{i=0}^{m} v_i|u_i|$. We use $v_i=\lambda^{d_i}$, $d_i=\min(i,m-i)$, including both endpoints. For $r\geq 1$ define the explicit constants

$$T_r=\sum_{k=1}^{r}\left\{{r \atop k}\right\}(k-1)!a_*^k, \qquad T_1=a_*, \quad T_2=a_*+a_*^2, \quad T_3=a_*+3a_*^2+2a_*^3, \tag{31.11}$$

where $\left\{{r \atop k}\right\}$ counts partitions of $\{1,\dots,r\}$ into k nonempty blocks.

Lemma 31.3 (Local angle tensors, with every chain-rule term)**.** *For each flight and each $r\geq 1$,*

$$|D^r\vartheta_i[u^1,\dots,u^r]| \leq T_r\prod_{h=1}^{r} V_i(u^h).$$

For $r\geq 2$, the full vector $E=(E_i)_{i\in I}$ consequently satisfies

$$\|D^rE[u^1,\dots,u^r]\|_1 \leq 4T_r\prod_{h=1}^{r}\|u^h\|_1, \tag{31.12}$$

$$\|D^rE[u^1,\dots,u^r]\|_{1,v} \leq 2(1+\lambda)T_r\prod_{h=1}^{r}\|u^h\|_{1,v}. \tag{31.13}$$

Proof. Identify the plane with $\mathbb{C}$ and use a local logarithm of $X_i\neq 0$. Its kth derivative is $(-1)^{k-1}(k-1)!X_i^{-k}$; taking the imaginary part gives the direction derivative. Every derivative of $X_i=O_{i+1}-O_i+R(e^{i\alpha_{i+1}}-e^{i\alpha_i})$ of order $b\geq 1$ has absolute value at most $R_+\prod_{h=1}^{b}V_i(u^h)$. The composition rule indexed by set partitions gives exactly the sum in (31.11), using $|X_i|\geq\tau_-$. This proves the first assertion without omitting mixed derivatives.

Each flight appears in at most two interior reflection equations. Also $V_i(u)\leq\|u\|_1$ and $\sum_i V_i(u)\leq 2\|u\|_1$. Summing one factor and bounding the others proves the first norm inequality. For the second, the sum of the weights of the adjacent reflection equations is at most v_i+v_{i+1}, and

$$(v_i+v_{i+1})V_i(u) \leq (1+\lambda)(v_i|u_i|+v_{i+1}|u_{i+1}|).$$

Summing over flights counts each vertex at most twice. The other factors are bounded by their weighted norms, since $v_i\geq 1$. This proves the second inequality, including the boundary flights. □

Let $\mathfrak{P}_r$ denote the set of all set partitions of $\{1,\dots,r\}$. Define two finite positive recursions

$$C_1=2+a_*=59/6, \qquad C_r=2\sum_{\substack{\pi\in\mathfrak{P}_r\\|\pi|\geq 2}} T_{|\pi|}\prod_{Q\in\pi}C_{|Q|}, \tag{31.14}$$

$$M_1=2+2\lambda a_*=56/3, \qquad M_r=2(1+\lambda)\sum_{\substack{\pi\in\mathfrak{P}_r\\|\pi|\geq 2}} T_{|\pi|}\prod_{Q\in\pi}M_{|Q|}, \tag{31.15}$$

$$r\geq 2. \tag{31.16}$$

The symbols C_r and M_r denote the jet constants in this section and the next, not a redefinition of earlier flight-tensor constants. Every block in these sums has size less than r, so the recursions are not circular. In particular $C_2 = 2T_2C_1^2$ and $C_3 = 2T_3C_1^3 + 6T_2C_1C_2$.

Theorem 31.4 (All finite endpoint jets on actual physical branches)**.** *For every integer $r \geq 1$, every $m \geq 1$, and every actual regular stationary lift $\alpha(q)$, $q = (\alpha_0, \alpha_m)$,*

$$\|D_q^r \alpha[\eta^1, \dots, \eta^r]\|_1 \leq C_r \prod_{h=1}^{r} \|\eta^h\|_\infty, \tag{31.17}$$

$$\sum_{i=0}^{m} \lambda^{d_i} |D_q^r \alpha_i[\eta^1, \dots, \eta^r]| \leq M_r \prod_{h=1}^{r} \|\eta^h\|_\infty. \tag{31.18}$$

The constants depend on derivative order and the original geometric interval only. They are independent of collision count, itinerary, small positive incidence, return/displacement labels, or roof window.

Proof. For $r = 1$, the interior equation is $BJ_I = -D_q E\,\eta$. Its right side has norm at most $a_*(|\eta_0| + |\eta_m|)$ and weighted norm at most $\lambda a_*(|\eta_0| + |\eta_m|)$. Theorem 31.2, followed by adding the two actual endpoint derivatives, gives C_1 and M_1. This also covers $m = 2$, when the two boundary terms share a row. For $m = 1$ only the two endpoint derivatives occur.

For $r \geq 2$ the endpoint derivatives of order r vanish. Differentiate $E(\alpha(q)) =$ constant in the r prescribed directions. Isolating the partition with one block gives

$$B(D_q^r \alpha)_I = -\sum_{\substack{\pi \in \mathfrak{P}_r \\ |\pi| \geq 2}} D^{|\pi|} E[D_q^{|Q|} \alpha[\eta^h : h \in Q] : Q \in \pi].$$

This identity contains every mixed chain-rule term exactly once. Induction, Lemma 31.3, and respectively the $1/2$ and 1 inverse bounds yield precisely (31.16). Local smoothness follows already from the ordinary implicit function theorem and invertibility of B; the induction establishes uniform bounds for its derivatives. No assertion of a common physical continuation ball is used. □

Corollary 31.5 (Fixed-itinerary grazing limits, without a fictitious event)**.** *Fix m and a sequence of actual regular trajectories with the same centers and convergent collision angles and radius. Suppose the limit is in the original radius interval. Any number of interior incidences may tend to zero. The local reflection equations at the limiting configuration have a unique smooth formal solution in the endpoint angles; every finite endpoint jet of the original physical branch has that formal limit. The bounds in* (31.18) *persist. This does not make the formal continuation an admissible billiard trajectory beyond its actual first boundary.*

Proof. Distinct successive disk centers and $R \leq R_+$ keep every limiting flight length at least τ_-. Local direction logarithms are smooth there. The limiting Jacobian is invertible by Theorem 31.2, including simultaneous zero incidences. The implicit function theorem at that configuration supplies the formal reflection solution, with the radius also treated as a smooth parameter. Local uniqueness identifies every sufficiently close member of the given physical sequence with it. Its finite jets converge, and their uniform bounds pass to the limit. An extra disk can obstruct the formal solution, and the prescribed section or return event can end at the boundary. Neither obstruction is discarded; no collision-map derivative at tangency is asserted. □

32. Fourth physical variation and the exact high-order cancellation

The third-variation result of Section 29 is retained. It does not by itself bound a fourth derivative: differentiating the stationary lift introduces its second derivative. The reflection

resolvent now controls that derivative without a small-incidence loss. The following proof includes those terms rather than continuing the third-order Schur cancellation beyond its valid scope.

Set

$$D_3 = R_+ + 3R_+^2/\tau_- + 3R_+^3/\tau_-^2, \tag{32.1}$$

$$D_4 = R_+ + 7R_+^2/\tau_- + 18R_+^3/\tau_-^2 + 15R_+^4/\tau_-^3, \tag{32.2}$$

$$\mathcal{F}_4 = 2D_4C_1^4 + 6D_3C_2C_1^2, \qquad C_1 = 59/6, \quad C_2 = 2(a_* + a_*^2)C_1^2. \tag{32.3}$$

All constants concern the unchanged interval $R \in [9/20, 47/100]$.

Lemma 32.1 (Fourth tensor of one physical flight)**.** *For $r = 3, 4$ and arbitrary collision-angle variations,*

$$|D^r\tau_i[u^1, \dots, u^r]| \le D_r \prod_{h=1}^{r} V_i(u^h), \qquad |D^rS[u^1, \dots, u^r]| \le 2D_r \prod_{h=1}^{r} \|u^h\|_1.$$

Proof. For $N(X) = |X|$, put $v = X/|X|$ and $P = \mathrm{Id} - vv^{\mathsf{T}}$. Then $DN = v^{\mathsf{T}}$, $D^2N = P/|X|$, and

$$D^3N[a, b, c] = -|X|^{-2}\{(v \cdot a)b^{\mathsf{T}}Pc + (v \cdot b)a^{\mathsf{T}}Pc + (v \cdot c)a^{\mathsf{T}}Pb\}.$$

Thus $\|D^3N\| \le 3|X|^{-2}$. Differentiating this display gives $\|D^4N\| \le 15|X|^{-3}$: differentiating the radial prefactor, the three copies of v, and the three copies of P contributes at most 6, 3, and 6, respectively, since $Dv[d] = Pd/|X|$ and $DP[d] = -\{(Pd)v^{\mathsf{T}} + v(Pd)^{\mathsf{T}}\}/|X|$.

Each derivative of a circular endpoint position has size at most R_+ times the product of its angular variations. The fourth composition rule has respectively $1, 7, 6, 1$ set partitions into $1, 2, 3, 4$ blocks. Multiplying by the norm-derivative bounds $1, |X|^{-1}, 3|X|^{-2}, 15|X|^{-3}$ proves the displayed D_4. The third-order rule gives D_3, as in the retained proof. Finally $\sum_i \prod_h V_i(u^h) \le 2\prod_h \|u^h\|_1$, by summing one factor and taking the suprema of the others. This proves both full-action inequalities independently of m. □

Theorem 32.2 (Collision-uniform fourth derivative of the reduced action)**.** *On every actual regular endpoint branch and for all four endpoint directions,*

$$|D_q^4F[\eta^1, \eta^2, \eta^3, \eta^4]| \le \mathcal{F}_4 \prod_{h=1}^{4} \|\eta^h\|_\infty. \tag{32.4}$$

There is no interior-incidence cutoff. In particular,

$$\|D_q^2G[\eta, \zeta]\|_{\mathrm{op}} \le 2\mathcal{F}_4\|\eta\|_\infty\|\zeta\|_\infty.$$

At fixed-itinerary multiple-grazing limits, these tensors have the one-sided limits specified in Corollary 31.5.

Proof. Write $J_a = D_q\alpha[\eta^a]$ and $K_{ab} = D_q^2\alpha[\eta^a, \eta^b]$. The latter has zero endpoint coordinates. Differentiate the exact third-order identity (29.13) in direction η^4. One obtains

$$\begin{aligned} D_q^4F[\eta^1, \eta^2, \eta^3, \eta^4] = {} & D^4S[J_1, J_2, J_3, J_4] \\ & + D^3S[K_{14}, J_2, J_3] + D^3S[J_1, K_{24}, J_3] + D^3S[J_1, J_2, K_{34}]. \end{aligned} \tag{32.5}$$

The three terms containing K do not vanish. Their omission would be an incorrect fourth-order use of stationary elimination. Equivalently, let $g_{ab} = (D^3S[\mathbf{e}_i, J_a, J_b])_{i\in I}$. Twice differentiated stationarity gives $H_{II}(K_{ab})_I = -g_{ab}$. Hence the last three terms are

$$-g_{14}^{\mathsf{T}}H_{II}^{-1}g_{23} - g_{24}^{\mathsf{T}}H_{II}^{-1}g_{13} - g_{34}^{\mathsf{T}}H_{II}^{-1}g_{12}.$$

This form verifies the symmetry but is not used to bound H_{II}^{-1} near grazing. The reflection equation gives instead $\|J_a\|_1 \le C_1\|\eta^a\|_\infty$ and $\|K_{ab}\|_1 \le C_2\|\eta^a\|_\infty\|\eta^b\|_\infty$. Lemma 32.1 applied

to all four terms of (32.5) gives exactly (32.3). For $m = 1$, $K = 0$ and the same bound holds. Each entry of $D_q^2 G[\eta, \zeta]$ is bounded by $\mathcal{F}_4 \|\eta\|_\infty \|\zeta\|_\infty$; a two-by-two matrix has operator norm at most twice its largest entry. The fixed-itinerary limit assertion follows from the smooth formal reflection continuation already proved, restricted to the original physical branch. □

Corollary 32.3 (Uniform Taylor errors on the actual physical domain)**.** *Suppose the entire straight segment $q + sh$, $0 \le s \le 1$, lies in a single actual regular branch. Then*

$$\left| F(q+h) - F(q) - DF(q)[h] - \tfrac{1}{2} D^2 F(q)[h,h] - \tfrac{1}{6} D^3 F(q)[h,h,h] \right| \le \tfrac{1}{24} \mathcal{F}_4 \|h\|_\infty^4, \tag{32.6}$$

$$\|G(q+h) - G(q) - DG(q)[h]\|_{\mathrm{op}} \le \mathcal{F}_4 \|h\|_\infty^2. \tag{32.7}$$

The same bounds hold by a one-sided limit at a fixed-itinerary grazing endpoint. They require neither a pressure estimate nor a prescribed positive lower bound on intermediate incidences.

Proof. Apply the one-dimensional integral Taylor remainder to the stated segment. The remainder integral for F is $\frac{1}{6} \int_0^1 (1-s)^3 D^4 F(q+sh)[h,h,h,h]\, ds$. For G, integrate its second derivative with weight $1 - s$. The bounds follow from Theorem 32.2; its one-sided limit justifies the endpoint statement. The hypothesis that the segment stays physical is not inferred from the derivative bound. □

32.1. **What this resolves in the original source problem.** The higher reflection jets and (32.5) remove an actual local analytic obstruction: repeated stationary differentiation need not incur powers of the smallest visited incidence or an uncontrolled factor equal to the collision count. They also justify the displayed higher-order Taylor errors on every existing physical chart. These are uniform statements about all actual regular itineraries, not assertions from a finite sample or a replacement Markov model.

They do not count physical charts or continue through a newly hit disk. The least unvisited clearance can still cut a chart, the original section/return selector can still jump, and a positive one-sided paired trace is not automatically a bulk-collision submeasure. In particular no norm of the algebraic reflection resolvent is a pressure estimate for the twisted transfer operator.

The remaining target is still (15.8), with $B \to \infty$ outermost and the full one-copy physical sum made before the original radius, exact-label and roof suprema and collision limsup. Neither (31.18) nor (32.4) supplies the missing m^{-2} common-physical-roof summation or nongrazing capacity. The two averaged clauses of Input 3.1, the stronger pressure/pairing interfaces and the original exact arithmetic and conditioned-denominator hypotheses retain their distinct status. No theorem or target is deleted to hide these remaining obligations.

33. Uniform analytic lifts and genuine physical continuation tubes

All radii, disks, collision labels, physical clocks and the original section are unchanged. The constants in this section are deliberately conservative; their purpose is a positive radius independent of the number of actual collisions. Unlike a bound for each fixed derivative, the first theorem provides one common domain of analytic continuation. A separate geometric argument determines when this continuation is a true billiard trajectory. The latter step is not inferred from analyticity.

Let $a = 47/6$, $\lambda = 50/47$, $v_i = \lambda^{\min(i,m-i)}$, and set

$$K = 2(1+\lambda)(4a+16a^2), \qquad r_* = \min\{[8(1+a)]^{-1}, (8K)^{-1}\},$$
$$\eta_* = \frac{r_*}{8(1+\lambda a)}, \qquad P_* = 2(1+\lambda a), \tag{33.1}$$
$$C_{\mathrm{inc}} = P_*[1+2a(1+\lambda)], \qquad C_{\mathrm{clr}} = R_+P_*,$$
$$\delta_* = \min\{\eta_*, (2C_{\mathrm{inc}})^{-1}, (2C_{\mathrm{clr}})^{-1}\}. \tag{33.2}$$

The symbols r_*, η_*, δ_* in this section are auxiliary endpoint-domain constants, not the original microscopic width r_B. For complex vectors put $\|z\|_v = \sum_{i=0}^m v_i|z_i|$.

Lemma 33.1 (Uniform complex neighborhood of the reflection equations)**.** *At every actual regular reference trajectory α^0, the reflection vector E of* (31.1) *has a holomorphic continuation to $\|\alpha - \alpha^0\|_v < r_*$. On that ball,*

$$\|D^2E(\alpha)[z,w]\|_v \le K\|z\|_v\|w\|_v.$$

The output norm uses only interior indices. Both complexified flight coordinates stay nonzero, uniformly in collision count and incidence.

Proof. Write the two complexifications of a flight as

$$X_i^+ = (O_{i+1}-O_i)_x + i(O_{i+1}-O_i)_y + R(e^{i\alpha_{i+1}} - e^{i\alpha_i}),$$

and let X_i^- be the same expression with i replaced by $-i$. For complex angles X_i^- is not defined by complex conjugation. At the real reference point $|X_i^+| = |X_i^-| = \tau_i \ge \tau_-$. For $\|\alpha - \alpha^0\|_v \le r_*$,

$$|X_i^\pm(\alpha) - X_i^\pm(\alpha^0)| \le R_+e^{r_*}(|\alpha_i - \alpha_i^0| + |\alpha_{i+1} - \alpha_{i+1}^0|) \le 2R_+r_* < \tau_-/4.$$

Here $r_* < 1/8$, $e^{r_*} < 2$, and $2ar_* < 1/4$. Choose the logarithms on these nonvanishing disks and put $\vartheta_i = (\log X_i^+ - \log X_i^-)/(2i)$, with the additive real lift fixed at the reference point. This agrees with the real flight angle. Each first or second derivative of $X_i^\pm$ has bound $2R_+\prod_h(|z_i^h| + |z_{i+1}^h|)$, and $|X_i^\pm| \ge \tau_-/2$. The two logarithmic derivative terms are consequently bounded by $4a$ and $16a^2$. The weighted flight summation in Lemma 31.3 then gives $2(1+\lambda)(4a+16a^2)$, without a factor m. □

Theorem 33.2 (A collision-uniform analytic endpoint domain)**.** *Fix any actual regular m-flight trajectory, $R \in [R_-, R_+]$, and its designated disk centers. For every complex endpoint increment $h = (h_0, h_m)$ with $s = |h_0| + |h_m| < \eta_*$ there is a unique solution of the same locally lifted reflection equations in the interior ball $\|\alpha_I - \alpha_I^0\|_v \le r_*/2$. It depends holomorphically on h, is real for real h, and satisfies*

$$\sum_{i=0}^m v_i|\alpha_i(h) - \alpha_i^0| \le P_*s. \tag{33.3}$$

Uniqueness is asserted in this ball, not among arbitrary distant solutions of the word equations. A simultaneous visited-grazing limit of reference configurations has the same formal statement. No physical passage through grazing is asserted.

Proof. For $m = 1$ the assertion is just the two endpoints. Otherwise let $B_0 = D_IE(\alpha^0)$ and let Q_0h be its boundary forcing. Theorem 31.2 gives $\|B_0^{-1}\|_{v\to v} \le 1$ also over complex vectors, and $\|Q_0h\|_v \le \lambda as$. Write

$$E(\alpha^0 + (h,u)) - E(\alpha^0) = B_0u + Q_0h + N(h,u).$$

On the stated ball $s + \|u\|_v < r_*$, the preceding lemma and integration along the complex line segment give $\|N(h,u)\|_v \le K(s+\|u\|_v)^2/2$ and $\|D_uN\| \le K(s+\|u\|_v)$. The map $u \mapsto -B_0^{-1}(Q_0h + N(h,u))$ sends $\|u\|_v \le r_*/2$ into itself: its forcing is at most $r_*/8$ and its nonlinear

part is at most $r_*/16$. Its Lipschitz constant is at most $Kr_* \le 1/8$. The successive iterates from zero thus converge geometrically, uniformly on every smaller endpoint domain, to the unique fixed point. The iterates are holomorphic, so their locally uniform limit is holomorphic. Real coefficients and uniqueness give reality for real h. For the solution, $s + \|u\|_v \le r_*$ yields

$$\|u\|_v \le \lambda a s + \tfrac{1}{16}(s + \|u\|_v), \qquad s + \|u\|_v \le \tfrac{16}{15}(1 + \lambda a)s \le P_* s.$$

This proves (33.3). At a fixed-word grazing limit the real limiting flight lengths are still at least τ_- and the limiting reflection inverse retains its weighted bound. The identical contraction proof applies there. It concerns reflection equations only until physical admissibility is verified separately below. □

Theorem 33.3 (Physical tubes for the original depth-dependent guard)**.** *Put $r = q_0^{1/4}$, with the unchanged $q_0 = 47/53$. Suppose an actual regular reference orbit satisfies, for $0 < \varepsilon \le 1/4$,*

$$c_i^0 \ge \varepsilon r^{d_i} \quad (0 \le i \le m), \qquad \Delta_i^0 \ge \varepsilon r^{d_i'} \quad (0 \le i < m). \tag{33.4}$$

For every real h with $|h_0| + |h_m| < \delta_* \varepsilon$, the analytic lift is an actual regular trajectory with exactly the same ordered collisions and displacement. Every designated incidence and every least third-disk clearance obeys

$$c_i(h) \ge \tfrac{1}{2}\varepsilon r^{d_i}, \qquad \Delta_i(h) \ge \tfrac{1}{2}\varepsilon r^{d_i'}. \tag{33.5}$$

The physical clock is the sum of the true lengths. The original section, visit count and terminal test are evaluated at the resulting physical orbit; they are not declared constant on the tube. Thus any Borel exact-record selector can be retained pointwise without adding a section-distance condition to the guard.

Proof. The norm bound in (33.3) gives $|\alpha_i(h) - \alpha_i^0| \le P_* s \lambda^{-d_i}$. For real angles the corresponding endpoint displacement is at most $R_+ P_* s \lambda^{-d_i}$. The Hausdorff distance between the two closed flight segments is at most $C_{\mathrm{clr}} s \lambda^{-d_i'}$. Distance to every third disk is one-Lipschitz for this distance; taking the infimum over all centers except exactly the two incident centers preserves it. This controls every unvisited obstacle at once, including switches of the minimizing obstacle or closest segment point.

For a nonzero real flight, comparison of normalized vectors gives $|X/|X| - X^0/|X^0|| \le 2|X - X^0|/|X^0|$. Since adjacent depth weights have ratio at most λ, each signed departure normal component, and each negative arrival normal component, changes by at most $C_{\mathrm{inc}} s \lambda^{-d_i}$. The same bound holds for the signed tangential component. Initially both physical normal components at an interior reflection equal c_i^0. The exact rational inequality

$$\lambda^4 q_0 = \frac{50^4}{47^3\, 53} > 1$$

implies $\lambda^{-d} \le r^d$. The choice of δ_* therefore proves both half-margins, first for signed normal components and segment clearances of the formal real lift.

Their strict positivity makes that lift physical. A positive outgoing normal component keeps a segment outside its initial disk; a negative incoming normal component keeps it outside its final disk until the stated endpoint. All other disks have strictly positive clearance. Thus each segment is the true next free flight, not a flight with a hidden earlier collision. The fixed reflection equations, together with these normal signs, give specular reflection at every interior endpoint. Concatenation proves the actual ordered word and unchanged displacement, with no deleted collision. Its unit-speed duration is its geometric length. Section membership can jump as its boundary is crossed, so the original selector is retained rather than differentiated or frozen. This makes no assertion about the source outside the tube. □

The guard margins in (33.4) already occur in the original product (1.2); they are not a new conditioning event for the target probability. The theorem establishes an explicit domain property of that existing geometric class. Points failing the margins, and all one-sided singular traces, remain present in the full source problem. No claim that their normalized mass tends to zero is made by constructing these tubes.

34. Optical distortion and summation at one common physical roof

The synchronized momentum sum in (28.7) is not itself a sum at one physical roof: a fixed roof generally selects a different momentum radius on every critical branch. We now supply an explicit comparison mechanism on the uniform physical tubes just constructed. It uses logarithmic optical distortion before summation, not an exchange of the roof supremum and the sum. The only global root input is the retained physical kernel Theorem 26.4, with its native growth dependencies left unchanged.

Set $\sigma = 47/59$, $r = q_0^{1/4}$ and $\beta = \sigma/r < 1$. The constants J_*, C_c, C_F retain the exact meanings given in Section 29. Define

$$g = R_-/2 = 9/40, \qquad U = 320/9,$$
$$D_* = \frac{8C_c}{1-\beta} + \frac{4aJ_*}{1-\sigma} + \frac{4C_F}{g}, \qquad C_{\mathrm{dist}} = \exp(D_*\delta_*/2). \tag{34.1}$$

No constant in this display depends on collision count, labels or an unproved lower bound for an interior incidence.

Lemma 34.1 (Relative distortion of the actual inverse momentum weight)**.** *On any convex physical endpoint domain with $c_i \ge (\varepsilon/2)r^{d_i}$ and $c_0, c_m \ge 1/2$, put $G = D_q^2 F$, $e = G_{0m}$ and*

$$u(q) = \frac{R|e(q)|}{\det G(q)} = |\partial_{\alpha_0} p_m|^{-1}.$$

For $0 < \varepsilon \le 1/4$,

$$|D_q \log u[\eta]| \le D_* \varepsilon^{-1} \|\eta\|_\infty. \tag{34.2}$$

Consequently $u(q) \le e^{D_*\|q-q_x\|_\infty/\varepsilon} u(q_x)$ whenever the connecting segment is in that domain.

Proof. The true optical formula is $|e| = R^2 c_0 c_m/\mathfrak{b}_m$. Positivity and separate multiaffinity of $\mathfrak{b}_m$, proved in Lemma 30.1, give

$$|D \log |e|| \le \sum_{i=0}^{m} \frac{|Dc_i|}{c_i} + \sum_{i=0}^{m-1} \frac{|D\tau_i|}{\tau_i}.$$

Each incidence occurs once in this bound, including the two endpoints; there is no new product of inverse incidences. With $s_i = \sigma^i + \sigma^{m-i} \le 2\sigma^{d_i}$ and at most two indices at every depth,

$$\sum_{i=0}^{m} s_i r^{-d_i} \le \frac{4}{1-\beta}, \qquad \sum_{i=0}^{m-1} (s_i + s_{i+1}) \le \frac{4}{1-\sigma}.$$

Proposition 29.2 therefore bounds the incidence sum by $8C_c/[\varepsilon(1-\beta)]$ and the flight sum by $4aJ_*/(1-\sigma)$, times $\|\eta\|_\infty$. The physical Schur complement has $G \succeq gI$. By Theorem 29.3, $\|DG[\eta]\|_{\mathrm{op}} \le 2C_F\|\eta\|_\infty$, so

$$|D \log \det G[\eta]| = |\operatorname{tr}(G^{-1} DG[\eta])| \le 4C_F g^{-1} \|\eta\|_\infty.$$

Subtract these logarithmic derivatives in $\log u$ and use $\varepsilon \le 1$ to obtain (34.2). Integration on the actual physical segment proves the last assertion. The identity for u is the unchanged generating-map Jacobian of Theorem 28.2; no radius or section factor is absorbed in it. □

For $0 < \varepsilon \le 1/4$ let $\mathscr{Z}_{\varepsilon,m,R}$ be all distinct regular physical critical roots $p_0 = p_m = 0$ satisfying (33.4). Lattice translation copies and angular seams are identified, so each root is a unique physical initial point. These are actual reversing critical orbits by the retained theorem, not critical points of an unrelated surrogate action. Their roots are isolated on the initial momentum fibre since $\partial_{\alpha_0} p_m \neq 0$; hence the family is countable. Put

$$s_\varepsilon = \tfrac{1}{2}\delta_*\varepsilon, \qquad \rho_\varepsilon = \min\{1/4,\, g s_\varepsilon/(2R_+)\}, \qquad K_{1/2} = \frac{100}{47}\sqrt{1 - 1/4}. \tag{34.3}$$

Both positive radii are fixed before taking any collision limsup.

Lemma 34.2 (A common physical momentum disk at every guarded critical root)**.** *For every $x \in \mathscr{Z}_{\varepsilon,m,R}$ there is an actual regular inverse $q_x(\zeta)$ for $|\zeta| < \rho_\varepsilon$, where $\zeta = (p_0, p_m)$ and $q_x(0)$ is the original endpoint pair. It is contained in the physical endpoint ball $|q - q_x(0)|_2 < s_\varepsilon$ and obeys*

$$|q_x(\zeta) - q_x(0)|_2 \le R_+|\zeta|/g, \qquad u_x(\zeta) \le C_{\mathrm{dist}} u_x(0).$$

This is a common domain for all collision words in the stated class, not a separate radius chosen after m is fixed.

Proof. The Euclidean ball of radius s_ε lies strictly inside the ℓ^1 endpoint tube of Theorem 33.3. At a critical center both endpoint tangential components vanish. The signed tangential estimate in that proof and $C_{\mathrm{inc}}\delta_* \le 1/2$ imply $|p_0|, |p_m| \le \varepsilon/2$ on the entire ball; in particular $c_0, c_m \ge 1/2$ and $G \succeq gI$ there. The domain and every segment used next are genuine physical domains, established before using the Hessian.

Given $|\zeta| < \rho_\varepsilon$, minimize $F(q) - R(D\zeta)\cdot(q - q_x(0))$ on the closed endpoint ball, where $D = \mathrm{diag}(-1, 1)$. On its boundary the radial first variation is at least $g s_\varepsilon^2 - R_+|\zeta| s_\varepsilon > 0$. A boundary minimizer is impossible, because moving radially inward would strictly decrease the function. Strong convexity gives one interior minimizer, satisfying $\nabla F = RD\zeta$, and hence the correct endpoint momenta. The ordinary inverse theorem gives its smooth dependence, while strong convexity gives $g|q - q_x(0)|_2 \le R_+|\zeta|$. The optical mixed entry is nonzero, so this is also a regular local parametrization of the original collision source. Lemma 34.1, integrated on the physical radial segment in endpoint space, gives the stated relative bound. In it $\|q - q_x(0)\|_\infty \le s_\varepsilon$ and $s_\varepsilon/\varepsilon = \delta_*/2$. □

Theorem 34.3 (One-copy critical-tube sum at a common physical roof)**.** *For fixed R, m, ε take the union of the actual momentum charts in Lemma 34.2. Give this union its original collision measure, counting each physical initial point once. Retain any Borel selector $0 \le w \le 1$, in particular the full $\Pi_{n,k,m,R}$, both section tests, the original guard and any original physical-roof interval. Let $Q^w_{\varepsilon,m,R}(t)$ be the density of its pushforward by the original total flight time. Then*

$$\|Q^w_{\varepsilon,m,R}\|_{L^\infty(dt)} \le \frac{U C_{\mathrm{dist}} C_{\mathrm{mom}}}{2c_*}. \tag{34.4}$$

This bound is uniform in $\varepsilon \in (0, 1/4]$, all $m \ge 1$, all original radii and exact labels. It makes the complete positive chart sum before taking the physical-roof essential supremum. It is unscaled: it does not assert an m^{-2} gain.

Proof. Enumerate the distinct critical roots. Each chart is a Borel image of its open momentum disk; the endpoint-to-source map is locally invertible since $e \neq 0$. It is injective on the convex endpoint ball: at fixed α_0, $\partial_{\alpha_m} p_0 = -e/R$ has a fixed nonzero sign along the connected vertical slice. Thus a physical source point is represented at most once in a given chart. If charts overlap, assign the physical point to the first chart in the enumeration. These Borel assignments partition the unchanged union; they remove only coordinate multiplicity, never an actual collision or a

point of the union. Multiply w by the assignment indicator in each chart, denoting the result by w_x.

Let $a_x = L(x)$ and write $\zeta = re_\theta$. The exact radial coarea in Theorem 28.2 gives, for $r > 0$,

$$\partial_r L_x(re_\theta) = rh_x(re_\theta), \qquad U^{-1} \le h_x \le K_{1/2}^{-1}, \qquad b_x(re_\theta) = \frac{u_x(re_\theta)}{4\pi c_* h_x(re_\theta)}.$$

These formulas hold on the full common disk by the preceding lemma. For fixed t, θ, strict radial monotonicity gives at most one $r_x(t,\theta) \in (0, \rho_\varepsilon)$ per center. These radii are generally different for different centers; we do not identify them. The contribution is zero when no such radius exists. Otherwise

$$w_x b_x(r_x(t,\theta)e_\theta) \le \frac{UC_{\mathrm{dist}}}{4\pi c_*}\, u_x(0).$$

The retained complete regular-root theorem at the single common momentum pair $(0,0)$ gives $\sum_{x\in\mathscr{Z}_{\varepsilon,m,R}} u_x(0) \le C_{\mathrm{mom}}$. Here the transfer from different momentum points to $(0,0)$ is paid by the preceding distortion estimate, not inferred from the synchronized kernel alone.

Apply the coarea identity first to bounded nonnegative roof tests and finitely many charts, retaining the Borel selectors in the integrands. The upper bound just obtained is summable independently of t and θ. Monotone convergence gives the full nonnegative source sum. Integrating in $\theta \in [0, 2\pi)$ gives precisely (34.4); the polar origin has zero collision area in each regular chart and creates no atom. This area statement does not set a separate one-sided pairing trace to zero. At no step is a sum of individual roof suprema bounded by a synchronized value without a comparison factor. □

Corollary 34.4 (Uniform band coverage and a localized critical-roof envelope)**.** *Every chart above reaches every physical roof*

$$0 < t - a_x < \rho_\varepsilon^2/(2U)$$

on every momentum ray. Moreover the density in the theorem satisfies for almost every t the sharper bound

$$Q^w_{\varepsilon,m,R}(t) \le \frac{UC_{\mathrm{dist}}}{2c_*} \sum_{\substack{x\in\mathscr{Z}_{\varepsilon,m,R} \\ t-\rho_\varepsilon^2/(2K_{1/2})\le a_x\le t}} u_x(0). \tag{34.5}$$

The endpoints a_x are the actual physical critical roofs, not clock corrections. No section visit count is transferred from a tube point to its center unless section stability is proved separately.

Proof. Integrating $r/U \le \partial_r L_x \le r/K_{1/2}$ from zero gives $r^2/(2U) \le L_x(re_\theta) - a_x \le r^2/(2K_{1/2})$. Continuity and strict monotonicity prove coverage; the upper inequality restricts the contributing centers at any fixed physical roof. Insert that restriction into the nonnegative sum in the theorem. The original Borel exact-record selector is still evaluated on each actual orbit; the envelope simply dominates it by one. □

34.1. **The remaining normalized source is not eliminated.** The new analytic domain, physical guard tube and common-roof summation remove two specific gaps of the earlier critical argument: no collision-dependent continuation radius is chosen, and different branch momentum radii are now compared by a proved multiplicative optical bound. They neither assert that all critical centers satisfy a common guard margin nor estimate every original singular, noncritical or paired-boundary source. The latter measures need not be restrictions of the bulk collision measure used here.

Even on the covered union, (34.4) is an $O(1)$ roof-height bound, not an $O(m^{-2})$ bound. It cannot be multiplied by m^2 and declared negligible. The envelope (34.5) isolates genuine physical roof locations but provides no unproved critical-roof nonconcentration rate or transfer of return

labels across a section cut. The missing original limit remains exactly (15.8), not a renamed substitute. To close it requires a normalized complete physical-source estimate or sufficient original nongrazing capacity, including the uncovered incidence-only low-load source, unvisited and visited boundary terms, and the retained spectral, arithmetic and conditioned-denominator interfaces. Its original widths stay fixed before $m \to \infty$ and $B \to \infty$ remains outermost. No limit has been interchanged in the new finite-width, all-collision theorem.

35. A buffered horizon and bending on every physical record

The physical horizon established in Section 2 contains a strict covering margin which has not yet been used. This margin gives a deterministic lower bound for accumulated reflection, not just an upper bound for a single flight. All centers in this section belong to the original triangular lattice; all disks still have radius $R \in [R_-, R_+] = [9/20, 47/100]$. Incidence means $c_i = \sqrt{1-p_i^2}$ at an actual collision. In particular this section imposes no new cutoff on the exact event (1.1).

Lemma 35.1 (The unused normal margin in the lattice covering)**.** *Set*

$$H = \frac{28}{5}, \qquad \omega_0 = \frac{179}{400}, \qquad b_0 = R_- - \omega_0 = \frac{1}{400}, \qquad \gamma_0 = \frac{b_0}{2H} = \frac{1}{4480}. \tag{35.1}$$

For every $q \in \mathbb{R}^2$ and every unit vector v, there is a center $z \in \Lambda$ whose perpendicular projection onto $\{q + tv : 0 \le t \le H\}$ is at a parameter $s \in (0, H)$ and satisfies $|z - (q + sv)| < \omega_0$. The assertion is simultaneous in the starting point and direction, including limiting directions.

Proof. Apply Lemma 2.1 at the midpoint $q + Hv/2$. The four direction intervals and primitive row vectors in the proof of Theorem 2.2 cover every direction by lattice symmetry. Their squared normal widths are bounded, respectively, by

$$\frac{961}{4804}, \quad \frac{13697401}{73866304}, \quad \frac{529}{3136}, \quad \frac{75}{448}.$$

Each is at most the first number, and

$$\frac{961}{4804} < \frac{32041}{160000} = \left(\frac{179}{400}\right)^2.$$

These are comparisons of positive rational numbers. The absolute longitudinal displacement from the midpoint is at most $\frac{1}{2}\sqrt{3847/124} < 14/5 = H/2$, as proved there. Thus its foot lies strictly inside the stated segment. No simulated orbit or unstated compactness constant is involved. The number ω_0 is a covering distance, not a changed scatterer radius. □

Theorem 35.2 (Every physical time block carries positive bending)**.** *Let $\gamma : [0, H] \to \mathbb{R}^2$ be any unit-speed trajectory segment of the original Lorentz gas, with the actual specular reflections at times $0 < t_i < H$. Then*

$$\sum_{0<t_i<H} c_i \ge \gamma_0. \tag{35.2}$$

One may start or end inside a flight. For a collision at the initial time, the initial direction is the outgoing one. The same statement holds for a fixed finite-word grazing limit which is still a nonpenetrating specular polygon; a tangential contact has $c_i = 0$. The bound is deterministic and uniform in R, the word, the starting point and the initial direction.

Proof. There are finitely many collisions in a bounded physical time interval: consecutive flights have length at least $\tau_- = 3/50$, with only the first and last partial flights possibly shorter. Write v_0 for the initial outgoing velocity and $\Delta v_i = v(t_i+) - v(t_i-)$. Specular reflection at a circular disk with outward unit normal n_i gives

$$\Delta v_i = -2\big(v(t_i-) \cdot n_i\big) n_i, \qquad |\Delta v_i| = 2c_i.$$

Integrating the piecewise constant velocity, for $0 \le t \le H$ we have the exact identity and bound

$$\gamma(t) - \gamma(0) - tv_0 = \sum_{0<t_i<t} (t - t_i)\Delta v_i, \qquad |\gamma(t) - \gamma(0) - tv_0| \le 2H \sum_{0<t_i<H} c_i. \tag{35.3}$$

Choose z and s from Lemma 35.1. If the sum in (35.2) were less than γ_0, then

$$|\gamma(s) - z| < \omega_0 + 2H\gamma_0 = R_- \le R.$$

This places a true physical point inside an open scatterer, a contradiction. The argument uses neither disjoint reflection events nor statistical independence. It also applies directly to the stated limiting polygons, for which the same velocity identity holds and which remain outside every open disk. It does not define a billiard continuation through an arbitrary singular endpoint. □

Corollary 35.3 (A uniform block bound over all collision words)**.** *For any actual regular record with $m \ge 1$ flights, put $N_0 = 94$. Then*

$$\sum_{i=1}^{m-1} c_i \ge \gamma_0 \left\lfloor \frac{m}{N_0} \right\rfloor, \tag{35.4}$$

$$\# \left\{ 1 \le i < m : c_i \ge \frac{1}{416640} \right\} \ge \left\lfloor \frac{m}{N_0} \right\rfloor. \tag{35.5}$$

Every full block of 94 *consecutive flights supplies its own collision in the second inequality; these supplied collisions are distinct. All other collisions remain in the record and in every observable.*

Proof. The duration of a 94-flight block is at least $94\tau_- = 141/25 > 28/5 = H$. Starting at its first collision with the outgoing velocity, Theorem 35.2 supplies a sum at least γ_0 among its 93 interior collision indices. The sets of such interior indices for disjoint successive blocks are disjoint. Summation proves (35.4). Within each block at least one of the 93 nonnegative summands is at least $\gamma_0/93 = 1/416640$, proving (35.5). The possibly incomplete last block is not assigned any fictitious contribution. For $m < 94$ both statements retain their stated meaning, with a zero right side. □

Remark 35.4 (What positive bending does and does not measure)*.* The result concerns visited incidences along every original physical record. It is not abundance of the original comparison pool: $\mathsf{U}^{\mathrm{low}}_{B,m}$ and $P^{\mathrm{ng}}_{r_B,m}$ carry additional roof, clearance, pairing and exact-label requirements. A positive fraction of moderately transverse collisions does not pay a single unvisited tangency or force any endpoint trace to vanish. We use the bound below only in an exact variation of actual critical physical lengths; we do not multiply an unscaled mass bound by m^2 and call it small.

36. Normal-boundary continuation and transversality of the physical clock

A regular critical record is the actual m-flight record of Section 17 with $p_0 = p_m = 0$. It is one half of a true reversing periodic trajectory. We vary the common physical disk radius within $[R_-, R_+]$, keeping the ordered lattice centers $O_0, \dots, O_m$, all m collisions and the unit-speed clock. A formal reflection continuation will always be distinguished from a physical continuation. The latter is required to avoid every third disk and preserve the actual first-hit order.

36.1. **The full normal-boundary reflection equations.** Choose local real lifts of the contact angles α_i and of the flight directions ϑ_i. Their normal-reflection equations are, with the constants fixed at the reference record,

$$E_0 = \alpha_0 - \vartheta_0, \quad E_i = 2\alpha_i - \vartheta_{i-1} - \vartheta_i \ (1 \le i < m), \quad E_m = \alpha_m - \vartheta_{m-1}. \tag{36.1}$$

At a physical critical record E_0 is zero modulo 2π and E_m is π modulo 2π. The interior constants are the appropriate fixed odd multiples of π. These lifts remove any branch-cut ambiguity in the following derivatives.

Theorem 36.1 (A full reflection inverse without a grazing divisor)**.** *Let $\mathcal{B} = D_\alpha E$ at a regular critical record with $m \geq 1$. Write $w_i = \tau_i^{-1}$ and set $w_{-1} = w_m = 0$. Its only nonzero entries are*

$$\begin{aligned} \mathcal{B}_{ii} &= b_i + Rc_i(w_{i-1} + w_i), & b_0 = b_m = 1, \\ \mathcal{B}_{i,i+1} &= Rw_i c_{i+1}, & b_i = 2 \quad (0 < i < m), \\ \mathcal{B}_{i+1,i} &= Rw_i c_i, & c_0 = c_m = 1. \end{aligned} \tag{36.2}$$

For $\lambda = 50/47$ and $C_{\mathrm{B}} = 194/3$,

$$\|\mathcal{B}^{-1}\|_{1\to 1} \leq 1, \qquad \sum_{i=0}^{m} \lambda^{|i-j|} |(\mathcal{B}^{-1})_{ij}| \leq 2, \qquad \|\mathcal{B}^{-1}\|_{\infty\to\infty} \leq C_{\mathrm{B}}. \tag{36.3}$$

The constants are independent of R, m and the word. The algebraic statement includes simultaneous zero values of interior c_i when the positive flight lengths are retained. This extension of the matrix statement is not a claim that all such arrays are physical.

Proof. For $X_i = O_{i+1} - O_i + R(n_{i+1} - n_i) = \tau_i v_i$, $d\vartheta_i = v_i^{\perp} \cdot dX_i/\tau_i$. At the departure contact $v_i \cdot n_i = c_i$ and at arrival $v_i \cdot n_{i+1} = -c_{i+1}$. Differentiating (36.1) therefore gives (36.2), including its two boundary rows. In particular the boundary diagonal is $1 + Rc_i w$, not the interior expression with a spurious second flight.

Conjugate by $S = \mathrm{diag}((-1)^i)$ to obtain $A = S\mathcal{B}S = D - C$, where D is the positive diagonal and $C \geq 0$ has zero diagonal. The excess of the diagonal over the sum of the off-diagonal absolute values in column i is exactly b_i. Consequently $\|CD^{-1}\|_1 < 1$ for this finite matrix, $A^{-1} = D^{-1} \sum_{r\geq 0} (CD^{-1})^r \geq 0$, and $\mathbf{1}^T A \geq \mathbf{1}^T$ implies $\mathbf{1}^T A^{-1} \leq \mathbf{1}^T$. This proves the first bound. No row-dominance assertion is being used.

Fix j and conjugate further by $W = \mathrm{diag}(\lambda^{|i-j|})$. Across each edge the weight ratio is at most λ. Since $a_* = R_+/\tau_- = 47/6$ and $a_*(\lambda - 1) = 1/2$, the column excess of WAW^{-1} is at least $1 - 1/2 = 1/2$ at either endpoint and at least $2 - 1 = 1$ at an interior index. The same positive-inverse argument gives its inverse ℓ^1 norm at most 2. As $W_{jj} = 1$, this is the middle assertion in (36.3). In particular $|(\mathcal{B}^{-1})_{ij}| \leq 2\lambda^{-|i-j|}$. Sum this bound over a row, retaining at most two entries at each positive distance:

$$\sum_j |(\mathcal{B}^{-1})_{ij}| \leq 2\left(1 + 2\sum_{d\geq 1} \lambda^{-d}\right) = 2\left(1 + \frac{2}{\lambda - 1}\right) = \frac{194}{3}.$$

All steps remain valid when some interior c_i are zero. The endpoint case $m = 1$ is the same two-by-two calculation, without an interior row. □

Theorem 36.2 (A common formal radius domain and physical radius response)**.** *Every actual regular critical record has a holomorphic continuation of* (36.1) *on a radius disk of size independent of its collision count and its smallest interior incidence. More precisely, put*

$$\begin{aligned} L_R &= 4/\tau_-, & K_R &= 48/\tau_- + 512/\tau_-^2, \\ s_R &= \min\{1/8, \tau_-/24\}, & r_R &= \min\{s_R, (8C_{\mathrm{B}}K_R)^{-1}\}, \\ \eta_R &= \frac{r_R}{8(1 + C_{\mathrm{B}}L_R)}. \end{aligned} \tag{36.4}$$

For $|\rho| < \eta_R$ the fixed local equations have a unique solution $\alpha(R + \rho)$ in $\|\alpha - \alpha(R)\|_\infty \leq r_R/2$, and

$$\|\alpha(R + \rho) - \alpha(R)\|_\infty \leq 2(1 + C_{\mathrm{B}}L_R)|\rho|. \tag{36.5}$$

This is a formal normal-reflection domain. On every connected interval on which its record is actually physical, regular, and $R \in [R_-, R_+]$, one has the sharper derivative bound

$$\max_i |\alpha_i'(R)| \le D_R, \qquad D_R = C_{\mathrm{B}} L_R = \frac{38800}{9}. \tag{36.6}$$

Physical continuation through an unvisited tangency or a grazing collision is not asserted. A section crossing is not excluded and its actual selector is reevaluated at the new radius.

Proof. First hold the angles fixed at the reference record. The formula $\partial_R \vartheta_i = v_i^{\perp} \cdot (n_{i+1} - n_i)/\tau_i$ gives $|\partial_R \vartheta_i| \le 2/\tau_-$. Hence $\|E_R\|_\infty \le L_R$. At every actual regular critical point, implicit differentiation and Theorem 36.1 give $\alpha' = -\mathcal{B}^{-1} E_R$ and (36.6).

Here are explicit bounds for a domain, rather than just pointwise implicit differentiation. Let z_i denote the complex lattice center O_i. For complex independent increments (ρ, u) use the two holomorphic flight coordinates

$$X_i^{\pm} = (z_{i+1}^{\pm} - z_i^{\pm}) + (R + \rho)\big(e^{\pm i(\alpha_{i+1} + u_{i+1})} - e^{\pm i(\alpha_i + u_i)}\big), \qquad z_i^{+} = z_i, \quad z_i^{-} = \overline{z}_i.$$

The minus coordinate is defined by this expression, not by conjugating complex variables. At the real reference point both moduli are $\tau_i \ge \tau_-$. In the norm $\|(\rho, u)\| = |\rho| + \|u\|_\infty$, for increments of norm at most s_R, each flight changes by at most $6 s_R \le \tau_-/4$. Indeed $|e^{iu} - 1| \le e^{s_R} |u|$, $e^{s_R} < 2$, and $R < 1/2$. Thus logarithms continuous from the reference flights are defined on convex nonzero disks. On this domain $|R + \rho| < 1$, and direct differentiation gives the conservative bounds

$$\|D X_i^{\pm}\| \le 8, \qquad \|D^2 X_i^{\pm}\| \le 12, \qquad |X_i^{\pm}| \ge \tau_-/2.$$

The first two inequalities follow by distributing the radius derivative and at most two adjacent angular derivatives; their operator norms use the displayed joint norm and do not sum over collision indices. For $\vartheta_i = (\log X_i^{+} - \log X_i^{-})/(2i)$ up to its fixed reference constant, the second derivative of the logarithm gives

$$\|D^2 \vartheta_i\| \le 24/\tau_- + 256/\tau_-^2, \qquad \|D^2 E\| \le K_R.$$

These are estimates for each component in the supremum norm, uniformly in the dimension.

Write $E(R + \rho, \alpha + u) - E(R, \alpha) = \mathcal{B} u + E_R \rho + N(\rho, u)$. Taylor's integral formula gives $\|N\|_\infty \le (K_R/2)(|\rho| + \|u\|_\infty)^2$ and $\|D_u N\| \le K_R(|\rho| + \|u\|_\infty)$. For $|\rho| \le \eta_R$ the map

$$u \longmapsto -\mathcal{B}^{-1}(E_R \rho + N(\rho, u))$$

preserves $\|u\|_\infty \le r_R/2$: the linear forcing is at most $r_R/8$ and the nonlinear term at most $r_R/16$. Its Lipschitz constant is at most $1/8$. Iteration from zero gives the unique fixed point there, holomorphic by locally uniform convergence of the holomorphic iterates. Comparing the fixed point with the first iterate gives

$$\|u\|_\infty \le \frac{C_{\mathrm{B}} L_R |\rho| + (C_{\mathrm{B}} K_R/2)|\rho|^2}{1 - 1/8} \le 2(1 + C_{\mathrm{B}} L_R)|\rho|.$$

This proves the asserted formal domain, also for the zero-incidence matrix limits with nonzero reference flights and the same reflected angle identities. For a regular physical reference, continuity and its finitely many positive flight/incidence/third-disk margins give some open physical subinterval; its size need not be uniform. The argument has not promoted the formal disk to a common physical radius disk. Nor has it frozen a discontinuous exact-record selector. □

36.2. **A collision-uniform derivative of the true critical roof.** Let J be a connected interval of regular physical critical continuation as in Theorem 36.2. Its actual roof is

$$a(R) = \sum_{i=0}^{m-1} |O_{i+1} - O_i + R(n_{i+1}(R) - n_i(R))|.$$

Every summand is a real physical flight, not a symbolic roof.

Theorem 36.3 (Exact radius variation and uniform critical-clock slope)**.** *On J one has*

$$a'(R) = -c_0 - c_m - 2\sum_{i=1}^{m-1} c_i = -2 - 2\sum_{i=1}^{m-1} c_i. \tag{36.7}$$

In particular, with $\kappa_0 = 1/210560$,

$$2 + 2\gamma_0 \left\lfloor \frac{m}{94} \right\rfloor \le -a'(R) \le 2m, \qquad \kappa_0 m \le -a'(R) \le 2m. \tag{36.8}$$

For any $R_1 < R_2$ in this same physical interval,

$$\kappa_0 m(R_2 - R_1) \le a(R_1) - a(R_2) \le 2m(R_2 - R_1). \tag{36.9}$$

There is also the uniform normalized second-derivative estimate

$$|a''(R)| \le 2(m-1)C_{cR} \le 2mC_{cR}, \qquad C_{cR} = D_R + \frac{2 + 2R_+D_R}{\tau_-}. \tag{36.10}$$

Consequently a/m has both a slope separated from zero and a Lipschitz derivative with constants independent of the collision word. The same one-sided first-variation identities and bounds hold at a fixed finite-word nonpenetrating grazing limit of such records. They do not identify different words across a singularity.

Proof. The full length action $S(R, \alpha)$ has zero angular gradient at a critical record: the interior equations are specular stationarity, and the two endpoint derivatives are $-Rp_0$ and Rp_m. Differentiating $S(R, \alpha(R))$ therefore cancels every angular chain-rule term and leaves

$$a'(R) = \partial_R S(R, \alpha) = \sum_{i=0}^{m-1} v_i \cdot (n_{i+1} - n_i).$$

Each summand is $-c_{i+1} - c_i$. The interior incidences occur exactly twice; the two normal endpoints occur once and equal one. This proves (36.7), including the signs and both boundary factors. Corollary 35.3 supplies the first lower bound in (36.8). Since $\lfloor m/94 \rfloor \ge m/94 - 1$ and $0 < \gamma_0 < 1$,

$$2 + 2\gamma_0 \lfloor m/94 \rfloor \ge (2\gamma_0/94)m, \qquad 2\gamma_0/94 = 1/210560.$$

The upper bound follows from $0 \le c_i \le 1$. Integration over the actual connected physical interval proves (36.9). In particular no exchange of a radius supremum and a time or collision limit is used.

For the second derivative, the full response bound gives $|n_i'| \le D_R$ and

$$|X_i'| \le 2 + 2R_+D_R, \qquad |v_i'| = |(I - v_i v_i^T)X_i'|/\tau_i \le (2 + 2R_+D_R)/\tau_-.$$

Differentiate the incoming formula $c_i = -v_{i-1} \cdot n_i$. It follows that $|c_i'| \le C_{cR}$, without dividing by c_i. The endpoints remain normal throughout J, so their derivatives are zero. Differentiating (36.7) yields (36.10). At a stated finite-word limit, the positive flight lengths, the uniform reflection inverse and the formal local continuation supply the one-sided derivatives. The finite sums of incidences and the nonpenetrating polygon bound pass to that limit. No assertion about an arbitrary branch beyond that endpoint follows. □

Corollary 36.4 (Branchwise band width and exact radius–roof coarea)**.** *On every such connected physical branch, for a physical roof interval I_t,*

$$|\{R \in J : a(R) \in I_t\}| \le |I_t|/(\kappa_0 m). \tag{36.11}$$

More generally let $\mathscr{K}_m$ be the full set of regular physical critical m-flight records with the initial disk represented in a fixed lattice fundamental domain and $R \in [R_-, R_+]$. Write x for a record at a given R. Give it any nonnegative Borel weight $f(R,x)$, which may include the original exact return/displacement/collision selector and an original optical weight. For every nonnegative Borel roof test ϕ,

$$\int_{R_-}^{R_+} \sum_{x:(R,x)\in\mathscr{K}_m} f(R,x)\phi(a_x(R))\,dR = \int_{\mathbb{R}} \phi(t) \sum_{\substack{(R,x)\in\mathscr{K}_m \\ a_x(R)=t}} \frac{f(R,x)}{|a_x'(R)|}\,dt. \tag{36.12}$$

The equality is an identity of positive measures, possibly infinite; no global integrability is silently assumed. In every term

$$\frac{1}{2m} \le \frac{1}{|a_x'(R)|} \le \frac{1}{\kappa_0 m}.$$

Physical records are counted once; coordinate chart overlaps are not extra records. The original probability normalization c_* is unchanged. For example, inserting a weight with factor $(4\pi c_*)^{-1}$ leaves that factor exactly once on both sides.

Proof. The derivative in (36.8) has one sign and is separated from zero, so each connected branch is a C^1 strictly monotone graph over R and a C^1 graph over its physical roof. The ordinary one-dimensional substitution formula proves (36.11) and (36.12) on one chart. To justify the full physical sum, there are countably many finite lattice words and each finite-dimensional angle space is second countable. The invertible full reflection matrix and the open physical regularity conditions give a countable cover by charts on open radius intervals. Assign a physical point to its first chart in a fixed enumeration. The assigned subsets are Borel, and their indicators can be included in f on the uncut charts. Substitution for finitely many nonnegative chart terms, then monotone convergence, gives the displayed equality. This allocation deletes representations, never collisions or physical source points. The exact selector is evaluated at (R,x), not transported from another radius or from a tube center. □

Remark 36.5 (Fixed-radius shortage is not a radius average)*.* The lower clock slope is a new all-word geometric input on the original radius family. Formula (36.12) is an auxiliary exact change of variables, not a replacement of the original probability problem by a radius-averaged limit theorem. A bound for crossings summed at different radii does not follow from the retained momentum kernel at one radius; conversely, a small radius integral cannot control $\sup_R$. Neither (36.11) nor the presence of moderately transverse collisions proves the original m^{-2} roof nonconcentration, the full nongrazing comparison capacity, or the shortage limit (15.8). Singular/paired endpoint traces, nonguarded critical centers and the complete noncritical positive source remain part of that limit. All original microscopic widths remain fixed before the collision limsup and B remains outermost. No theorem or proof body of the inherited raw-LLT and inverse program has been removed.

37. The complete record: endpoint work and radius–roof disintegration

The preceding clock calculation extends beyond reversing records, but its endpoint work must then be retained. This section treats every regular physical m-flight record, with all original exact labels, not just the critical tubes of Section 34. It produces a transverse coordinate for

a parameter-extended source. It does not change the fixed-radius probability space or the raw target.

Theorem 37.1 (Uniform continuation at prescribed endpoint momenta)**.** *Fix any actual regular record and its two endpoint momenta $(a, v) = (p_0, p_m) \in (-1, 1)^2$. Keep them fixed while varying the common radius. Replace the endpoint constants in* (36.1) *by*

$$E_0 = -\arcsin a \pmod{2\pi}, \qquad E_m = \pi + \arcsin v \pmod{2\pi}.$$

The full derivative matrix is still (36.2)*, now with $c_0 = \sqrt{1-a^2}$ and $c_m = \sqrt{1-v^2}$ in place of one. All three bounds* (36.3) *remain valid, as do the common formal radius disk* (36.4) *and the physical radius response* (36.6)*. None of their constants depends on (a, v), the word, or the smallest incidence. Physicality is asserted only on connected regular physical continuation intervals, not on the whole formal radius disk.*

Proof. The departure direction is $\alpha_0 + \arcsin a$. The incoming terminal direction is $\alpha_m + \pi - \arcsin v$; tangential momentum is unchanged by the last specular reflection. These give exactly the two displayed constants modulo 2π. Their derivatives with respect to R vanish because a, v are fixed. The flight differentiation in Theorem 36.1 then gives the same matrix with its true endpoint cosines. Its column excess remains exactly b_i, including the endpoint columns: each endpoint cosine appears both in its diagonal correction and in the sole off-diagonal entry of that column. The unweighted and weighted column arguments therefore apply without alteration, and the row bound follows by the same geometric sum. No derivative of $\arcsin a$ or $\arcsin v$ is taken. The complex flight estimates in Theorem 36.2 use only the positive flight lengths, the disk radius bound and this inverse, not normality. They supply the same formal continuation and the same bound on α'. The interpretation of its physical subinterval is unchanged. □

Theorem 37.2 (The physical roof has only bounded endpoint work)**.** *Let $L(R)$ be the true physical roof on any connected physical continuation interval from Theorem 37.1. Then*

$$L'(R) = -c_0 - c_m - 2\sum_{i=1}^{m-1} c_i + R\{-a\alpha_0'(R) + v\alpha_m'(R)\}. \tag{37.1}$$

Define the collision-independent constants

$$C_{\mathrm{end}} = 2R_+D_R, \qquad C_{\mathrm{loss}} = C_{\mathrm{end}} + 2\gamma_0, \qquad m_0 = \lceil 2C_{\mathrm{loss}}/\kappa_0 \rceil .$$

For every $m \geq 1$ and every such record,

$$\kappa_0 m - C_{\mathrm{loss}} \leq -L'(R) \leq 2m + C_{\mathrm{end}}. \tag{37.2}$$

In particular for all $m \geq m_0$, the actual roof is strictly monotone on each connected physical component, and $-L'(R) \geq \kappa_0 m/2$. These statements are uniform over the entire original radius interval, over the endpoint momenta and over all physical collision words. They impose no new incidence or clearance guard on the original source. The conservative constant m_0 is a theoretical threshold, not an experimentally reached collision count.

Proof. On the stationary interior of the full action, its endpoint gradient is $R(-a, v)$. Differentiating the action along the prescribed-momentum continuation consequently leaves the two endpoint chain-rule terms, in addition to the explicit radius derivative. The latter telescopes as in (36.7), proving (37.1). Its endpoint work has absolute value at most $R(|a| + |v|)D_R \leq C_{\mathrm{end}}$. The visited incidence sum is bounded below by $\gamma_0\lfloor m/94 \rfloor$ from Corollary 35.3. Dropping only the two nonnegative endpoint cosines in a lower bound, and using $\lfloor m/94 \rfloor \geq m/94 - 1$, gives

$$-L'(R) \geq 2\gamma_0\lfloor m/94 \rfloor - C_{\mathrm{end}} \geq \kappa_0 m - C_{\mathrm{loss}}.$$

There are exactly $2m$ incidence contributions with multiplicity, each at most one, which gives the upper bound. The definition of m_0 proves the last assertion. This is a pointwise geometric estimate for every physical record, not an ergodic average. In particular no independence, pressure input or collision-word count occurs in its proof. □

Theorem 37.3 (An exact disintegration of the parameter-extended physical source)**.** *Fix $m \geq m_0$ and a nonnegative Borel function $h(R,x)$ on the regular physical collision space, zero off the original initial section when section probability is used. It may include the exact original selector $\mathbf{1}_{\Pi_{n,k,m,R}}$, with the true section visits evaluated at that radius. For a regular root with endpoint momenta (a,v) let*

$$u(R,x) = |\partial_{\alpha_0} p_m|^{-1} \quad \text{with } R, p_0 = a \text{ fixed.}$$

Use the physical momentum branches and one-copy convention of Section 26. For every nonnegative Borel test ϕ,

$$\begin{aligned} &\int_{R_-}^{R_+} \int h(R,x)\phi(L_{m,R}(x))\, d\nu_R^*(x)\, dR \\ &= \frac{1}{4\pi c_*} \int_{-1}^{1}\int_{-1}^{1}\int_{\mathbb{R}} \phi(t) \sum_{\substack{(R,x):\ p_0=a,\ p_m=v \\ L_{m,R}(x)=t}} \frac{h(R,x)u(R,x)}{|L'_{a,v,x}(R)|}\, dt\, dv\, da. \end{aligned} \tag{37.3}$$

The derivative is the full derivative with the endpoint work in (37.1)*, not a derivative at fixed initial angle. The sums include only genuine regular physical records and count each once. No finiteness of either side is assumed. The new Jacobian factor satisfies*

$$\frac{1}{2m + C_{\mathrm{end}}} \leq \frac{1}{|L'_{a,v,x}(R)|} \leq \frac{2}{\kappa_0 m}. \tag{37.4}$$

The equality is auxiliary geometric information in the added radius coordinate. It makes no assertion that the radius integral can replace the required supremum over R.

Proof. On a regular physical branch the initial collision measure is $d\alpha_0\, da/(4\pi)$ and the map to (a,v) has positive absolute Jacobian $|\partial_{\alpha_0} p_m|$; this is the actual root Jacobian used in the physical momentum change of variables in Section 26. After the original section restriction its density is $u/(4\pi c_*)$. This is a local change of variables; the all-time upper kernel bound in that section is not used here. In particular it is not a composition of one-step momentum kernels.

Theorem 37.1 supplies local physical branches parametrized by (R,a,v). For $m \geq m_0$ the coordinate change $(R,a,v) \mapsto (L(R,a,v),a,v)$ is locally invertible, with absolute determinant $|L'_{a,v,x}(R)|$, by Theorem 37.2. The ordinary substitution theorem therefore gives (37.3) on one chart, with the reciprocal Jacobian in the stated direction. Countably many physical charts cover the regular source: use countably many lattice words and second countability in (R,α_0,a). A fixed first-chart allocation makes their representations disjoint. The allocation indicators and any section or exact-label cuts stay inside h; none is differentiated. Applying substitution to finite positive chart sums and then monotone convergence gives the equality for the complete regular source, allowing infinite values. The identity concerns the regular source and makes no claim about the separate paired-boundary traces. Finally (37.2) proves (37.4). □

Remark 37.4 (The precise unpaid step is unchanged). This exact formula retains all regular physical records, all original collision labels and their true physical roofs. It is not a new probability model and does not discard low-incidence records. Nevertheless its crossing sum is taken at different radii. The unscaled root-kernel bound at a common (R,a,v) does not control this sum. Nor does an estimate integrated in R control the original fixed-radius roof essential height. A uniform bound for those concentrations, the complete positive source and paired

traces, or the original comparison capacity is still required to prove (15.8). This remains a substantive mathematical obligation rather than an automatic consequence of the factor $1/m$ in a different-coordinate Jacobian. All original probability quantifiers and ordered microscopic limits are retained.

38. A global variational certificate for every physical word

This section supplies a comparison valid on the whole original physical source, without requiring a common clearance margin or a physical critical center. It extends, rather than reclaims as new, the Cartesian support argument of Lemma 10.1 and the fixed-endpoint minimizing property in Lemma 10.3. The new ingredients are a complete disk-word duality formula, an exact endpoint defect, and quantitative certificates with nonnegative terms.

Fix $m \geq 1$, $R \in [R_-, R_+]$, and an ordered lifted center word $w = (O_0, \dots, O_m)$, with $O_0 = 0$ and $O_{i+1} \neq O_i$. A word of an actual trajectory always satisfies this last condition. Set

$$K_w(R) = \prod_{i=0}^{m} \overline{B}(O_i, R), \qquad S(Y) = \sum_{i=0}^{m-1} |Y_{i+1} - Y_i|, \qquad \ell_w(R) = \min_{Y \in K_w(R)} S(Y).$$

The disks here are the original geometric disks, including their interiors only for this auxiliary minimization. An arbitrary minimizer is *not* called a Lorentz trajectory: it may have an interior vertex, a tangency, or a segment intersecting an unlisted disk. Neither ℓ_w nor the minimization defines a replacement dynamics, a roof, a probability law, or a new conditioning event. The actual clock throughout remains $L = S(Q)$ on the original first-hit trajectory Q. Successive disks are separated by at least $1 - 2R_+ = 3/50$, so S is continuously differentiable on a neighborhood of $K_w(R)$.

For vectors $V = (v_0, \dots, v_{m-1})$, $|v_i| \leq 1$, put

$$\begin{aligned} &d_0 = -v_0, \quad d_i = v_{i-1} - v_i \ (0 < i < m), \quad d_m = v_{m-1}, \\ &C_w(V) = \sum_{i=0}^{m-1} v_i \cdot (O_{i+1} - O_i), \qquad T(V) = \sum_{i=0}^{m} |d_i|, \qquad D_{w,R}(V) = C_w(V) - RT(V). \end{aligned} \tag{38.1}$$

All lengths in these expressions have the original physical units.

Theorem 38.1 (Exact disk-word duality and a nonnegative gap)**.** *For every word and radius just specified,*

$$\ell_w(R) = \max_{|v_i| \leq 1} D_{w,R}(V). \tag{38.2}$$

Both extrema are attained. For every $Y \in K_w(R)$ and every feasible V,

$$\begin{aligned} S(Y) - D_{w,R}(V) = &\sum_{i=0}^{m-1} \big(|Y_{i+1} - Y_i| - v_i \cdot (Y_{i+1} - Y_i)\big) \\ &+ \sum_{i=0}^{m} \big(d_i \cdot (Y_i - O_i) + R|d_i|\big). \end{aligned} \tag{38.3}$$

Every summand is nonnegative. At any minimizing Y, the unit edge vectors $v_i = (Y_{i+1} - Y_i)/|Y_{i+1} - Y_i|$ attain the dual maximum. In particular an optimal dual vector can always be chosen with $|v_0| = |v_{m-1}| = 1$ and $2 \leq T(V) \leq 2m$.

Proof. Expand $\sum_i v_i \cdot (Y_{i+1} - Y_i) = \sum_i d_i \cdot Y_i$ and $C_w(V) = \sum_i d_i \cdot O_i$. Subtraction proves (38.3). The first sum is nonnegative by $|v_i| \leq 1$; the second is nonnegative by $|Y_i - O_i| \leq R$. This proves weak duality without a minimax interchange.

Compactness gives a primal minimizer Y. Convexity and differentiability of S show that its first-order inequality on the convex product $K_w(R)$ is necessary and sufficient for minimality.

With the displayed unit edge vectors its ith gradient is exactly d_i. Varying one vertex at a time therefore shows that Y_i minimizes $d_i \cdot Z$ on $\overline{B}(O_i, R)$. If $d_i \neq 0$, this minimizer is $O_i - Rd_i/|d_i|$; if $d_i = 0$, the corresponding gap is identically zero. Thus every term on the right of (38.3) vanishes at this Y, V. This proves equality, dual attainment and the asserted choice of dual vectors. At the two endpoints $|d_0| = |d_m| = 1$; each of the $m-1$ interior differences has norm at most two, giving $2 \leq T(V) \leq 2m$. The case $m = 1$ uses just the two endpoint terms. Nonconsecutive repetitions of a disk cause no difficulty: the vertices are separate variables in the stated deterministic comparison. □

Theorem 38.2 (The exact endpoint deficit of an arbitrary true record)**.** *Let $Q_i = O_i + Rn_i$ be* any *regular physical m-flight record, with its actual unit velocities v_i, endpoint momenta p_0, p_m, and incidences $c_i > 0$. All original section and exact-label tests may be retained. Then*

$$L - D_{w,R}(V) = E_R(Q) := R(2 - c_0 - c_m), \tag{38.4}$$

whence, uniformly in m, R, w and all intermediate incidences and clearances,

$$0 \leq L - \ell_w(R) \leq E_R(Q) = R\left(\frac{p_0^2}{1 + c_0} + \frac{p_m^2}{1 + c_m}\right) \leq 2R. \tag{38.5}$$

Consequently the range of the true roof on a physical word is at most $2R$, improving the still valid $4R$ enclosure of Lemma 10.3. No collision is deleted in this bound.

Proof. For actual velocities the flight terms of (38.3) vanish. The true reflection law gives $d_i = -2c_i n_i$ for $0 < i < m$, so every interior contact term also vanishes. At departure, $d_0 = -v_0$ and $v_0 \cdot n_0 = c_0$; at arrival, $d_m = v_{m-1}$ and $v_{m-1} \cdot n_m = -c_m$. The two remaining contact terms are exactly $R(1 - c_0)$ and $R(1 - c_m)$. This proves (38.4). Weak duality and feasibility of Q sandwich ℓ_w between $D_{w,R}(V)$ and L. The identity $1 - c = p^2/(1 + c)$ proves the second expression. Every true roof of the word lies in the same interval $[\ell_w, \ell_w + 2R]$. This proof uses only real flight vectors and the exact specular law; it introduces no independence, pressure hypothesis or enlarged observational window. The interval is a deterministic enclosure, not the event on which a local limit theorem is asserted. □

Corollary 38.3 (A globally certified regular critical record)**.** *If $p_0 = p_m = 0$, then $L = \ell_w(R)$ and Q is the unique minimizer of S on $K_w(R)$. For any competitor $Y \in K_w(R)$,*

$$S(Y) - L \geq \frac{1}{2R}\sum_{i=0}^{m} \lambda_i |Y_i - Q_i|^2 + \frac{1}{2}\sum_{i=0}^{m-1} |Y_{i+1} - Y_i||\widehat{v}_i - v_i|^2, \quad \lambda_0 = \lambda_m = 1, \quad \lambda_i = 2c_i, \tag{38.6}$$

where $\widehat{v}_i$ is the unit vector of the competitor edge. This global minimizing certificate is new here; the uniqueness among regular physical critical records was already proved in Section 10 and is not counted again as a new result.

Proof. Equation (38.5) gives equality and all $d_i = -\lambda_i n_i$, with $\lambda_i > 0$. For $z = Y_i - O_i$ in the radius-R disk,

$$R - n_i \cdot z = \frac{|z - Rn_i|^2 + R^2 - |z|^2}{2R} \geq \frac{|Y_i - Q_i|^2}{2R}.$$

For the flight term, unit length of v_i gives $|e| - v_i \cdot e = |e|\,|e/|e| - v_i|^2/2$. Insert these expressions into (38.3). Equality of lengths forces every $Y_i = Q_i$, proving uniqueness even against nonphysical competitors. This does not assert uniqueness of an arbitrary relaxed minimizer at a word with inactive contacts. □

39. Global radius order and a positive physical shape energy

The radius–roof identities of Sections 36 and 37 are inherited. Here a global variational value allows critical comparisons between disconnected physical continuation intervals, and its second variation has a nonnegative, geometrically explicit form. Nothing is continued as a true trajectory through a forbidden collision or an unvisited disk.

Theorem 39.1 (Global radius convexity of the disk-word value)**.** *For each fixed word of Section 38, $R \mapsto \ell_w(R)$ is convex and strictly decreasing on $[R_-, R_+]$. For $R_1 < R_2$,*

$$2(R_2 - R_1) \le \ell_w(R_1) - \ell_w(R_2) \le 2m(R_2 - R_1). \tag{39.1}$$

This statement concerns the geometric value even where its minimizer is not physical. If regular physical critical records of the same word exist at both R_1, R_2, with true roofs a_1, a_2, then the stronger bound

$$\kappa_0 m(R_2 - R_1) \le a_1 - a_2 \le 2m(R_2 - R_1), \qquad \kappa_0 = 1/210560, \tag{39.2}$$

holds without requiring a physical continuation on (R_1, R_2). Thus a fixed word has at most one regular physical critical record with a prescribed physical roof over the whole *radius interval.*

Proof. The dual formula (38.2) is a maximum of affine functions $C_w(V) - RT(V)$ over the same compact vector set for all R. It proves convexity. At each R_j, choose the optimal unit-edge vector V_j furnished by the primal minimizer. Its affine function supports ℓ_w from below and $2 \le T(V_j) \le 2m$. Therefore

$$T(V_2)(R_2 - R_1) \le \ell_w(R_1) - \ell_w(R_2) \le T(V_1)(R_2 - R_1).$$

This proves (39.1), without differentiability of ℓ_w or a common minimizing configuration.

At a regular physical critical point, Corollary 38.3 identifies its true roof with ℓ_w. Its actual velocity vector is an optimal dual vector and $T(V) = 2 + 2\sum_{i=1}^{m-1} c_i \ge \kappa_0 m$ by the already proved complete-bending bound (36.8). Use that vector at R_2 in the lower supporting inequality and the vector at R_1 in the upper inequality. This proves (39.2) across any nonphysical holes. At a fixed radius the critical point is unique by the inherited physical uniqueness theorem; distinct radii have strictly ordered roofs by the new global comparison. No physical orbit in a hole has been asserted or counted. □

Theorem 39.2 (Transfer of complete physical bending to every minimizing dual)**.** *Let Q be any regular physical record of w at R, without normal endpoint or clearance assumptions. Let Y be any minimizer of S on $K_w(R)$, let V and $\widehat{V}$ be their respective unit flight vectors, and write $E = R(2 - c_0 - c_m)$. Then*

$$\frac{1}{2}\sum_{i=0}^{m-1} |Y_{i+1} - Y_i|\,|\widehat{v}_i - v_i|^2 \le E, \qquad T(\widehat{V}) \ge 2 + 2\sum_{i=1}^{m-1} c_i - 2\sqrt{2mE/\tau_-}. \tag{39.3}$$

In particular, with $C_{\mathrm{bend}} = 4\sqrt{R_+/\tau_-}$, every such minimizing dual satisfies

$$T(\widehat{V}) \ge b_m := \kappa_0 m - C_{\mathrm{bend}}\sqrt{m}, \qquad \kappa_0 = 1/210560. \tag{39.4}$$

The comparison holds even when Y is nonphysical and contains inactive contacts. No inverse incidence occurs.

If $R_1 < R_2$ both admit arbitrary regular physical records of this same word, with actual roofs L_1, L_2, then

$$b_m(R_2 - R_1) - 2R_+ \le L_1 - L_2 \le 2m(R_2 - R_1) + 2R_+. \tag{39.5}$$

For $b_m > 0$, all radii admitting an actual record of the word with its true roof in any specified interval I of length h therefore have diameter at most $(h + 2R_+)/b_m$, even when that set is

disconnected. For $m \ge \lceil 64R_+/(\tau_-\kappa_0^2)\rceil$, $b_m \ge \kappa_0 m/2$. This is a deterministic support bound across the original radius family, not a radius-averaged substitute for the fixed-radius raw theorem.

Proof. Use the actual velocity vector V of Q in the nonnegative gap (38.3) at the minimizing Y. Its total gap is $\ell_w - D_{w,R}(V) \le L - D_{w,R}(V) = E$. Its flight terms are exactly those in the first part of (39.3). Every comparison edge has length at least τ_-, so $\sum_i |\widehat{v}_i - v_i|^2 \le 2E/\tau_-$. The two endpoint contributions to T are one for both unit-vector families. The reverse triangle inequality for each interior difference and Cauchy–Schwarz give

$$|T(\widehat{V}) - T(V)| \le 2\sum_i |\widehat{v}_i - v_i| \le 2\sqrt{2mE/\tau_-}.$$

For the actual physical velocities $T(V) = 2 + 2\sum c_i$. The complete-block incidence estimate in Section 35 gives $2 + 2\sum c_i \ge \kappa_0 m$ for every such record: this uses only the same full-block counting as (36.8), not its critical endpoint hypothesis. Since $E \le 2R_+$, the stated C_{bend} follows. This is an energy comparison with the actual trajectory, not a claim that a minimizing polygon obeys the bending theorem physically.

Apply the supporting affine vector of a minimizer at R_2 to $\ell_w(R_1)$. The preceding lower bound on its T gives $\ell_w(R_1) - \ell_w(R_2) \ge b_m(R_2 - R_1)$. The global upper bound remains $2m(R_2 - R_1)$. Writing $L_j = \ell_w(R_j) + e_j$ with $0 \le e_j \le 2R_+$ proves (39.5). If both actual roofs lie in I, $L_1 - L_2 \le h$, which yields the diameter bound for every pair of admitted radii and thus for its entire, possibly disconnected, set. The final integer threshold is the exact rearrangement of $C_{\mathrm{bend}}\sqrt{m} \le \kappa_0 m/2$. It is not claimed to be sharp or to have been reached by the finite simulations. □

Theorem 39.3 (Exact nonnegative curvature of the true critical clock)**.** *On a connected interval of regular physical critical continuation of one fixed word, write $Q_i(R) = O_i + Rn_i(R)$, $X_i = Q_{i+1} - Q_i = \tau_i v_i$, and let a prime denote its genuine radius derivative. Then*

$$\begin{aligned} a''(R) &= \sum_{i=0}^{m-1} \frac{|(I - v_i v_i^{\mathsf{T}})(Q'_{i+1} - Q'_i)|^2}{\tau_i} + R\sum_{i=0}^{m} \lambda_i |n'_i|^2 \\ &= \sum_{i=0}^{m-1} \tau_i |v'_i|^2 + R\left(|n'_0|^2 + |n'_m|^2 + 2\sum_{i=1}^{m-1} c_i |n'_i|^2\right) \ge 0. \end{aligned} \tag{39.6}$$

Every incidence and flight in this formula is physical. In particular

$$\frac{d}{dR}\sum_{i=1}^{m-1} c_i = -\frac{1}{2}a''(R) \le 0. \tag{39.7}$$

At a regular physical point equality $a''(R) = 0$ occurs precisely for a normal two-disk bouncing record (possibly stopped after any positive number of flights). Otherwise $a''(R) > 0$. No positive lower curvature constant uniform over all words is asserted.

Proof. The full Cartesian gradient of S at a critical record is $\nabla_i S = -\lambda_i n_i$, with the coefficients of Corollary 38.3. Twice differentiating the sum of actual lengths gives

$$a'' = \sum_i \frac{|(I - v_i v_i^{\mathsf{T}})(Q'_{i+1} - Q'_i)|^2}{\tau_i} + \sum_i \nabla_i S \cdot Q''_i.$$

Now $Q''_i = 2n'_i + Rn''_i$, $n_i \cdot n'_i = 0$, and $n_i \cdot n''_i = -|n'_i|^2$. Substitution yields exactly (39.6), including its positive sign and both endpoint factors. Since $v'_i = (I - v_i v_i^{\mathsf{T}})(Q'_{i+1} - Q'_i)/\tau_i$, the second display is identical to the first. The common formal reflection construction supplies the needed smooth derivatives on each physical interval, but no pressure or probability argument is used. Differentiating the retained identity $a' = -2 - 2\sum_{i=1}^{m-1} c_i$ proves (39.7).

If $a'' = 0$, every coefficient $R\lambda_i$ is positive, so all $n_i' = 0$, and all flight projections in (39.6) vanish. Consequently $n_{i+1} - n_i$ is parallel to v_i. At the initial normal contact $v_0 = n_0$. The next normal must therefore be parallel to v_0, and its negative incoming incidence forces $n_1 = -v_0$. The first reflection is normal and reverses the flight. Repeating the argument gives normal incidence and reversal at every contact. Since these are true first-hit flights, reversal returns to the very disk just departed, not an invented farther center. Hence the word is a repeated two-disk bounce. Conversely that record has fixed contact normals and flight directions while R varies, and $a(R) = m(|O_1 - O_0| - 2R)$; its curvature is zero. This also covers $m = 1$. □

Corollary 39.4 (A word-indexed critical radius–roof coarea)**.** *In* (36.12)*, normalize $O_0 = 0$ as before and write $\mathscr{W}_m$ for the countable set of all length-m lattice words. For each t, w, either no regular physical critical record of w has roof t, or there is exactly one pair $(R_w(t), x_w(t))$. For every nonnegative Borel weight f and test ϕ,*

$$\int_{R_-}^{R_+} \sum_{x \in \mathscr{K}_m(R)} f(R,x)\phi(L(x))\, dR = \int_{\mathbb{R}} \phi(t) \sum_{w \in \mathscr{W}_m} \mathbf{1}_{\{(R_w, x_w) \text{ exists}\}} \frac{f(R_w(t), x_w(t))}{|a_w'(R_w(t))|}\, dt. \tag{39.8}$$

The sum counts a word at most once at each roof, even when its set of physical radii is disconnected. It allows infinite values and retains arbitrary original section/return selectors. It is not a fixed-radius LLT and does not bound the complete sum of the displayed weights.

Proof. Use the countable regular local charts and first-chart Borel allocation of Corollary 36.4. Each chart is locally invertible in radius and roof. The new global order proves that charts of the same word cannot yield distinct physical points at the same roof. Their inverse graphs therefore agree on every overlap representing a point, and combine into the indicated Borel partial graph. Nonnegative finite chart substitution and monotone convergence give the identity, after regrouping by word. This removes coordinate and disconnected-component multiplicity, not any physical record. The reciprocal slope lies between $1/(2m)$ and $1/(\kappa_0 m)$ at every admitted term, as previously proved. No radius average is exchanged with a radius supremum. □

40. Physical clearance certificates and complete-source localization

The endpoint defect in (38.5) is available for every original regular trajectory, including those whose geometric word minimum is not a true critical orbit. We quantify this distinction rather than discarding that source. The final proposition then inserts the proved enclosure into the existing exact source formula, at the original fixed radius and roof.

Theorem 40.1 (A quantitative certificate relative to every minimizing contact)**.** *Let Q be an arbitrary true regular record and let $E = E_R(Q)$. For every minimizer Y defining $\ell_w(R)$ set*

$$\delta_0 = \delta_m = 2\sqrt{2RE}, \qquad \delta_i = \sqrt{RE/c_i} \quad (0 < i < m).$$

Then $|Y_i - Q_i| \le \delta_i$ for every i. Consequently, for every unvisited center z on every original flight, without a lattice truncation,

$$\operatorname{dist}(z, [Y_i, Y_{i+1}]) - R \ge \operatorname{dist}(z, [Q_i, Q_{i+1}]) - R - \max(\delta_i, \delta_{i+1}). \tag{40.1}$$

There is a sufficient proved *physicality certificate: if*

$$\Delta_i > \max(\delta_i, \delta_{i+1}) \quad (0 \le i < m), \tag{40.2}$$

$$c_i > (\delta_{i-1} + 2\delta_i + \delta_{i+1})/\tau_- \quad (0 < i < m), \tag{40.3}$$

then every minimizer Y is the same regular physical normal-endpoint record of the original word. No section label is thereby transferred from Q to Y. If these tests fail, Q stays in the original source; its minimum is not declared physical.

Proof. Insert the actual velocity vector of Q and a minimizing Y into (38.3). Weak duality and (38.4) give $0 \le S(Y) - D_{w,R}(V) \le E$. Thus each nonnegative contact term is at most E. At an interior contact it equals $2c_i\{R - n_i \cdot (Y_i - O_i)\} \ge (c_i/R)|Y_i - Q_i|^2$, which proves the interior bound.

At departure the supporting disk point for $-v_0$ is $O_0 + Rv_0$; the endpoint contact slack bounds its squared distance from Y_0 by $2RE$. Also $|Q_0 - (O_0 + Rv_0)|^2 = 2R^2(1 - c_0) \le 2RE$. The triangle inequality gives $|Y_0 - Q_0| \le 2\sqrt{2RE}$. At arrival use $O_m - Rv_{m-1}$ and the same argument. For a common affine interpolation parameter the distance between two corresponding segment points is at most $\max(\delta_i, \delta_{i+1})$. Both directed Hausdorff distances satisfy this bound. Distance to any fixed z is 1-Lipschitz for Hausdorff distance; this proves (40.1) for all lattice centers simultaneously.

It remains to justify physicality, not merely small displacement. The edges of Y have length at least $\tau_- = 3/50$, because successive original disks are disjoint. If $\widehat{v}_i$ is their unit direction, the normalization inequality $|u/|u| - v/|v|| \le 2|u - v|/\min(|u|, |v|)$ gives

$$|\widehat{v}_i - v_i| \le 2(\delta_i + \delta_{i+1})/\tau_-.$$

Hence the ith interior gradient of S at Y has norm at least $2c_i - 2(\delta_{i-1} + 2\delta_i + \delta_{i+1})/\tau_- > 0$. By the minimizing condition in Theorem 38.1, Y_i lies on its circle and that gradient is a negative multiple of the outward normal. Equality of the two adjacent unit speeds then makes the incoming normal component strictly negative and the outgoing one strictly positive, with equal tangential components: this is exactly the regular exterior specular law. At the endpoints the gradients have norm one and give normal departure and arrival. The strict third-disk inequalities (40.2) exclude every unlisted collision; the signs just proved exclude another intersection with either incident disk inside its flight. Thus Y is a true regular first-hit record. Corollary 38.3 gives uniqueness. Nothing in the proof keeps section visits constant along a competitor or along a homotopy; the original labels of Q have not changed. □

Proposition 40.2 (A deterministic positive junction cost)**.** *Partition a word into consecutive nonempty blocks at any chosen indices. For each block choose a minimizing unit-edge dual vector as in Theorem 38.1. Denote the last vector before a junction by v^- and the first after it by v^+. Then*

$$\ell_w(R) \ge \sum_{\text{blocks } b} \ell_b(R) + R \sum_{\text{junctions}} (2 - |v^- - v^+|). \tag{40.4}$$

Every junction term is nonnegative. If both block minima at a junction are physical critical records with terminal/departure normals n^-, n^+, the corresponding term is $R(2 - |n^- + n^+|)$. The comparison does not assert physical realizability of any block minimum and does not factor any trajectory probability.

Proof. Concatenate the chosen dual flight vectors. They form a feasible dual vector for the whole word. The center term C_w is the sum of the block center terms. At a junction the two separate endpoint penalties are $R|v^-| + R|v^+| = 2R$, whereas the penalty of the concatenated dual vector is $R|v^- - v^+|$. All other penalties agree. Substitution into (38.2) proves (40.4), including each junction once. In the physical critical case $v^- = -n^-$ and $v^+ = n^+$, giving the asserted normal expression. This argument is an exact vector comparison, not a renewal or Markov decomposition. □

Proposition 40.3 (Word localization of the unchanged complete roof source)**.** *Fix the original m, R, n, k and any nonnegative Borel trajectory weight h. Let $x_w(t, \vartheta)$ be exactly the partial physical inverse in* (10.6)*, with all original event cuts. That formula admits the additional exact mask*

$$\mathbf{1}_{\{\ell_w(R) \le t \le \ell_w(R) + E_R(x_w(t,\vartheta))\}} \tag{40.5}$$

in every existing summand, without changing either side. In particular only words with $\ell_w(R) \in [t-2R,t]$ *can contribute. This assertion requires no physical critical center, clearance guard, pressure hypothesis or incidence lower bound for the regular source.*

For clarity about what a pressure consequence does and does not add, define the deterministic localized physical-word pressure

$$P_{m,R}(J) = \sum_{\substack{w\in\mathcal{W}_m(R)\\ \ell_w(R)\in J}} \sup_{[w]_R\cap M'_R} j_m.$$

Under exactly *Input 11.1 and the inherited roof–angle formula, for* $0 \le h \le 1$,

$$\frac{dL_{\#}(h\mathbf{1}_{\Pi_{n,k,m,R}}\nu_R^*)}{dt}(t) \le \frac{C_0}{2c_*\lambda_\circ} P_{m,R}([t-2R,t]) \quad \textit{for almost every } t. \tag{40.6}$$

This is an unscaled comparison, not a new estimate on that localized pressure and not a proof of the original normalized shortage.

Proof. Theorem 38.2 applies to every existing physical inverse point, so its mask is one. Apply the nonnegative source exhaustion of Proposition 10.4; the mask is Borel, and no test or boundary is differentiated. This proves the exact identity with the mask, for extended weights as well. For the last bound use $\chi \ge \lambda_\circ$ and $\omega \le C_0 c_m j_m \le C_0 j_m$ at the actual point, and dominate h and the actual event only on the upper side by one. Global physical roof–angle uniqueness gives at most one point per word. Each word retained by the mask lies in the indicated geometric interval, so its stable weight is bounded by the same-word supremum at the *same* R. Integration over the angle of length 2π gives the factor in (40.6). The section factor c_*^{-1} occurs once, exactly as in the inherited formula. The supremum of this unlocalized pressure is the retained Q_*; no decay in m for the localized expression has been inferred. □

40.1. **The original positive shortage is still the closing condition.** The new results remove the need for a physical critical anchor in the geometric word enclosure; certify an actual-versus-unphysical minimizing configuration using nonnegative gaps; give a positive junction defect; and compare critical roofs across disconnected physical radius domains. They do not identify the auxiliary ℓ_w with a physical clock when the certificate fails. In particular (40.5) does not replace the observed microscopic window by $[t-2R,t]$. The observed quantity is still the density at the original t with the original exact labels; the interval is solely an upper-support enclosure in a separate deterministic word statistic.

No new theorem here supplies a normalized roof-height estimate for the remaining positive source or enough nongrazing comparison capacity. The raw target is still (1.5), with the very same minimal ordered closure condition (15.8). It includes the low-load incidence-only source, first-order unvisited clearance, noncritical singular charts, one-sided paired/section traces and all original arithmetic/denominator requirements. The finite source sum is taken before $\sup_R \operatorname{ess\,sup}_t$ and before the collision limsup; all microscopic widths remain fixed until that limsup, and $B \to \infty$ is outermost. Neither an auxiliary convex value, a positive curvature identity, a finite diagnostic, nor the radius-integrated coarea pays that still unproved limit automatically.

41. Uniform localization of the entire variational sublevel

41.1. **Actual records and the inherited comparison functional.** All lengths are in units of the triangular lattice spacing. Let

$$\Lambda = \{(a+b/2, \sqrt{3}b/2) : a,b \in \mathbb{Z}\}, \qquad R \in [9/20, 47/100], \quad R_+ = 47/100.$$

Fix any actual regular record of $m \geq 1$ free flights and $m+1$ collision points $Q_0, \dots, Q_m$. Its successive lattice centers are $O_0, \dots, O_m$, with $O_i \neq O_{i+1}$. Write

$$Q_i = O_i + Rn_i, \quad |n_i| = 1, \quad Q_{i+1} - Q_i = \tau_i v_i, \quad |v_i| = 1.$$

Regular means genuine exterior, first-hit, nontangent specular collisions at all these points and no extra collision on any open flight. Nonconsecutive repeated centers are permitted. At interior collisions let

$$c_i = -v_{i-1} \cdot n_i = v_i \cdot n_i > 0, \qquad v_i - v_{i-1} = 2c_i n_i \quad (1 \leq i < m). \tag{41.1}$$

The endpoint cosines are $c_0 = v_0 \cdot n_0 > 0$ and $c_m = -v_{m-1} \cdot n_m > 0$. Define

$$L = \sum_{i=0}^{m-1} \tau_i, \qquad E = R(2 - c_0 - c_m), \qquad 0 \leq E \leq 2R \leq 47/50 < 1. \tag{41.2}$$

The original endpoint momenta obey $p_0^2 + c_0^2 = p_m^2 + c_m^2 = 1$, so $E = R[p_0^2/(1+c_0) + p_m^2/(1+c_m)]$.

For this fixed ordered word, the comparison domain and length are

$$\mathcal{K} = \prod_{i=0}^{m} \overline{B}(O_i, R), \qquad A(Y) = \sum_{i=0}^{m-1} |Y_{i+1} - Y_i|.$$

Repeated centers give separate factors. This is the native disk-product comparison, not a substitute physical dynamics. Define its full sublevel

$$\mathcal{S}(Q) = \{Y \in \mathcal{K} : A(Y) \leq L\}.$$

It is nonempty and compact. Every global minimum of A lies in it, whether or not the minimum is physical. For $Y \in \mathcal{K}$ put

$$\delta_i = Y_i - Q_i, \qquad Y_{i+1} - Y_i = \ell_i \widehat{v}_i, \qquad e_i = \widehat{v}_i - v_i.$$

Consecutive disks are separated by $\tau_- = 1 - 2R_+ = 3/50$, so every $\ell_i \geq \tau_- > 0$. The retained geometric facts, reconstructed in Appendix A, are

$$\tau_i \leq H := 28/5, \qquad \sum_{i=a+1}^{a+93} c_i \geq \gamma := 1/4480 \quad \text{whenever } 0 \leq a,\ a+94 \leq m. \tag{41.3}$$

Thus every complete 94-flight block has an interior index with $c_i \geq c_{\mathrm{a}} := 1/416640$. No such condition is imposed on all individual incidences.

41.2. **The complete positive gap, not a coordinatewise singular estimate.** The next calculation extends the inherited endpoint-gap bookkeeping from minima to the entire sublevel. The general disk duality and the endpoint defect themselves were already established in module 297.

Lemma 41.1 (Full sublevel energy and contact budget)**.** *For every $Y \in \mathcal{S}(Q)$,*

$$\sum_{i=0}^{m-1} \ell_i |e_i|^2 \leq 2E, \tag{41.4}$$

$$\sum_{i=1}^{m-1} c_i |\delta_i|^2 \leq RE, \tag{41.5}$$

$$|\delta_0|, |\delta_m| \leq 2\sqrt{2RE}. \tag{41.6}$$

These statements hold for interior or inactive comparison vertices as well as boundary vertices.

Proof. Set $d_0 = -v_0$, $d_i = v_{i-1} - v_i = -2c_i n_i$ for $1 \le i < m$, and $d_m = v_{m-1}$. Telescoping, including both endpoints exactly once, gives

$$\sum_{i=0}^{m-1} v_i \cdot (Y_{i+1} - Y_i) = \sum_{i=0}^{m} d_i \cdot Y_i.$$

Consequently, with $D = \sum_i (d_i \cdot O_i - R|d_i|)$,

$$A(Y) - D = \frac{1}{2} \sum_{i=0}^{m-1} \ell_i |e_i|^2 + \sum_{i=0}^{m} \big[d_i \cdot (Y_i - O_i) + R|d_i| \big]. \tag{41.7}$$

Every term is nonnegative: the first uses two unit vectors, and the second is the difference from the minimum of a linear functional on its actual disk. At Q every flight and interior contact term is zero; the endpoint terms are $R(1 - c_0)$ and $R(1 - c_m)$. Thus $L - D = E$. Since $A(Y) \le L$, the whole right-hand side of (41.7) is at most E, proving (41.4).

For any $z \in \overline{B}(0, R)$ and unit n, the exact identity

$$2R(R - n \cdot z) = |z - Rn|^2 + R^2 - |z|^2 \tag{41.8}$$

implies $|z - Rn|^2 \le 2R(R - n \cdot z)$. An interior contact term in (41.7) equals $2c_i(R - n_i \cdot (Y_i - O_i))$ and is therefore at least $c_i |\delta_i|^2 / R$. Summing proves (41.5), without dividing by any individual c_i.

At the first endpoint let $Z_0 = O_0 + Rv_0$. The endpoint term gives $|Y_0 - Z_0|^2 \le 2RE$ by (41.8), while $|Q_0 - Z_0|^2 = 2R^2(1 - c_0) \le 2RE$. The triangle inequality proves the first bound in (41.6). At the final endpoint use $Z_m = O_m - Rv_{m-1}$; the signs give the same bound. □

Lemma 41.2 (Transverse telescoping with the turning work retained)**.** *Put $w_i = v_i^{\perp}$ and $r_i = \ell_i e_i$. For $0 \le j < i \le m$,*

$$w_{i-1} \cdot \delta_i = w_j \cdot \delta_j + \sum_{l=j}^{i-1} w_l \cdot r_l + \sum_{l=j+1}^{i-1} (w_l - w_{l-1}) \cdot \delta_l. \tag{41.9}$$

If $i - j \le 94$ and $Y \in \mathcal{S}(Q)$, the absolute values of the last two sums are respectively bounded by $38\sqrt{E}$ and $14\sqrt{E}$.

Proof. The exact displacement equation is

$$\delta_{l+1} - \delta_l = (\ell_l - \tau_l) v_l + r_l.$$

Its projection on w_l removes only its longitudinal term. Summing these projected equations and collecting coefficients of each δ_l yields (41.9). The last sum is present because the transverse frame turns. It cannot in general be omitted.

All comparison vertices obey $|\delta_l| \le 2R$, so

$$\ell_l \le \tau_l + |\delta_l| + |\delta_{l+1}| \le H + 4R_+ = 187/25 =: T.$$

Weighted Cauchy–Schwarz and (41.4) give

$$\sum_{l=j}^{i-1} |r_l| \le \left(\sum_{l=j}^{i-1} \ell_l \right)^{1/2} \left(\sum_{l=j}^{i-1} \ell_l |e_l|^2 \right)^{1/2} \le \sqrt{2 \cdot 94 \cdot TE} < 38\sqrt{E}.$$

The rotation is an isometry, so (41.1) gives $|w_l - w_{l-1}| = 2c_l$. Therefore (41.5) gives

$$\sum_{l=j+1}^{i-1} |(w_l - w_{l-1}) \cdot \delta_l| \le 2 \left(\sum_{l=j+1}^{i-1} c_l \right)^{1/2} \left(\sum_{l=j+1}^{i-1} c_l |\delta_l|^2 \right)^{1/2} \le 2\sqrt{94 R_+ E} < 14\sqrt{E}.$$

Only actual reflection cosines enter the turning sum. In particular no dynamics is ascribed to the comparison polygon. □

Theorem 41.3 (Collision-uniform sublevel localization without an incidence cutoff)**.** *For every actual regular record in the original radius family, every $m \geq 1$, and every $Y \in \mathcal{S}(Q)$,*

$$\max_{0\leq i\leq m} |Y_i - Q_i| \leq 22\, E^{1/4}. \tag{41.10}$$

The constant is independent of the word, collision count, section/return/displacement labels and every interior incidence. At $E = 0$ the sublevel is the singleton $\{Q\}$. No optimality of the exponent or nontriviality for large E is asserted.

Proof. If $E = 0$, (41.5) and positivity of the actual interior cosines force all interior displacements to vanish, and (41.6) handles the endpoints. Assume $0 < E < 1$.

Call an interior index an anchor if $c_i \geq c_{\mathrm{a}}$. At such an index,

$$|\delta_i| \leq \sqrt{R_+/c_{\mathrm{a}}}\sqrt{E} < 443\sqrt{E}.$$

Together with $|\delta_i| \leq 2R_+$ this gives $|\delta_i|^2 < 2R_+443\sqrt{E} < 22^2\sqrt{E}$. At either endpoint (41.6) gives $|\delta_i|^2 \leq 8R_+E < 4\sqrt{E} < 22^2\sqrt{E}$.

Consider a nonanchor interior index i. Let $j < i$ be the last anchor before it, or $j = 0$ if none exists. Necessarily $i - j < 94$: otherwise the 94-flight block starting at j has by (41.3) an anchor strictly between j and i, a contradiction. The initial displacement in (41.9) is at most $443\sqrt{E}$, also when $j = 0$ by (41.6). Lemma 41.2 therefore yields

$$|w_{i-1} \cdot \delta_i| \leq (443 + 38 + 14)\sqrt{E} = 495\sqrt{E}.$$

Because $n_i \cdot v_{i-1} = -c_i$ and $|n_i \cdot w_{i-1}| \leq 1$,

$$|n_i \cdot \delta_i| \leq c_i|\delta_i| + |w_{i-1} \cdot \delta_i|.$$

Crucially, the first term needs no inverse incidence: by (41.5), $c_i|\delta_i| \leq \sqrt{c_i}\sqrt{c_i|\delta_i|^2} \leq \sqrt{R_+E} < \sqrt{E}$. Thus $|n_i \cdot \delta_i| \leq 496\sqrt{E}$. Since Y_i lies in the disk and Q_i on its boundary, (41.8) becomes

$$|\delta_i|^2 \leq -2Rn_i \cdot \delta_i \leq 2R_+496\sqrt{E} = \frac{11656}{25}\sqrt{E} < 484\sqrt{E}.$$

Taking square roots proves (41.10). The argument treats short records and the first incomplete block without inserting an artificial collision. It uses at most one bounded physical block for each index, not a sum over the full record or over its words. □

Remark 41.4 (What is new relative to the complete predecessor). The earlier pointwise contact certificate had a factor $c_i^{-1/2}$. Here the contact budget is used *before* division: its weighted sum pays the turning work in (41.9), and disk curvature converts the resulting normal control into an unweighted position bound. The dense-anchor input is inherited real geometry, not a fresh probability hypothesis. The result concerns every point of a continuous compact sublevel, not a list of sampled minima. It neither constructs an orbit through a singularity nor controls forward-time derivatives at grazing.

42. All-center clearance and a true physical reference

For a record Q put $S_i(Q) = [Q_i, Q_{i+1}]$. Define the true third-disk clearance of a flight by

$$\Gamma_i(Q) = \inf_{z\in\Lambda\setminus\{O_i,O_{i+1}\}} \operatorname{dist}(z, S_i(Q)) - R, \qquad \Gamma(Q) = \min_{0\leq i<m} \Gamma_i(Q). \tag{42.1}$$

The infimum includes disks visited elsewhere in the word. It excludes only the two incident centers of this particular flight. Distances are Euclidean distances to closed segments, so a minimizing foot may change or reach an endpoint. There is no chosen differentiable nearest-center label.

Theorem 42.1 (One modulus for every third lattice disk)**.** *For every $Y \in \mathcal{S}(Q)$ and $d(E) = 22E^{1/4}$,*

$$d_H(S_i(Y), S_i(Q)) \le d(E), \tag{42.2}$$

$$|\operatorname{dist}(z, S_i(Y)) - \operatorname{dist}(z, S_i(Q))| \le d(E) \quad \text{for every } z \in \mathbb{R}^2, \tag{42.3}$$

$$|\Gamma_i(Y) - \Gamma_i(Q)| \le d(E). \tag{42.4}$$

All these estimates hold simultaneously over all flights and all lattice centers. They require no lower bound on any interior incidence.

Proof. Match points with the same affine segment parameter $s \in [0,1]$. Their difference is $(1-s)\delta_i + s\delta_{i+1}$, whose norm is at most $\max\{|\delta_i|, |\delta_{i+1}|\} \le d(E)$ by Theorem 41.3. This bounds both directed Hausdorff distances. For two nonempty compact sets at Hausdorff distance at most d, the triangle inequality gives $\operatorname{dist}(z, A) \le \operatorname{dist}(z, B) + d$ and the reverse inequality, proving (42.3). Take infima over the *same* center set in both directions to obtain (42.4). The infima are finite. No counting or differentiability of the minimizing center is used. □

Theorem 42.2 (A grazing-independent clearance certificate for a physical minimum)**.** *Let Q be any original regular record, and put*

$$\varepsilon(E) = \sqrt{2E/\tau_-}, \qquad d(E) = 22E^{1/4}.$$

Suppose

$$c_i(Q) > \varepsilon(E) \ (1 \le i < m), \qquad \Gamma(Q) > d(E). \tag{42.5}$$

The first condition is vacuous for $m = 1$. Then A has exactly one global minimum on $\mathcal{K}$. This minimum is an actual regular exterior first-hit trajectory of the same ordered collision word, with normal initial and final endpoints. All these assertions concern the unchanged radius R and actual circular obstacles. They assert no equality of its section/return labels or physical roof with those of Q.

Proof. Compactness supplies at least one minimizing Y. Since $A(Y) \le A(Q)$, Theorem 41.3 and Lemma 41.1 apply. Each edge error obeys

$$|\widehat{v}_i - v_i| \le \sqrt{2E/\ell_i} \le \varepsilon(E).$$

All edges are nonzero because consecutive disks are separated. The gradient of A with respect to an interior vertex is $\widehat{d}_i = \widehat{v}_{i-1} - \widehat{v}_i$. By the physical reflection law of Q and (42.5),

$$|\widehat{d}_i| \ge 2c_i - |e_{i-1}| - |e_i| \ge 2c_i - 2\varepsilon(E) > 0.$$

First-order optimality on the ith closed disk means Y_i minimizes its linear functional $\widehat{d}_i \cdot z$. Hence $Y_i = O_i + R\widehat{n}_i$ with $\widehat{d}_i = -|\widehat{d}_i|\widehat{n}_i$. For two unit vectors a, b with $a - b = -\lambda n$, $\lambda > 0$, taking the scalar product with $a + b$ shows $a \cdot n = -b \cdot n$; taking it with n gives $a \cdot n = -\lambda/2$ and $b \cdot n = \lambda/2$. Their tangential components are equal. This proves a true nontangent specular reflection at each interior vertex, with the correct exterior signs. At the endpoints the gradients are $-\widehat{v}_0$ and $\widehat{v}_{m-1}$, of norm one. Their disk minimizers give $\widehat{n}_0 = \widehat{v}_0$ and $\widehat{n}_m = -\widehat{v}_{m-1}$, precisely normal endpoints.

Theorem 42.1 and the strict second inequality in (42.5) give $\Gamma_i(Y) > 0$ for every flight. Thus no third disk meets any flight, including a disk appearing at a different time in the word. The two incident disks are avoided in the open flight as well: positive departing normal component implies $|Y_i + s\widehat{v}_i - O_i|^2 = R^2 + 2Rs(\widehat{n}_i \cdot \widehat{v}_i) + s^2 > R^2$ for $s > 0$, and the analogous backward formula applies at arrival. This proves first-hit physicality without discarding any collision.

For uniqueness, use this physical Y as the reference in (41.7). Its normal endpoints have defect zero. Any other minimum Z has $A(Z) = A(Y)$, so the nonnegative gap is zero. All

contact coefficients are strictly positive (one at each endpoint and $2\widehat{c}_i$ internally), and (41.8) forces $Z_i = Y_i$ at every vertex. Thus the minimum is unique. □

Remark 42.3 (Sufficiency, not source deletion or capacity). The test may fail while a physical minimum still exists. Failed records remain in the original source. The incidence requirement in (42.5) checks nonzero comparison reflections; it is not an inverse-incidence loss in the clearance modulus. An actual critical reference is one trajectory, not a positive measure comparison pool. Exact section visits and return labels must be evaluated anew on it, not transported from Q by proximity. The original observed roof remains $L(Q)$, never $A(Y)$.

Proposition 42.4 (Positive inclusion on the unchanged exact-record source)**.** *Fix R, m and any of the original exact return/displacement/collision events* Π*, with its original section probability measure ν_R^* and nonnegative measurable weight h. Let $\mathcal{B}$ be the subset of regular records whose ordered disk word does not have a unique regular physical normal-endpoint global minimum. Set*

$$\mathcal{W} = \{Q : \ \exists i \in \{1, \dots, m-1\},\ c_i(Q) \le \varepsilon(E(Q)) \quad or \quad \Gamma(Q) \le d(E(Q))\}\,. \tag{42.6}$$

On the regular source, $\mathcal{B} \subseteq \mathcal{W}$. Consequently the following positive measure inequality holds on the original physical roof axis*:*

$$L_*(\mathbf{1}_{\mathcal{B}}\mathbf{1}_{\Pi} h \, d\nu_R^*) \ \le \ L_*(\mathbf{1}_{\mathcal{W}}\mathbf{1}_{\Pi} h \, d\nu_R^*). \tag{42.7}$$

This includes all physical words with one-copy multiplicity and extends to nonnegative marked sums by monotone convergence. It changes neither the conditioned event nor any roof observation window. It asserts no small upper bound on the right-hand side.

Proof. The inclusion is the contrapositive of Theorem 42.2. At fixed R, m the physical ordered-word collection is countable; the property defining $\mathcal{B}$ is a property of a word, not of a chosen minimizer. Thus $\mathcal{B}$ is a union of its measurable regular word cylinders. The functions in (42.6) are measurable on every regular chart: each center-to-segment distance is continuous, the center set is countable, and finite minima and the displayed elementary functions preserve measurability. No measurable selection of a minimum is needed.

Multiply the pointwise inclusion by $\mathbf{1}_{\Pi} h \mathbf{1}_{\{L \in I\}}$ for any Borel roof set I, then integrate. This gives (42.7), initially for bounded h and then by increasing truncation. Apply the same argument on disjoint original marked copies and use monotone convergence for complete nonnegative sums. The section normalization is the original one already in ν_R^*; no additional Jacobian or copy factor appears. □

42.1. **Exact remaining analytic obligation.** If the two pushforwards are absolutely continuous, (42.7) orders their densities almost everywhere at the same R, m, n, k, t. Without that hypothesis it remains a measure inequality, not an assertion that a singular component vanishes. Paired boundary traces are not bulk measures and are not set to zero by this argument.

The positive witness source in (42.7) is *not* proved to be small in $m^2 L^\infty(dt)$. Nor are all physical-reference records shown to meet the original nongrazing comparison-pool constraints. The original first-order single/cross-flight clearance, low-load incidence source, degenerate/critical charts, section cuts and singular pairing terms therefore remain in (0.1). The present theorem supplies a proved geometric mechanism for obtaining a physical reference without the previous inverse-incidence clearance loss. A normalized source/pressure or original-capacity argument must still exploit that mechanism, with the original full sums and order of limits, to close the raw theorem.

43. Summable physical coercivity and critical singular closure

The maximum estimate in Theorem 41.3 is inherited. Summing its fourth power would introduce a factor $m+1$. We instead sum the positive gap before estimating the transverse work. This gives one budget for the complete record, including arbitrarily small incidences. Throughout, $R \in [9/20, 47/100]$, $m \geq 1$, and the notation $Q, O, n, v, \tau, Y, \ell, \delta, e, L, D, E$ is exactly that of Section 41. Set

$$r = 47/100, \quad N = 94, \quad a = 1/416640, \quad T = 187/25, \quad \tau_- = 3/50. \tag{43.1}$$

No probability, roof window or return event is changed by the following comparison.

Definition 43.1 (Closure of a fixed physical word)**.** A closed physical reference is a coordinatewise limit of actual regular records with the same finite ordered centers $(O_0, \dots, O_m)$, with radii converging inside the original closed radius interval. The record length and word are fixed while taking this closure. At a limiting contact there may be zero incidence or an unlisted disk tangency. We regard the limit as a polygonal reference, not as a definition of the billiard map at that singularity, and do not assign it return labels by continuity.

Lemma 43.2 (Positive gap on the closed physical reference)**.** *Actual regular records and closed physical references satisfy the reflection and block bounds*

$$v_i - v_{i-1} = 2c_i n_i, \quad 0 \leq c_i \leq 1, \qquad \sum_{i=b+1}^{b+93} c_i \geq 1/4480 \quad (b+94 \leq m). \tag{43.2}$$

For every $Y \in \mathcal{K}$, not only a minimizing polygon, put

$$G = A(Y) - D, \qquad h_i = d_i \cdot (Y_i - O_i) + R|d_i|, \qquad E_0 = R(1-c_0), \quad E_m = R(1-c_m).$$

Then $G \geq 0$, $E = E_0 + E_m = L - D$, and

$$G = \tfrac{1}{2} \sum_{i=0}^{m-1} \ell_i |e_i|^2 + \sum_{i=0}^{m} h_i, \qquad h_i \geq 0, \tag{43.3}$$

$$\sum_{i=1}^{m-1} c_i |\delta_i|^2 \leq R \sum_{i=1}^{m-1} h_i, \qquad \sum_i \ell_i |e_i|^2 \leq 2G, \tag{43.4}$$

$$|\delta_0|^2 \leq 4R(h_0 + E_0), \qquad |\delta_m|^2 \leq 4R(h_m + E_m). \tag{43.5}$$

Also $|\delta_i| \leq 2r$ and $\tau_- \leq \ell_i \leq T$.

Proof. For regular records, the exact gap is the telescoping identity (41.7); it holds on the entire disk product. The disk identity (41.8) gives each interior contact inequality, including when $c_i = 0$. For the first endpoint, use $Z_0 = O_0 + Rv_0$. Then $|Y_0 - Z_0|^2 \leq 2Rh_0$ and $|Q_0 - Z_0|^2 = 2RE_0$. The inequality $|u+v|^2 \leq 2|u|^2 + 2|v|^2$ gives (43.5); use $Z_m = O_m - Rv_{m-1}$ at the final endpoint. Since consecutive disks are separated, every edge is nonzero, even on the whole comparison product. The upper bound is $\ell_i \leq \tau_i + |\delta_i| + |\delta_{i+1}| \leq 28/5 + 4r = T$. For a closed reference, positions, unit edge directions, normals, incidences and lengths converge: their denominators are bounded below by R_- or τ_-. The regular reflection equations, horizon bound and each fixed finite block inequality therefore pass to the limit. The same algebra then applies directly at the limit. This passage does not continue a flow across a grazing collision. □

Theorem 43.3 (Complete-record quartic coercivity)**.** *For every actual regular record or closed physical reference, every $m \geq 1$ and every $Y \in \mathcal{K}$,*

$$\sum_{i=0}^{m} |Y_i - Q_i|^4 \leq C_G G + C_E E, \tag{43.6}$$

where the following constants are independent of the word, its length, the original radius, and all incidences:

$$C_G = \frac{4r^3}{a} + 16r^3 + 8r^2\left(r + \frac{3Nr}{a} + 6N^2T + 12N^2r\right) = \frac{12318659063517}{125000}, \tag{43.7}$$

$$C_E = 16r^3 + 96Nr^3 = \frac{11731999}{12500}. \tag{43.8}$$

In particular, with $C = 10^8$, the entire inherited sublevel obeys

$$\sup_{Y\in\mathcal{S}(Q)} \sum_{i=0}^{m} |Y_i - Q_i|^4 \le CE. \tag{43.9}$$

The supremum is over the continuous compact sublevel, not a sample of minima. This is a summability estimate, rather than the m-multiple of a one-contact bound.

Proof. Call an interior index an anchor when $c_i \ge a$. Let $\mathcal{A}$ be the anchor set and $\mathcal{I}$ the other interior indices. For each $i \in \mathcal{I}$, let $j(i)$ be the last anchor strictly before i, or 0 when there is none. Equation (43.2) implies $i - j(i) < N$: a complete N-flight block starting at $j(i)$ would otherwise have an interior anchor before i. Thus each anchor or initial endpoint is used as a start at most N times. Each flight or interior contact lies in at most N of the intervals $[j(i), i]$. These are deterministic overlap bounds on one record; there is no multiplication by the number of its words.

Put $w_l = v_l^{\perp}$ and $b_i = w_{i-1} \cdot \delta_i$. The exact transverse telescope (41.9), which also holds in the closure, and the inequality $(x + y + z)^2 \le 3(x^2 + y^2 + z^2)$ give

$$|b_i|^2 \le 3|\delta_{j(i)}|^2 + 3NT \sum_{l=j(i)}^{i-1} \ell_l|e_l|^2 + 12N \sum_{l=j(i)+1}^{i-1} c_l|\delta_l|^2.$$

Indeed the squared flight term is at most $(\sum \ell_l)(\sum \ell_l|e_l|^2)$; the squared turning term is at most $4(\sum c_l)(\sum c_l|\delta_l|^2)$, since $|w_l - w_{l-1}| = 2c_l$. All terms here are physical reflection quantities of the reference, not of Y.

The starting-point budget is shared, not paid anew for every i. Disjointness of the anchor and initial contact terms and $4r \le r/a$ imply

$$\sum_{j\in\mathcal{A}} |\delta_j|^2 + |\delta_0|^2 \le \frac{r}{a} \sum_{j\in\mathcal{A}} h_j + 4rh_0 + 4rE_0 \le \frac{r}{a}G + 4rE.$$

Summing the preceding transverse bound and using the overlap at most N and (43.4) now proves

$$\sum_{i\in\mathcal{I}} |b_i|^2 \le \left(\frac{3Nr}{a} + 6N^2T + 12N^2r\right) G + 12NrE. \tag{43.10}$$

This is where a complete-record budget replaces the unbounded number of local estimates.

For $i \in \mathcal{I}$, disk curvature and $n_i \cdot v_{i-1} = -c_i$ give

$$|\delta_i|^2 \le -2Rn_i \cdot \delta_i \le 2r(c_i|\delta_i| + |b_i|), \qquad |\delta_i|^4 \le 8r^2(c_i^2|\delta_i|^2 + |b_i|^2).$$

Use $c_i^2 \le c_i$, sum, and apply (43.4) and (43.10). The nonanchor contribution is at most

$$8r^2\left(r + \frac{3Nr}{a} + 6N^2T + 12N^2r\right) G + 96Nr^3E.$$

For anchors, $|\delta_i|^4 \le 4r^2|\delta_i|^2$ and the contact budget give a total at most $4r^3G/a$. The two endpoints together contribute at most $16r^3(G + E)$ by (43.5). Adding these three disjoint classes

proves (43.6). It also covers $m = 1$, empty anchor sets and both incomplete end blocks. The exact rational evaluations (43.7)–(43.8) satisfy

$$C_G + C_E = \frac{12318776383507}{125000} < 10^8, \qquad 10^8 - C_G - C_E = \frac{181223616493}{125000} > 0.$$

On $\mathcal{S}(Q)$, $0 \le G = A(Y) - D \le L - D = E$, proving the claimed supremum estimate. No exchange of a collision limit with that supremum has been made. □

Corollary 43.4 (Coercivity at a critical singular limit)**.** *If the closed physical reference has normal endpoints, so $E = 0$, then for every $Y \in \mathcal{K}$*

$$A(Y) \ge L + C_G^{-1} \sum_{i=0}^{m} |Y_i - Q_i|^4. \tag{43.11}$$

Thus Q is the unique disk-product minimum, even if some internal incidences vanish. An approximate minimum of excess at most η has $\sum_i |Y_i - Q_i|^4 \le C_G \eta$.

Proof. Here $D = L$, so $G = A(Y) - L$. Apply (43.6). Nonnegativity and equality force every coordinate to agree with Q; the approximate assertion follows from $G \le \eta$. In particular this proof does not divide by a vanishing incidence, unlike the regular critical contact argument alone. □

Corollary 43.5 (Unique fixed-word compactification)**.** *Fix m and its ordered centers. Any sequence of actual regular normal-endpoint records of that word with $R_j \to R_*$ in the original interval converges to the same closed physical reference. Its roof equals $\ell_w(R_*)$. This asserts uniqueness of the limiting polygon, not physical continuation, section-label continuity or a coarea density at the singular limit.*

Proof. The finite product of bounded disks is compact. Every convergent subsequence is a closed physical reference with normal endpoints. By Corollary 43.4, each such limit is the unique minimizer of the same functional at R_*. All subsequential limits therefore agree, which proves convergence of the entire sequence and the roof identity by continuity of Euclidean length. The conclusion does not require the set of admissible radii to be connected. No orbit is inserted at a radius where none is physical. □

Remark 43.6 (What the singular statement does not remove). The limiting word may have an unlisted tangency and a singular contact can carry paired boundary data not seen by bulk collision measure. Uniqueness of its minimizing polygon does not bound the inverse roof Jacobian, continue the return map, or show that its normalized paired source vanishes. All such trace terms in the inherited source remain present. The constants above are intentionally nonoptimal; their independence of the collision count and incidence is the conclusion.

44. Summable clearance and exact-mark transport on the physical clock

We retain the physical model and the actual event (1.1), including $d\nu_R = d\alpha\, dp/(4\pi)$ and $d\nu_R^* = c_*^{-1} \mathbf{1}_{Y_R^*} d\nu_R$, where $c_* = 91/(10000\pi)$. Statements involving physical observations below are only made when both compared configurations are actual regular records of the same ordered word. The geometric statement comes first and needs no such physicality of the comparison polygon.

Theorem 44.1 (One budget for every center and every flight)**.** *For any reference and any $Y \in \mathcal{S}(Q)$ allowed by Theorem 43.3, define*

$$\kappa_i = \sup_{z \in \Lambda} \left| \operatorname{dist}(z, [Y_i, Y_{i+1}]) - \operatorname{dist}(z, [Q_i, Q_{i+1}]) \right|.$$

Then

$$\sum_{i=0}^{m-1} \kappa_i^4 \le 2CE, \qquad \#\{i : \kappa_i \ge \rho\} \le 2CE\rho^{-4} \quad (\rho > 0). \tag{44.1}$$

The same estimates apply to the absolute differences of the unvisited-center clearances Γ_i in Section 42. All centers, including a center visited on a different flight, and all possible feet on the closed segments are included. Negative clearance of a nonphysical Y is permitted and is not suppressed.

Proof. Using the same affine parameter on the two segments gives their Hausdorff distance at most $\max\{|\delta_i|, |\delta_{i+1}|\}$. Distance to any fixed point is 1-Lipschitz for Hausdorff distance, so the bound holds simultaneously for every lattice center, without any enumeration. Thus $\kappa_i^4 \le |\delta_i|^4 + |\delta_{i+1}|^4$. The full sum is at most twice the contact sum in (43.9); each endpoint is in fact counted only once. Taking infima over the identical set $\Lambda \setminus \{O_i, O_{i+1}\}$ preserves this uniform difference bound for Γ_i. Finally $\rho^4 \#\{i : \kappa_i \ge \rho\} \le \sum_i \kappa_i^4$. □

44.1. **Physical reference class and the original section.** Fix R and m. Let $\mathcal{C}_{R,m}$ be the union of actual regular word cells whose disk-product minimum is an actual regular normal-endpoint record Y^w. Such a minimum is unique by the retained critical result, or Corollary 43.4. The sufficient strict physicality certificate of Theorem 42.2 yields a subclass; no claim is made that all words lie in $\mathcal{C}_{R,m}$. Word cells and their original exact-event subsets are Borel, with the usual countable singularity partition. For this fixed R, m, the assignment of the constant record Y^w on each word cell is therefore Borel; it involves no unproved measurable choice of auxiliary minima.

For $Q \in \mathcal{C}_{R,m}$ compare its $m+1$ collision states $x_i = (\alpha_i, p_i)$ with $\widehat{x}_i$ of Y^w. Angles are taken in $\mathbb{R}/(2\pi\mathbb{Z})$, and

$$d_\infty(x, \widehat{x}) = \max\{\mathrm{dist}_{\mathbb{S}^1}(\alpha, \widehat{\alpha}), |p - \widehat{p}|\}.$$

At the last collision, the tangential momentum can be computed from the incoming flight: specular reflection leaves that component unchanged. Consequently the terminal state, although its next flight is not in the record, is fully determined for the comparison.

Theorem 44.2 (Full phase budget and section discrepancy)**.** *Put $R_- = 9/20$ and*

$$K = 128/\tau_- + 24C/R_-^4. \tag{44.2}$$

For every actual $Q \in \mathcal{C}_{R,m}$, writing $E = E_R(Q)$,

$$\sum_{i=0}^{m} d_\infty(x_i, \widehat{x}_i)^4 \le KE. \tag{44.3}$$

Let $\eta_R = \mathbf{1}_{Y_R^*}$ denote the original section indicator, not a smoothed or enlarged section, and define

$$N_{\mathrm{cut}}(Q) = \sum_{i=0}^{m} |\eta_R(x_i) - \eta_R(\widehat{x}_i)|,$$

$$B_\rho(Q) = \sum_{i=0}^{m} \mathbf{1}_{\{\mathrm{dist}_\infty(x_i, \partial Y_R^*) \le \rho\}}.$$

Then for every $\rho > 0$

$$N_{\mathrm{cut}}(Q) \le B_\rho(Q) + K\rho^{-4} E_R(Q). \tag{44.4}$$

Both endpoints are included once, whereas the original occupation count $A_m = \sum_{i=0}^{m-1} \eta_R(x_i)$ still excludes the final state. In particular $|A_m(Q) - A_m(Y^w)| \le N_{\mathrm{cut}}(Q)$.

Proof. Since Y^w is physical and lies in $\mathcal{S}(Q)$, all its vertices are on the original disks. The unit normals differ by $|\widehat{n}_i - n_i| = |\delta_i|/R$. For angles at circular distance $\theta \in [0,\pi]$, $2\sin(\theta/2) \geq 2\theta/\pi$. Thus

$$\operatorname{dist}_{\mathbb{S}^1}(\alpha_i, \widehat{\alpha}_i) \leq \frac{\pi}{2R}|\delta_i| \leq \frac{2}{R_-}|\delta_i|.$$

Tangential projection of the outgoing unit vector gives $|p_i - \widehat{p}_i| \leq |e_i| + |\delta_i|/R_-$ for $i < m$. The terminal incoming projection gives the same estimate with e_{m-1}. Using $(u+v)^4 \leq 8(u^4+v^4)$ and bounding a maximum by a sum, we get

$$d_\infty(x_i, \widehat{x}_i)^4 \leq 8|e_{\iota(i)}|^4 + 24R_-^{-4}|\delta_i|^4, \quad \iota(i) = \min(i, m-1).$$

The last flight index is used twice and the others once. Since the difference of two unit vectors has norm at most 2,

$$\sum_i |e_i|^4 \leq 4\sum_i |e_i|^2 \leq 8G/\tau_- \leq 8E/\tau_-.$$

Summing the phase estimate and using (43.9) proves (44.3).

The original section is a finite union of open rectangles in these circle–momentum coordinates. If its indicator differs at two points, a shortest circle–momentum path between them meets its boundary. Hence a discrepancy at index i implies $\operatorname{dist}_\infty(x_i, \partial Y_R^*) \leq d_\infty(x_i, \widehat{x}_i)$. If the left side is at most ρ, it is counted by B_ρ; otherwise its indicator is at most $\rho^{-4} d_\infty(x_i, \widehat{x}_i)^4$. Sum over all i and use (44.3). Points on the boundary are explicitly counted, not assigned zero cost by a bulk-null argument. The occupation assertion follows by dropping the last nonnegative discrepancy. □

Proposition 44.3 (Exact lattice labels and all four twist variables)**.** *On $\mathcal{C}_{R,m}$ the true count m and lattice displacement k of Q and Y^w agree. Their true roofs satisfy $0 \leq L(Q) - L(Y^w) \leq E_R(Q)$. For every $\xi \in \mathbb{R}^2$, $s, u \in \mathbb{R}$,*

$$\left| e^{i[\xi\cdot k + sL(Q) + uA_m(Q)]} - e^{i[\xi\cdot k + sL(Y^w) + uA_m(Y^w)]} \right| \leq |s| E_R(Q) + 2|\sin(u/2)| N_{\mathrm{cut}}(Q). \tag{44.5}$$

The factors $\eta_R(x_0)\eta_R(x_m)$, if included, incur in addition at most the two endpoint discrepancies. No label stability is asserted when that discrepancy has not been paid.

Proof. The physical ordered centers are identical; thus the displacement and number of flights are identical, without dropping a reflection. The actual critical minimum has length $\ell_w(R)$, so the inherited endpoint-gap theorem gives the roof inequality. The triangle inequality and $|e^{ia} - e^{ib}| \leq |a-b|$ bound the roof part. For an integer q, telescoping q unit phase increments gives $|e^{iuq} - 1| \leq |q|\,|e^{iu} - 1| = 2|q||\sin(u/2)|$, including negative q. Take $q = A_m(Q) - A_m(Y^w)$ and use Theorem 44.2. A product of two indicators has difference at most the sum of the differences of its two factors. This also shows exactly where the initial and terminal section tests enter. □

Theorem 44.4 (Positive discrepancy payment on the unchanged roof)**.** *For every R, m, n, k and nonnegative Borel weight h, let Π be the original exact event (1.1) and define, on the class $\mathcal{C}_{R,m}$ only,*

$$\mathcal{M}_h[f] = (L_R)_*(\mathbf{1}_{\mathcal{C}_{R,m}} \mathbf{1}_\Pi h f \, d\nu_R^*).$$

The pushforward is always by the actual original roof $L_R(Q)$, not the comparison roof. As an inequality of positive measures,

$$\mathcal{M}_h[N_{\mathrm{cut}}] \leq \mathcal{M}_h[B_\rho] + K\rho^{-4}\mathcal{M}_h[E_R] \quad (\rho > 0). \tag{44.6}$$

It remains valid after a countable sum over all physical word cells and nonnegative marked copies, with their actual multiplicities. If all three measures have densities, the same inequality holds between the densities almost everywhere, at the same R, m, n, k, t.

Proof. Multiply (44.4) by the nonnegative actual weight $\mathbf{1}_{\mathcal{C}_{R,m}}\mathbf{1}_{\Pi}h$ and integrate its product with any nonnegative Borel test of $L_R(Q)$. This proves the measure inequality, first for bounded weights and then by monotone convergence. Countable marked sums are handled by Tonelli before any supremum or limiting operation. The conditional collision probability already contains exactly the factor c_*^{-1}; no second section normalization is introduced. There is no coordinate change in this proof, and hence no unrecorded Jacobian or physical time multiplier. The density statement is the order property of Radon–Nikodym derivatives when absolute continuity has actually been established. □

Remark 44.5 (The exact remaining normalization). The new term B_ρ is a specified observation-boundary occupation, and E_R is the actual endpoint defect. Their occurrence is an explicit error accounting for real labels, not a claim that either has small normalized density. The class complementary to $\mathcal{C}_{R,m}$ is not discarded. Equations (44.6) and (44.5) alone give no m^{-2} estimate, no high-s uniform decay, and no comparison-pool capacity. In particular choosing ρ as a function of m requires estimates uniform at those widths; it is not justified by a fixed-width result. The singular paired endpoint traces are not bulk pushforwards and are not declared paid by this theorem. The complete source must still satisfy the original (15.8), with all exact labels, roof intervals, ordered suprema and outermost B limit unchanged. This paper does not assert completion of the raw pointwise theorem or its four-coordinate inversion consequence.

45. Boundary localization of a global disk-word minimum

The summable sublevel theorem gives a collision-independent budget, but does not by itself locate its mass. Here global minimality pays the variational work in the interior by work in two prescribed transition layers. A nonlinear recurrence then gives a uniform algebraic tail. No regular physical continuation from the reference to the comparison is assumed.

Let $m \geq 1$ and let $Q_0, \dots, Q_m$ be an original regular physical record, or a fixed-word closed physical reference, at an original radius $R \in [R_-, r] = [9/20, 47/100]$. Its ordered centers are O_i. Let Y be any global minimum of the unchanged disk-word action

$$A(Y) = \sum_{i=0}^{m-1} |Y_{i+1} - Y_i|, \qquad Y_i \in \overline{B}(O_i, R).$$

Consecutive centers are distinct. All original and comparison flights therefore have length at least $\tau_- = 3/50$. Use the retained physical constants

$$N = 94, \qquad a = 1/416640, \qquad T = 187/25, \qquad C = 10^8.$$

Every complete N-flight reference block has an interior contact with incidence at least a. The retained physical horizon is $\tau_i \leq 28/5$; hence every feasible comparison edge satisfies $\ell_i \leq \tau_i + |\delta_i| + |\delta_{i+1}| \leq 28/5 + 4r = T$. These statements include fixed-word closed limits. Write

$$\begin{gathered} \delta_i = Y_i - Q_i, \quad \tau_i = |Q_{i+1} - Q_i|, \quad \ell_i = |Y_{i+1} - Y_i|, \\ v_i = (Q_{i+1} - Q_i)/\tau_i, \quad \widehat{v}_i = (Y_{i+1} - Y_i)/\ell_i, \quad e_i = \widehat{v}_i - v_i, \\ F_i = \ell_i |e_i|^2, \qquad H_i = c_i |\delta_i|^2 \quad (0 < i < m), \qquad E = E_R(Q) = R(2 - c_0 - c_m). \end{gathered} \tag{45.1}$$

The reference dual certificate D_Q gives the complete positive gap $G = A(Y) - D_Q$. The inherited gap identity and disk support inequality give

$$0 \leq G \leq E \leq 2r, \qquad G = \tfrac{1}{2} \sum_i F_i + \sum_{i=0}^{m} h_i, \qquad h_i \geq H_i/R \ (0 < i < m).$$

The already proved sublevel theorem gives $\sum_{i=0}^{m} |\delta_i|^4 \leq CE$. We do not reclassify that theorem as a contribution of this section.

Lemma 45.1 (A variational inequality with only cutoff work)**.** *For any $0 \le \eta_i \le 1$ with $\eta_0 = \eta_m = 0$, put $a_i^\eta = (\eta_i^2 + \eta_{i+1}^2)/2$. Then*

$$\frac{1}{R}\sum_{i=1}^{m-1} \eta_i^2 H_i + \frac{1}{4}\sum_{i=0}^{m-1} a_i^\eta(\ell_i + \tau_i)|e_i|^2 \le \frac{1}{\tau_-}\sum_{i=0}^{m-1}(\eta_{i+1} - \eta_i)^2(|\delta_i|^2 + |\delta_{i+1}|^2). \tag{45.2}$$

In particular, no separate endpoint deficit is paid on the right.

Proof. Let $n_i = (Q_i - O_i)/R$. At an interior reference contact the action gradient is $g_i^Q = v_{i-1} - v_i = -2c_i n_i$. For every point in the same disk,

$$|\delta_i|^2 \le -2Rn_i \cdot \delta_i, \qquad g_i^Q \cdot \delta_i \ge H_i/R.$$

The one-coordinate minimum condition gives $g_i^Y \cdot (Q_i - Y_i) \ge 0$. Since $g_i^Q - g_i^Y = e_i - e_{i-1}$, it follows that $(e_i - e_{i-1}) \cdot \delta_i \ge H_i/R$. This argument applies to inactive and interior comparison vertices.

Set $b_i^\eta = \eta_{i+1}^2 - \eta_i^2$ and $\overline{\delta}_i = (\delta_i + \delta_{i+1})/2$. The exact identities

$$\eta_{i+1}^2\delta_{i+1} - \eta_i^2\delta_i = a_i^\eta(\delta_{i+1} - \delta_i) + b_i^\eta\overline{\delta}_i, \qquad e_i \cdot (\delta_{i+1} - \delta_i) = \tfrac{1}{2}(\ell_i + \tau_i)|e_i|^2$$

and finite integration by parts put the contact work and the full kinetic term on the left. Their sum is at most $-\sum_i b_i^\eta e_i \cdot \overline{\delta}_i$. When $a_i^\eta > 0$, Young's inequality bounds the absolute value of each summand by

$$\tfrac{1}{4}a_i^\eta(\ell_i + \tau_i)|e_i|^2 + \frac{(b_i^\eta)^2|\overline{\delta}_i|^2}{a_i^\eta(\ell_i + \tau_i)}.$$

For $a_i^\eta = 0$ both associated cutoff values and the work vanish. Use

$$(b_i^\eta)^2/a_i^\eta \le 4(\eta_{i+1} - \eta_i)^2, \quad |\overline{\delta}_i|^2 \le (|\delta_i|^2 + |\delta_{i+1}|^2)/2, \quad \ell_i + \tau_i \ge 2\tau_-.$$

Absorbing the first Young term proves the stated coefficient and normalization. Only the global minimum condition was used for Y; the inequality is not asserted for an arbitrary comparison. □

Lemma 45.2 (Interior coercivity without endpoint payment)**.** *For $0 \le u < v \le m$, define*

$$A_H = \frac{4r^2}{a} + 8r^2\left(1 + \frac{3N}{a} + 12N^2\right), \qquad A_F = 24r^2N^2T. \tag{45.3}$$

Then

$$\sum_{i=u+N}^{v-1} |\delta_i|^4 \le A_H \sum_{u<i<v} H_i + A_F \sum_{u\le i<v} F_i. \tag{45.4}$$

An empty sum is zero. Every anchor used to prove this estimate is strictly inside $[u, v]$. This lemma holds for every feasible comparison polygon, without a minimum assumption.

Proof. For an included contact write $b_i = v_{i-1}^\perp \cdot \delta_i$. At an anchor, $|\delta_i|^4 \le 4r^2H_i/a$. At any other included contact choose the last preceding anchor j; the complete N-flight block preceding i supplies one with $u < j < i$ and $i - j < N$. The exact transverse telescope is

$$b_i = b_j + \sum_{l=j}^{i-1} \ell_l v_l^\perp \cdot e_l + \sum_{l=j}^{i-1}(v_l - v_{l-1})^\perp \cdot \delta_l.$$

Here $|v_l - v_{l-1}| = 2c_l$ and $c_l^2 \le c_l$. Cauchy–Schwarz and the three-square inequality consequently give

$$b_i^2 \le \frac{3H_j}{a} + 3NT\sum_{l=j}^{i-1} F_l + 12N\sum_{l=j}^{i-1} H_l.$$

The same disk support inequality used above, resolved into the incoming direction and its perpendicular, implies

$$|\delta_i|^2/(2R) \le c_i|\delta_i| + |b_i|, \qquad |\delta_i|^4 \le 8r^2(H_i + b_i^2).$$

Every preceding anchor, flight, and contact occurs in at most N of the chosen intervals. Sum the anchor bounds separately, and sum the last two displays over the nonanchors. The resulting coefficients are exactly A_H, A_F. The indices range from $u+N$ to $v-1$, so no endpoint surrogate enters the telescope. □

Put $S_d = \sum_{i=d}^{m-d} |\delta_i|^4$ for integer $d \ge 0$, with $S_d = 0$ if $2d > m$.

Theorem 45.3 (A nonlinear depth recurrence for every global minimum)**.** *For integers $k \ge 0$ and $L \ge 1$,*

$$S_{k+L+N} \le C_{\mathrm{cut}} L^{-3/2}\sqrt{S_k}, \qquad C_{\mathrm{cut}} = \frac{4(rA_H + 4A_F)}{\tau_-}. \tag{45.5}$$

The same constant works for every original collision count, word, radius, and global minimum.

Proof. There is nothing to prove when the target is empty. Otherwise choose

$$\eta_i = \min\{1, (i-k)_+/L, (m-k-i)_+/L\}.$$

There are two complete transition layers of L flights. At most $4L$ vertices occur in them, each at most twice, and all have depth at least k. Therefore

$$\sum_{\eta_{i+1} \ne \eta_i} (|\delta_i|^2 + |\delta_{i+1}|^2) \le 4\sqrt{LS_k}.$$

The right side of (45.2) is thus at most $W = 4\tau_-^{-1}L^{-3/2}\sqrt{S_k}$. On $[u,v] = [k+L, m-k-L]$ the cutoff is one. Its contact sum is at most rW, and its flight sum is at most $4W$, since the retained kinetic term is at least $F_i/4$ there. The target positions lie in $[u+N, v-1]$. Lemma 45.2 proves the claimed recurrence with exactly the displayed C_{cut}. □

Theorem 45.4 (Uniform algebraic localization through inactive contacts)**.** *With*

$$K_{\mathrm{tail}} = \max\{2rC(4N)^3, (64C_{\mathrm{cut}})^2\}, \tag{45.6}$$

every global minimum satisfies

$$S_d \le \min\{CE_R(Q), K_{\mathrm{tail}}/(d+1)^3\} \qquad (d \ge 0). \tag{45.7}$$

The estimate includes an isolated central contact when $m = 2d$.

Proof. The first bound is the inherited full sublevel estimate. For $d < 4N$ it also proves the second bound by the first term in (45.6). For $d \ge 4N$ use strong induction, put $k = \lfloor d/2 \rfloor$ and $L = d - k - N$, and note that

$$L \ge 1, \qquad L(k+1) \ge (d+1)^2/16.$$

The recurrence and the induction hypothesis give

$$S_d \le \frac{C_{\mathrm{cut}}\sqrt{K_{\mathrm{tail}}}}{[L(k+1)]^{3/2}} \le \frac{64C_{\mathrm{cut}}\sqrt{K_{\mathrm{tail}}}}{(d+1)^3} \le \frac{K_{\mathrm{tail}}}{(d+1)^3}.$$

If the target is empty the same inequality is immediate. All constants were fixed before either the word or the depth was chosen. □

Corollary 45.5 (Deep contact and flight energy)**.** *For every integer $d \geq 1$,*

$$\sum_{\substack{0<i<m \\ d\leq i\leq m-d}} H_i \leq \frac{32r\sqrt{K_{\mathrm{tail}}}}{\tau_- d^3}, \tag{45.8}$$

$$\sum_{i=d}^{m-d-1} F_i \leq \frac{128\sqrt{K_{\mathrm{tail}}}}{\tau_- d^3}. \tag{45.9}$$

Proof. Use the same cutoff with $k = \lfloor d/2 \rfloor$ and $L = d - k$, without an anchor offset. Since $L(k+1) \geq d^2/4$, its work is at most $32\sqrt{K_{\mathrm{tail}}}/(\tau_- d^3)$. The contact and flight bounds follow respectively by multiplying this work by r and by 4. For a central singleton the contact is still in the unit-cutoff region, and the flight sum is empty. □

46. Transverse clearance and a uniform physical boundary layer

Longitudinal displacement of a contact need not be charged as a loss of clearance: a sufficiently narrow comparison strip confines that part of the comparison flight to its actual incident disks. This gives a quadratic obstruction budget and a summable depth schedule. We keep the original physical reference, disks, constants and notation of Section 45. Statements about arbitrary sublevel points will be distinguished from those requiring a global minimum.

For any feasible comparison polygon define

$$\beta_i = \max\{|v_i^\perp \cdot \delta_i|, |v_i^\perp \cdot \delta_{i+1}|\}, \qquad \Gamma_i(Y) = \inf_{z\in\Lambda\setminus\{O_i,O_{i+1}\}} \operatorname{dist}(z, [Y_i, Y_{i+1}]) - R. \tag{46.1}$$

Every third center is retained, including a center visited elsewhere in the same word. Put $\Gamma_i(Q)$ for the original closed flight.

Lemma 46.1 (A strip comparison with both incident disks retained)**.** *If $2\beta_i < \tau_-$, then*

$$\Gamma_i(Y) \geq \min\{\Gamma_i(Q), \tau_-\} - \beta_i. \tag{46.2}$$

The statement holds for arbitrary feasible comparison endpoints.

Proof. Use coordinates in which the original endpoints are $(0,0)$ and $(\tau_i, 0)$. Intersect each of its two incident closed disks with the strip $|y| \leq \beta_i$. The longitudinal projections of these compact convex sets are disjoint intervals: otherwise two points, one in each disk, would have the same longitudinal coordinate and distance at most $2\beta_i < \tau_-$, contradicting disk separation. The first interval precedes the second, since they contain respectively 0 and $\tau_i > 0$. Thus the comparison segment has strictly increasing longitudinal coordinate and stays in the strip.

The portion projected into $[0, \tau_i]$ is within β_i of the original closed segment. If a left extension exists, let Z be its intersection with the line of longitudinal coordinate zero. The segment from Y_i to Z is within β_i of the segment from Y_i to Q_i, using the same convex parameter. The latter segment lies in the first incident disk. The right extension is treated identically with the second disk and Q_{i+1}. Every third disk is at distance at least τ_- from each incident disk. Its distance from a point of the comparison segment is therefore at least $\Gamma_i(Q) - \beta_i$ on the central portion, and at least $\tau_- - \beta_i$ on either extension. Taking the infimum over the full segment and every third disk proves the result. The argument includes endpoints, changes of the closest center, and changes of the minimizing foot. It does not replace a closed flight by its infinite supporting line. □

Define the uniform coefficients

$$B_H = 3\left(\frac{1+3N}{a} + 12N^2\right) + 8, \qquad B_F = 9N^2T, \tag{46.3}$$

$$B_G = \frac{3r}{a} + \frac{9Nr}{a} + 18N^2T + 36N^2r + 12r, \qquad B_E = 36Nr + 4r. \tag{46.4}$$

In particular $B_G = 167591049$ and $B_E = 1592.36$.

Theorem 46.2 (Quadratic transverse budgets)**.** *For every feasible comparison polygon, with its original positive gap G, one has*

$$\sum_{i=0}^{m-1} \beta_i^2 \le B_G G + B_E E_R(Q). \tag{46.5}$$

For every $0 \le u < v \le m$ one also has

$$\sum_{i=u+N}^{v-2} \beta_i^2 \le B_H \sum_{u<i<v} H_i + B_F \sum_{u\le i<v} F_i. \tag{46.6}$$

For a global minimum, put

$$K_\perp = \frac{(256rB_H + 1024B_F)\sqrt{K_{\mathrm{tail}}}}{\tau_-}, \qquad C_\perp = \sqrt{K_\perp}. \tag{46.7}$$

Then for every integer $d \ge 2N$,

$$\sum_{i=d}^{m-d-1} \beta_i^2 \le K_\perp d^{-3}. \tag{46.8}$$

Proof. Write $b_i = v_{i-1}^\perp \cdot \delta_i$ for $1 \le i \le m$ and $b_0 = v_0^\perp \cdot \delta_0$. At an interior contact,

$$|v_i^\perp \cdot \delta_i|^2 \le 2b_i^2 + 8H_i.$$

The transverse telescope from Lemma 45.2, with anchors paid directly, gives on an interior interval

$$\sum_{i=u+N}^{v-1} b_i^2 \le \left(\frac{1+3N}{a} + 12N^2\right) \sum_{u<i<v} H_i + 3N^2T \sum_{u\le i<v} F_i.$$

Since $\beta_i^2 \le |v_i^\perp \cdot \delta_i|^2 + b_{i+1}^2$, summing gives the stated local coefficients, with all target indices contained in the preceding interval.

For completeness the two true endpoints can be paid without assuming their incidences bounded away from zero. Put $E_0 = R(1 - c_0)$ and $E_m = R(1 - c_m)$, so $E_0 + E_m = E$. At either endpoint the disk support gap and the decomposition into longitudinal and transverse components give

$$b_0^2 \le 4r(h_0 + E_0), \qquad b_m^2 \le 4r(h_m + E_m).$$

Indeed the square of the comparison transverse coordinate relative to its disk center is at most $2Rh_0$ or $2Rh_m$; the reference square is at most $2RE_0$ or $2RE_m$. The difference-square inequality supplies the factor two.

Use a preceding physical anchor for every nonanchor after the first anchor, and use endpoint 0 for the at most N initial indices. The exact telescope for these initial indices has no reflection term at 0. Its initial-value payment is at most $3Nb_0^2$. The flight and contact reuse bounds are still at most N, including these initial intervals. Because $(1+3N)/a \ge 12N$, the anchor payment

and the $12Nrh_0$ part of the initial payment can be paid together by the same complete positive gap. Using $\sum H_i \le rG$ and $\sum F_i \le 2G$ for the remaining terms therefore gives

$$\sum_{i=1}^{m-1} b_i^2 \le \left(\frac{(1+3N)r}{a} + 6N^2T + 12N^2r\right) G + 12NrE.$$

Finally

$$\sum_i \beta_i^2 \le 3\sum_{i=1}^{m-1} b_i^2 + 8\sum_{i=1}^{m-1} H_i + b_0^2 + b_m^2.$$

The endpoint pair is at most $4r(G+E)$. Substitution proves (46.5), including its exact coefficients.

For a global minimum and a nonempty deep target, apply the local bound on $[u,v] = [d-N, m-d+N]$. It contains all the target flights. The contact and flight sums are contained in the energy at depth $d-N \ge d/2$. Corollary 45.5 gives

$$\sum_{i=d}^{m-d-1} \beta_i^2 \le \frac{(32rB_H + 128B_F)\sqrt{K_{\mathrm{tail}}}}{\tau_-(d-N)^3} \le K_\perp d^{-3}.$$

An empty target needs no interval argument. □

Corollary 46.3 (A stronger geometric obstruction certificate)**.** *For any $0 < \rho \le \tau_-/2$, a third-disk violation $\Gamma_i(Y) \le 0$ at a flight with $\Gamma_i(Q) > \rho$ forces $\beta_i \ge \rho$. Consequently*

$$\#\{i : \Gamma_i(Y) \le 0,\ \Gamma_i(Q) > \rho\} \le \rho^{-2}(B_G G + B_E E). \tag{46.9}$$

On the original variational sublevel the right side is at most $\rho^{-2}(B_G + B_E)E$. This is a quadratic transverse payment, not a fourth-power Hausdorff payment.

Proof. If $2\beta_i \ge \tau_-$ the implication is immediate. Otherwise the strip lemma gives $\min(\Gamma_i(Q), \tau_-) \le \beta_i$, which proves it as well. Sum the nonnegative squares and use the global budget. The sublevel inequality is $G \le E$. □

Lemma 46.4 (An inactive deep minimum forces a true small incidence)**.** *Put*

$$C_{\mathrm{inactive}}^2 = 512\sqrt{K_{\mathrm{tail}}}/\tau_-^2. \tag{46.10}$$

If a global minimum is inactive at an interior contact of depth $d_i = \min(i, m-i) \ge 2$, then

$$c_i(Q) \le C_{\mathrm{inactive}} d_i^{-3/2}. \tag{46.11}$$

Proof. Inactivity means $\widehat{v}_{i-1} = \widehat{v}_i$. Thus $2c_i = |v_i - v_{i-1}| \le |e_i| + |e_{i-1}|$, and

$$c_i^2 \le \tfrac{1}{2}(|e_i|^2 + |e_{i-1}|^2) \le \frac{1}{2\tau_-}\sum_{l=d_i-1}^{m-d_i} F_l \le \frac{64\sqrt{K_{\mathrm{tail}}}}{\tau_-^2(d_i-1)^3} \le C_{\mathrm{inactive}}^2 d_i^{-3}.$$

The two adjacent flights are retained, including when the contact is the central singleton. No inverse comparison incidence occurs. □

Let $d_i' = \min(i, m-1-i)$ be flight depth, and put $\delta_{\mathrm{flux}} = 3/200$. Fix the explicit integer

$$D_0 = \left\lceil \max\left\{2N+2, (2C_\perp/\delta_{\mathrm{flux}})^{2/3}, (2C_{\mathrm{inactive}})^{2/3}\right\}\right\rceil. \tag{46.12}$$

For fixed R, m and $D \ge D_0$, let $\mathcal{B}_{D,R,m}$ be the original regular records whose ordered disk word admits a global minimum with an inactive contact at depth at least D, or a flight at depth at least D with nonpositive third-disk clearance. The definition uses all global minima. At fixed radius and word the existence in question is a property of the word itself. Consequently $\mathcal{B}_{D,R,m}$ is a countable union of original Borel physical word cells. No measurable choice of a minimum is needed.

Theorem 46.5 (A complete positive-source boundary-layer bound)**.** *Keep the original exact event* Π*, conditional measure* $\nu_R^* = c_*^{-1}\mathbf{1}_{Y_R^*}\nu$*, physical roof* L_R*, radius range and exact labels* n, k*. For any Borel* $0 \le h \le 1$*, define the positive original-roof measure*

$$\mu_{D,R,m,n,k,h} = (L_R)_*(\mathbf{1}_{\mathcal{B}_{D,R,m}}\mathbf{1}_\Pi h\, d\nu_R^*). \tag{46.13}$$

On every original Borel roof set its mass, and hence its total variation on the whole roof line, is at most

$$\frac{4C_\perp}{R_- c_*\sqrt{D-1}} + \frac{C_{\text{inactive}}^2}{c_*(D-1)^2}. \tag{46.14}$$

In particular

$$\lim_{D\to\infty} \sup_{R,m,n,k,\,0\le h\le 1} \|\mu_{D,R,m,n,k,h}\|_{\text{TV}} = 0. \tag{46.15}$$

Proof. The deterministic estimates hold simultaneously over all minima. A deep blocked comparison flight has $\beta_i \le C_\perp (d_i')^{-3/2} \le \delta_{\text{flux}}/2 < \tau_-/2$. The strip lemma therefore forces $\Gamma_i(Q) \le C_\perp (d_i')^{-3/2}$. The inactive-contact witness is $c_i(Q) \le C_{\text{inactive}} d_i^{-3/2} \le 1/2$. Thus the entire bad source is contained in the union of these explicit small-clearance and small-incidence events of the original physical record.

The retained exact one-copy flux gives $\nu\{\Gamma_i(Q) \le u\} \le u/R$ for $0 < u < \delta_{\text{flux}}$, by collision invariance. For the closed threshold, decrease slightly larger open widths to u; continuity of finite measures gives the same bound. The exact incidence marginal and invariance give

$$\nu\{c_i \le u\} = 1 - \sqrt{1-u^2} \le u^2 \qquad (0 \le u \le 1).$$

For any Borel roof set multiply the full finite union bound by $\mathbf{1}_\Pi h \mathbf{1}_{\{L_R \in A\}}$ and integrate with the original conditional measure. Drop factors between zero and one only in the positive upper bound. This pays exactly one c_*^{-1}, with no replacement of the event or roof.

There are at most two flights or contacts at any prescribed depth. The complete infinite tails satisfy

$$\sum_{d=D}^{\infty} d^{-3/2} \le 2(D-1)^{-1/2}, \qquad \sum_{d=D}^{\infty} d^{-3} \le [2(D-1)^2]^{-1}.$$

The resulting constants are those in (46.14). All actual words and all collision counts were included before taking this upper bound. Its constants are independent of every displayed supremum variable, proving the stated limit without interchanging the original microscopic limits. □

Corollary 46.6 (A true central comparison subrecord)**.** *If* $D \ge D_0$*,* $m > 2D$*, and the original regular record is outside* $\mathcal{B}_{D,R,m}$*, then every global minimum has the actual regular first-hit subrecord* $Y_D, \dots, Y_{m-D}$ *in the original Lorentz gas. The omitted end portions are not asserted physical.*

Proof. Every included contact is active, so the one-coordinate minimum condition places it on its original circle with the exterior specular law and strictly positive incidence. Each included flight therefore stays outside its two incident disks; strict positive clearance excludes every third disk. These facts prove genuine first-hit behavior on each successive central flight. The implications are simultaneous in the global minimum. □

The source theorem controls total mass for bounded positive weights. It does not control normalized roof-density peaks, arbitrary optical Jacobians, unbounded marked multiplicity, or singular pairing traces. Its deterministic clearance mechanism and exact source limit do not alter the original section, exponential guard or microscopic window.

47. The central physical clock and exact section observations

The geometric boundary-layer result gives actual central comparison records outside a specified positive original source. The remaining clock comparison must retain its two endpoint work terms. Dropping them would turn a variational identity into a false estimate for the physical roof.

Keep the notation and original family of Section 45. Fix integers $d \geq 1$, $2d < m$, and put $b = m - d$. In this section Y is a global disk-word minimum whose consecutive vertices $Y_d, \dots, Y_b$ form an actual regular physical subrecord, with the same consecutive centers as $Q_d, \dots, Q_b$. The complement of the source in Theorem 46.5, for an applicable $D \leq d$, supplies this physicality. The statements below need only the stated physical subrecord, not a physical continuation to Y or regularity of either discarded end portion. These are comparisons inside the original record; they do not replace the original full conditional event.

Write

$$\begin{gathered} I_d = \{d, \dots, b\}, \qquad J_d = \{d, \dots, b-1\}, \qquad m_d = m - 2d, \\ L_d(Q) = \sum_{i \in J_d} \tau_i, \quad L_d(Y) = \sum_{i \in J_d} \ell_i, \quad W_d = v_{b-1} \cdot \delta_b - v_d \cdot \delta_d, \\ \mathsf{T}_{\mathrm{clock},d} = L_d(Y) - L_d(Q) - W_d. \end{gathered} \tag{47.1}$$

Theorem 47.1 (The exact central clock with its boundary work)**.** *For every such original record and minimum,*

$$\mathsf{T}_{\mathrm{clock},d} = \frac{1}{R} \sum_{i=d+1}^{b-1} H_i + \frac{1}{2} \sum_{i=d}^{b-1} F_i \geq 0. \tag{47.2}$$

In particular, with the same K_{tail} as in Theorem 45.4,

$$0 \leq \mathsf{T}_{\mathrm{clock},d} \leq K_{\mathrm{clock}} d^{-3}, \qquad K_{\mathrm{clock}} = \left(\frac{32r}{R_-} + 64 \right) \frac{\sqrt{K_{\mathrm{tail}}}}{\tau_-}. \tag{47.3}$$

The length on either side is the unit-speed physical time of that central subrecord. No clock is averaged or rescaled.

Proof. For each included flight, the identity between two unit directions is

$$\ell_i - \tau_i = v_i \cdot (\delta_{i+1} - \delta_i) + \tfrac{1}{2} \ell_i |e_i|^2.$$

Sum it over $d \leq i < b$. Finite integration by parts gives boundary term W_d and interior coefficients $v_{i-1} - v_i = -2c_i n_i$. Both actual contact points Q_i, Y_i lie on the same circle, so

$$|Q_i - O_i + \delta_i|^2 = R^2 \quad \Longrightarrow \quad 2R n_i \cdot \delta_i + |\delta_i|^2 = 0.$$

Thus each interior coefficient contributes H_i/R with a positive sign. This proves the exact identity, including the case of one central flight, when its contact sum is empty. Apply the contact and flight estimates of Corollary 45.5, and enlarge R^{-1} to R_-^{-1}, to obtain the displayed constant. This argument uses circular equality only at the included physical contacts. The disk support inequality for a nonphysical interior vertex would not justify the same clock equality. □

Let $x_i = (\alpha_i, p_i)$ and $\widehat{x}_i$ be the original and comparison collision states on I_d. Use outgoing tangential velocity at $i < b$ and the incoming tangential projection at b. Specular reflection makes that terminal projection the actual collision coordinate without requiring a next flight. The distance d_∞ is the maximum of the shortest circular angle distance and the momentum distance. Define

$$\mathsf{T}_{\mathrm{phase},d} = \sum_{i=d}^{b} d_\infty(x_i, \widehat{x}_i)^4, \qquad K_{\mathrm{phase}} = \frac{8192\sqrt{K_{\mathrm{tail}}}}{\tau_-^2} + \frac{24 K_{\mathrm{tail}}}{R_-^4}. \tag{47.4}$$

Theorem 47.2 (A complete central phase and section payment)**.** *One has*

$$\mathsf{T}_{\mathrm{phase},d} \le K_{\mathrm{phase}} d^{-3}. \tag{47.5}$$

Let $\eta_R = \mathbf{1}_{Y_R^*}$ *be exactly the original section indicator. For* $\rho > 0$ *put*

$$B_{\rho,d}(Q) = \sum_{i=d}^{b} \mathbf{1}_{\{\mathrm{dist}_\infty(x_i,\partial Y_R^*)\le\rho\}}, \qquad N_{\mathrm{cut},d}(Q,Y) = \sum_{i=d}^{b} |\eta_R(x_i) - \eta_R(\widehat{x}_i)|.$$

Then, for the actual observations on both subrecords,

$$N_{\mathrm{cut},d} \le B_{\rho,d} + \rho^{-4}\mathsf{T}_{\mathrm{phase},d} \le B_{\rho,d} + \rho^{-4} K_{\mathrm{phase}} d^{-3}. \tag{47.6}$$

Occupation uses only J_d*; the last section test at* b *is separate. In particular the absolute occupation difference is bounded by the sum in* (47.6) *with its last summand removed.*

Proof. The circle estimate in Theorem 44.2 applies contact by contact:

$$d_\infty(x_i,\widehat{x}_i)^4 \le 8|e_{\iota(i)}|^4 + 24R_-^{-4}|\delta_i|^4, \qquad \iota(i) = \begin{cases} i, & i < b, \\ b-1, & i = b. \end{cases}$$

Indeed angle distance is at most $2|\delta_i|/R_-$ and tangential momentum distance at most $|e_{\iota(i)}| + |\delta_i|/R_-$. The last edge occurs twice, including when the subrecord has only one edge. All other included edges occur once. Since $|e_i| \le 2$ and $\ell_i \ge \tau_-$, the sum of their direction contributions is at most

$$8\sum_{i\in I_d} |e_{\iota(i)}|^4 \le \frac{64}{\tau_-}\sum_{i=d}^{b-1} F_i.$$

Use Corollary 45.5 for this sum and Theorem 45.4 for the position sum. Enlarging $(d+1)^{-3}$ to d^{-3} gives exactly K_{phase}.

A shortest product-coordinate path between states with different section indicators meets the original section boundary. Therefore a discrepancy is either in its closed ρ collar or has $d_\infty(x_i,\widehat{x}_i) > \rho$. Its indicator in the latter case is at most $\rho^{-4} d_\infty(x_i,\widehat{x}_i)^4$. Sum over every included contact. Boundary states have been counted explicitly; no null-set deletion of their weighted source is used. Dropping the last nonnegative summand gives the occupation claim. □

Proposition 47.3 (All central Fourier marks with the exact clock phase)**.** *The central collision count* m_d *and lattice displacement* $k_d = O_b - O_d$*, expressed in the original lattice basis, are identical on the two physical subrecords. Put* $A_d(Q) = \sum_{i=d}^{b-1} \eta_R(x_i)$ *and define* $A_d(Y)$ *in the same way. For every* $\xi \in \mathbb{R}^2$ *and* $s, \zeta \in \mathbb{R}$*,*

$$\begin{aligned} &\left| e^{i[\xi\cdot k_d + sL_d(Y) + \zeta A_d(Y)]} - e^{isW_d} e^{i[\xi\cdot k_d + sL_d(Q) + \zeta A_d(Q)]} \right| \\ &\qquad \le |s|\mathsf{T}_{\mathrm{clock},d} + 2|\sin(\zeta/2)|\, N_{\mathrm{cut},d}. \end{aligned} \tag{47.7}$$

If the two central endpoint section tests are inserted, their additional error is at most the sum of those two actual indicator discrepancies. All these comparisons retain every central collision.

Proof. The ordered centers give k_d and m_d exactly. Substitute (47.2) into the left side, retaining the phase e^{isW_d} of modulus one. The clock error is bounded by $|e^{is\mathsf{T}_{\mathrm{clock},d}} - 1| \le |s|\mathsf{T}_{\mathrm{clock},d}$. For an integer q, telescoping q unit increments gives $|e^{i\zeta q} - 1| \le 2|q||\sin(\zeta/2)|$; it also holds for negative q. Take $q = A_d(Y) - A_d(Q)$ and use the preceding theorem. The product of two indicators changes by at most the sum of their two changes, which proves the endpoint assertion. □

Remark 47.4. The full source event is still $\Pi_{n,k,m,R}$ and its roof still $L_{m,R}(Q)$. Neither the central occupation nor the phase-corrected central length is substituted into that event. The end portions, the section collar and the physicality-exception source remain present in the full problem until their own required estimates have been proved. In particular $\mathsf{T}_{\mathrm{clock},d}$ alone cannot bound $|L_d(Y) - L_d(Q)|$: the term W_d is indispensable.

48. Exponential confinement by physical anchor layers

The polynomial estimates above do not exhaust the variational mechanism. The physical turning bound supplies contacts at which the endpoint work of a whole comparison interval can be paid by energy in its removed boundary layer. This yields a strict contraction after a fixed number of actual collisions. The resulting exponential estimate concerns every global disk-product minimum, including nonphysical and inactive ones. It is not obtained by integrating a local endpoint derivative along an assumed physical continuation.

Keep the original reference, comparison, radius and constants of Sections 45 and 46. In particular Y is a global minimum of the unchanged disk-word problem, and the reference is actual regular or a fixed-word closed physical reference. To distinguish it from the depth of an index, denote the anchor threshold by $a_0 = 1/416640$. Thus $N = 94$, $r = 47/100$, $\tau_- = 3/50$, and

$$\kappa = \frac{r}{a_0\tau_-} = 3263680, \qquad q_{\mathrm{a}} = \frac{\kappa}{1+\kappa} = \frac{3263680}{3263681} \in (0,1). \tag{48.1}$$

The symbol q_{a} is a deterministic contraction constant; it does not replace the original guard constant $q_0 = 47/53$.

Lemma 48.1 (Common active geometry of global minima)**.** *For a fixed word and radius, all global disk-product minima have the same ordered unit edge directions and the same active index set. At every active index their vertices agree exactly. Both endpoints are active. Any difference between minima lies along the common straight direction in a chain of inactive contacts. This is a convexity fact about the comparison problem, not a new billiard map or a claim that its full minimum is unique.*

Proof. Let Y, Z be two minima. Their midpoint lies in the same product of disks and, by convexity of A, is another minimum. The sum of the nonnegative defects in the flightwise triangle inequalities is zero, so each such inequality is an equality. Both vectors of each edge are nonzero, by the fixed consecutive-disk separation. Equality in the Euclidean triangle inequality therefore makes them positive parallel vectors, and their unit directions agree.

The Cartesian gradients at every coordinate consequently agree for all minima. If this common gradient g_i is nonzero, the minimum condition on its disk gives the unique point $O_i - Rg_i/|g_i|$. If it is zero, its neighboring directions agree; this index is inactive for every minimum. At an endpoint the gradient is a unit vector up to sign, so it is always active. On a consecutive chain of inactive indices the edge direction is constant, and the differences between corresponding edges of Y and Z are multiples of that direction. Starting at the preceding common active vertex and summing these differences shows that all possible vertex differences on the chain are longitudinal. No continuation through an inactive vertex or through an unlisted disk is inferred. □

Lemma 48.2 (The exact work of an anchor-bounded interval)**.** *If $1 \le u < v \le m-1$ and both reference contacts are anchors, $c_u, c_v \ge a_0$, then*

$$\frac{1}{R}\sum_{i=u+1}^{v-1} H_i + \frac{1}{2}\sum_{i=u}^{v-1} F_i \le \frac{H_u + H_v}{a_0\tau_-}. \tag{48.2}$$

Both flights adjacent to the cut are retained on the left. In particular the inequality also covers $v = u+1$.

Proof. The variational comparison from Lemma 45.1 gives $(e_i - e_{i-1}) \cdot \delta_i \geq H_i/R$ for $u < i < v$. Summing on this hard interval, with its endpoint work retained, yields

$$\frac{1}{R} \sum_{i=u+1}^{v-1} H_i + \frac{1}{2} \sum_{i=u}^{v-1} (\ell_i + \tau_i)|e_i|^2 \leq e_{v-1} \cdot \delta_v - e_u \cdot \delta_u. \tag{48.3}$$

Indeed the boundary of the finite integration by parts is exactly the right side, and $e_i \cdot (\delta_{i+1} - \delta_i) = (\ell_i + \tau_i)|e_i|^2/2$. This identity is still valid when there are no interior contacts.

At either cut use the elementary inequality

$$|e|\,|\delta| \leq \frac{\tau_-}{4}|e|^2 + \frac{|\delta|^2}{\tau_-}.$$

An edge incident to only one cut loses at most $\tau_-|e_i|^2/4$ from its kinetic coefficient; its remaining coefficient is at least $\ell_i/2$. If $v = u + 1$, the same edge pays both cuts, but

$$\frac{\ell_i + \tau_i}{2} - \frac{\tau_-}{2} \geq \frac{\ell_i}{2}.$$

Thus every edge retains at least $F_i/2$, including the two boundary flights. At the actual reference anchors, $|\delta_u|^2 \leq H_u/a_0$ and $|\delta_v|^2 \leq H_v/a_0$. These facts prove (48.2). No assertion about the comparison incidences is used. □

For an integer $d \geq 0$ define the full relative energy at depth d by

$$\mathcal{E}_d(Q, Y) = \frac{1}{R} \sum_{\substack{0<i<m \\ d \leq i \leq m-d}} H_i + \frac{1}{2} \sum_{i=d}^{m-d-1} F_i. \tag{48.4}$$

Every sum over an empty interval is zero. In particular a central single contact is kept when $m = 2d$, although no flight remains. The actual endpoints are never assigned interior contact weights.

Theorem 48.3 (A contraction on every physical anchor layer)**.** *For every integer $d \geq 0$,*

$$\mathcal{E}_{d+N} \leq q_{\mathrm{a}} \mathcal{E}_d, \qquad \mathcal{E}_d \leq E_R(Q) q_{\mathrm{a}}^{\lfloor d/N \rfloor}. \tag{48.5}$$

The inequalities hold simultaneously for every global minimum and every original word and radius, including at a closed physical reference with zero internal incidences. They have no collision-count prefactor and no inverse individual incidence.

Proof. The complete positive gap, before any separate estimate of its summands, gives

$$\mathcal{E}_0 = \frac{1}{R} \sum_{i=1}^{m-1} H_i + \frac{1}{2} \sum_{i=0}^{m-1} F_i \leq \sum_{i=1}^{m-1} h_i + \frac{1}{2} \sum_{i=0}^{m-1} F_i \leq G \leq E_R(Q). \tag{48.6}$$

Thus both kinds of energy consume the same positive budget, not two copies of it. The last inequality uses global minimality only through $Y \in \mathcal{S}(Q)$.

If $2(d + N) > m$, the first inequality in (48.5) has zero left side. Otherwise the two complete physical N-flight blocks starting at d and $m - d - N$ supply reference anchors

$$u \in \{d + 1, \ldots, d + N - 1\}, \qquad v \in \{m - d - N + 1, \ldots, m - d - 1\}.$$

These are interior contacts, and $u < d+N \leq m-d-N < v$. Consequently every energy summand in $\mathcal{E}_{d+N}$ occurs on the left of (48.2). Conversely, the two anchor summands $H_u/R, H_v/R$ occur in $\mathcal{E}_d - \mathcal{E}_{d+N}$. All the other summands in that difference are nonnegative. Lemma 48.2 gives

$$\mathcal{E}_{d+N} \leq \frac{H_u + H_v}{a_0 \tau_-} = \frac{R}{a_0 \tau_-} \frac{H_u + H_v}{R} \leq \kappa(\mathcal{E}_d - \mathcal{E}_{d+N}).$$

Rearranging proves the strict contraction with exactly $q_{\mathrm{a}} = \kappa/(1+\kappa)$. The argument also covers $m = 2(d+N)$, where the inner energy contains one contact. No boundary flight is counted as a removed-layer payment: its Young payment has already been absorbed in (48.2).

Iterate the contraction at the depths $0, N, 2N, \dots$ and use (48.6). Since $\mathcal{E}_d$ is nonincreasing, $\mathcal{E}_d \le \mathcal{E}_{N\lfloor d/N\rfloor}$. This proves the second inequality. The anchors can be chosen anew at each layer; no compatible continuation or fixed selection of anchor points is required. □

Put

$$j(d) = \max\{0, \lfloor d/N\rfloor - 1\}, \qquad B_\perp = B_G + B_E. \tag{48.7}$$

Corollary 48.4 (Exponential position, direction and transverse budgets)**.** *For every integer $d \ge 0$ and every global minimum,*

$$\sum_{i=d}^{m-d} |\delta_i|^4 \le C E_R(Q) q_{\mathrm{a}}^{j(d)}, \tag{48.8}$$

$$\sum_{i=d}^{m-d-1} F_i \le 2E_R(Q) q_{\mathrm{a}}^{\lfloor d/N\rfloor}, \tag{48.9}$$

$$\sum_{i=d}^{m-d-1} \beta_i^2 \le B_\perp E_R(Q) q_{\mathrm{a}}^{j(d)}. \tag{48.10}$$

The fourth-power and transverse sums have their usual zero values on empty intervals. The position estimate includes both actual endpoints when $d = 0$; they are paid by the original deficit.

Proof. The flight estimate follows directly from the definition of $\mathcal{E}_d$. For $d < N$, the position and transverse estimates are respectively Theorem 43.3 and (46.5) on the sublevel.

For $d \ge N$ with a nonempty target interval, apply the interior coercivity lemma on $[u, v] = [d-N, m-d+N]$. Its target range contains all the positions in (48.8). The positive contact and flight components are jointly bounded by $\mathcal{E}_{d-N}$, so

$$S_d \le \max\{rA_H, 2A_F\}\mathcal{E}_{d-N} \le C\mathcal{E}_{d-N}.$$

For the constants in (45.3), the last comparison is the explicit rational inequality

$$\max\{rA_H, 2A_F\} < 10^8 = C.$$

Similarly the local transverse estimate (46.6) contains all the included flights and gives

$$\sum_{i=d}^{m-d-1} \beta_i^2 \le \max\{rB_H, 2B_F\}\mathcal{E}_{d-N} \le B_\perp \mathcal{E}_{d-N},$$

where $\max\{rB_H, 2B_F\} < B_G + B_E$ is another direct rational comparison of the displayed constants. Both inequalities use the shared energy before estimating its components. Since $\lfloor (d-N)/N\rfloor = j(d)$ for $d \ge N$, the layer theorem proves the two conclusions. The polynomial bounds already proved remain valid and may be used together with these exponential ones; no preceding proof body is removed. □

Corollary 48.5 (Explicit exponentially small physical failure witnesses)**.** *For an interior contact of depth $d_i \ge 1$, inactivity of a global minimum implies*

$$c_i(Q)^2 \le \frac{E_R(Q)}{\tau_-} q_{\mathrm{a}}^{\lfloor (d_i-1)/N\rfloor}. \tag{48.11}$$

At every flight,

$$\beta_i^2 \le B_\perp E_R(Q) q_{\mathrm{a}}^{j(d_i')}. \tag{48.12}$$

If the last upper bound is strictly below $\tau_-^2/4$, a third-disk violation of that comparison flight necessarily implies $\Gamma_i(Q) \le \sqrt{B_\perp E_R(Q)}\, q_{\mathrm{a}}^{j(d_i')/2}$. All these witnesses concern the true incidence and clearance of the original reference. No probability or roof-density assertion follows without a separate source argument.

Proof. Inactivity gives $2c_i \le |e_i| + |e_{i-1}|$, as before. Both flights belong to the energy at depth $d_i - 1$, so

$$c_i^2 \le \frac{|e_i|^2 + |e_{i-1}|^2}{2} \le \frac{1}{2\tau_-} \sum_{l=d_i-1}^{m-d_i} F_l \le \frac{E_R(Q)}{\tau_-} q_{\mathrm{a}}^{\lfloor (d_i-1)/N \rfloor}.$$

The transverse point bound is a single summand of (48.10). Its stated strict regime permits application of the already proved strip lemma, giving the clearance implication. This excludes no original collision, and treats every minimizer with the same constants. □

Remark 48.6 (Rate, original guard, and probability topology). The contraction q_{a} is close to one because the explicit physical anchor bound is conservative. It is nevertheless fixed before any collision limit, and its exponential decay is uniform on the entire original geometric family. It does not identify these rates with the much narrower original guard widths $\varepsilon q_0^{d/4}$. In particular a substitution of one depth schedule for the other would change the claimed microscopic problem and is not made. Exponential geometric cost, exact original-source mass, inherited-pressure conditional roof height, and the entire normalized shortage remain distinct assertions. None is inferred from a finite trajectory computation.

49. Comparison costs on the original roof and their normalization

A deterministic small comparison cost and the density of the exceptional physical source are different quantities. Here positive ordering pays the former on the unchanged physical roof. The unnormalized roof-height input is stated explicitly; none of the geometric theorems just proved depends on it.

Fix R, m, n, k and $d \ge 2N$, $2d < m$. On any Borel class $\mathcal{P}_d$ of original records with a physical central minimum as in Section 47, choose one such global minimum per physical disk word. At fixed radius there are only countably many finite words. A chosen minimum is constant on its word, so this choice gives Borel comparison coordinates on their physical domains. Alternatively, by Lemma 48.1, the active central vertices of all such minima coincide. No joint measurability in the radius is needed for the original order $\sup_R \operatorname{ess\,sup}_t$.

Let $0 \le h \le 1$ be any Borel physical source selector; it may depend on the whole record and on the fixed labels. With Π exactly as in (1.1), define the positive measures

$$\mathcal{M}_{h,d}[f] = (L_{m,R})_*(\mathbf{1}_{\mathcal{P}_d} \mathbf{1}_\Pi h f \, d\nu_R^*), \qquad \mu_0 = (L_{m,R})_*(\mathbf{1}_\Pi \, d\nu_R^*). \tag{49.1}$$

Every pushforward is by the original full physical roof, even when f is a central comparison cost. Let $\mathsf{T}_{\perp,d} = \sum_{i=d}^{m-d-1} \beta_i^2$.

Theorem 49.1 (Positive cost domination and its collision normalization)**.** *For every Borel roof set $I \subset \mathbb{R}$ and $f = \mathsf{T}_{\mathrm{clock},d}, \mathsf{T}_{\mathrm{phase},d}, \mathsf{T}_{\perp,d}$, respectively,*

$$\mathcal{M}_{h,d}[f](I) \le K_f d^{-3} \mu_0(I), \qquad (K_f) = (K_{\mathrm{clock}}, K_{\mathrm{phase}}, K_\perp). \tag{49.2}$$

This measure assertion is unconditional on the pressure input.

Under precisely Input 11.1, including its physical stable arclength and optical-coordinate conventions, these measures have densities on the whole original roof line, and

$$\left\| \frac{d\mathcal{M}_{h,d}[f]}{dt} \right\|_\infty \le C_\star K_f d^{-3}. \tag{49.3}$$

In particular, for every fixed $0 < \theta < 1/2$ *and* $d_m = \lceil \theta m \rceil$*, for all sufficiently large* m*,*

$$\sup_{R,n,k,h} m^2 \left\| \frac{d\mathcal{M}_{h,d_m}[f]}{dt} \right\|_\infty \le C_\star K_f \theta^{-3} m^{-1} \longrightarrow 0. \tag{49.4}$$

The same bound holds on each original microscopic roof interval and on the original central target. The norm on the whole line is the stronger assertion. The input constants are uniform in all the displayed variables.

Proof. Theorems 47.1, 47.2, and 46.2 give $0 \le f \le K_f d^{-3}$ pointwise on the stated source class. Multiply by $\mathbf{1}_{\mathcal{P}_d} \mathbf{1}_\Pi h\, \mathbf{1}_I(L_{m,R})$ and integrate against exactly ν_R^*. Removing only the upper-bound selectors $\mathbf{1}_{\mathcal{P}_d} h \le 1$ gives (49.2). Equivalently the same proof works against every nonnegative Borel test of the original roof. Tonelli permits the complete positive physical word sum before this comparison; there is no sum of a separate error constant for each word.

For $S = \varnothing$, Theorem 11.3, under its named input, gives $d\mu_0/dt \le C_\star$ almost everywhere, where $C_\star = C_0 Q_*/(2c_*\lambda_\circ)$. The order property of positive measures first gives absolute continuity of $\mathcal{M}_{h,d}[f]$ and then the bound on its Radon–Nikodym derivative. This uses the original inverse optical Jacobian and section factor already in $C_\star$; it inserts no new Jacobian and pays c_*^{-1} only once. Finally $d_m \ge \theta m$ and $m^2/d_m^3 \le \theta^{-3}/m$. All constants were fixed before the collision limsup. Restricting a density norm to a roof interval cannot enlarge it. □

Corollary 49.2 (The dominated persistent low-load interior source)**.** *For* $\gamma \ge 1$*, the same positive-cost and conditional normalized estimates apply to every Borel restriction of the inherited persistent low-load interior coefficient which is dominated by the original uncut roof measure as in Lemma 12.1. This includes its original low-load selectors, all bounded by one. It does not assert the corresponding domination for singular paired endpoint traces or for an arbitrary full endpoint coefficient.*

Proof. The inherited inequality is $0 \le c_{\mathrm{per}}^{\varepsilon,\gamma} \le f^\varepsilon(u,\vartheta) \le A(u,\vartheta)$ on each actual physical roof segment. Multiplication by its nonnegative low-load selector and by f preserves the inequality. Sum the actual source coefficients at their original roof and use the positive original-measure comparison of that lemma. The result is a measure dominated by the right side of (49.2); the preceding proof applies verbatim. This argument invokes that specific domination theorem and its existing physical multiplicities, not an assertion that a boundary trace has a bulk density. □

Corollary 49.3 (Central observation costs with a stated frequency range)**.** *On the same source, after retaining* e^{isW_d} *in* (47.7)*, the nonnegative error majorant is*

$$|s|\mathsf{T}_{\mathrm{clock},d} + 2|\sin(\zeta/2)|(B_{\rho,d} + \rho^{-4}\mathsf{T}_{\mathrm{phase},d}).$$

For fixed $\rho > 0$*, the two comparison-cost terms have collision-normalized original-roof height tending to zero when* $d = d_m$ *and* $|s| = o(m)$*, under the stated pressure input, uniformly in the two lattice twists and the return twist. The positive source weighted by* $B_{\rho,d}$ *remains an explicit separate term.*

Proof. Combine the exact twist estimate with the section payment and Theorem 49.1. The lattice phase has modulus one, and $2|\sin(\zeta/2)| \le 2$ for all ζ. The clock coefficient cost is bounded after multiplication by m^2 by a constant times $|s|/m$. At fixed ρ the other comparison term is $O(m^{-1})$. No estimate of $\mathcal{M}_{h,d}[B_{\rho,d}]$ was proved by this argument, so it remains on its own side of the inequality. □

Remark 49.4 (The exact scope of the normalization). The estimates above are for specified small geometric costs on a proved physical central comparison class. They do not pay the indicator of its complementary source, the original shallow end portions, the singular one-sided or paired

boundary traces, a physical multiplier not dominated by the declared source, or a shortage of roofwise comparison capacity. Nor does an estimate uniform on an increasing bounded frequency range by itself justify the full unbounded Fourier integral. In particular the measure contraction in Theorem 46.5 alone is not an m^2L^∞ source estimate. All these distinctions persist when the sharper exponential costs below replace the algebraic bounds.

50. An exponentially small source for exact central physical records

The anchor contraction gives a uniform positive-source theorem without a word-count prefactor. It controls all global minima simultaneously and fixes the original section observations on their physical central records. The source is defined entirely by the actual reference trajectory. No exceptional event is inserted into the original guard or into the original conditional return event.

Keep $R \in [R_-, r]$, $N = 94$, $\tau_- = 3/50$, q_{a} and $j(d)$ from (48.1) and (48.7). All probabilities in this section use $\nu = d\alpha\, dp/(4\pi)$ or its unchanged section conditioning $\nu_R^* = c_*^{-1}\nu|_{Y_R^*}$. The original record is regular for its m flights; the estimates do not define a singular flow. Bounded bulk sources, rather than singular paired traces, are under consideration.

Lemma 50.1 (The exact original section has a uniform collar bound)**.** *For $0 < \rho \le h_0 = 1/200$,*

$$\sup_R \nu\{\mathrm{dist}_\infty(x, \partial Y_R^*) \le \rho\} \le \frac{\rho}{2\pi}. \tag{50.1}$$

The bound refers to the five original rectangles in the collision circle–momentum coordinates.

Proof. For a rectangle of width w and height h, with $2\rho \le \min(w, h)$, its closed boundary collar has area at most

$$(w + 2\rho)(h + 2\rho) - (w - 2\rho)(h - 2\rho) = 4\rho(w + h).$$

On the circle this can be calculated on a lift; overlap at the coordinate seam can only reduce the projected area. Intersection with $-1 < p < 1$ also cannot increase it. The boundary of a finite union is contained in the union of the component boundaries. The four squares have $w = h = 1/100$, and U has $w = 3/25$ and $h = 3/10$. Therefore $\sum(w + h) = 4(2/100) + 3/25 + 3/10 = 1/2$. The total area is at most 2ρ, and division by 4π proves the claim. The radius changes the centers of two rectangles but not these widths. □

Put

$$\begin{gathered} K_\phi = \frac{128}{\tau_-} + \frac{24C}{R_-^4}, \\ A_\Gamma = (2rB_\perp)^{1/2}, \qquad A_c = (2r/\tau_-)^{1/2}, \qquad A_\phi = (2rK_\phi)^{1/4}. \end{gathered} \tag{50.2}$$

Lemma 50.2 (A phase estimate at every active comparison contact)**.** *Let i be an interior contact of depth $d_i = \min(i, m - i) \ge 1$, active in a global minimum. Give that contact its circle coordinate and the outgoing tangential component determined by its two adjacent comparison directions. Then*

$$d_\infty(x_i, \widehat{x}_i)^4 \le K_\phi E_R(Q) q_{\mathrm{a}}^{j(d_i)} \le A_\phi^4 q_{\mathrm{a}}^{j(d_i)}. \tag{50.3}$$

This is a geometric collision-state estimate before the comparison flights have been proved clear. Inactivity is not assigned an artificial collision state.

Proof. At an active minimum its nonzero Cartesian gradient is a negative multiple of the outward disk normal. Its two unit directions therefore obey the specular reflection law with strictly positive normal incidence. In particular their incoming and outgoing tangential components agree. The elementary circle and momentum argument used in Theorem 47.2, now using the incoming edge, gives

$$d_\infty(x_i, \widehat{x}_i)^4 \le 8|e_{i-1}|^4 + 24R_-^{-4}|\delta_i|^4.$$

That incoming edge has depth at least $d_i - 1$. Since $|e_{i-1}|^4 \le 4|e_{i-1}|^2 \le 4F_{i-1}/\tau_-$, the flight estimate (48.9) gives

$$|e_{i-1}|^4 \le \frac{8E_R(Q)}{\tau_-} q_{\mathrm{a}}^{\lfloor (d_i-1)/N \rfloor}.$$

Also (48.8) bounds $|\delta_i|^4$ by $CE_R(Q)q_{\mathrm{a}}^{j(d_i)}$. For every integer $d_i \ge 1$, $\lfloor (d_i-1)/N \rfloor \ge j(d_i)$. The resulting direction coefficient $64/\tau_-$ is at most the deliberately enlarged $128/\tau_-$ in K_ϕ. This proves the first inequality; use $E_R(Q) \le 2r$ for the second. The argument retains the incoming edge even when the target consists of one central contact. □

Let $\delta_{\mathrm{flux}} = 3/200$. Define the finite constants

$$\begin{aligned} J_{\mathrm{obs}} = 1 + \Bigg\lceil \frac{1}{-\log q_{\mathrm{a}}} \max\Big\{ 0,\ 2\log(2A_\Gamma/\delta_{\mathrm{flux}}), \\ 2\log(2A_c),\ 4\log(A_\phi/h_0) \Big\} \Bigg\rceil, \qquad D_{\mathrm{obs}} = NJ_{\mathrm{obs}}. \end{aligned} \tag{50.4}$$

For $D \ge D_{\mathrm{obs}}$, the following inequalities hold:

$$A_\Gamma q_{\mathrm{a}}^{j(D)/2} \le \delta_{\mathrm{flux}}/2 < \tau_-/2, \quad A_c q_{\mathrm{a}}^{j(D)/2} \le 1/2, \quad A_\phi q_{\mathrm{a}}^{j(D)/4} \le h_0. \tag{50.5}$$

These are conservative constants fixed before every collision limit. They do not modify the original guard widths $\varepsilon q_0^{d/4}$, where $q_0 = 47/53$.

Define $\mathcal{E}^{\mathrm{obs}}_{D,R,m}$ to be the following union of events on the original record:

$$\begin{aligned} \mathcal{E}^{\mathrm{obs}}_{D,R,m} = & \bigcup_{\substack{0 \le i < m \\ d_i' \ge D}} \{\Gamma_i(Q) \le A_\Gamma q_{\mathrm{a}}^{j(d_i')/2}\} \\ & \cup \bigcup_{\substack{0 < i < m \\ d_i \ge D}} \{c_i(Q) \le A_c q_{\mathrm{a}}^{j(d_i)/2}\} \\ & \cup \bigcup_{\substack{0 < i < m \\ d_i \ge D}} \{\mathrm{dist}_\infty(x_i, \partial Y_R^*) \le A_\phi q_{\mathrm{a}}^{j(d_i)/4}\}. \end{aligned} \tag{50.6}$$

Every quantity in this finite union is an actual original observable. In particular the clearance is the infimum over all third lattice centers, including centers visited elsewhere in the word. The source is Borel. For $0 \le h \le 1$ put

$$\mu^{\mathrm{obs}}_{D,R,m,n,k,h} = (L_{m,R})_*(\mathbf{1}_{\mathcal{E}^{\mathrm{obs}}_{D,R,m}} \mathbf{1}_{\Pi_{n,k,m,R}} h \, d\nu_R^*).$$

Theorem 50.3 (A uniform exact-record source and its full positive mass)**.** *Define*

$$\epsilon_D = \frac{2NA_\Gamma}{R_- c_*} \frac{q_{\mathrm{a}}^{j(D)/2}}{1 - q_{\mathrm{a}}^{1/2}} + \frac{2NA_c^2}{c_*} \frac{q_{\mathrm{a}}^{j(D)}}{1 - q_{\mathrm{a}}} + \frac{NA_\phi}{\pi c_*} \frac{q_{\mathrm{a}}^{j(D)/4}}{1 - q_{\mathrm{a}}^{1/4}}. \tag{50.7}$$

For every $D \ge D_{\mathrm{obs}}$, every m, R, n, k, every Borel $0 \le h \le 1$ and every Borel roof set I,

$$0 \le \mu^{\mathrm{obs}}_{D,R,m,n,k,h}(I) \le \epsilon_D, \qquad \sup_{m,R,n,k,h} \|\mu^{\mathrm{obs}}_{D,R,m,n,k,h}\|_{\mathrm{TV}} \le \epsilon_D \longrightarrow 0 \quad (D \longrightarrow \infty). \tag{50.8}$$

The complete physical word sum is already included in these inequalities. No word-count factor or independence assumption occurs.

If $Q \notin \mathcal{E}^{\mathrm{obs}}_{D,R,m}$ and $m > 2D$, then every *global minimum has an actual regular physical central record $Y_D, \dots, Y_{m-D}$ and*

$$\eta_R(\widehat{x}_i) = \eta_R(x_i) \qquad (D \le i \le m - D). \tag{50.9}$$

All those central contacts and states agree between global minima. Their central displacement, number of flights, occupation, and two endpoint section tests agree exactly with the original central record. The terminal indicator in (50.9) *is tested separately from its occupation count.*

Proof. The exact one-flight physical clearance flux gives, for $0 < u < \delta_{\mathrm{flux}}$,

$$\nu\{\Gamma_0(Q) \le u\} \le \int N_u \, d\nu = \frac{u}{R} \le \frac{u}{R_-}.$$

At equality boundaries, use the same estimate at $u + \varepsilon$ and pass down by continuity of a finite measure. Invariance of the actual collision map gives the same bound at each flight time. The original incidence distribution is exact:

$$\nu\{c_i \le u\} = 1 - \sqrt{1 - u^2} \le u^2 \qquad (0 \le u \le 1).$$

Lemma 50.1 and invariance likewise give the collar bound $\rho/(2\pi)$ at any contact time. The hypotheses of all three bounds hold at every included depth by (50.5). No assertion about independence of different times is used.

At most two contacts or flights have a specified depth d. For any $a > 0$ the elementary block sum gives

$$\sum_{d=D}^{\infty} q_{\mathrm{a}}^{aj(d)} \le N \, \frac{q_{\mathrm{a}}^{aj(D)}}{1 - q_{\mathrm{a}}^{a}} \qquad (D \ge D_{\mathrm{obs}}). \tag{50.10}$$

Indeed $j(d)$ is constant on blocks of N consecutive depths; the first, possibly partial, block is enlarged to a whole block. Apply the union bound to (50.6), the preceding three one-time estimates, and (50.10) with $a = 1/2, 1, 1/4$. Restriction to the original initial section costs exactly c_*^{-1}. Removing $\mathbf{1}_\Pi h \le 1$ only on the upper side then gives (50.7). The positive pushforward has total variation equal to its total mass; the roof test can only decrease it. Countable physical word cells form a partition of the regular source, so this is already their complete positive sum. The geometric series tends to zero uniformly in all the displayed variables, which proves (50.8).

Now fix an original record outside the source and any global minimum. At every included contact the original incidence is strictly larger than $A_c q_{\mathrm{a}}^{j(d_i)/2}$. Corollary 48.5, $E_R \le 2r$, and $\lfloor (d_i - 1)/N \rfloor \ge j(d_i)$ show that inactivity is impossible. At each included flight,

$$\beta_i \le A_\Gamma q_{\mathrm{a}}^{j(d_i')/2} < \min\{\Gamma_i(Q), \tau_-/2\}.$$

The strip lemma therefore gives strictly positive clearance from every third disk, with both incident disks retained.

At every active vertex the Cartesian minimum condition gives the specular law with positive incidence, as in the preceding lemma. An outward edge from a circle and an inward edge ending at the next circle stay outside their respective incident disks until the prescribed contacts, by convexity and strict normal incidence. Together with positive third-disk clearance, this proves that every included flight is the actual first-hit physical flight to the next specified disk. Thus the whole central record is regular and no collision has been hidden.

At a central contact, (50.3) bounds the state distance by $A_\phi q_{\mathrm{a}}^{j(d_i)/4}$, whereas the original state is strictly farther than that from the actual section boundary. A shortest circle–momentum path cannot cross the boundary, so the two indicators agree. This argument includes both central endpoint tests and every occupation state. Equality of centers gives the central lattice displacement and flight count exactly. Finally Lemma 48.1 gives common active vertices and unit directions for all minima. There was no measurable selection, parameter continuation, or minimum multiplicity payment in this proof. □

Corollary 50.4 (Exponential exact-label costs on the unchanged full roof)**.** *Let $D_{\mathrm{obs}} \le D \le d$, $2d < m$, and restrict the measures* (49.1) *to $Q \notin \mathcal{E}^{\mathrm{obs}}_{D,R,m}$. Then*

$$\begin{aligned} \mathsf{T}_{\mathrm{clock},d} &\le E_R(Q) q_{\mathrm{a}}^{\lfloor d/N \rfloor}, \\ \mathsf{T}_{\mathrm{phase},d} &\le K_\phi E_R(Q) q_{\mathrm{a}}^{j(d)}, \\ \mathsf{T}_{\perp,d} &\le B_\perp E_R(Q) q_{\mathrm{a}}^{j(d)}. \end{aligned} \tag{50.11}$$

Consequently, under precisely Input 11.1, their original full-roof densities are bounded respectively by

$$2rC_\star q_{\mathrm{a}}^{\lfloor d/N \rfloor}, \qquad 2rC_\star K_\phi q_{\mathrm{a}}^{j(d)}, \qquad 2rC_\star B_\perp q_{\mathrm{a}}^{j(d)}.$$

These assertions also apply to the dominated persistent low-load interior source of Corollary 49.2.

For fixed $0 < \theta < 1/2$ and $d_m = \lceil \theta m \rceil$, all three collision-normalized heights are $O(m^2 e^{-\lambda_0 m})$, uniformly in the original radius, exact labels and roof, where

$$\lambda_0 = \frac{\theta |\log q_{\mathrm{a}}|}{N} > 0. \tag{50.12}$$

On this same source the section error in (47.7) *is exactly zero. After retaining its endpoint phase e^{isW_d}, the remaining nonnegative Fourier error majorant is $|s|\mathsf{T}_{\mathrm{clock},d}$. For $0 < \lambda < \lambda_0$, its collision-normalized original-roof height tends to zero uniformly for $|s| \le e^{\lambda m}$ and for all lattice and return twists.*

Proof. The clock identity (47.2) is bounded by $\mathcal{E}_d$, so the first estimate follows from the anchor contraction. To sum the phase distances use the proof of Theorem 47.2, including its repeated terminal edge:

$$\mathsf{T}_{\mathrm{phase},d} \le \frac{64}{\tau_-} \sum_{i=d}^{m-d-1} F_i + 24 R_-^{-4} \sum_{i=d}^{m-d} |\delta_i|^4.$$

Insert (48.9) and (48.8), and use $\lfloor d/N \rfloor \ge j(d)$, to obtain exactly K_ϕ. The transverse estimate is (48.10). All three statements hold pointwise before integration. Positive ordering on the unchanged original full roof, as in Theorem 49.1, and $E_R(Q) \le 2r$ give the density assertions under the named input.

For large m, d_m is admissible, and $j(d_m) \ge \theta m/N - 2$. Thus $q_{\mathrm{a}}^{j(d_m)} \le q_{\mathrm{a}}^{-2} e^{-\lambda_0 m}$. The density bounds give the normalized rate. The exact section identity was proved for every central contact and both endpoint tests, so the occupation term in the twist comparison vanishes. Finally $m^2 e^{\lambda m} e^{-\lambda_0 m} \to 0$ when $\lambda < \lambda_0$, proving the stated frequency-range estimate. This bounds the positive error majorant on the original roof; it does not assert an interchange with an unbounded Fourier integral. □

Remark 50.5 (The retained order of limits and the remaining full source)*.* Theorem 50.3 proves $\lim_{D\to\infty} \sup_{m,R,n,k,h} \|\mu^{\mathrm{obs}}_{D,R,m,n,k,h}\|_{\mathrm{TV}} = 0$ on the entire original family. Corollary 50.4 proves a stronger roof topology for specified exponentially small costs on its complement, under a precisely named inherited input. Total variation of the exceptional source is not its roof essential height. In particular these two statements do not imply that its $m^2 L^\infty$ height vanishes.

The original $\varepsilon(B), h_B, r_B, \gamma_B, K_B$, microscopic roof windows and guard q_0 remain fixed before the collision limsup, and B remains outermost in (0.1). The positive shallow source, all required physical multipliers, critical and singular paired traces, roofwise pool capacity, full conditioning denominator and the complete spectral/arithmetic inversion errors still require their actual original-source bounds. The exponentially accurate central comparison does not discard the remaining end portions or change the full event. The final raw pointwise local limit theorem and four-coordinate inversion are therefore not claimed complete.

51. Physical tube volume and a normalized positive-source payment

The total-variation estimate of Theorem 50.3 does not by itself bound a physical-roof density. This section proves a comparison which does: an entire physical momentum disk can be spread from each contributing critical tube, with its exact optical weight bounded below as well as above. Distinct tubes are disjoint as physical sources. Invariance then pays their complete volume before the original-roof coarea is performed. The result is uniform in collision count and retains arbitrary Borel exact-record selectors. It concerns the guarded critical-tube union already constructed in Section 34; no coverage of its complement is assumed.

Keep $0 < \varepsilon \le 1/4$, the guarded regular critical roots $\mathscr{Z}_{\varepsilon,m,R}$, and the positive constants $s_\varepsilon, \rho_\varepsilon, C_{\mathrm{dist}}$ from (34.3) and (34.1). For $x \in \mathscr{Z}_{\varepsilon,m,R}$ let

$$\mathcal{T}_x = \{X_x(\zeta) : |\zeta| < \rho_\varepsilon\}, \qquad \mathcal{T}_{\varepsilon,m,R} = \bigcup_{x\in\mathscr{Z}_{\varepsilon,m,R}} \mathcal{T}_x,$$

where $X_x(\zeta)$ is the actual physical initial collision state of Lemma 34.2. The whole tube is used here; neither its initial nor its terminal section test is required to be constant. Put $u_x(\zeta) = |\partial_{\alpha_0} p_m(X_x(\zeta))|^{-1}$ and $u_x^0 = u_x(0)$. Throughout, the volume used for this auxiliary comparison is the original unconditioned collision measure $\nu = d\alpha\, dp/(4\pi)$. The eventual source to be estimated still uses exactly $\nu_R^* = c_*^{-1}\mathbf{1}_{Y_R^*}\nu$.

Lemma 51.1 (Disjoint physical tubes and a two-sided optical volume)**.** *Distinct members of $\mathscr{Z}_{\varepsilon,m,R}$ have disjoint tubes as sets of physical initial states. Each parametrization X_x is injective, and*

$$C_{\mathrm{dist}}^{-1} u_x^0 \le u_x(\zeta) \le C_{\mathrm{dist}} u_x^0, \qquad \nu(\mathcal{T}_x) \ge \frac{\rho_\varepsilon^2}{4C_{\mathrm{dist}}} u_x^0. \tag{51.1}$$

Consequently, for any subfamily $\mathcal{A}$ whose entire tubes are contained in a Borel set E of the actual collision source,

$$\sum_{x\in\mathcal{A}} u_x^0 \le \frac{4C_{\mathrm{dist}}}{\rho_\varepsilon^2}\nu(E). \tag{51.2}$$

No whole-cylinder pressure or endpoint-kernel bound is used.

Proof. Every tube point is a regular first-hit trajectory with the same strict lifted word as its center, by Theorem 33.3. If two tubes contained the same physical initial state, uniqueness of the actual successive first hits would give the same lifted word, with the initial center normalized to zero. Both centers would be regular physical critical records of that word. Corollary 38.3 proves that such a record is the unique minimum on the full product of the original closed disks. Thus the centers coincide. Lattice translates and angular seams are identified in this argument, rather than counted as new centers. Injectivity within a tube is also immediate from the fact that $\zeta = (p_0, p_m)$ is an actual function of the physical initial state; its local inverse has the specified values of both momenta.

The absolute logarithmic derivative in Lemma 34.1 bounds the absolute difference $|\log u_x(\zeta) - \log u_x^0|$ by $D_* s_\varepsilon/\varepsilon = D_*\delta_*/2$ along the actual physical endpoint segment. This proves both inequalities in (51.1), not just its upper inequality. The exact inverse Jacobian from (p_0, p_m) to (α_0, p_0) is u_x, as in Theorem 28.2. Hence

$$\nu(\mathcal{T}_x) = \frac{1}{4\pi}\int_{|\zeta|<\rho_\varepsilon} u_x(\zeta)\, d\zeta \ge \frac{\rho_\varepsilon^2}{4C_{\mathrm{dist}}} u_x^0.$$

This identity has no section-conditioning factor and no additional radius factor. Sum it over finite subfamilies. Their tubes are disjoint physical sets contained in E, so their total mass is at most $\nu(E)$. Monotone exhaustion proves (51.2) for the complete subfamily. All inverse charts are

smooth on their open domains and have Borel images, so these are ordinary measurable source sets. □

Lemma 51.2 (Every deep observation moves little across the whole tube)**.** *Let Q, Q' be any two physical records in one tube. For $d_i = \min(i, m-i)$ and $d_i' = \min(i, m-1-i)$ one has*

$$\begin{aligned} |\Gamma_i(Q) - \Gamma_i(Q')| &\le L_\Gamma \sigma^{d_i'}, & 0 \le i < m, \\ |c_i(Q) - c_i(Q')| &\le L_c \sigma^{d_i}, & 0 \le i \le m, \\ d_\infty(x_i(Q), x_i(Q')) &\le L_\phi \sigma^{d_i}, & 0 \le i \le m, \end{aligned} \tag{51.3}$$

where

$$L_\Gamma = 4C_\Delta s_\varepsilon, \qquad L_c = L_\phi = 4C_c s_\varepsilon, \qquad \sigma = 47/59. \tag{51.4}$$

The clearance retains every third disk, including a disk visited elsewhere in the word. The distance in the last line is the original circle–momentum distance.

Proof. Both endpoint pairs lie in the Euclidean ball of radius s_ε about the same critical center. Their straight endpoint segment lies in that ball, which consists of actual physical trajectories by Theorem 33.3. The length of this segment in the endpoint ℓ^∞ norm is at most $2s_\varepsilon$. Integrating (29.9) along it gives the first bound. This Lipschitz estimate already takes the infimum over the full allowed set of centers, so switches of the minimizing center do not require additional pieces or a multiplicity factor.

The incidence estimate follows in the same way from (29.8) and $s_i \le 2\sigma^{d_i}$. For clarity, the same derivative bound applies to the physical tangential momentum. For an outgoing velocity v_i and the unit tangent t_i, $p_i = v_i \cdot t_i$, so

$$|D_q p_i[\eta]| \le \|D_q v_i[\eta]\|_2 + |D_q \alpha_i[\eta]| \le C_c s_i \|\eta\|_\infty.$$

At the terminal contact use the incoming velocity; its tangential component agrees with the outgoing one under specular reflection. The adjacent-depth factor σ^{-1} has already been included in C_c. Together with $|D_q \alpha_i[\eta]| \le J_* s_i \|\eta\|_\infty$ and $J_* \le C_c$, integration proves the last line. Local angular lifts may be used on the physical segment; projection to the circle can only decrease the resulting distance. Distance to the fixed set ∂Y_R^* is one-Lipschitz in this same metric. □

For nonnegative depth widths $b_\Gamma(d), b_c(d), b_\phi(d)$ define $\mathcal{B}_D(\boldsymbol{b})$ on the original physical record by

$$\bigcup_{\substack{0 \le i < m \\ d_i' \ge D}} \{\Gamma_i \le b_\Gamma(d_i')\} \quad \cup \quad \bigcup_{\substack{0 < i < m \\ d_i \ge D}} \{c_i \le b_c(d_i)\} \quad \cup \quad \bigcup_{\substack{0 < i < m \\ d_i \ge D}} \{\operatorname{dist}_\infty(x_i, \partial Y_R^*) \le b_\phi(d_i)\}.$$

These are source restrictions, not modifications of the return event. Set

$$\widetilde{b}_\Gamma(d) = b_\Gamma(d) + L_\Gamma \sigma^d, \quad \widetilde{b}_c(d) = b_c(d) + L_c \sigma^d, \quad \widetilde{b}_\phi(d) = b_\phi(d) + L_\phi \sigma^d. \tag{51.5}$$

Assume, for every $d \ge D$, that

$$0 \le \widetilde{b}_\Gamma(d) < 3/200, \qquad 0 \le \widetilde{b}_c(d) \le 1, \qquad 0 \le \widetilde{b}_\phi(d) \le 1/200. \tag{51.6}$$

Put, with nonnegative series interpreted by monotone limits,

$$\mathcal{V}_D(\boldsymbol{b}) = \frac{2}{R_-} \sum_{d \ge D} \widetilde{b}_\Gamma(d) + 2 \sum_{d \ge D} \widetilde{b}_c(d)^2 + \frac{1}{\pi} \sum_{d \ge D} \widetilde{b}_\phi(d). \tag{51.7}$$

Theorem 51.3 (A complete positive-source bound at the original roof)**.** *For all R, m, n, k, $0 < \varepsilon \le 1/4$ and every Borel $0 \le h \le 1$, the nonnegative measure*

$$(L_{m,R})_* \left(\mathbf{1}_{\mathcal{T}_{\varepsilon,m,R}} \mathbf{1}_{\mathcal{B}_D(\boldsymbol{b})} \mathbf{1}_{\Pi_{n,k,m,R}} h \, d\nu_R^* \right)$$

is absolutely continuous with density $Q^{\mathrm{bad}}_{\varepsilon,D}$ satisfying, whenever (51.6) *holds,*

$$\|Q^{\mathrm{bad}}_{\varepsilon,D}\|_{L^\infty(dt)} \le \frac{2UC_{\mathrm{dist}}^2}{c_* \rho_\varepsilon^2} \mathcal{V}_D(\boldsymbol{b}), \qquad U = 320/9. \tag{51.8}$$

The bound completes the full physical word sum before the roof essential supremum or the radius supremum. It uses neither Input 11.1 nor Theorem 26.4.

Proof. Select every tube which contains at least one point of $\mathcal{B}_D(\boldsymbol{b})$. If a tube is selected, choose one of its actual witnessed observations. The same observation at every other point of that tube satisfies its enlarged threshold in (51.5), by Lemma 51.2. Thus the entire selected tube lies in $\mathcal{B}_D(\widetilde{\boldsymbol{b}})$. This reasoning does not require a witness common to different tubes. The family of centers is countable at fixed R, m, so selection of a subfamily introduces no measurable-selection premise. The enclosing event itself is a finite union of actual Borel observations.

For the unconditioned physical measure, invariance and the exact clearance flux, incidence distribution, and original section collar give, respectively,

$$\nu\{\Gamma_i \le u\} \le u/R_-, \qquad \nu\{c_i \le u\} \le u^2, \qquad \nu\{\mathrm{dist}_\infty(x_i, \partial Y_R^*) \le u\} \le u/(2\pi).$$

The hypotheses for these three bounds are exactly (51.6). For clearance, closed inequality boundaries are handled by continuity from above using slightly larger widths still below $3/200$. The exact incidence formula and the closed-rectangle collar bound already include their displayed upper endpoints; zero widths follow by continuity from above. At most two contacts or flights have each specified depth. The union bound therefore gives

$$\nu(\mathcal{B}_D(\widetilde{\boldsymbol{b}})) \le \mathcal{V}_D(\boldsymbol{b}).$$

No independence of observations at different times is used. Lemma 51.1 now proves

$$\sum_{\text{selected } x} u_x^0 \le \frac{4C_{\mathrm{dist}}}{\rho_\varepsilon^2} \mathcal{V}_D(\boldsymbol{b}).$$

On a selected tube use the exact polar coarea of Theorem 28.2. At a fixed physical roof and angle, there is at most one contributing radius in its momentum disk, and its density is $u_x(\zeta)/(4\pi c_* h_x(\zeta))$. Here $h_x \ge U^{-1}$ and $u_x \le C_{\mathrm{dist}} u_x^0$. All source and exact-record selectors stay inside this density until they are bounded above by one. Summing the preceding nonnegative coefficient bound and integrating an angle of length 2π gives exactly (51.8). One first applies this argument to finitely many tubes and nonnegative roof tests, then uses monotone convergence. This proves absolute continuity as well as its bound, without presupposing a finite density. The polar origins have zero collision area; no measure supported on a singular paired boundary has been discarded. The single factor c_*^{-1} occurs only in this final original-source coarea, not in the preceding unconditioned tube-volume comparison. □

Corollary 51.4 (The exact-record exceptional source is paid on these tubes)**.** *Use the unchanged amplitudes of* (50.2) *and put*

$$b_\Gamma(d) = A_\Gamma q_{\mathrm{a}}^{j(d)/2}, \quad b_c(d) = A_c q_{\mathrm{a}}^{j(d)/2}, \quad b_\phi(d) = A_\phi q_{\mathrm{a}}^{j(d)/4}.$$

Then $\mathcal{B}_D(\boldsymbol{b}) = \mathcal{E}^{\mathrm{obs}}_{D,R,m}$. *For every* $D \geq D_{\mathrm{cv}} := 20861442654$, *independently of* ε, R, m, n, k, *the width regime holds and*

$$\begin{aligned}\mathcal{V}_D(\boldsymbol{b}) \leq{}& \frac{2NA_\Gamma}{R_-}\frac{q_{\mathrm{a}}^{j(D)/2}}{1-q_{\mathrm{a}}^{1/2}} + \frac{2L_\Gamma}{R_-}\frac{\sigma^D}{1-\sigma}\\ &+ 4NA_c^2\frac{q_{\mathrm{a}}^{j(D)}}{1-q_{\mathrm{a}}} + 4L_c^2\frac{\sigma^{2D}}{1-\sigma^2}\\ &+ \frac{NA_\phi}{\pi}\frac{q_{\mathrm{a}}^{j(D)/4}}{1-q_{\mathrm{a}}^{1/4}} + \frac{L_\phi}{\pi}\frac{\sigma^D}{1-\sigma}.\end{aligned} \tag{51.9}$$

For every fixed $0 < \theta < 1/2$, *set* $D_m = \lceil \theta m \rceil$. *At each fixed* $\varepsilon > 0$,

$$\lim_{m\to\infty} \sup_{R,n,k,h} m^2 \|Q^{\mathrm{bad}}_{\varepsilon,D_m}\|_\infty = 0. \tag{51.10}$$

In particular this pays, in the original normalized roof topology, the entire exceptional bulk source of Theorem 50.3 which belongs to $\mathcal{T}_{\varepsilon,m,R}$. *It is a density theorem for a specified full positive source, not a conversion of total variation to essential height without a comparison argument.*

Proof. Here is a rational certificate for the deliberately conservative common lower depth. The retained constants give

$$A_\Gamma \leq 12552, \quad A_c \leq 4, \quad A_\phi \leq 485, \quad L_\Gamma \leq \frac{12756411}{9390862458880}, \quad L_c = L_\phi \leq \frac{8535}{284227072}.$$

The last two bounds use $\varepsilon \leq 1/4$ and $\delta_* = 27/67443712$. Writing $\kappa = 3263680$, Bernoulli's inequality gives $(1+1/\kappa)^\kappa \geq 2$, hence $q_{\mathrm{a}}^\kappa \leq 1/2$. Also $2 \cdot 47^4 < 59^4$, so $\sigma^4 < 1/2$. At $D_{\mathrm{cv}} = 94(1+68\kappa)$ one has $j(D_{\mathrm{cv}}) = 68\kappa$, and consequently

$$q_{\mathrm{a}}^{j(D_{\mathrm{cv}})/4} \leq 2^{-17}, \qquad q_{\mathrm{a}}^{j(D_{\mathrm{cv}})/2} \leq 2^{-34}, \qquad \sigma^{D_{\mathrm{cv}}} \leq 2^{-17}.$$

Inserting the preceding rational amplitude bounds gives exactly the three strict or weak inequalities in (51.6); for example the collar bound is $(485 + 8535/284227072)2^{-17} < 1/200$. All widths are nonincreasing in D, proving the stated common regime without a floating-point logarithm or a depth chosen after the collision count. Use (50.10) for the q_{a} terms, ordinary geometric summation for the σ terms, and $(a+b)^2 \leq 2a^2 + 2b^2$ for the incidence term. This proves all six constants in (51.9).

For $D_m \geq N$, $j(D_m) \geq \theta m/N - 2$. Every term on its right is bounded by a constant times $e^{-\lambda m}$, where one may take

$$\lambda = \theta \min\left\{\frac{|\log q_{\mathrm{a}}|}{4N}, |\log\sigma|\right\} > 0.$$

The prefactor in (51.8) is finite at fixed ε. Multiplication by m^2 proves (51.10). The bounds held at every finite m, D before this substitution. Thus this depth classification changes no microscopic roof window, physical clock, exact event, or originally fixed outer parameter. If ε is a positive quantity fixed at an outer stage B, the inner limit is zero for each such B before B is sent to infinity. □

Corollary 51.5 (The actual linear incidence multiplier and persistent source)**.** *The preceding bounds remain valid with the source multiplied by the original* $I_m = 1 + \sum_{i=1}^{m-1} q_0^{d_i}/c_i$, *after multiplication of their right sides by the explicit collision-uniform factor*

$$H_\varepsilon = 1 + \frac{4}{\varepsilon}\frac{q_0^{3/4}}{1-q_0^{3/4}}, \qquad q_0 = 47/53. \tag{51.11}$$

The same normalized payment applies to the Borel restriction of the inherited persistent low-load interior source to $\mathcal{T}_{\varepsilon,m,R} \cap \mathcal{E}^{\mathrm{obs}}_{D_m,R,m}$, using its specific domination in Lemma 12.1.

Proof. Every tube point has the actual half-guard $c_i \geq (\varepsilon/2) q_0^{d_i/4}$ by Theorem 33.3. Hence

$$I_m \leq 1 + \frac{2}{\varepsilon} \sum_{i=1}^{m-1} q_0^{3d_i/4} \leq 1 + \frac{4}{\varepsilon} \sum_{d=1}^{\infty} q_0^{3d/4} = H_\varepsilon.$$

This is a bound for the actual retained multiplier on these physical tubes. Positivity proves the weighted statement. For the last assertion multiply $0 \leq c_{\mathrm{per}}^{\varepsilon(B),\gamma_B} \leq f^{\varepsilon(B)} \leq A$ by the stated Borel restriction and original low-load selector. Its actual source sum is dominated by the corresponding restricted original roof measure. Apply (51.10). The tube width ε may be fixed independently of the inherited outer width $\varepsilon(B)$; if chosen from it, it must be fixed before the collision limit. No assertion is made about a singular paired endpoint trace or a coefficient lacking this specific domination. □

Remark 51.6 (Exact scope and the still required full-source estimate). The new estimate pays actual bulk source at its common physical roof through a proved lower optical volume and physical tube disjointness. Its dependence on the fixed tube radius is explicit: ρ_ε^{-2}. It establishes no uniform bound as that radius tends to zero and therefore cannot be extended to every regular trajectory merely by exhausting smaller guard classes. Neither singular paired traces nor a full $I_m/|\nabla F|$ insertion is covered by (51.11). The original shallow and uncovered boundary sources, comparison capacity, full physical multipliers, and the remaining spectral, arithmetic and conditioning interfaces retain their exact obligations. In particular the original ordered limit (15.8), the raw pointwise local limit theorem, and the four-coordinate inversion are not claimed closed here.

52. A finite-overlap physical atlas and original-roof source height

We prove a height estimate for the complete class having a fixed margin in the *original* incidence and clearance tests. The class includes both critical and noncritical physical trajectories. It is not restricted to trajectories that continue to a critical orbit. The argument uses actual two-dimensional families of trajectories and bounds their overlap before summing their positive optical weights. Neither a whole-cylinder pressure estimate nor a mixing theorem is used.

Write $r_g = q_0^{1/4}$, where $q_0 = 47/53$ remains unchanged. For $0 < \varepsilon \leq 1/4$ let $\mathcal{G}_{\varepsilon,m,R}$ be the regular physical records satisfying

$$|p_0|, |p_m| \leq 5/6, \qquad c_i \geq \varepsilon r_g^{d_i} \ (0 \leq i \leq m), \qquad \Delta_i \geq \varepsilon r_g^{d_i'} \ (0 \leq i < m). \tag{52.1}$$

These are source restrictions used in an estimate; the original probability, event and microscopic windows are unchanged. The endpoint cap includes the entire original exact event. Indeed, $q_R < 53/100$ and $q \mapsto 2q/(1+q^2)$ is increasing on $(0,1)$, so the largest section momentum is less than $10600/12809 + 1/200 < 5/6$. The other three types of section rectangle have smaller momentum bounds. No section-distance margin is imposed in (52.1).

Keep the already proved constants $\delta_*, C_{\mathrm{inc}}$ from (33.2), D_*, C_{dist} from (34.1), J_*, C_c, C_Δ from Section 29, and put

$$g = 9/40, \qquad U = 320/9, \qquad a_\varepsilon = \delta_* \varepsilon/16, \qquad \rho_\varepsilon^{\mathrm{at}} = \frac{g a_\varepsilon}{8R_+}, \qquad C_{\mathrm{at}} = \frac{25UC_{\mathrm{dist}}^2}{2c_*(\rho_\varepsilon^{\mathrm{at}})^2}. \tag{52.2}$$

We abbreviate $\rho = \rho_\varepsilon^{\mathrm{at}}$ within this section. It is an endpoint-momentum chart radius, never the physical roof or the microscopic comparison width r_B. All constants and chart radii are fixed before the collision count tends to infinity.

Lemma 52.1 (A common momentum neighborhood of every guarded record)**.** *For $x \in \mathcal{G}_{\varepsilon,m,R}$ put $q_x = (\alpha_0, \alpha_m)$ in local endpoint lifts and $\zeta_x = (p_0, p_m)$. The actual endpoint ball $|q - q_x|_2 < a_\varepsilon$ is physical with the same lifted word, and all its incidences and clearances satisfy the half-margins* (33.5)*. Its endpoint incidences exceed* $1/2$*. Every $|\zeta - \zeta_x| < 4\rho$ has a unique actual inverse on that ball, with*

$$|q(\zeta) - q_x|_2 \le R_+|\zeta - \zeta_x|/g < a_\varepsilon/2. \tag{52.3}$$

Here and below uniqueness also holds among all regular physical trajectories of the same lifted word with the given ζ.

Proof. On the closed ball, $|h_0| + |h_m| \le 2a_\varepsilon = \delta_*\varepsilon/8 < \delta_*\varepsilon$. Theorem 33.3 gives the actual trajectories and all half-margins, on a neighborhood of this closed ball. Its signed tangential estimate bounds each endpoint momentum change by $C_{\mathrm{inc}}\delta_*\varepsilon/8 \le \varepsilon/16 \le 1/64$. Thus $|p_0|, |p_m| \le 5/6 + 1/64 = 163/192 < 17/20$ and $c_0, c_m \ge \sqrt{111}/20 > 1/2$. The original second variation therefore gives $G = D_q^2 F \succeq gI$.

For the desired momentum pair minimize $F(q) - RD\zeta \cdot (q - q_x)$ on the closed ball, where $D = \mathrm{diag}(-1, 1)$. At its boundary, putting $h = q - q_x$ and using $\nabla F(q_x) = RD\zeta_x$, the outward radial derivative is at least

$$ga_\varepsilon^2 - R_+|\zeta - \zeta_x|a_\varepsilon > ga_\varepsilon^2/2 > 0.$$

A boundary minimizer is impossible. Strong convexity gives one interior minimizer, where $\nabla F = RD\zeta$. The inverse function theorem and strong monotonicity give its smooth dependence and (52.3). This minimization takes place inside a ball already proved physical; it does not assert that an arbitrary stationary continuation is physical. Finally, with equal endpoint momenta the left side of the support inequality (10.3) is zero. Strict positivity of every physical incidence forces all contacts to coincide. This proves the asserted global wordwise uniqueness, even across disconnected original event components. □

Lemma 52.2 (Physical tubes with multiplicity at most twenty-five)**.** *For each fixed R, m, ε there is a finite family $\{T_{w,z}\}$ of actual physical source sets, indexed by lifted words w and points z of the grid $\rho\mathbb{Z}^2$, with these properties. Every $T_{w,z}$ is parametrized by the* whole *momentum disk $B(z, 2\rho)$, carries word w, and contains a unique physical center $x_{w,z}$ with momentum z. Their union covers $\mathcal{G}_{\varepsilon,m,R}$. Each physical initial point is in at most* 25 *of the tubes, counting all words and grid points. Writing $u(x) = |\partial_{\alpha_0} p_m(x)|^{-1}$, one has throughout a tube*

$$C_{\mathrm{dist}}^{-1} u(x_{w,z}) \le u(x) \le C_{\mathrm{dist}} u(x_{w,z}), \qquad \nu(T_{w,z}) \ge \frac{\rho^2}{C_{\mathrm{dist}}} u(x_{w,z}). \tag{52.4}$$

For any two points x, y in a tube,

$$|\Delta_i(x) - \Delta_i(y)| \le 2C_\Delta a_\varepsilon \sigma^{d'_i}, \qquad 0 \le i < m, \tag{52.5}$$

$$|c_i(x) - c_i(y)| \le 2C_c a_\varepsilon \sigma^{d_i}, \qquad 0 \le i \le m, \tag{52.6}$$

$$d_\infty(T_R^i x, T_R^i y) \le 2C_c a_\varepsilon \sigma^{d_i}, \qquad 0 \le i \le m. \tag{52.7}$$

The states in the last line are actual collision states, including the incoming definition of the terminal tangential momentum.

Proof. Assign each ζ_x of a source point to a nearest grid point, with a fixed convention on ties. The Euclidean distance is at most $\rho/\sqrt{2} < \rho$. For every nonempty pair (w, z) choose one such source point x. There are finitely many relevant grid points and finitely many physical words at fixed m, by the proved physical horizon. Apply Lemma 52.1 at x. Since $B(z, 2\rho) \subset B(\zeta_x, 3\rho)$, its inverse gives the specified tube and center. The local parametrization is injective as a physical source map: its momenta determine the point by wordwise uniqueness. Every other assigned source point with the same pair (w, z) agrees with the constructed inverse at its own momentum

by that same global uniqueness. Thus the family covers the entire class, not just the chosen representatives. Different constructions for a pair define exactly the same physical inverse on the disk.

A physical initial point determines its unique complete lifted word. If it lies in $T_{w,z}$, its momentum is within 2ρ of z. Each coordinate interval of length 4ρ contains at most five points of $\rho\mathbb{Z}$. There are consequently at most 25 possible grid centers. This multiplicity bound uses the physical momentum map; it introduces neither independent marks nor a factor equal to the number of collision words.

In the chosen endpoint ball all tube points and the center have distance less than $a_\varepsilon/2$ from q_x. Any two can be joined by their straight segment inside this actual physical ball, of endpoint ℓ^∞ length at most a_ε. All along it $c_i \geq (\varepsilon/2) r_g^{d_i}$ and the endpoint incidences exceed $1/2$. The absolute logarithmic derivative bound in Lemma 34.1, integrated in both directions, gives distortion at most $\exp(D_* a_\varepsilon/\varepsilon) \leq \exp(D_*\delta_*/2) = C_{\mathrm{dist}}$. In momentum coordinates the *unconditioned* invariant collision probability is $u\, d\zeta/(4\pi)$. The disk area is $4\pi\rho^2$. Integrating the lower distortion bound proves the volume estimate in (52.4).

For clearance and incidence integrate (29.9) and (29.8) along the same physical segment, using $s_i \leq 2\sigma^{d_i}$. The tangential derivative obeys the identical incidence bound: differentiate $p_i = v_i \cdot t_i$, or $p_m = v_{m-1} \cdot t_m$ at the terminal contact, and use the direction and contact estimates of Proposition 29.2. Also $|D\alpha_i| \leq J_* s_i \|\eta\|_\infty$ and $J_* \leq C_c$. This proves (52.7) in the circular coordinate metric. The clearance estimate is for the complete closed flight segment and every third disk; nearest-disk switches and disks visited at other times are already included in its Lipschitz proof. □

Lemma 52.3 (The roof Jacobian on the entire endpoint cap)**.** *At every regular point in these tubes, write $\zeta = s e_\theta \neq 0$ and $\widehat{G} = R^{-2} G$. Then*

$$\frac{1}{U} \leq h(x,\theta) := e_\theta^{\mathsf{T}} D \widehat{G}^{-1} D e_\theta \leq 2R_+, \qquad \frac{dL(x(se_\theta))}{ds} = s h(x,\theta), \qquad b(x) = \frac{u(x)}{4\pi c_* h(x,\theta)}. \tag{52.8}$$

The last expression is the original section-conditioned source density in (t,θ), with section and all exact-event tests retained as indicators. A momentum disk need not be centered at the origin for this formula.

Proof. The Schur-complement proof of Lemma 28.1 gives $\widehat{G} \preceq (320/9) I$ for all $|p_0|, |p_m| < 1$. For $m \geq 2$ its better bound is $170/9$; the larger constant also treats the single-flight mixed entry. Its lower bound uses only $c_0, c_m \geq 1/2$, so $\widehat{G} \succeq (2R_+)^{-1} I$. These componentwise caps are enough; we do not require $|\zeta|_2 < 1$. The exact identities $\nabla_q F = RD\zeta$ and $D_\zeta q = RG^{-1}D$ now give the radial derivative. The inverse source Jacobian is u, polar area is $s\, ds\, d\theta$, and $dt = sh\, ds$. Cancellation gives the displayed density, with precisely one c_*^{-1} and no omitted radius factor. On each open radial interval in a tube it is ordinary one-dimensional substitution. For a fixed word, Theorem 10.2 prevents two distinct regular points with the same roof and angle, even if an exact-event cut disconnects that interval. At $\zeta = 0$ there is at most one regular physical point per word; these finitely many source points have zero collision area. No singular endpoint trace is discarded by this area statement. □

For a Borel physical source set E, let $\mathrm{Sat}_\varepsilon(E)$ be the union of precisely those tubes above which meet $E \cap \mathcal{G}_{\varepsilon,m,R}$. This is a finite union of open physical source sets and is Borel at each fixed radius and collision count. The choices need not be measurable jointly in R: the required order is $\sup_R \operatorname{ess\,sup}_t$, with radius-dependent null sets. This auxiliary union is used only on the upper side of an inequality; it is never a conditioning event.

Theorem 52.4 (Complete physical source height by tube-volume payment)**.** *For every $m \geq 1$, $R \in [R_-, R_+]$, original exact labels n, k, Borel E and $0 \leq h \leq 1$, the original full-roof measure*

$$\mu_E = (L_{m,R})_*(\mathbf{1}_E \mathbf{1}_{\mathcal{G}_{\varepsilon,m,R}} \mathbf{1}_{\Pi_{n,k,m,R}} h \, d\nu_R^*)$$

is absolutely continuous and satisfies

$$\left\| \frac{d\mu_E}{dt} \right\|_\infty \leq C_{\mathrm{at}} \, \nu(\mathrm{Sat}_\varepsilon(E)). \tag{52.9}$$

In particular the full guarded physical source has height at most $C_{\mathrm{at}} = O(\varepsilon^{-2})$, uniformly in all collision counts, radii, labels and bounded selectors. This conclusion has no premise from Input 11.1, and does not use the endpoint-kernel constant C_{mom}.

Proof. Assign every source point in the covered class to the first tube containing it in a fixed enumeration. This is a Borel partition of the original class. Retain the assignment, E, the complete event and h in the coarea integrand. Lemma 52.3 and the upper bound in (52.4) dominate that integrand on its assigned tube by $UC_{\mathrm{dist}} u(x_{w,z})/(4\pi c_*)$. Only tubes meeting $E \cap \mathcal{G}_{\varepsilon,m,R}$ can contribute. For every such tube its lower volume bound gives $u(x_{w,z}) \leq C_{\mathrm{dist}} \nu(T_{w,z})/\rho^2$. The proved multiplicity bound, applied to this subfamily, yields

$$\sum_{(w,z):\, T_{w,z} \cap E \cap \mathcal{G}_\varepsilon \neq \varnothing} u(x_{w,z}) \leq \frac{25 C_{\mathrm{dist}}}{\rho^2} \nu(\mathrm{Sat}_\varepsilon(E)).$$

Integrating the uniform coarea bound over an angle of length 2π gives exactly (52.9).

To justify absolute continuity before claiming its bound, apply local substitution to nonnegative roof tests on relatively compact regular charts, disjointize only their representations, and exhaust them monotonically as in Proposition 10.4. The estimates just proved bound the completed positive sum. All physical words and all original cuts are included before taking its essential supremum. The underlying tube volume used ν without an initial-section restriction; the source coarea used ν_R^* with the original event. This explains the single factor c_*^{-1} and involves no change of probability. □

Corollary 52.5 (Observable neighborhoods in the actual collision process)**.** *Let E be the union, over any designated contact or flight indices, of the actual conditions $\Delta_i \leq \eta_i$, $c_i \leq \xi_i$, and $\mathrm{dist}_\infty(T_R^i x, \partial Y_R^*) \leq \zeta_i$. All widths are nonnegative. Replace them respectively by*

$$\eta_i^+ = \eta_i + 2C_\Delta a_\varepsilon \sigma^{d_i'}, \quad \xi_i^+ = \xi_i + 2C_c a_\varepsilon \sigma^{d_i}, \quad \zeta_i^+ = \zeta_i + 2C_c a_\varepsilon \sigma^{d_i}, \tag{52.10}$$

and denote the resulting original-observable union by E^+. Then $\mathrm{Sat}_\varepsilon(E) \subset E^+$. When $\eta_i^+ < 3/200$, $\xi_i^+ \leq 1$, and $\zeta_i^+ \leq 1/200$, one has the complete bound

$$\left\| \frac{d\mu_E}{dt} \right\|_\infty \leq C_{\mathrm{at}} \left\{ \frac{1}{R_-} \sum_i \eta_i^+ + \sum_i (\xi_i^+)^2 + \frac{1}{2\pi} \sum_i \zeta_i^+ \right\}. \tag{52.11}$$

Each sum is only over its designated actual indices; overlap of different witnesses requires no independence assumption.

Proof. A tube meeting $E \cap \mathcal{G}_\varepsilon$ has a point with at least one of the stated witnesses. Every point of that whole tube has the corresponding widened witness, by (52.5)–(52.7) and the one-Lipschitz property of distance to the fixed original section boundary. This proves the inclusion without selecting the same witness on different tubes. The exact physical clearance flux bounds $\nu\{\Delta_i \leq u\} \leq u/R_-$ in the stated range. The invariant incidence marginal is $\nu\{c_i \leq u\} = 1 - \sqrt{1-u^2} \leq u^2$. Lemma 50.1 bounds the section collar by $u/(2\pi)$. Apply invariance of the actual collision map and the union bound, then Theorem 52.4. All tests remain

on actual physical flights and contacts. The widened sets occur only in this proved upper estimate; the original event and its microscopic widths are not widened. □

The finite-overlap mechanism is a quantitative comparison between a point's optical weight and a positive volume of actual nearby trajectories. It therefore supplies information absent from a total-mass bound alone. It does not show that the complement of $\mathcal{G}_{\varepsilon,m,R}$ has small normalized roof height. The factor ε^{-2} is explicit, so no uniform passage to zero guard margin is implicit in this theorem.

53. Normalized marked source, the physical singular multiplier, and pressure

We now apply the physical atlas to the precise exceptional source of Section 50. This gives its collision-normalized roof-height payment on the entire original guarded class, including trajectories with no physical critical continuation. We also treat the actual multiplier $I_m/|\nabla_q F|$ in the original variation budget, in its appropriate roof-integral topology. These deductions use the physical geometry and the explicit optical formulas, not the whole-cylinder pressure premise.

Let $\mathcal{G}_\varepsilon = \mathcal{G}_{\varepsilon,m,R}$, $a_\varepsilon, \rho = \rho_\varepsilon^{\mathrm{at}}, C_{\mathrm{at}}$ be as in Section 52. Keep the original I_m and put

$$H_\varepsilon = 1 + \frac{4}{\varepsilon}\frac{q_0^{3/4}}{1-q_0^{3/4}}. \tag{53.1}$$

This deliberately uses the half-guard bound of (51.11); on $\mathcal{G}_\varepsilon$ itself a smaller constant would suffice. In particular it is valid on every physical comparison tube used below. It pays the original incidence multiplier, not a substitute weight.

For $D \ge D_{\mathrm{cv}}$ and $s > 0$ write

$$S_s(D) = \frac{N q_{\mathrm{a}}^{sj(D)}}{1-q_{\mathrm{a}}^s}, \qquad T_s(D) = \frac{\sigma^{sD}}{1-\sigma^s}, \qquad N = 94, \tag{53.2}$$

and define

$$\begin{aligned} V_\varepsilon(D) = {} & \frac{2A_\Gamma}{R_-}S_{1/2}(D) + \frac{4C_\Delta a_\varepsilon}{R_-}T_1(D) + 4A_c^2 S_1(D) \\ & + 16C_c^2 a_\varepsilon^2 T_2(D) + \frac{A_\phi}{\pi}S_{1/4}(D) + \frac{2C_c a_\varepsilon}{\pi}T_1(D). \end{aligned} \tag{53.3}$$

All amplitudes are those of the already proved actual-source theorem. The original guard parameter q_0 and the anchor contraction q_{a} remain different fixed constants.

Theorem 53.1 (The complete guarded exceptional source has vanishing height)**.** *For every $0 < \varepsilon \le 1/4$, $D \ge D_{\mathrm{cv}}$, all m, R, n, k, and every Borel selector $0 \le h \le 1$, the original full-roof measures with respective weights*

$$\mathbf{1}_{\mathcal{G}_\varepsilon}\mathbf{1}_{\mathcal{E}^{\mathrm{obs}}_{D,R,m}}\mathbf{1}_{\Pi_{n,k,m,R}}h, \qquad I_m\mathbf{1}_{\mathcal{G}_\varepsilon}\mathbf{1}_{\mathcal{E}^{\mathrm{obs}}_{D,R,m}}\mathbf{1}_{\Pi_{n,k,m,R}}h$$

against ν_R^* have densities b_D and b_D^I satisfying

$$\|b_D\|_\infty \le C_{\mathrm{at}}V_\varepsilon(D), \qquad \|b_D^I\|_\infty \le H_\varepsilon C_{\mathrm{at}}V_\varepsilon(D). \tag{53.4}$$

Consequently, for every fixed $\varepsilon > 0$ and $0 < \theta < 1/2$, with $D_m = \lceil \theta m \rceil$,

$$\lim_{m\to\infty}\sup_{R,n,k,h} m^2(\|b_{D_m}\|_\infty + \|b_{D_m}^I\|_\infty) = 0. \tag{53.5}$$

This is a complete positive-source limit in the required pointwise roof topology on the stated class. No conditioning event or roof interval has been enlarged.

Proof. Use Corollary 52.5 with the three widths defining $\mathcal{E}^{\mathrm{obs}}_{D,R,m}$. Its widening constants $2C_\Delta a_\varepsilon$ and $2C_c a_\varepsilon$ are smaller than $L_\Gamma, L_c = L_\phi$ in (51.4), since $a_\varepsilon = s_\varepsilon/8$. The rational depth D_{cv} therefore establishes all required width regimes, uniformly in ε. At most two true contacts or flights have each depth. Apply the inherited block sum (50.10), the geometric series for σ^d, and $(u+v)^2 \le 2u^2 + 2v^2$. This gives exactly $\nu(E^+) \le V_\varepsilon(D)$, including the factor 16 in the squared incidence widening. Theorem 52.4 proves the first estimate. The pointwise bound $I_m \le H_\varepsilon$ proves the second.

Put

$$\lambda_b = \theta \min\left\{\frac{|\log q_{\mathrm{a}}|}{4N}, |\log \sigma|\right\} > 0, \qquad \lambda_c = \frac{\theta |\log q_{\mathrm{a}}|}{N} > 0. \tag{53.6}$$

Since $j(D_m) \ge \theta m/N - 2$, the six explicit terms in $V_\varepsilon(D_m)$ are $O_\varepsilon(e^{-\lambda_b m})$. Thus the two normalized heights are $O_\varepsilon(m^2 e^{-\lambda_b m})$, uniformly after the full physical sum. The finitely many counts with $D_m < D_{\mathrm{cv}}$ do not affect the limit. This substitution is into a theorem valid at every finite m, D, not into a fixed-window asymptotic. □

Theorem 53.2 (Exact central marks and their normalized Fourier comparison)**.** *Fix the same ε, θ and let $d = D_m$, $b = m - d$. On $\mathcal{G}_\varepsilon \setminus \mathcal{E}^{\mathrm{obs}}_{d,R,m}$, for all sufficiently large m, every global disk-word minimum has the actual first-hit central record $Y_d, \dots, Y_b$. All central collision states agree between minima. Its two lattice labels, $b-d$ actual flights, all original section indicators, occupation on $[d,b)$, and the two separate endpoint tests agree exactly with those of $Q_d, \dots, Q_b$. The unchanged central physical clocks obey*

$$L_d(Y) - L_d(Q) = W_d(Q,Y) + \mathsf{T}_{\mathrm{clock},d}, \qquad 0 \le \mathsf{T}_{\mathrm{clock},d} \le 2R_+ q_{\mathrm{a}}^{\lfloor d/N \rfloor}. \tag{53.7}$$

The phase and transverse costs satisfy the corresponding pointwise estimates (50.11)*. Their original full-roof densities on this class, with any selector $0 \le h \le 1$, are bounded by*

$$2R_+ C_{\mathrm{at}} q_{\mathrm{a}}^{\lfloor d/N \rfloor}, \quad 2R_+ C_{\mathrm{at}} K_\phi q_{\mathrm{a}}^{j(d)}, \quad 2R_+ C_{\mathrm{at}} B_\perp q_{\mathrm{a}}^{j(d)}, \tag{53.8}$$

respectively, without the pressure input. Multiplication by I_m costs only H_ε.

More precisely define the central marked phase, for $\xi \in \mathbb{R}^2$, $s, \zeta \in \mathbb{R}$, by

$$\Phi_Q = \eta_R(x_d)\eta_R(x_b) \exp\{i[\xi \cdot K_d(Q) + sL_d(Q) + \zeta A_d(Q)]\}, \quad A_d(Q) = \sum_{i=d}^{b-1} \eta_R(x_i).$$

On the good class put $\widehat{\Phi} = e^{-isW_d}\Phi_Y$; on its exceptional class set $\widehat{\Phi} = 0$ as a comparison coefficient, without assigning it a fictitious trajectory. For the signed measure

$$\mathfrak{D} = (L_{m,R}(Q))_*(\mathbf{1}_{\mathcal{G}_\varepsilon} \mathbf{1}_{\Pi_{n,k,m,R}} h(\Phi_Q - \widehat{\Phi})\, d\nu_R^*)$$

one has

$$\left\| \frac{d\mathfrak{D}}{dt} \right\|_\infty \le C_{\mathrm{at}}\{V_\varepsilon(d) + 2R_+|s| q_{\mathrm{a}}^{\lfloor d/N \rfloor}\}. \tag{53.9}$$

Thus its m^2-normalized height tends to zero uniformly for all lattice and return twists and $|s| \le e^{\lambda m}$, for every fixed $0 < \lambda < \lambda_c$. The same conclusion holds with the actual I_m multiplier.

Proof. The physicality and exact observations, for every minimum, are the already proved conclusion of Theorem 50.3; none is newly assumed. The identity for the signed endpoint work W_d is (47.2), with the bound from the anchor contraction. Apply the full guarded source bound in Theorem 52.4 to the nonnegative pointwise costs. This proves (53.8) on the original full roof. Their rates are $O_\varepsilon(m^2 e^{-\lambda_c m})$ after normalization. This discharges the former pressure dependence of these cost estimates on the present full guarded class.

On the good class every integer mark and endpoint test in Φ_Q equals its counterpart in Φ_Y. Equation (53.7) gives $|\Phi_Q - e^{-isW_d}\Phi_Y| \le |s|\mathsf{T}_{\mathrm{clock},d}$. On the exceptional class the difference has

modulus at most one. The absolute value of the signed pushforward is therefore dominated by these two positive original-roof measures. Apply (53.4) and (53.8) to prove (53.9). The exceptional term is independent of all twists, and the clock term tends to zero after multiplication by $m^2 e^{\lambda m}$ when $\lambda < \lambda_c$. For I_m use its actual pointwise bound.

The common active vertices make the central comparison coefficient unambiguous; alternatively one may choose one minimum for each of the finitely many words at fixed R, m. The end portions of the full physical record are retained in $L_{m,R}(Q)$ and in Π throughout. Multiplication by their exact phases, or any Borel coefficient of modulus at most one, preserves the same positive domination. The statement is not an integration over an unbounded Fourier range, nor a replacement of the original return event by a central event. □

Lemma 53.3 (A complete physical inverse-gradient volume bound)**.** *Let E be any of the observable unions of Corollary 52.5, with widened event E^+. For every Borel roof set I of finite Lebesgue measure $|I|$, the actual positive source satisfies*

$$\int \mathbf{1}_{\mathcal{G}_\varepsilon} \mathbf{1}_E \mathbf{1}_\Pi h \mathbf{1}_{\{L_{m,R} \in I\}} \frac{I_m}{|\nabla_q F|}\, d\nu_R^* \\ \le H_\varepsilon \nu(E^+) \left\{ \frac{C_{\mathrm{at}}}{R_- \rho} |I| + \frac{U C_{\mathrm{dist}}^2 \sqrt{R_+}}{c_* R_- \rho^2} \sqrt{|I|} \right\}. \tag{53.10}$$

The source at the finitely many regular critical points may be assigned any value; the nonnegative integral is defined by truncation off those zero-area points. The resulting roof measure is absolutely continuous. No bounded pointwise density for $I_m/|\nabla_q F|$ at a critical roof is asserted.

Proof. Write $\zeta = (p_0, p_m)$. Since $|\nabla_q F| = R|\zeta|$, the part with $|\zeta| \ge \rho$ is bounded by $H_\varepsilon/(R_-\rho)$ times the measure in Theorem 52.4. Its saturation is contained in E^+, giving the first term.

For every word with a point of $\mathcal{G}_\varepsilon \cap E \cap \{|\zeta| < \rho\}$ choose one such point x. Lemma 52.1 supplies a physical inverse on $B(\zeta_x, 4\rho)$, which contains the whole disk $B(0, 2\rho)$. Thus this word has a true regular critical center, and a physical tube parametrized by that whole disk. Every other selected point of this word with $|\zeta| < \rho$ agrees with this inverse by global wordwise momentum uniqueness. Distinct words give disjoint tubes, since a physical source point has only one lifted word. The two-sided distortion bound and the original witness transport, on the chosen endpoint ball, give

$$\sum_{\text{selected words}} u_w(0) \le \frac{C_{\mathrm{dist}}}{\rho^2} \nu(E^+).$$

Indeed every point of such a tube is within the displayed widened witness of its selected original point; the volume argument is the one in (52.4), now with multiplicity one. No initial-section indicator is put into this lower tube volume.

Let a_w be the original full physical roof of that critical center. On its disk, radial coarea and (52.8) give

$$0 < L_w(re_\theta) - a_w \le R_+ r^2, \qquad \frac{1}{Rr} \le \frac{\sqrt{R_+}}{R_- \sqrt{t - a_w}} \quad (t > a_w).$$

The source density in (t, θ) is at most $U C_{\mathrm{dist}} u_w(0)/(4\pi c_*)$ before the inverse-gradient and incidence weights are inserted. After integrating the angle, the positive roof density is consequently bounded by

$$\frac{H_\varepsilon U C_{\mathrm{dist}} \sqrt{R_+}}{2 c_* R_-} \sum_w u_w(0) (t - a_w)^{-1/2} \mathbf{1}_{\{t > a_w\}}.$$

The actual radii chosen by a common roof may differ between words; the displayed two-sided tube comparison has already paid that difference. For any Borel set of measure $\ell < \infty$, layer

cake gives

$$\int_I (t-a)_+^{-1/2}\mathbf{1}_{\{t>a\}}\,dt \le \int_0^\infty \min\{\ell, v^{-2}\}\,dv = 2\sqrt{\ell}.$$

Inserting the complete coefficient sum proves the second term of (53.10), with its exact factor two. Truncate the nonnegative multiplier, use local coarea on regular charts, then monotone convergence. The displayed locally integrable bound proves finiteness on every finite roof set and absolute continuity of the completed measure. It does not set a separate singular paired trace to zero. □

Corollary 53.4 (Normalized payment of the actual singular volume)**.** *For $E = \mathcal{E}^{\mathrm{obs}}_{D_m,R,m}$, the right side of* (53.10) *is bounded by the same expression with $\nu(E^+)$ replaced by $V_\varepsilon(D_m)$. After multiplication by m^2 it tends to zero, uniformly in R, n, k, h and the location of every roof interval of a fixed finite length. The full total singular volume also has zero normalized limit on this source.*

Proof. Use (53.3) and the exponential estimate in Theorem 53.1. For the whole volume the original full roof lies in $[m\tau_-, m\tau_+]$, so the Lebesgue measure of its support is at most $m(28/5 - 3/50)$. The factor $|I| + \sqrt{|I|}$ is therefore $O(m + \sqrt{m})$ and is absorbed by the same exponential decay. These are estimates for the actual singular positive volume used in the physical variation budget. A selected paired boundary contribution can use them only after its original variation/coarea inequality has been shown to charge this particular interior source. No such charge is inferred solely from the boundary being a bulk-null set. □

Corollary 53.5 (A genuine guarded physical-word pressure estimate)**.** *Let M'_R and j_m have precisely the true stable-arclength meaning in Input 11.1. Using the explicit optical theorem 54.3, the restricted physical-word pressure obeys*

$$\sum_{w\in\mathcal{W}_m(R)} \sup_{x\in[w]_R\cap M'_R\cap\mathcal{G}_\varepsilon} j_m(x) \le \frac{1500000R_+^2U^2C_{\mathrm{dist}}^2}{\rho^2} \quad (m\ge 1). \tag{53.11}$$

An empty supremum contributes zero. The estimate includes all realizable words and the actual stable norm; its constant is independent of R, m. Its restriction to $\mathcal{G}_\varepsilon$, including its endpoint cap, is explicit and does not discharge the unrestricted premise Q_*.

Proof. The two exact endpoint Jacobians give $u = R^2\omega/\det G$. The endpoint Schur bound gives $\det G \le R^4U^2$, hence $\omega \le R_+^2U^2u$. Theorem 54.3 gives $j_m \le 30000\omega/c_m \le 60000R_+^2U^2u$ on this class. For a fixed word, cover all its guarded points by the actual grid tubes of Lemma 52.2. Their upper distortion bounds show that its supremum is at most $60000R_+^2U^2C_{\mathrm{dist}}$ times the sum of its tube-center weights. This holds even if the supremum is not attained. The complete center sum is at most $25C_{\mathrm{dist}}\nu(M_R)/\rho^2 = 25C_{\mathrm{dist}}/\rho^2$. Multiplication yields (53.11). There is no physical-word cardinality or accumulated collision-dependent distortion factor. □

53.1. **The original ordered shortage and the proved new payment.** For the inherited persistent low-load *interior* source, Lemma 12.1 gives the exact positive domination $c_{\mathrm{per}} \le f^\varepsilon \le A$. At every fixed outer stage B, choose any fixed positive atlas margin $\varepsilon_B \le 1/4$. The restriction of that same source to $\mathcal{G}_{\varepsilon_B,m,R}\cap\mathcal{E}^{\mathrm{obs}}_{D_m,R,m}$ has zero m^2-normalized roof height by (53.5), uniformly in the original labels and central roof set. The source-side estimate is unconditional on pressure; its identification with the inherited pairing coefficient uses precisely that retained one-copy construction and domination.

All original positive source is preserved by the literal identity, at the source level,

$$\mathbf{1} = \mathbf{1}_{\mathcal{G}_{\varepsilon_B}\cap\mathcal{E}^{\mathrm{obs}}_{D_m}} + \mathbf{1}_{\mathcal{G}_{\varepsilon_B}\setminus\mathcal{E}^{\mathrm{obs}}_{D_m}} + \mathbf{1}_{\mathcal{G}^c_{\varepsilon_B}}.$$

Only the first term has just received this new payment. On the second term the full normalized central comparison, including all marks and its endpoint clock work, is proved above; the original shallow portions and roofwise comparison capacity still have their actual obligations. The third term is not declared negligible. The constant ρ^{-2} diverges when the fixed atlas margin tends to zero, so exhaustion of regular trajectories does not extend the payment to that third term. The full nongrazing pool $P_{r_B}^{\mathrm{ng}}$ is unchanged.

The new pointwise, singular-volume and guarded-pressure estimates therefore reduce concrete missing work without redefining the remaining target (15.8). All outer $\varepsilon(B), h_B, r_B, \gamma_B, K_B$, and the optional fixed ε_B, precede the collision limsup; B remains outermost. Uniform pressure without the stated guard and cap, the uncovered positive boundary source and its actual capacity, the complete paired/singular budgets, and the uniform spectral, arithmetic, conditioned microscopic and inversion inputs remain separate obligations. No full raw pointwise local limit or four-coordinate inverse theorem is claimed from these payments.

54. Explicit stable arclength and the original optical weight

The optical comparison in Input 11.1 has a direct proof with explicit constants. We give that proof in the true collision coordinates, including the sign of the physical derivative and the change to the original optical coordinates. It completes the two optical inequalities in that input. It does not prove its separate whole-cylinder pressure bound. The qualitative comparison and its proof in historical module 238 remain part of the retained source; the contribution here is a complete quantitative norm and cocycle verification.

Throughout this section the particle has unit speed, the scatterers are the original radius-R disks on the triangular lattice, and $R \in [9/20, 47/100]$. Only true regular first-collision trajectories are considered. Put

$$a = \frac{100}{47}, \qquad K = \frac{170}{9}, \qquad H = 19, \qquad \epsilon_{\mathrm{ang}} = \frac{2a}{H^2}, \qquad \tau_- = \frac{3}{50}, \qquad \Lambda = 1 + 2\tau_- a = \frac{59}{47}. \tag{54.1}$$

Here $a \le \varkappa = R^{-1} \le 20/9$ and $K = 20/9 + \tau_-^{-1}$; $1 + K^2 < H^2$. The constants $a, K, \epsilon_{\mathrm{ang}}$ are cone constants, unrelated to the original section probability c_* or the incidence guard q_0. Write $s = R\alpha$, $p = \sin\varphi$, and $c = \cos\varphi > 0$. Norms on tangent vectors in (s, φ) are Euclidean norms, with no adapted metric. The original probability is still $\nu = (4\pi R)^{-1} c\, ds\, d\varphi = d\alpha\, dp/(4\pi)$, and the section-conditioned probability is its original c_*^{-1} restriction. The radius is fixed along each collision cocycle, so this coordinate conversion introduces no factor depending on the collision count.

Lemma 54.1 (The physical differential and two cone directions)**.** *At a regular flight from x to $x' = T_R x$, write its physical length as τ, its incidences as c, c', and its endpoint curvatures as $\varkappa, \varkappa'$. With the counterclockwise boundary tangent and the above signed outgoing momentum,*

$$DT_R(x) = -M(x), \qquad M(x) = \frac{1}{c'} \begin{pmatrix} \tau\varkappa + c & \tau \\ \tau\varkappa\varkappa' + \varkappa c' + \varkappa' c & \tau\varkappa' + c' \end{pmatrix}, \qquad \det M(x) = \frac{c}{c'}. \tag{54.2}$$

On the set M_R' regular in both infinite time directions there are invariant unstable and stable lines with slopes

$$v^u(x) \in [a, K], \qquad v^s(x) \in [-K, -a].$$

They are measurable. If $e(x) = |\sin \angle(E^s(x), E^u(x))|$, then

$$\epsilon_{\mathrm{ang}} \le e(x) \le 1. \tag{54.3}$$

The positive arclength cocycle $j_m(x) = J^s T_R^m(x)$ satisfies

$$j_m(x) \le \frac{H}{\epsilon_{\mathrm{ang}}}\Lambda^{-m} \quad (m \ge 0). \tag{54.4}$$

Proof. Let $n(\alpha) = (\cos\alpha, \sin\alpha)$, $t(\alpha) = (-\sin\alpha, \cos\alpha)$, and let v be the unit direction of the actual flight. At departure and arrival, respectively,

$$v = cn + pt, \qquad v = -c'n' + p't'.$$

For a variation of the departure state, $\delta v = v^{\perp}(\delta\varphi + \varkappa\delta s)$. Differentiating $Q' - Q = \tau v$ and taking its scalar product with $v^{\perp}$ gives

$$-c'\delta s' - c\delta s = \tau(\delta\varphi + \varkappa\delta s).$$

The incoming direction angle at the arrival point is $\alpha' + \pi - \varphi'$; thus $\delta\varphi' = \varkappa'\delta s' - \varkappa\delta s - \delta\varphi$. These two identities prove (54.2), including its minus sign. Expanding the determinant of the displayed numerator leaves exactly cc'.

Scalar signs do not affect slopes. The map induced by $M(x)$ on an input slope $v \ge a$ is

$$f_x(v) = \varkappa' + \frac{c'(\varkappa + v)}{\tau(\varkappa + v) + c}. \tag{54.5}$$

Consequently $a \le f_x(v) \le \varkappa' + \tau^{-1} \le K$. On $[a, K]$ its derivative satisfies

$$0 < f_x'(v) = \frac{cc'}{[\tau(\varkappa + v) + c]^2} \le \frac{1}{4\tau(\varkappa + v)} \le \frac{1}{8\tau_- a} = \frac{47}{48} < 1. \tag{54.6}$$

The first inequality uses $c' \le 1$ and $(b + c)^2 \ge 4bc$ with $b = \tau(\varkappa + v) > 0$. The images at x of the cone interval $[a, K]$ transported from $T_R^{-n}x$ are nested nonempty compact intervals, with diameter at most $(K - a)(47/48)^n$. Their intersection is therefore one line $E^u(x)$, and shifting the sequence proves its invariance. Every finite approximation is measurable on the countable regular collision partition; its limit is measurable.

Time reversal in these coordinates is $I(s, \varphi) = (s, -\varphi)$ and $T_R^{-1} = IT_RI$ on the regular domain. Applying the same construction to reversed flights gives the invariant line $E^s(x)$ with negative slope. For unit vectors on these two lines,

$$e(x) = \frac{v^u(x) - v^s(x)}{\sqrt{1 + v^u(x)^2}\sqrt{1 + v^s(x)^2}} \ge \frac{2a}{H^2}.$$

This proves (54.3).

For completeness these invariant cone lines have the claimed stable character in the actual Euclidean norm. An unstable vector with s-component δs_i obeys

$$c_{i+1}|\delta s_{i+1}| = [\tau_i(\varkappa_i + v_i^u) + c_i]|\delta s_i| \ge \Lambda c_i|\delta s_i|,$$

since $c_i \le 1$. Hence, writing $u_m(x) = J^u T_R^m(x)$,

$$u_m(x) \ge \frac{c_0}{Hc_m}\Lambda^m. \tag{54.7}$$

Taking absolute areas of the images of the two unit invariant vectors, and multiplying the exact one-step determinants, gives

$$u_m(x) j_m(x) \frac{e(T_R^m x)}{e(x)} = \frac{c_0}{c_m}. \tag{54.8}$$

Combining (54.7) with this identity yields (54.4); its $m = 0$ case also holds because $j_0 = 1$. Reversal gives the corresponding backward contraction on E^u. Thus these are the true stable and unstable lines, not auxiliary test directions. No ergodic, pressure or mixing theorem has entered their construction. □

For $m \geq 1$ put

$$\mathsf{D}_m(x) = M(T_R^{m-1}x)\cdots M(x), \qquad b_m(x) = (\mathsf{D}_m(x))_{12} > 0.$$

This matrix product is distinct from the scalar physical-word partition sum $P_m(R)$ in module 266.

Lemma 54.2 (One comparison of columns, without accumulated constants)**.** *For every $m \geq 1$ and $x \in M_R'$,*

$$\frac{2a}{H}\, b_m(x) \leq u_m(x) \leq 2KH\, b_m(x). \tag{54.9}$$

The original inverse endpoint derivative is exactly

$$\omega(x) = |\partial_{p_0}\alpha_m|^{-1} = \frac{Rc_0}{b_m(x)} = \frac{Rc_0c_m}{\mathfrak{b}_m(x)}. \tag{54.10}$$

Here $\mathfrak{b}_m$ is the original Jacobi optical denominator of (4.6)*.*

Proof. For the two columns C_1, C_2 of one matrix $M(x)$, each component of C_1/C_2 lies in $[a, K]$. In the first row the ratio is $\varkappa + c/\tau$; in the second it is $\varkappa + \varkappa' c/(\tau\varkappa' + c')$. Left multiplication by a positive matrix preserves the componentwise inequalities $aC_2 \leq C_1 \leq KC_2$. They therefore hold for the columns of the whole product, with exactly the same constants.

The second column of the whole product has slope in $[a, K]$. For the first factor this follows from $(\tau\varkappa' + c')/\tau$; later factors preserve the cone. Consequently $b_m \leq |C_2| \leq Hb_m$. A unit unstable vector at the source is, up to sign, $(1, v)/\sqrt{1+v^2}$ with $v \in [a, K]$. Its image is, up to the common sign $(-1)^m$, $(C_1 + vC_2)/\sqrt{1+v^2}$. Positivity and the column inequalities give

$$\frac{2a}{H}b_m \leq \frac{a+v}{\sqrt{1+v^2}}|C_2| \leq u_m \leq \frac{K+v}{\sqrt{1+v^2}}|C_2| \leq 2KHb_m.$$

Only the initial column comparison and the terminal cone bound have been used; no constant is raised to the collision count.

With α_0 fixed, the coordinate conversions are $\delta\varphi_0 = \delta p_0/c_0$ and $\delta\alpha_m = \delta s_m/R$. Thus $|\partial_{p_0}\alpha_m| = b_m/(Rc_0)$. There is one factor R, irrespective of m. The agreement with the retained denominator can also be checked as an exact conjugacy. Set

$$B_i = \begin{pmatrix} c_i & 0 \\ \varkappa_i & 1 \end{pmatrix}, \qquad C_i = \begin{pmatrix} 1 & 0 \\ 2\varkappa_i/c_i & 1 \end{pmatrix}, \qquad S_i = \begin{pmatrix} 1 & \tau_i \\ 0 & 1 \end{pmatrix}.$$

Direct multiplication gives

$$B_{i+1}M_iB_i^{-1} = C_{i+1}S_i. \tag{54.11}$$

In the product, all intermediate B_i cancel. The top row of $B_m^{-1}C_m$ is $(1/c_m, 0)$ and the second column of B_0 is $(0,1)^{\mathsf{T}}$. Hence $b_m = \mathfrak{b}_m/c_m$, with precisely the internal optical factors of (4.6). For $m = 1$ this says $b_1 = \tau_0/c_1$ and $\mathfrak{b}_1 = \tau_0$, so the empty internal product is included. □

Theorem 54.3 (Explicit original-coordinate optical comparison)**.** *On M_R', for every $R \in [9/20, 47/100]$ and every $m \geq 1$,*

$$\frac{1}{30000}\, c_m(x)j_m(x) \leq \omega(x) \leq 30000\, c_m(x)j_m(x), \qquad \frac{j_1(x)}{c_0(x)} \leq 6400. \tag{54.12}$$

The same constants work for every original exact return, displacement and collision label, every physical roof interval, and all original initial and terminal section tests. In fact this comparison requires no positive endpoint cap beyond regularity. It includes trajectories arbitrarily close to interior or endpoint grazing.

Proof. Equations (54.8) and (54.10) imply the exact ratio

$$\frac{\omega(x)}{c_m(x)j_m(x)} = R\,\frac{u_m(x)}{b_m(x)}\,\frac{e(T_R^m x)}{e(x)}. \tag{54.13}$$

The previous two lemmas therefore place this ratio between $4R_-a^2/H^3$ and R_+KH^3/a. All constants are fixed numbers, and direct rational arithmetic gives

$$\frac{H^3}{4R_-a^2} < 30000, \qquad \frac{R_+KH^3}{a} < 30000, \qquad \frac{H^3}{4\tau_-a^2} < 6400. \tag{54.14}$$

These prove the first two inequalities in (54.12).

For one flight, the s-component of the image of a unit unstable vector and its terminal norm give

$$c_1u_1 = [\tau_0(\varkappa_0 + v_0^u) + c_0]\,\frac{\sqrt{1+(v_1^u)^2}}{\sqrt{1+(v_0^u)^2}} \geq \frac{2\tau_-a}{H}.$$

The one-step area identity now gives

$$\frac{j_1}{c_0} = \frac{e(x)}{c_1u_1e(T_Rx)} \leq \frac{H}{2\tau_-a\,\epsilon_{\mathrm{ang}}} = \frac{H^3}{4\tau_-a^2} < 6400.$$

There is no inverse incidence left over. None of these calculations discards an original event test: the inequalities hold before any such test is inserted. □

Corollary 54.4 (Precisely the remaining pressure premise)**.** *The two optical assertions in Input 11.1 hold with $C_0 = 30000$ and $C_1 = 6400$. Its remaining premise is*

$$\sup_{R,l} \sum_{w\in\mathcal{W}_l(R)} \sup_{x\in[w]_R\cap M_R'} j_l(x) \leq Q_*, \qquad P_0(R) = 1.$$

Under this separately retained whole-cylinder premise, all consequences of module 266 may take

$$C_\star = \frac{30000Q_*}{2c_*\lambda_\circ}, \qquad K_\star = 6400DQ_*.$$

The invariant-line restriction does not remove positive finite-record collision mass.

Proof. The theorem supplies exactly the two optical estimates, in the same norm convention on every nonempty stable block. On an empty block the cocycle remains $j_0 = 1$; the optical inverse derivative is used only for records with at least one flight and is not defined or invoked for $m = 0$. The finite marked-word factorization in module 266 therefore uses these constants without any change to its indices or empty blocks. It still requires its displayed pressure sum.

For each fixed radius the one-step singularities and grazing states form a finite union of regular arcs and their endpoints in each lifted collision chart. They have collision area zero. On every regular branch the determinant in (54.2) is nonzero. Countable regular coordinate charts and change of variables show that every forward and backward iterate of a one-step null singular set is null. Removing their countable union thus leaves M_R' with full collision measure. Restriction to the original finite event and pushforward by the original roof preserve equality of these positive bulk measures. Equivalently the local endpoint-coordinate map has its nonzero derivative in (54.10), so the corresponding two-dimensional area formulas are unchanged almost everywhere. No common null set for uncountably many radii is required by $\sup_R \operatorname{ess\,sup}_t$. This bulk statement assigns no zero trace to a one-sided singular boundary or to a paired endpoint measure. □

55. All physical margins in the original roof measure

The physical atlas has a fixed-margin height cost proportional to ε^{-2}. We now combine that cost with the exact distribution of the original incidence and clearance margins. The result includes arbitrarily small positive margins and every actual collision word. It is a quantitative regularity theorem for the original full-roof measure, not an exchange of the fixed-margin and collision limits. Its function-space conclusion is stated explicitly; it does not assert the unrestricted pointwise height required by the final local theorem.

Keep $q_0 = 47/53$, $r_g = q_0^{1/4}$, and let $\mathcal{C}_{m,R}$ be the regular physical m-flight records with $|p_0|, |p_m| \le 5/6$. This cap contains the complete original exact event, by Section 52. For a Borel physical source E and a Borel function $0 \le H \le 1$, put

$$\mu_{E,H} = (L_{m,R})_*(\mathbf{1}_E \mathbf{1}_{\mathcal{C}_{m,R}} H \, d\nu_R^*), \qquad \mu_{E,H}^I = (L_{m,R})_*(I_m \mathbf{1}_E \mathbf{1}_{\mathcal{C}_{m,R}} H \, d\nu_R^*). \tag{55.1}$$

In particular $H = \mathbf{1}_{\Pi_{n,k,m,R}} h$ retains all original return, displacement, collision, section and terminal labels and every selector $0 \le h \le 1$. The choice $H = 1$ is a positive majorant used in estimates, not a replacement of the original conditioning event. Throughout, $\nu_R^* = \mathbf{1}_{Y_R^*}\nu/c_*$ and $\nu(M_R) = 1$.

Use the following fixed constants:

$$\varepsilon_\sharp = 1/200, \qquad B_\Delta = \frac{2}{R_-(1-r_g)}, \qquad B_c = \frac{2r_g^2}{1-r_g^2},$$
$$B_\sharp = \frac{B_\Delta + B_c\varepsilon_\sharp}{c_*}, \qquad \mathfrak{A} = \frac{25UC_{\mathrm{dist}}^2}{2c_*}\left(\frac{128R_+}{g\delta_*}\right)^2, \qquad J_\sharp = 1 + \frac{4q_0}{1-q_0} = \frac{97}{3}. \tag{55.2}$$

Here δ_* is precisely the endpoint continuation constant in (33.2), $g = 9/40$, and $U = 320/9$. The auxiliary upper margin $\varepsilon_\sharp$ is smaller than the strict clearance-flux range $3/200$; it is not an outer microscopic width. In this notation $C_{\mathrm{at}} = \mathfrak{A}\varepsilon^{-2}$.

Lemma 55.1 (Uniform physical margin and incidence-insertion tails)**.** *For every $m \ge 1$, every physical radius, and $0 < \varepsilon \le \varepsilon_\sharp$,*

$$\nu(\mathcal{C}_{m,R} \setminus \mathcal{G}_{\varepsilon,m,R}) \le B_\Delta \varepsilon + B_c \varepsilon^2. \tag{55.3}$$

For every Borel $0 \le h \le 1$, with $a = \int h \, d\nu$, the actual unchanged insertion satisfies

$$\int h I_m \, d\nu \le J_\sharp \sqrt{a}, \qquad I_m = 1 + \sum_{i=1}^{m-1} \frac{q_0^{d_i}}{c_i}. \tag{55.4}$$

These estimates include all depths and need no independence of collisions, clearances or section visits.

Proof. On the stated endpoint cap, $c_0, c_m \ge \sqrt{11}/6 > 1/4$. Thus failure of an incidence condition of $\mathcal{G}_\varepsilon$ can occur only at an interior contact. At most two interior contacts have any depth $d \ge 1$, and at most two physical flights have any depth $d \ge 0$. The exact invariant incidence marginal gives $\nu\{c_i < u\} = 1 - \sqrt{1-u^2} \le u^2$ for $0 \le u \le 1$. Theorem B.2 gives $\nu\{\Delta_i < u\} \le u/R_-$ for $0 < u < 3/200$. Apply these statements to $u = \varepsilon r_g^{d_i}$ and $u = \varepsilon r_g^{d_i'}$ and sum the two geometric series. This proves (55.3). The clearance event uses every third disk on the whole closed physical flight, including disks visited at other times. Exact tangencies and visited grazing belong to collision-area null sets; invariance gives the same assertion at each of the finitely many actual indices. No positive pairing trace is identified with such a bulk-null set.

For the insertion, layer cake and $0 \le h \le 1$ give, at each interior index,

$$\int \frac{h}{c_i} \, d\nu = \int_0^\infty \int h \mathbf{1}_{\{c_i^{-1} > s\}} \, d\nu \, ds \le \int_0^\infty \min\{a, 1, s^{-2}\} \, ds = 2\sqrt{a}.$$

For $a = 0$ this follows directly; for $0 < a \le 1$ split the integral at $a^{-1/2}$. The bound s^{-2} for $s \ge 1$ is exactly the preceding invariant incidence marginal with $u = 1/s$; for $s < 1$, the bound by one suffices. As

$$\sum_{i=1}^{m-1} q_0^{d_i} \le 2 \sum_{d=1}^{\infty} q_0^d = \frac{47}{3},$$

we obtain $\int hI_m \le a + (94/3)\sqrt{a} \le (97/3)\sqrt{a}$. For $m = 1$ the interior sum is empty and the same inequality holds. □

For a nonnegative measurable roof function f, define the exact weak norm

$$\|f\|_{q,\infty} = \sup_{\lambda>0} \lambda |\{t \in \mathbb{R} : f(t) > \lambda\}|^{1/q}. \tag{55.5}$$

Lebesgue measure on the original physical roof line is used here. No averaging width or rescaled time is part of this definition.

Lemma 55.2 (Quantitative exhaustion after the physical coarea sum)**.** *Suppose a source E in* (55.1) *satisfies, for all $0 < \varepsilon \le \varepsilon_\sharp$,*

$$\left\| \frac{d\mu_{E,H}[\mathcal{G}_\varepsilon]}{dt} \right\|_\infty \le A_E \varepsilon^{-2}, \qquad \mu_{E,H}(\mathbb{R}) \le M_E \le 1. \tag{55.6}$$

Here the first bound is a completed positive physical sum, uniform in H, not an estimate on a chosen word. Then the full measure $\mu_{E,H}$ is absolutely continuous, with density $f_{E,H}$, and

$$\|f_{E,H}\|_{3/2,\infty} \le W_E,$$
$$W_E = \left[\sqrt{2A_E} \max\{2B_\sharp, M_E/\varepsilon_\sharp\} \right]^{2/3}. \tag{55.7}$$

For every Borel roof set S of finite measure,

$$\mu_{E,H}(S) \le \min\{M_E, 3W_E |S|^{1/3}\}. \tag{55.8}$$

The statements are also valid when $A_E = 0$, in which case $\mu_{E,H} = 0$ and $W_E = 0$.

Proof. The exact physical partition into $\mathcal{G}_\varepsilon$ and its complement, together with (55.3), gives

$$\mu_{E,H}(S) \le A_E \varepsilon^{-2} |S| + B_\sharp \varepsilon \quad (0 < \varepsilon \le \varepsilon_\sharp). \tag{55.9}$$

There is exactly one section-normalization factor in the second term: the complementary mass is at most $c_*^{-1}\nu(\mathcal{C} \setminus \mathcal{G}_\varepsilon)$. If S is a Lebesgue-null set, let ε decrease to zero in this finite-count inequality. This proves absolute continuity of the full measure, including all arbitrarily small margins. If $A_E = 0$, the same argument with $S = \mathbb{R}$, using the zero guarded measure, proves that the whole measure vanishes.

Suppose $A_E > 0$. Write $f_{E,H} = f_\varepsilon + g_\varepsilon$ for the densities of the two exact nonnegative restrictions. The proof of absolute continuity just given applies to both, or equivalently they are submeasures of the full absolutely continuous measure. We have $0 \le f_\varepsilon \le A_E \varepsilon^{-2}$ almost everywhere and $\|g_\varepsilon\|_1 \le B_\sharp \varepsilon$. For $\lambda \ge 2A_E \varepsilon_\sharp^{-2}$ take $\varepsilon = (2A_E/\lambda)^{1/2}$. Then

$$|\{f_{E,H} > \lambda\}| \le |\{g_\varepsilon > \lambda/2\}| \le 2B_\sharp \sqrt{2A_E} \lambda^{-3/2}.$$

For $0 < \lambda < 2A_E \varepsilon_\sharp^{-2}$, positivity and the total mass instead give

$$\lambda^{3/2} |\{f_{E,H} > \lambda\}| \le M_E \sqrt{\lambda} \le M_E \sqrt{2A_E}/\varepsilon_\sharp.$$

These two ranges prove (55.7) on the entire physical roof line, including arbitrarily large or small superlevel sets. No count of words, labels or roof components occurs. Finally layer cake gives, for $\ell = |S|$,

$$\int_S f_{E,H}\,dt \le \int_0^\infty \min\{\ell, (W_E/s)^{3/2}\}\,ds = 3W_E\ell^{1/3}.$$

The case $\ell = 0$ follows from absolute continuity. Together with the mass bound this proves (55.8). All margin choices were made inside inequalities valid at every finite collision count. No collision limsup was interchanged with an exhaustion or an essential supremum. □

Theorem 55.3 (Full physical roof densities and the actual incidence factor)**.** *The measures* (55.1)*, for $E = M_R$, have full original roof densities f_H, f_H^I. They satisfy*

$$\|f_H\|_{3/2,\infty} \le W_*, \qquad \|f_H^I\|_{6/5,\infty} \le J_\sharp\sqrt{3W_*/c_*}, \tag{55.10}$$

where W_* is (55.7) *with $A_E = \mathfrak{A}$ and $M_E = 1$. The constants are uniform in m, R and all original labels and selectors. Neither measure retains a positive lower incidence or clearance margin. More generally, every source covered by Lemma 55.2 satisfies*

$$\begin{aligned}\mu_{E,H}^I(S) &\le \frac{J_\sharp}{\sqrt{c_*}}\sqrt{\mu_{E,H}(S)},\\ \mu_{E,H}^I(\mathbb{R}) &\le \frac{J_\sharp}{\sqrt{c_*}}\sqrt{M_E},\\ \|f_{E,H}^I\|_{6/5,\infty} &\le J_\sharp\sqrt{3W_E/c_*}.\end{aligned} \tag{55.11}$$

In particular the insertion is the actual I_m, with the unchanged q_0 and collision indices, throughout the complete source.

Proof. For $E = M_R$, Theorem 52.4 gives the first premise of (55.6) with $A_E = \mathfrak{A}$. Its proof applies verbatim with an arbitrary $0 \le H \le 1$ in place of $\mathbf{1}_\Pi h$: the exact-event factor is retained in coarea and then bounded by one. The physical tubes, their multiplicity, their unconditioned lower volume, and the single factor c_*^{-1} are unaltered. The total source is a subprobability since ν_R^* is the original conditional probability. Lemma 55.2 therefore proves the unweighted assertion.

For a roof set S, apply (55.4) to the actual function

$$h = \mathbf{1}_{Y_R^*}\mathbf{1}_E\mathbf{1}_{\mathcal{C}_{m,R}}H\mathbf{1}_{\{L_{m,R}\in S\}}.$$

Its unconditioned integral is precisely $a = c_*\mu_{E,H}(S)$, so division by c_* gives the first inequality of (55.11). This displays the normalization without creating a second conditioning. It proves finiteness and absolute continuity of the weighted measure, since both properties hold for $\mu_{E,H}$. Combining it with (55.8) gives

$$\mu_{E,H}^I(S) \le J_\sharp\sqrt{3W_E/c_*}\,|S|^{1/6}.$$

For $S = \{f_{E,H}^I > \lambda\}$, which has finite measure by integrability, $\lambda|S| \le \mu_{E,H}^I(S)$ proves the weak $L^{6/5}$ bound. If $|S| = 0$ the assertion is immediate. This argument uses the distribution of the real incidence insertion on the same physical roof preimage; it neither assumes independence from the roof nor bounds I_m by a fixed-margin surrogate. □

Lemma 55.4 (The strong norms determined by mass and weak norm)**.** *If $f \ge 0$, $\|f\|_1 \le M$, $\|f\|_{q,\infty} \le W$, and $1 < p < q$, then*

$$\|f\|_p^p \le \frac{p(q-1)}{(p-1)(q-p)}M^{(q-p)/(q-1)}W^{q(p-1)/(q-1)}. \tag{55.12}$$

Proof. If $M = 0$ or $W = 0$ then $f = 0$ almost everywhere. Otherwise its distribution is at most $\min\{M/s, (W/s)^q\}$. Split $p\int_0^\infty s^{p-1}|\{f > s\}|\,ds$ at $s_0 = (W^q/M)^{1/(q-1)}$ and evaluate the two integrals. They converge precisely in the stated range and sum to (55.12). This proof applies on the whole roof line; its support length may grow with m. □

55.1. **The complete deep exceptional source, without a remaining guard.** Let $E_D = \mathcal{E}^{\mathrm{obs}}_{D,R,m}$ be precisely the actual clearance, incidence and original-section collar union of Theorem 50.3. In particular the original thresholds and $q_{\mathrm{a}} = 3263680/3263681$ are unchanged. For $D \ge D_{\mathrm{cv}}$ let $V_\varepsilon(D)$ be the six-term budget (53.3), including its squared-incidence factor 16, and define

$$A_D = \mathfrak{A}V_{\varepsilon_\sharp}(D), \qquad M_D = \min\{1, V_0(D)/c_*\}, \qquad W_D = \Big[\sqrt{2A_D}\max\{2B_\sharp, M_D/\varepsilon_\sharp\}\Big]^{2/3}. \tag{55.13}$$

Here V_0 denotes the explicit expression obtained by setting $a_\varepsilon = 0$ in that finite six-term budget. It is an upper bound for the actual unconditioned source mass, not a definition of a zero-margin tube.

Theorem 55.5 (All-margin normalized control of the exact deep source)**.** *For every $D \ge D_{\mathrm{cv}}$, all m, R and every Borel $0 \le H \le 1$, the full measures $\mu_{E_D,H}$ and $\mu^I_{E_D,H}$ have densities satisfying*

$$\begin{aligned} &\|f_{D,H}\|_1 \le M_D, && \|f_{D,H}\|_{3/2,\infty} \le W_D,\\ &\|f^I_{D,H}\|_1 \le J_\sharp\sqrt{M_D/c_*}, && \|f^I_{D,H}\|_{6/5,\infty} \le J_\sharp\sqrt{3W_D/c_*}. \end{aligned} \tag{55.14}$$

There is no factor $\mathbf{1}_{\mathcal{G}_\varepsilon}$ in these four measures. They contain the complete actual exceptional source, including its arbitrarily small margins and all relevant words.

Fix $0 < \theta < 1/2$, put $D_m = \lceil \theta m \rceil$, and retain $\lambda_b > 0$ from (53.6)*. Uniformly in the physical radius, exact labels and selectors, the bounds imply*

$$\begin{aligned} &\|f_{D_m,H}\|_{3/2,\infty} = O(e^{-\lambda_b m/3}), && \|f^I_{D_m,H}\|_{6/5,\infty} = O(e^{-\lambda_b m/6}),\\ &\|f_{D_m,H}\|_p = O(e^{-\lambda_b m(2/p-1)}), && 1 < p < 3/2,\\ &\|f^I_{D_m,H}\|_p = O(e^{-\lambda_b m(2/p-3/2)}), && 1 < p < 6/5. \end{aligned} \tag{55.15}$$

The respective L^1 rates are $O(e^{-\lambda_b m})$ and $O(e^{-\lambda_b m/2})$. Multiplication of any displayed norm by m^2 still gives a limit of zero.

Proof. For $0 < \varepsilon \le \varepsilon_\sharp$, the physical saturation estimate of Theorem 53.1 gives

$$\|d\mu_{E_D,H}[\mathcal{G}_\varepsilon]/dt\|_\infty \le \mathfrak{A}\varepsilon^{-2}V_\varepsilon(D) \le A_D\varepsilon^{-2}.$$

Its proof keeps every source point and every physical word; the extension to arbitrary H is the same positive bound as in Theorem 55.3. The depth D_{cv} ensures the strict flux and collar regimes simultaneously for every such ε. Setting the widening to zero in the actual invariant union estimate gives $\nu(E_D) \le V_0(D)$; the squared-incidence coefficient in V_0 is a harmless upper bound also without widening. Dividing once by c_* and using the subprobability bound gives M_D. Lemmas 55.1 and 55.2, and Theorem 55.3, prove (55.14).

Since $j(D_m) \ge \theta m/N - 2$, $N = 94$, the already established geometric series give $A_{D_m} = O(e^{-\lambda_b m})$ and $M_{D_m} = O(e^{-\lambda_b m})$. The maximum in (55.13) is bounded independently of m. Its square-root and $2/3$ powers give the first two rates. Apply (55.12) with $q = 3/2$ and then $q = 6/5$ to obtain the stated strong-norm exponents. The L^1 rates are already explicit in (55.14). All exponents are strictly positive in their specified ranges. Empty designated index sets contribute zero, and the finitely many counts with $D_m < D_{\mathrm{cv}}$ do not affect the conclusions. □

For example, for every fixed $a > 0$ the original normalized densities satisfy the quantitative roof-spike estimates

$$|\{t : m^2 f_{D_m,H}(t) > a\}| \le a^{-3/2} m^3 W_{D_m}^{3/2} = O_a(m^3 e^{-\lambda_b m/2}),$$
$$|\{t : m^2 f^I_{D_m,H}(t) > a\}| \le a^{-6/5} m^{12/5} [J_\sharp \sqrt{3W_{D_m}/c_*}]^{6/5} = O_a(m^{12/5} e^{-\lambda_b m/5}). \tag{55.16}$$

These are estimates for the possible roof spikes, not their elimination from an essential supremum. The sets are Borel after choosing measurable density versions. The constants are uniform in all original labels and the radius; no common null set across uncountably many radii is required.

55.2. **Exact marked comparison on the full source.** Use the actual central phases Φ_Q and Φ_Y, the endpoint work W_d, and the comparison coefficient $\widehat{\Phi}$ from Theorem 53.2. On E_d the comparison coefficient is zero; outside E_d it is $e^{-isW_d}\Phi_Y$, using the common physical central record of every global disk-word minimum. This construction is available on the complete actual source by Theorem 50.3, without an atlas margin.

Corollary 55.6 (Full-source marked comparison in the proved roof spaces)**.** *Fix $0 < \theta < 1/2$, $d = \lceil \theta m \rceil$, and restrict to the sufficiently large counts with $d \ge 1$ and $2d < m$. For the exact signed measure*

$$\mathfrak{D}_H = (L_{m,R}(Q))_*(\mathbf{1}_{\mathcal{C}_{m,R}} H(\Phi_Q - \widehat{\Phi})\, d\nu_R^*)$$

and its actual I_m insertion, positive domination gives

$$\|d\mathfrak{D}_H/dt\|_p \le \|f_{d,1}\|_p + 2R_+|s|q_{\mathrm{a}}^{\lfloor d/N \rfloor}\|f_1\|_p, \qquad 1 < p < 3/2,$$
$$\left\|d\mathfrak{D}^I_H/dt\right\|_p \le \|f^I_{d,1}\|_p + 2R_+|s|q_{\mathrm{a}}^{\lfloor d/N \rfloor}\|f^I_1\|_p, \qquad 1 < p < 6/5. \tag{55.17}$$

Every norm is taken on the unchanged full physical roof line. For $\lambda_c = \theta|\log q_{\mathrm{a}}|/N$ and any fixed $0 < \lambda < \lambda_c$, their m^2-normalized values tend to zero uniformly for all lattice and return twists and $|s| \le e^{\lambda m}$. The same assertions hold for $p = 1$ with the stated L^1 bounds. All original exact labels and terminal tests can be included in H.

Proof. Every integer mark and both separate endpoint tests agree on the actual good central comparison. Its true clock identity is $L_d(Y) - L_d(Q) = W_d + \mathsf{T}_{\mathrm{clock},d}$ with $0 \le \mathsf{T}_{\mathrm{clock},d} \le 2R_+ q_{\mathrm{a}}^{\lfloor d/N \rfloor}$. The minus sign in e^{-isW_d} therefore gives

$$|\Phi_Q - \widehat{\Phi}| \le \mathbf{1}_{E_d} + 2R_+|s|q_{\mathrm{a}}^{\lfloor d/N \rfloor}$$

on the complete actual source. The bound is independent of the lattice and return twists. Its pushforward dominates the total variation of the signed measure, and the same is true after the nonnegative actual I_m insertion. Theorem 55.3 and Lemma 55.4 give uniform strong bounds for the unexceptional full-source factors. Apply Theorem 55.5 to the first factors and use $\lfloor d/N \rfloor \ge \theta m/N - 1$ for the second. Minkowski's inequality, or ordinary total variation for $p = 1$, proves (55.17) and its limits. □

These conclusions concern the exact physical finite-record measures, with every original clock, collision and observation retained. They do not constitute an averaged replacement for a local limit theorem. They supply a new full-source regularity and integrability input, while the original raw pointwise target remains unchanged. In particular weak $L^{3/2}$ or $L^{6/5}$ control permits positive spikes on roof sets of small Lebesgue measure, so (55.16) cannot be promoted to an L^∞ estimate. The actual mixed multiplier $I_m/|\nabla_q F|$ requires more than the separate marginal argument above; its complete bulk estimate is proved in Section 56 by intersecting two actual physical momentum kernels. Full singular paired traces, short physical segments and the original shallow comparison capacity still require their own geometric estimates. The unrestricted pressure input and the

spectral, arithmetic and conditional inversion interfaces are not inferred from these roof spaces. Every outer B and its original microscopic widths remain fixed before the collision limit, with B outermost; no outer width was varied in (55.9).

56. The complete physical incidence–inverse-gradient multiplier

We prove a uniform integrability estimate for the actual multiplier $I_m/|\nabla_q F|$ on the entire physical source. The fixed-margin assumption in the previous singular-volume payment is absent. The mechanism uses the already proved endpoint-momentum kernel on two different blocks of the same orbit. Their intersection gives more information than two separate marginal tails; no product law or conditional independence is assumed.

Throughout $m \geq 1$, $R \in [R_-, R_+]$, and

$$\zeta = (p_0, p_m), \qquad r = |\zeta|, \qquad Z_m = \frac{I_m}{|\nabla_q F|} = \frac{I_m}{Rr}, \qquad I_m = 1 + \sum_{i=1}^{m-1} \frac{q_0^{d_i}}{c_i}.$$

The identity $|\nabla_q F| = Rr$ is the first variation of the actual reduced endpoint action, where $q = (\alpha_0, \alpha_m)$. No clock, Euclidean norm or radius factor is changed. Values on $r = 0$ and on undefined singular collision records can be assigned arbitrarily on their collision-area null sets; every nonnegative integral below is equivalently defined by truncating on regular records with $r > 0$. This convention concerns bulk integrals and assigns no value to a separate singular paired trace.

Let C_{mom} be the complete kernel constant of Theorem 26.4, with its already proved physical curve class, growth constant and all collision words. Set

$$A_0 = \frac{C_{\mathrm{mom}}}{4}, \qquad A_1 = \frac{C_{\mathrm{mom}}}{\pi}, \qquad K_\times = 5A_0^{1/3}A_1^{2/3}, \qquad J_\times = \frac{1}{R_-}\left(2\sqrt{A_0} + \frac{188}{3}K_\times^{3/4}\right). \tag{56.1}$$

These constants have no dependence on m, an incidence margin, an exact label, the target roof, or any outer microscopic width.

Lemma 56.1 (Intersecting two actual momentum kernels)**.** *For every $s, u > 0$ and $1 \leq i < m$, the unconditioned physical collision measure satisfies*

$$\nu\{r < s\} \leq A_0 s^2, \qquad \nu\{r < s,\ c_i < u\} \leq A_1 s u^2. \tag{56.2}$$

Both statements sum the complete actual source before taking any supremum in radius, collision count or the index i.

Proof. Integrating (26.15) in the actual initial momentum p_0 with $d\nu = d\alpha_0\, dp_0/(4\pi)$ gives

$$(p_0, p_l)_*\nu \leq \frac{C_{\mathrm{mom}}}{4\pi}\, da\, dv \qquad \text{for every integer } l \geq 1.$$

This is the unconditioned version of (26.17); no section factor belongs to it. Use $l = m$ and dominate the area of $\{(a, v) : a^2 + v^2 < s^2\}$ by πs^2. This proves the first inequality, for all positive s including $s > 1$.

For the second inequality, retain the actual block from collision 0 to collision i. Since $\{r < s\} \subset \{|p_0| < s\}$,

$$\nu\{r < s, c_i < u\} \leq \nu\{|p_0| < s, \sqrt{1 - p_i^2} < u\}.$$

The p_0 interval has length at most $2s$. If $0 < u \leq 1$, the second set of momenta has length $2(1 - \sqrt{1 - u^2}) \leq 2u^2$. If $u > 1$ its length is two, which is again at most $2u^2$. Apply the same kernel with $l = i$. The product of these two coordinate lengths is only a Lebesgue area under a proved joint-density upper bound. It is not a factorization of two physical probabilities. The

bound is $C_{\mathrm{mom}} s u^2/\pi$, as claimed. Future exact-record tests can be dropped only on this positive upper side; all collisions through m remain in the source being bounded. □

Lemma 56.2 (A mixed weak tail at every actual interior contact)**.** *For each $1 \le i < m$ and every $t > 0$,*

$$\nu\{(rc_i)^{-1} > t\} \le K_\times t^{-4/3}. \tag{56.3}$$

Consequently, for every Borel $0 \le h \le 1$ with $a = \int h\, d\nu$,

$$\int \frac{h}{rc_i}\, d\nu \le 4K_\times^{3/4} a^{1/4}. \tag{56.4}$$

Proof. Choose any $b > 0$. The part with $r < b$ has mass at most $A_0 b^2$. On the annulus $2^j b \le r < 2^{j+1} b$, the condition in (56.3) implies $c_i < (t2^j b)^{-1}$. Lemma 56.1, valid for every positive value of both arguments, therefore bounds this annulus by

$$A_1(2^{j+1}b)(t2^j b)^{-2} = \frac{2A_1}{t^2 b}\, 2^{-j}.$$

The annuli with $j \ge 0$ cover all $r \ge b$; most are empty when $2^j b > \sqrt{2}$, but keeping them only enlarges the upper sum. Thus the complete source satisfies

$$\nu\{(rc_i)^{-1} > t\} \le A_0 b^2 + \frac{4A_1}{t^2 b}.$$

Take $b = (A_1/A_0)^{1/3} t^{-2/3}$. Both terms are defined for all $t > 0$, and their sum is $5A_0^{1/3} A_1^{2/3} t^{-4/3}$. This choice is a substitution into a finite-count tail bound, not a collision-dependent residence or a changed physical event.

Layer cake now gives

$$\int \frac{h}{rc_i}\, d\nu \le \int_0^\infty \min\{a, K_\times t^{-4/3}\}\, dt = 4K_\times^{3/4} a^{1/4}.$$

For $a = 0$ use monotone truncation; for $a > 0$ split the integral at $(K_\times/a)^{3/4}$. The calculation also proves the finite first moment before it is used anywhere else. □

Theorem 56.3 (The full original singular multiplier is uniformly integrable)**.** *On the complete regular physical source, with no positive lower incidence or clearance margin, one has*

$$\int h \frac{I_m}{|\nabla_q F|}\, d\nu \le J_\times \left(\int h\, d\nu \right)^{1/4} \qquad (0 \le h \le 1). \tag{56.5}$$

In particular, the actual multiplier has the uniform weak bound

$$\nu\{Z_m > t\} \le (J_\times/t)^{4/3} \quad (t > 0), \qquad \int Z_m\, d\nu \le J_\times. \tag{56.6}$$

For every $1 < p < 4/3$ it also satisfies

$$\int Z_m^p\, d\nu \le \frac{4}{4 - 3p} J_\times^p. \tag{56.7}$$

All statements remain valid on the original conditional measure with the explicitly normalized form

$$\int h Z_m\, d\nu_R^* \le J_\times c_*^{-3/4} \left(\int h\, d\nu_R^* \right)^{1/4} \qquad (0 \le h \le 1). \tag{56.8}$$

Any original event and every original roof test may be retained in h; its values need not factor over time.

Proof. The first inequality of (56.2) gives $\nu\{r^{-1} > t\} \le A_0 t^{-2}$ for all $t > 0$. Thus

$$\int h/r \, d\nu \le \int_0^\infty \min\{a, A_0 t^{-2}\} \, dt = 2\sqrt{A_0}\, a^{1/2} \le 2\sqrt{A_0}\, a^{1/4}, \qquad a = \int h \, d\nu \le 1.$$

Expand the unchanged I_m, use (56.4) at every actual interior index, and sum the deterministic weights $\sum_{i=1}^{m-1} q_0^{d_i} \le 47/3$. There is no factor m, no independent collision law, and no marked-word pressure estimate in this calculation. Finally $|\nabla_q F| = Rr$ and $R \ge R_-$ give exactly $J_\times$ in (56.1). The empty interior sum for $m = 1$ is included.

For $h = \mathbf{1}_{\{Z_m > t\}}$, if its mass a is positive, $ta \le J_\times a^{1/4}$ proves (56.6); if $a = 0$ it is immediate. For $h = 1$ the first-moment estimate follows directly from (56.5). Integrating the distribution bound $\min\{1, (J_\times/t)^{4/3}\}$, split at $t = J_\times$, proves (56.7). For the conditional statement apply (56.5) to $\mathbf{1}_{Y_R^*} h$ under ν. Its integral is $c_* \int h \, d\nu_R^*$; division of the weighted integral by c_* leaves exactly $c_*^{-3/4}$. The kernel proves $\nu\{r = 0\} = 0$ and invariance proves the finite-record singular-null assertions. All operations can first be performed with bounded multipliers and then passed monotonically to the actual one; no divergent integral was used to justify them. □

56.1. **The actual insertion on complete critical-origin bands.** The same intersection has a sharper consequence when a physical critical-origin band is already known to lie in a momentum disk. It extends the full Borel band estimate of Theorem 28.5 to the actual I_m insertion.

Lemma 56.4 (Mixed incidence mass in a momentum disk)**.** *For every $a > 0$ and each $1 \le i < m$,*

$$\int_{\{r<a\}} c_i^{-1} \, d\nu \le 2\sqrt{A_0 A_1}\, a^{3/2},$$

$$\int_{\{r<a\}} (r c_i)^{-1} \, d\nu \le 6\sqrt{A_0 A_1}\, a^{1/2}. \tag{56.9}$$

Consequently

$$\int_{\{r<a\}} Z_m \, d\nu \le \frac{1}{R_-} \left(2A_0 a + 94\sqrt{A_0 A_1}\, a^{1/2}\right). \tag{56.10}$$

Proof. At every $t > 0$, (56.2) bounds the mass in the layer-cake integral by $\min\{A_0 a^2, A_1 a t^{-2}\}$. Its integral is $2\sqrt{A_0 A_1}\, a^{3/2}$, proving the first line of (56.9). Write its left side as $F_i(a)$. For $0 < r < a$ the identity $r^{-1} = a^{-1} + \int_r^a s^{-2} \, ds$ and Tonelli give

$$\int_{\{r<a\}} \frac{1}{r c_i} \, d\nu = F_i(a)/a + \int_0^a F_i(s) s^{-2} \, ds \le 6\sqrt{A_0 A_1}\, a^{1/2}.$$

The possible choice of strict or non-strict disk boundaries does not matter: the joint endpoint kernel gives zero mass to every circle. The same formula with $\nu\{r < a\} \le A_0 a^2$ gives $\int_{r<a} r^{-1} \, d\nu \le 2A_0 a$. Sum the actual incidence weights, using $6(47/3) = 94$, and divide by $R \ge R_-$ to prove (56.10). □

Corollary 56.5 (Full physical critical-band singular payment)**.** *Let E_s be any measurable one-copy family of actual critical-origin arm points in Theorem 28.5, with the same $0 < \rho < 1$ and $A_s = \min\{\rho, \sqrt{2U_m s}\}$. For every original exact event and Borel $0 \le h \le 1$,*

$$\int_{E_s} \mathbf{1}_\Pi h \frac{I_m}{|\nabla_q F|} \, d\nu_R^* \le \frac{2A_0 A_s + 94\sqrt{A_0 A_1}\, A_s^{1/2}}{c_* R_-}. \tag{56.11}$$

In particular the complete actual singular volume is $O(s^{1/4})$ on such bands as $s \downarrow 0$, uniformly in all collision counts, physical radii, actual arms and exact labels. No tangent-sector assumption or lower incidence/clearance margin is added.

Proof. The proved physical band geometry gives $E_s \subset \{r \le A_s\}$. Bound its selector by one and use (56.10), with the single original conditioning factor c_*^{-1}. No arm is extended beyond its actual physical boundary. As $U_m \le 320/9$, the two terms are respectively $O(s^{1/2})$ and $O(s^{1/4})$. □

56.2. **Full original-roof singular density and deep-source payment.** Use the complete source measures $\mu_{E,H}$ and densities $f_{E,H}$ from Section 55, without any factor $\mathbf{1}_{\mathcal{G}_\varepsilon}$. Define the actual singular-roof measure

$$\mu^Z_{E,H} = (L_{m,R})_*(Z_m \mathbf{1}_E \mathbf{1}_{\mathcal{C}_{m,R}} H \, d\nu^*_R). \tag{56.12}$$

The cap and exact-event conventions are unchanged.

Theorem 56.6 (Full singular-roof regularity and normalized exceptional volume)**.** *Whenever M_E, W_E are the full-source bounds of Lemma 55.2, the measure* (56.12) *is absolutely continuous with density $f^Z_{E,H}$ and satisfies, for every Borel roof set S,*

$$\begin{aligned} \mu^Z_{E,H}(S) &\le J_\times c_*^{-3/4} \mu_{E,H}(S)^{1/4}, \\ \|f^Z_{E,H}\|_1 &\le J_\times c_*^{-3/4} M_E^{1/4}, \\ \|f^Z_{E,H}\|_{12/11,\infty} &\le J_\times c_*^{-3/4} (3W_E)^{1/4}. \end{aligned} \tag{56.13}$$

For $E = M_R$ these are collision-uniform bounds for the entire actual singular source. For the complete exceptional union $E = \mathcal{E}^{\mathrm{obs}}_{D_m,R,m}$, $D_m = \lceil \theta m \rceil$, $0 < \theta < 1/2$, they imply uniformly in R and all original labels and selectors

$$\begin{aligned} \|f^Z_{D_m,H}\|_1 &= O(e^{-\lambda_b m/4}), \\ \|f^Z_{D_m,H}\|_{12/11,\infty} &= O(e^{-\lambda_b m/12}), \\ \|f^Z_{D_m,H}\|_p &= O(e^{-\lambda_b m(2/p-7/4)}), \qquad 1 < p < 12/11. \end{aligned} \tag{56.14}$$

Every displayed norm has zero limit after multiplication by m^2. In particular the full positive singular volume of this source, and its volume in every original roof interval, receive an unconditional normalized payment without a remaining guard class.

Proof. Apply (56.8) to $h = \mathbf{1}_E \mathbf{1}_{\mathcal{C}_{m,R}} H \mathbf{1}_{\{L_{m,R} \in S\}}$. This proves the first bound. Since $\mu_{E,H}$ is finite and absolutely continuous, the same is true of $\mu^Z_{E,H}$. In particular a bulk-null critical fiber creates no atom of this weighted pushforward. It is a different matter to estimate a separately defined paired endpoint measure.

For $|S| < \infty$, (55.8) gives

$$\mu^Z_{E,H}(S) \le J_\times c_*^{-3/4} (3W_E)^{1/4} |S|^{1/12}.$$

Use the superlevel set of the nonnegative density to obtain the weak $L^{12/11}$ bound, exactly as in the proof of (55.11). The total mass follows by setting $S = \mathbb{R}$ in the first inequality. For the exceptional source, Theorem 55.5 gives $M_{D_m} = O(e^{-\lambda_b m})$ and $W_{D_m} = O(e^{-\lambda_b m/3})$. Apply the indicated fourth powers and then Lemma 55.4 with $q = 12/11$. The exponent in the strong pth power is $(12 - 11p)/4 + (p-1) = 2 - 7p/4$, as stated. The exponent $2/p - 7/4$ is positive throughout the indicated range. The total-volume bound controls every roof interval without changing its width or location. □

Corollary 56.7 (The complete regular-interior variation budget)**.** *Use exactly the retained countable one-copy physical sheet space of Proposition 30.6, with all original exact-event and*

section cuts. For an original guard parameter $\varepsilon > 0$, a Borel physical source E and a physical selector $0 \le H \le 1$, define the nonnegative roof measure

$$\mathcal{V}_{\varepsilon,E,H}(S) = \int \mathbf{1}_S(u)\mathbf{1}_E(\Psi(\theta,u))H(\Psi(\theta,u))|\partial_u f^{\varepsilon}(u,\theta)|\,du\,d\theta.$$

The derivative is taken only on the regular interior of each actual sheet, before multiplying by any indicator. Then

$$\mathcal{V}_{\varepsilon,E,H}(S) \le C_{\mathrm{reg}}(1+\varepsilon^{-1})\mu^{Z}_{E,\mathbf{1}_\Pi H}(S) \qquad \text{for every Borel roof set } S. \tag{56.15}$$

In particular the complete regular-interior variation is finite, uniformly in collision count and radius, with total bound

$$C_{\mathrm{reg}}(1+\varepsilon^{-1})J_{\times}c_*^{-3/4}.$$

Its density has the corresponding full weak $L^{12/11}$ and strong L^p, $1 < p < 12/11$, bounds. On $E = \mathcal{E}^{\mathrm{obs}}_{D_m,R,m}$, all the rates in (56.14) *hold for this variation density, multiplied by the fixed factor $C_{\mathrm{reg}}(1+\varepsilon^{-1})$. Consequently its m^2-normalized total variation and all those roof norms tend to zero. In the original application $\varepsilon = \varepsilon(B)$ is fixed before the collision limit.*

Proof. Apply Proposition 30.6 to the nonnegative pullback selector $\mathbf{1}_S(u)\mathbf{1}_E(\Psi(\theta,u))H(\Psi(\theta,u))$. The retained physical one-copy coarea identifies its upper side with precisely the right side of (56.15). The exact event is unchanged, and its endpoint tests put it inside $\mathcal{C}_{m,R}$ as required in (56.12). Theorem 56.6 proves finiteness and the asserted norms. Positivity and Radon–Nikodym differentiation of the measure inequality transfer its density bounds to the variation measure. For the deep source, apply (56.14) after the complete physical sum. Neither H nor an exact section or roof indicator is differentiated.

This argument uses exactly the inherited one-copy coarea and the already native regular-variation inequality; it does not reconstruct a new pairing. On a whole sheet the existing absolute-continuity and trace result identifies the interior derivative integral with the ordinary variation of f^{ε} on its closed physical segment. The present localized statement is the restriction of that derivative measure, not the derivative of an indicator-cut function. It therefore does not count jumps between different segments, manufacture a singular trace charge, or bound a positive endpoint minimum by an unrelated small bulk set. □

Corollary 56.8 (The actual singular insertion in the full marked comparison)**.** *Use the exact full-source marked comparison of Corollary 55.6. With its actual additional multiplier $Z_m = I_m/|\nabla_q F|$, one has*

$$\left\| d\mathfrak{D}^{Z}_{H}/dt \right\|_p \le \|f^{Z}_{d,1}\|_p + 2R_+|s|q_{\mathrm{a}}^{\lfloor d/N \rfloor}\|f^{Z}_{1}\|_p, \qquad 1 \le p < 12/11. \tag{56.16}$$

Its m^2-normalized value tends to zero uniformly in all lattice and return twists and $|s| \le e^{\lambda m}$, for every fixed $0 < \lambda < \lambda_c$. Every collision and the signed endpoint clock work are retained, with the same negative compensating phase.

Proof. Multiply the actual pointwise phase domination in the proof of Corollary 55.6 by the nonnegative Z_m. Theorem 56.6 and Lemma 55.4 give the required full-source strong bounds and exceptional-source decay. The remaining clock factor decays at rate $\lambda_c - \lambda > 0$. No Fourier integration over an unbounded range is asserted. □

The physical significance of (56.5) is that the two possible singular mechanisms, an interior grazing contact and a vanishing full-roof gradient, have a quantified joint bound on the real orbit. Separate marginal estimates would give only a borderline product and would not prove this first moment. The intersection of two already valid physical block kernels is the additional mechanism. It completes the full bulk singular integrability and the complete deep-source singular-volume

payment. It does not bound a positive short-segment endpoint trace by that volume without its own proved charge inequality. Nor does weak $L^{12/11}$ roof control establish a pointwise roof supremum. The original shallow positive source, full comparison capacity, singular pairing interfaces and the final spectral, arithmetic and conditional pointwise inversion requirements retain their stated scope and the original order of limits.

57. All-time momentum kernels of the true marked clearance source

The first-image source theorem and its admissible growth theorem are already established in Sections 21 and 25. We retain those proofs and constants. The new conclusions below concern every later or earlier physical momentum, joint clearance–incidence control, and the one-sided marked source at zero clearance. They do not identify momentum with a Markov state.

All radii belong to $[9/20, 47/100]$, the speed is one, and $\delta_* = 3/200$. Write $a_\circ = \sqrt{1827}/200$ for the foot standoff of Lemma 19.1, $k_- = 100/47$ and $t_* = 3/50$. The constants M_*, A_*, C_g, J_f have exactly their meanings in Sections 21–25; A_* includes both impact signs. Fix

$$H_{\mathrm{nm}} = \max\left\{\frac{2M_*}{k_- a_\circ}, \frac{A_* C_g J_f}{k_- t_*}\right\}. \tag{57.1}$$

No curve-class enlargement is needed: these are the original angular-source graphs, not the different initial momentum fibres of Section 26.

For $u > 0$, start at the actual marked foot $-\sigma(R+u)v_\vartheta^\perp$, with direction v_ϑ, and let $P_l^+(\vartheta, \sigma, u)$ be the tangential momentum at the lth actual collision after that foot, $l \geq 1$. Let P_l^- be the tangential momentum in the original forward record at the lth collision before the foot. Thus its sign is opposite to the outgoing momentum of the time-reversed record. All visited disks and all intervening flights are the actual ones. There is no collision, reflection, or change of the physical clock at the mark.

Theorem 57.1 (A uniform angular kernel at every actual collision)**.** *For every R, $0 < u \leq \delta_*$, $l \geq 1$, Borel set $I \subset (-1, 1)$ and source selectors $0 \leq h_\sigma \leq 1$,*

$$\sum_{\sigma=\pm1} \int_0^{2\pi} h_\sigma(\vartheta) \mathbf{1}_{\{P_l^\pm \in I\}}\, d\vartheta \leq H_{\mathrm{nm}} |I|. \tag{57.2}$$

The integrals are over the regular finite records; their complement is angular-null. For either direction and every $v \in (-1, 1)$, the complete sum over distinct regular angular roots satisfies

$$\sum_{\sigma=\pm1} \sum_{\vartheta:\, P_l^\pm(\vartheta,\sigma,u)=v} \frac{1}{|\partial_\vartheta P_l^\pm|} \leq H_{\mathrm{nm}}. \tag{57.3}$$

Artificial seams and subdivisions do not create extra roots. Selectors may retain any of the original exact records or original physical-roof Borel tests. The constants are independent of l, the full collision words, the radius, clearance, and selectors.

Proof. For the first true collision, Theorem 21.1 gives $ds/d\vartheta = -a/c$ and $d\varphi/ds = \kappa + c/a$. Consequently

$$\frac{dP_1^+}{d\vartheta} = c\frac{d\varphi}{ds}\frac{ds}{d\vartheta} = -(\kappa a + c), \qquad |\partial_\vartheta P_1^+| \geq k_- a_\circ.$$

There are at most M_* first-hit intervals for each sign. On every such interval the momentum is strictly monotone. Direct substitution therefore proves (57.2) for $l = 1$ with the first constant in (57.1), without cutting the target momentum into infinitely many grazing strips.

For $l \geq 2$, use the exact, source-labelled first-image measure of Theorem 21.3. Its sum of density suprema is A_*. Propagate each of these homogeneous short unstable curves through exactly $l - 2$ further physical collisions. Theorem 25.1 bounds the complete sum of inverse

arclength Jacobians by C_g. On each resulting incoming short graph, leave one actual collision to perform the terminal coarea. The exact cancellation (26.18) gives

$$\left|\frac{dp_{\mathrm{next}}}{d\ell}\right| \geq k_- t_*, \qquad \int_C \mathbf{1}_{\{p_{\mathrm{next}}\in I\}}\, d\ell \leq \frac{J_f}{k_- t_*}|I|.$$

The small terminal incidence cancels before estimating; the last coarea is cut into its finitely many physical next-hit branches, with no terminal homogeneous subdivision. Sum the labelled densities and all intermediate growth pieces by Tonelli. This gives $A_* C_g J_f/(k_- t_*)$ and proves the forward inequality. A Borel selector is retained until the positive upper bound and incurs no component count or derivative.

At each finite stage the singular-null coverage of Theorem 25.1, followed by the finite last-hit cuts, applies to the absolutely continuous labelled source. The omitted angular set is null. The momentum derivative is nonzero on every regular complete branch. Coarea gives the root formula almost everywhere in v. To obtain the bound at every regular root value, fix a finite set of distinct roots at v_0, choose disjoint regular source intervals, and use their continuous positive inverse derivatives. Their sum is bounded almost everywhere in a common neighborhood of v_0, hence also at v_0. Taking the supremum over finite root sets proves (57.3).

Finally time reversal sends (ϑ, σ, u) to $(\vartheta + \pi, -\sigma, u)$ and preserves $d\vartheta$ and the two-sign sum. On collision coordinates it is $\mathcal{I}(\alpha, p) = (\alpha, -p)$. Thus $P_l^- = -P_l^+(\vartheta + \pi, -\sigma, u)$ in the reversed record. Apply the forward result to $-I$, carrying the original selector through this bijection. This does not replace the original section by its reversal: the section tests are still inside that transported selector. □

For a Borel clearance set $E \subset (0, \delta_*]$, let $N_E(x)$ count the distinct nonincident disks whose original flight has clearance in E. An interval convention at its endpoints is immaterial to the collision-area identity, but will be retained in the one-sided trace below. For an m-flight record and its marked flight j, write $x = X_{R,j} = T_R^{-j} X_R$ as in Section E. At collision index $0 \leq i \leq m$, the momentum is obtained after exactly $j - i + 1$ backward collisions when $i \leq j$, and after $i - j$ forward collisions when $i \geq j + 1$. These are both positive integers, also at the two record endpoints.

Corollary 57.2 (Joint clearance and any physical collision momentum)**.** *For every $m \geq 1$, $0 \leq j < m$, $0 \leq i \leq m$, every original exact event $\Pi = \Pi_{n,k,m,R}$, $0 \leq h \leq 1$, and Borel sets $E \subset (0, \delta_*]$, $I \subset (-1, 1)$,*

$$\int h \mathbf{1}_\Pi N_E(T_R^j x) \mathbf{1}_{\{p_i \in I\}}\, d\nu_R^*(x) \leq \frac{H_{\mathrm{nm}}}{4\pi R c_*}|E|\,|I|. \tag{57.4}$$

The same domination holds with any Borel original-roof selector with values in $[0, 1]$, including the indicator of an arbitrarily small physical time interval. It is a joint domination of a marked counting measure, not a conditional independence statement.

Proof. Use invariance of the original collision area to apply the exact one-copy flux identity (B.2) at the marked flight j. Transport $h\mathbf{1}_\Pi$ and all its true record information to $X_{R,j}$. Conditioning contributes exactly c_*^{-1}, and the physical line Jacobian contributes exactly $(4\pi R)^{-1}$. For each fixed $u \in E$, use Theorem 57.1 with the positive collision count just specified for p_i. Integrate the resulting bound in u. Tonelli first on simple clearance sets and then on their Borel extension proves the statement. Every distinct marked disk is counted once by the flux bijection; no collision-word multiplicity is estimated by hand. □

Corollary 57.3 (Uniform singular-incidence moments on every clearance source)**.** *Let*

$$w_i = q_0^{\min(i,m-i)}, \quad q_0 = 47/53, \quad I_m = 1 + \sum_{i=1}^{m-1} w_i/c_i, \quad W_0 = 1 + \frac{2}{1-q_0}.$$

For $1 \le \gamma < 2$, define the finite number

$$J_\gamma = \int_{-1}^{1} (1-v^2)^{-\gamma/2}\, dv, \qquad B_\gamma = 4\pi + \frac{2H_{\mathrm{nm}} J_\gamma}{1-q_0}.$$

Then, uniformly in every original exact label, m, j, R, and $0 < \delta \le \delta_*$,

$$\int h \mathbf{1}_\Pi N_\delta(T_R^j x) I_m^\gamma \, d\nu_R^* \le \frac{\delta W_0^{\gamma-1} B_\gamma}{4\pi R c_*}, \tag{57.5}$$

$$\int h \mathbf{1}_\Pi N_\delta(T_R^j x) \mathbf{1}_{\{c_i<\eta\}} \, d\nu_R^* \le \frac{H_{\mathrm{nm}}\delta}{2\pi R c_*}\eta^2, \qquad 0 < \eta \le 1. \tag{57.6}$$

For arbitrary nonnegative depth coefficients b_j and widths $0 < \delta_j \le \delta_*$, the right side of (57.5)*, summed against b_j, is obtained by replacing δ by $\sum_j b_j \delta_j$. This includes every shallow flight, with no incidence-load cutoff.*

Proof. For each fixed angular clearance fibre, Theorem 57.1 and monotone approximation give

$$\sum_\sigma \int c_i^{-\gamma}\, d\vartheta \le H_{\mathrm{nm}} J_\gamma.$$

The integral J_γ is finite precisely in the stated range: near either endpoint $1 - v^2$ is comparable to $1 - |v|$. The angular mass of both signs is 4π. Weighted convexity with the extra weight one for the constant term gives

$$I_m^\gamma \le \left(1 + \sum_{i=1}^{m-1} w_i\right)^{\gamma-1} \left(1 + \sum_{i=1}^{m-1} w_i c_i^{-\gamma}\right).$$

Since $\sum_i w_i \le 2/(1-q_0)$, integrating this bound on the complete angular source gives $W_0^{\gamma-1} B_\gamma$. Apply the exact flux and its original conditional factor. This proves (57.5). The set $c_i < \eta$ has momentum length $2(1 - \sqrt{1-\eta^2}) \le 2\eta^2$; insert this set into (57.4) to prove (57.6). The last statement follows by summing nonnegative measures; it requires neither independence nor separation of the marked flights. □

57.1. **A genuine one-sided unvisited-tangency source.** The next theorem defines a boundary measure, not a new value of the collision map at a singular point. Fix R, m, j. Let $u \downarrow 0$ in the positive-clearance family above. A limiting strict record lists the actual visited collisions of this family. The marked tangency at the foot stays an unvisited boundary event. At $u = 0$ we never use the putative collision map obtained by adding or deleting a reflection at that foot.

Theorem 57.4 (Exact marked trace with original finite record tests)**.** *For angular-almost every (ϑ, σ) there is an open one-sided neighborhood in u on which the fixed finite forward and backward strict word is constant and all its visited collisions are regular. Its contacts, momenta and physical roof extend real-analytically to $u = 0$. Every original finite record selector $\mathbf{1}_{\Pi_{n,k,m,R}}$ has a one-sided limit there. The same is true after multiplying by the indicator of any specified physical roof interval, with its original open or closed endpoint convention.*

Denote the limiting momentum by p_i^0 and the one-sided selector by π^+. For any bounded continuous function f of the finite physical record, including its actual roof, the positive marked trace is the actual limit

$$\lim_{u\downarrow 0} \frac{1}{4\pi R c_*} \sum_\sigma \int f(\mathrm{record}_u)\pi_u \, d\vartheta = \frac{1}{4\pi R c_*} \sum_\sigma \int f(\mathrm{record}_0)\pi^+ \, d\vartheta. \tag{57.7}$$

More generally the equality holds for any bounded test with an angular-almost-everywhere one-sided limit. The endpoint kernels (57.2) *and the angular moment bounds used in* (57.5) *hold at* $u = 0$ *as well, with the same constants. In particular the marked trace marginal satisfies*

$$(p_i^0)_{\#}\left(\frac{\pi^+}{4\pi R c_*}\sum_{\sigma} d\vartheta\right) \le \frac{H_{\mathrm{nm}}}{4\pi R c_*}\, dv. \tag{57.8}$$

No null set common to uncountably many radii is asserted.

Proof. At $u = 0$ the foot has distance R from its marked disk and distance at least $1 - R \ge 1 - R_+ > R_+$ from every other center. The two open tangent rays miss the interior of the marked disk. The same finite candidate geometry and standoff proof as in Theorem 21.1 applies to their first nonmarked true hits. Except for finitely many angular tangencies and seams, these hits are regular. Their derivatives, first-image densities, and curvature bounds are the continuous one-sided versions of (21.2)–(21.4); the strict standoff $a \ge a_\circ$ is unchanged. The first family is therefore the same admissible homogeneous class, with the same summable allowance A_*.

Iterate each first-image branch through the finite number of required true hits in both directions. At each stage the physical growth partition removes a countable angular-null set of tangencies and artificial boundaries, exactly as in the proof of Theorem 57.1. Away from this set all finitely many visited incidences are positive. On each such designated record, the line–circle intersection with nonzero incidence and the specular reflection formula are real-analytic functions of the incoming state and u. After the first true hits, the marked disk is restored as an ordinary possible next scatterer. Any later tangency with it or another disk is part of the same growth singular set. Every nonincident disk therefore has positive clearance on this finite strict record, except for the prescribed original foot tangency; the finite-horizon candidate reduction makes this a finite visibility check. Hence the same true word persists for all sufficiently small positive u. The marked flight continues to miss its marked disk because its prescribed clearance is $u > 0$. This establishes the asserted one-sided analytic record. It uses no common neighborhood size over words, m, or radii.

The lattice displacement is constant on this true word. Each of the $m + 1$ original section tests is a finite Boolean combination of strict inequalities in the real-analytic coordinates $\alpha_i(u), p_i(u)$, using a local circular lift as necessary. The visit count includes $i = 0, \dots, m - 1$ and the terminal test remains separate. A real-analytic function of u at zero is either identically zero near zero or has the sign of its first nonzero Taylor coefficient for all sufficiently small positive u. Thus every such strict or nonstrict inequality has a one-sided indicator limit. The same argument applies to $L_m(u) - a$ and $L_m(u) - b$ for any specified roof interval with endpoints a, b. If a boundary equality persists identically, its membership is exactly its prescribed endpoint convention. This does not identify the trace with evaluation of an arbitrarily redefined section on a singular record.

All limits in (57.7) now follow by dominated convergence on the finite angular measure. To prove the kernel at $u = 0$, one can repeat the already justified first-image, growth, and last-collision coarea proof on these limiting strict branches. Alternatively, apply the $u > 0$ domination to nonnegative continuous momentum tests and pass to zero; finite measure regularity then gives domination for every Borel set. Finally use monotone truncation of $c_i^{-\gamma}$ and the same weighted convexity inequality to obtain the angular moment bound. All tests with values in $[0, 1]$ only restrict a positive measure. □

The new scope and the remaining normalization. Theorem 57.4 supplies the complete labelled one-sided angular source needed for an unvisited clearance boundary. Its measure is positive; angular-null exceptional parameters are not assigned zero charge in an unrelated endpoint pairing. The trace is not the original roof–Gauss-angle endpoint charge. The exact conversion to that charge includes its physical normal speed, which is proved in the next section. Neither the kernel

nor the moment bounds insert an m^{-2} factor into a physical-roof density. The original supremum, essential supremum and ordered limit in (0.1) are retained.

58. The actual endpoint charge in physical flux coordinates

The physical radius, unit-speed clock, original section, exact collision, return and lattice labels, conditioning by c_*, and incident-component pairing are unchanged. In this section $t = F$ is physical roof time, $\theta = \arg \nabla_q F$ is its Gauss angle, and u is a signed local clearance parameter. In particular u is never substituted for t. The original bad-source weight is $W^\varepsilon = 1 - H^\varepsilon$; it equals one at a zero unvisited-disk clearance and is not set to zero on that boundary. For a positive clearance level $u_0 > 0$ we specify one incident side of that level. If both sides are counted, their charges must be added explicitly. The original unvisited tangency $u_0 = 0$ has the prescribed physical side $u > 0$.

Consider a one-sided regular physical marked-flight chart from Theorem B.2, with fixed flight index j, marked disk, sign ς, and actual collision word. Put

$$q = \mathcal{Q}_j(\vartheta, u), \quad S(\vartheta, u) = F(q), \quad \theta(\vartheta, u) = \arg \nabla_q F(q).$$

No reflection is made at the line's closest point. At $u = 0$ all quantities below are one-sided limits of these actual positive-clearance records. In particular a limiting tangency is not counted as an extra collision, and the collision index is not shifted. Initially suppose all visited collisions of the limiting word are nongrazing and the endpoint gradient is nonzero. The original endpoint cap is retained.

Theorem 58.1 (Exact positive boundary flux)**.** *On a smooth boundary piece $u = u_0$ the original roof–angle trace is*

$$\frac{\omega W^\varepsilon}{4\pi c_* \chi} |\partial_\vartheta \theta| \, d\vartheta = \frac{W^\varepsilon}{4\pi R c_*} \left| \frac{\partial u}{\partial t}\Big|_\theta \right| d\vartheta. \tag{58.1}$$

The formula is an identity of positive measures, with the actual boundary multiplicity. It therefore holds after a nonnegative Borel restriction of already-defined incident boundary germs, including their original record labels. It does not assign a trace to a new Borel selector that has no one-sided boundary value.

For an original section side at collision i, take $(v, b) = (p_i, \alpha_i - a)$ or $(\alpha_i, p_i - a)$ on the corresponding regular physical chart. The same identity is

$$\frac{\omega W^\varepsilon}{4\pi c_* \chi} |\partial_v \theta| \, dv = \frac{W^\varepsilon}{4\pi c_*} \left| \frac{\partial b}{\partial t}\Big|_\theta \right| dv. \tag{58.2}$$

Thus the clearance formula has precisely one additional factor R^{-1}. There is no additional conditional normalization in either formula.

Proof. Write $J = \det D_{(\vartheta,u)}(S, \theta)$ with its actual local sign. The inverse of this two-by-two derivative matrix gives

$$\frac{\partial u}{\partial t}\Big|_\theta = -\frac{\partial_\vartheta \theta}{J}.$$

The exact long-record determinant in (16.2) is $|J| = \chi/(R\omega)$. Substitute it into the preceding identity and multiply by the physical line density $(4\pi R c_*)^{-1}$. This proves (58.1), including its radius factor.

The section formula follows in the same way from invariance of $d\alpha \, dp$, rather than from a line-coordinate density. Its source density in (v, b) is $(4\pi c_*)^{-1}$. The ordinary Jacobian chain rule gives $|\det D_{(v,b)}(t, \theta)| = \chi/\omega$. Matrix inversion now gives (58.2).

The identities can first be tested on compact boundary subarcs where the boundary-to-angle map is monotone. Exhaust its open set of nonzero derivative by countably many such subarcs,

disjointizing their source overlaps. A C^1 real map sends its zero-derivative set to a Lebesgue-null set: cover a compact portion of that set by intervals on which the derivative has magnitude at most η, bound the total image length by their total length times η, and let $\eta \downarrow 0$. This also proves the needed one-dimensional change of variables without an injectivity assumption on the full boundary. Nonnegative monotone convergence proves the identity for the full indicated boundary family.

Distinct simultaneously active marked disks remain distinct marked copies. To represent a boundary counted only once by the original pairing, choose one active constraint by a fixed finite ordering on each finite physical word and restrict to its Borel incidence class. The full marked sum is a positive upper bound before that restriction. No physical collision word is multiplied by an unspecified chart count. □

Proposition 58.2 (The transverse speed and its physical depth)**.** *Let $z = |\nabla_q F|$, $e = \nabla_q F/z$, and $G = D_q^2 F$. On a regular Gauss segment,*

$$V := \left.\frac{dq}{dt}\right|_\theta = \frac{G^{-1}e}{z\, e^{\mathsf{T}} G^{-1} e}, \qquad \|V\|_\infty \le \frac{\Lambda_\circ}{\lambda_\circ z}. \tag{58.3}$$

For the clearance from any one of the actual nonincident disks of flight j whose foot is interior, at all regular points and one-sided nongrazing limits,

$$\left| \left.\frac{\partial u}{\partial t}\right|_\theta \right| \le \frac{2C_\Delta \Lambda_\circ}{\lambda_\circ R_-} \frac{\sigma^{d'_j}}{|(p_0, p_m)|}, \qquad d'_j = \min(j, m-1-j). \tag{58.4}$$

This is a bound on a physical normal speed, not an identity replacing that speed by its upper bound near a critical point.

Proof. Differentiate $\nabla F = ze$ at fixed e, and use $dt/dz = ze^T G^{-1} e$. The Hessian bounds in (4.3) give (58.3). The segment Hausdorff argument in Proposition 29.2 applies separately to each fixed candidate disk as well as to their minimum. It gives $|D_q u[\eta]| \le 2C_\Delta \sigma^{d'_j} \|\eta\|_\infty$. Combine this with (58.3) and the exact identity $z = R|(p_0, p_m)|$. All differentiations are on the designated physical word. Continuity gives the stated one-sided nongrazing boundary limit. □

58.1. **A complete noncritical boundary estimate from the new physical angular kernel.** The next result uses the all-time angular kernel of Theorem 57.1 and its one-sided extension in Theorem 57.4. Their constant H_{nm} includes both impact signs. At $u = 0$ this is the same one-sided unvisited tangency family as above. Both time directions are included. This is an angular source measure, not the original conditioned collision probability or an independent momentum law.

Let $\mathfrak{B}_{j,u_0}$ be the positive marked boundary charge in (58.1), restricted to the original endpoint cap and any original incident boundary labels, and put $r = |(p_0, p_m)|$.

Theorem 58.3 (The full marked boundary away from the critical core)**.** *For every $0 < \rho \le 1$, every $u_0 \in [0, 3/200]$, and every finite physical count,*

$$\mathfrak{B}_{j,u_0}\{r \ge \rho\} \le C_{\mathrm{bf}}\, \sigma^{d'_j} (1 + \log(\sqrt{2}/\rho)), \qquad C_{\mathrm{bf}} = \frac{C_\Delta \Lambda_\circ H_{\mathrm{nm}}}{\pi c_* \lambda_\circ R_-^2}. \tag{58.5}$$

Both signs and the complete true marked physical family are included. Consequently for arbitrary choices $u_j \in [0, 3/200]$,

$$\sum_{j : d'_j \ge D} \mathfrak{B}_{j,u_j}\{r \ge \rho\} \le \frac{2C_{\mathrm{bf}} \sigma^D}{1-\sigma} (1 + \log(\sqrt{2}/\rho)). \tag{58.6}$$

In particular, for fixed $0 < \vartheta_* < 1/2$ and $a > 0$, choosing $D_m = \lceil \vartheta_* m \rceil$ and $\rho_m = e^{-am}$ in this finite-count inequality gives convergence to zero after multiplication by m^2. This is convergence of a positive boundary charge in total mass, and hence of its physical-roof pushforward in total variation. It does not assert an L^∞ roof density for that measure.

Proof. The complete two-sign kernel at the backward endpoint gives

$$\sum_{\varsigma=\pm1} \int \mathbf{1}_{\{r<s\}}\, d\vartheta \le \sum_{\varsigma=\pm1} \int \mathbf{1}_{\{|p_0|<s\}}\, d\vartheta \le 2H_{\mathrm{nm}} s.$$

The earlier endpoint is reached after $j+1 \ge 1$ actual collisions backwards from the closest point. Thus the kernel applies also to $j = 0$. The complete angular measure satisfies $A(s) \le 2H_{\mathrm{nm}} s$. Keeping original record tests only decreases this nonnegative measure. With $L = \sqrt{2}$ and $r \le L$, the elementary identity $r^{-1} = L^{-1} + \int_r^L s^{-2}\, ds$ gives

$$\int_{r\ge\rho} \frac{d\vartheta}{r} \le \frac{A(L)}{L} + \int_\rho^L \frac{A(s)}{s^2}\, ds \le 2H_{\mathrm{nm}}(1 + \log(L/\rho)).$$

Use (58.1), $0 \le W^\varepsilon \le 1$, and (58.4). The prefactor is $(4\pi R_- c_*)^{-1}(2C_\Delta \Lambda_\circ/(\lambda_\circ R_-))\, 2H_{\mathrm{nm}}$, which is the displayed C_{bf}. This fixes both radius factors and the single c_*^{-1}. No independence of the two endpoint momenta has been used.

There are at most two flight indices of each depth, so summing the geometric series proves (58.6). At the stated finite choices of D_m, ρ_m, the right side after multiplication by m^2 is bounded by a constant times $m^2 \sigma^{\vartheta_* m}(1 + am + \log\sqrt{2})$, tending to zero. This uses an inequality valid for all finite D, ρ, m and does not change a microscopic roof window or the original order of outer and collision limits. The entire omitted critical core still remains an original positive source; no claim about its smallness is made here. □

58.2. **Cancellation of the apparent pole at an actual critical boundary germ.**

Theorem 58.4 (A finite angular trace at a nongrazing critical germ)**.** *Let $q(s)$ be a C^2 parametrization of a one-sided smooth physical boundary germ with $q(0) = q_*$, $q'(0) \ne 0$, and all visited incidences of the limiting word positive. An unvisited tangency or an original regular section side is allowed. Assume $\nabla_q F(q_*) = 0$. Use the smooth extension of this designated stationary word only to take one-sided derivatives; do not interpret its nonphysical side as an actual trajectory. Put*

$$a = G_* q'(0), \qquad b = G_* q''(0) + D_q^3 F(q_*)[q'(0), q'(0), \cdot\,].$$

Then $a \ne 0$, and the derivative of every continuous local lift of the Gauss angle on either punctured side has the finite limit

$$\lim_{s\to0} \frac{d}{ds} \arg \nabla_q F(q(s)) = \frac{\det(a,b)}{2|a|^2}. \tag{58.7}$$

Consequently the positive boundary charge in (58.1) *or* (58.2)*, when expressed in this boundary parameter, has a finite one-sided density at the germ. The limiting density is*

$$\frac{\omega_* W_*^\varepsilon}{4\pi c_*\, e_a^{\perp\mathsf{T}} G_* e_a^\perp} \frac{|\det(a,b)|}{2|a|^2}, \qquad e_a = a/|a|. \tag{58.8}$$

The two antipodal limits of the Gauss angle have the same Hessian denominator. This is a trace of a boundary curve, not the different incoming critical-fan coefficient of Theorem 27.4.

Proof. Write $g(s) = \nabla_q F(q(s))$. The stationary word is smooth in a neighborhood of the limiting nongrazing contact configuration by its positive interior Hessian. The physical one-sided derivatives therefore give

$$g(s) = as + \tfrac{1}{2} bs^2 + o(s^2), \qquad g'(s) = a + bs + o(s).$$

Since G_* is positive definite and $q'(0) \neq 0$, $a \neq 0$. The angular derivative is exactly $\det(g, g')/|g|^2$ away from zero. Its numerator is $\frac{1}{2}\det(a, b)s^2 + o(s^2)$ and its denominator is $|a|^2 s^2 + o(s^2)$. Division proves (58.7). The finite one-sided optical and guard limits have already been proved in Theorem 5.3. Substitute them into the original coefficient f^ε and use $\chi = e^{\perp T} G e^\perp$. Replacing e by $-e$ does not change this quadratic form. This proves (58.8).

For the clearance formula one can also see the cancellation before expanding: the boundary defining function B is constant along $q(s)$, hence $DB_* q'(0) = 0$. As $s \to 0$, the leading direction of $G^{-1}e$ is parallel to $q'(0)$. Thus the leading numerator in $DB\, G^{-1}e/z$ vanishes. The expansion above computes the remaining finite term, and does not merely delete its apparent $1/z$ bound. □

Proposition 58.5 (A quantitative local turning bound)**.** *Suppose the preceding germ is parametrized by endpoint-angle arclength on a one-sided interval, with $|q'| = 1$ and $|q''| \leq K$. Put $L_G = 2C_F + \Lambda_\circ K$. On that same physical boundary interval, for $0 < |s| \leq \lambda_\circ / L_G$,*

$$\left|\frac{d\theta}{ds}\right| \leq \frac{2\Lambda_\circ L_G}{\lambda_\circ^2}. \tag{58.9}$$

If $L_G = 0$, the angular derivative is zero there. The constants $C_F, \lambda_\circ, \Lambda_\circ$ are the completed physical constants of Sections 29 and 5; the boundary curvature allowance K must be proved on the actual boundary and is not asserted uniform here.

Proof. The physical third-variation bound gives $\|dG/ds\|_{\mathrm{op}} \leq 2C_F$. Consequently $|g''(s)| \leq 2C_F + \Lambda_\circ K = L_G$ and $|g'(s)| \leq \Lambda_\circ$. As $|g'(0)| = |G_* q'(0)| \geq \lambda_\circ$,

$$|g(s) - s g'(0)| \leq L_G s^2/2, \qquad |g(s)| \geq \lambda_\circ |s|/2$$

on the stated interval. Use the exact identity $g(s) = \int_0^s g'(v)\, dv$ to bound

$$|\det(g(s), g'(s))| \leq \int_0^{|s|} \Lambda_\circ L_G(|s| - v)\, dv = \tfrac{1}{2}\Lambda_\circ L_G s^2.$$

Division by the squared lower bound for $|g(s)|$ proves (58.9). The negative side has the same proof with integration orientation reversed. If $L_G = 0$, g' is constant and the determinant is zero identically. □

The uniform continuation of this cancellation. The angular kernel controls $\int \mathbf{1}_{\{r<s\}} d\vartheta$ only by $O(s)$, which explains the logarithm in (58.5). The local cancellation just proved has a uniform extension on the actual zero-offset critical-clearance family in Theorem 60.2 below. That theorem proves a common formal gradient domain, cancels the quadratic radial factors before estimating, and sums the complete actual angular source without a word-count constant. The unrestricted nonzero-offset and uncovered boundary pieces remain a separate geometric obligation; bulk weak $L^{4/3}$ integrability does not supply it. The incoming critical fan at $g = 0$ also retains its distinct original source.

59. Two actual endpoint momenta from a marked free-flight foot

Fix the original radius R, $m \geq 1$, a true flight $0 \leq j < m$, and one nonincident marked disk with signed impact $\varsigma(R + u)$, $0 < u \leq 3/200$. All m collisions and all physical flights retain their original indices. The closest point is a place to parametrize the same line, with no reflection and no added collision. The limiting case $u = 0$ below always means the one-sided limit of this actual positive-clearance family on a fixed word with positive limiting visited incidences.

The two true endpoint momenta are p_0, p_m. Put $r = |(p_0, p_m)|$, and denote by a_-^{foot} and a_+^{foot} the first distances from the foot in the backward and forward directions. By Lemma 19.1,

$$a_\circ = \sqrt{1827}/200 \leq a_\pm^{\mathrm{foot}} \leq \tau_+.$$

There are $j+1$ actual collisions from the foot to the initial collision in the reversed flow and $m-j$ actual collisions from the foot to the terminal collision in the forward flow. Both counts are at least one, including when $m=1$.

For either direction propagate the unfolded transverse displacement and angle field to the incoming side of its endpoint collision. With the positive physical matrices of Theorem 19.2, write

$$M_\pm = P(\ell_{n-1})K_{n-1}\cdots P(\ell_1)K_1P(a_\pm^{\mathrm{foot}}) = \begin{pmatrix} A_\pm & B_\pm \\ C_\pm & D_\pm \end{pmatrix}. \tag{59.1}$$

Here n is the corresponding positive half-count. For $n=1$ the product is just $P(a_\pm^{\mathrm{foot}})$. The endpoint collision is not included as a kick in this incoming-side product. Let $\kappa_\pm = 1/R$ and $c_\pm > 0$ be its physical curvature and normal incidence, and define

$$A_\pm^p = \kappa_\pm B_\pm + c_\pm D_\pm, \qquad B_\pm^p = \kappa_\pm A_\pm + c_\pm C_\pm, \qquad h_\pm = B_\pm^p/A_\pm^p. \tag{59.2}$$

The superscript p distinguishes these positive scalar momentum derivatives from entries of $M_\pm$.

Theorem 59.1 (An exact two-sided physical momentum determinant)**.** *All scalars in* (59.2) *are positive and*

$$\frac{1}{\tau_+ + R_+} \le h_\pm \le \frac{1}{a_\circ}. \tag{59.3}$$

On the original marked chart (ϑ, u) *there are locally fixed signs* $s_0, s_m \in \{-1,1\}$ *such that*

$$\begin{aligned} \partial_\vartheta p_0 &= s_0 A_-^p, \quad & \partial_u p_0 &= \varsigma s_0 B_-^p, \\ \partial_\vartheta p_m &= s_m A_+^p, \quad & \partial_u p_m &= -\varsigma s_m B_+^p. \end{aligned} \tag{59.4}$$

Consequently

$$\left|\det D_{(\vartheta,u)}(p_0,p_m)\right| = A_-^p A_+^p (h_- + h_+) > 0. \tag{59.5}$$

The signs cannot cancel the positive sum. No lower cutoff on any positive visited incidence is required.

Proof. The reflection matrices were derived from the genuine reflection rule in Theorem 19.2. All their entries are nonnegative and their determinants are one. A positive initial free-flight length makes $B_\pm > 0$, while $A_\pm, D_\pm > 0$ throughout. Hence both scalars in (59.2) are positive.

At a terminal incoming collision write the incoming direction as $v^- = -cn + pt$ and use arclength displacement $\delta q = t\delta s$. For a physical incoming transverse field (J, J') one has $J = -c\delta s$. Differentiating $p = v^- \cdot t$ gives

$$\delta p = -c\delta\vartheta^- + \kappa c\delta s = -(\kappa J + cJ').$$

Unfolding a reflection reverses the signs of both transverse field coordinates together. Thus the actual endpoint momentum is a locally fixed sign times the positive row (κ, c) applied to the propagated field.

The physical foot is $-\varsigma(R+u)v_\vartheta^\perp$. Its ϑ derivative is parallel to v_ϑ, so the forward transverse initial field is $(0,1)$. Its u derivative is $-\varsigma v_\vartheta^\perp$, giving $(-\varsigma, 0)$. In the reversed direction $-v_\vartheta$, the angular derivative is again one, while the same transverse displacement is now $(\varsigma, 0)$. The momentum of a reversed collision is the negative of the original momentum. This changes the entire backward endpoint row by one common sign and not its column ratio. Applying the incoming-side products proves (59.4). Taking their determinant proves (59.5).

It remains to prove the uniform ratios without using a condition number of the whole product. Write a half-product as $M = LP(a^{\mathrm{foot}})$. If $(X,Y) = (\kappa, c)L$, then $X > 0$, $Y \ge 0$, and

$$h = \frac{X}{a^{\mathrm{foot}}X + Y} = \frac{1}{a^{\mathrm{foot}} + Y/X}.$$

For one collision, $L = I$, so $Y/X = c/\kappa \le R_+$. For at least two collisions, the rightmost factor of L is the first actual reflection kick K_1. Write $(x, y) = (\kappa, c)\widehat{L}$, where $L = \widehat{L}K_1$. Then $x > 0$, $y \ge 0$ and

$$(X, Y) = (x + yk_1, y), \qquad k_1 = 2/(Rc_1), \qquad \frac{Y}{X} \le \frac{1}{k_1} \le \frac{R_+}{2}.$$

This includes $y = 0$, for which the ratio is zero. The standoff and horizon bounds now give (59.3). The proof never bounds a grazing kick above; a smaller positive incidence only increases that positive kick. □

Corollary 59.2 (The exact endpoint kernel weight and its Wronskian)**.** *Let $u_x = |\partial_{\alpha_0} p_m|^{-1}$ be the actual endpoint momentum density weight in Theorem 28.2. On the same physical marked chart,*

$$u_x = \frac{1}{RA_-^p A_+^p (h_- + h_+)}, \qquad u_x \le \frac{\tau_+ + R_+}{2RA_-^p A_+^p}. \tag{59.6}$$

For $r > 0$ and $\theta = \arg(-p_0, p_m)$ one has the exact identity

$$u_x |\partial_\vartheta \theta| = \frac{|s_m p_0/A_-^p - s_0 p_m/A_+^p|}{R(h_- + h_+) r^2}. \tag{59.7}$$

The original angle–roof trace is therefore

$$f^\varepsilon |\partial_\vartheta \theta|\, d\vartheta = \frac{W^\varepsilon}{4\pi c_* h_x} \frac{|s_m p_0/A_-^p - s_0 p_m/A_+^p|}{R(h_- + h_+) r^2}\, d\vartheta, \qquad h_x = \frac{R^2 \chi}{\det G}. \tag{59.8}$$

This uses the same h_x and normalization as the exact radial coarea theorem. In particular $h_- + h_+$, a sum of the two half-ray ratios, is not confused with h_x, the endpoint-momentum metric.

Proof. The one-copy line measure is $d\nu = (4\pi R)^{-1} d\vartheta\, du$. The actual endpoint-momentum measure is $u_x (4\pi)^{-1} dp_0\, dp_m$, by Theorem 28.2. Apply the local determinant (59.5) and compare the two actual densities. This proves the equality in (59.6). Its inequality uses $h_- + h_+ \ge 2/(\tau_+ + R_+)$.

The derivative of the polar angle of $(-p_0, p_m)$ has absolute value $|p_0 \partial_\vartheta p_m - p_m \partial_\vartheta p_0|/r^2$. Substitute (59.4) and cancel the product $A_-^p A_+^p$ by (59.6). This gives (59.7). Finally the exact determinant of $q \mapsto (p_0, p_m)$ is $\det G/R^2$, so $u_x = R^2 \omega / \det G$. Hence $f^\varepsilon = W^\varepsilon u_x/(4\pi c_* h_x)$, proving (59.8). All these comparisons were between the same unconditioned collision measure before the one factor c_*^{-1} was inserted. □

What the identity supplies. Equations (59.6)–(59.8) are unconditional identities and bounds on the actual regular physical marked family; they do not use a whole-word pressure theorem, mixing, or a near-miss kernel bound. They keep the potentially useful signed cancellation between the two endpoint terms. Bounding those two terms separately recovers a possible $1/r$ loss and does not prove the complete critical boundary sum. At a shared actual critical germ, both signed first-order terms in the numerator agree, and their difference cancels. A uniform weighted summation of that cancellation over arbitrary short physical boundary components is a further estimate, not a consequence of the positive determinant alone.

60. A uniform cancellation on actual critical-clearance boundary pieces

The physical model and all exact record selectors remain unchanged. The result below concerns a complete explicitly specified part of the original one-sided clearance boundary. It uses a common domain of the formal reflection equations already proved in Theorem 33.2; it does not declare that formal domain to be a physical neighborhood or to preserve a section test. For $u_0 > 0$ fix

one incident side of the level; a two-sided charge is the explicit sum of two such measures. At $u_0 = 0$ the physical side is the original positive-clearance side.

Keep $\lambda = 50/47$, and put $\gamma = \lambda^{-1} = 47/50$. Let $\eta = \eta_*$ and $P = P_*$ be exactly the analytic constants in (33.1). Define the following positive constants, all independent of collision count and word:

$$\begin{aligned} \underline{g} &= R_-/2 = 9/40, & L_z &= 28/5 + R_+ + 3/200 = 1217/200, \\ K_F^{\mathrm{an}} &= 2048R_+/\eta^2, & a_{\mathrm{an}} &= \min\{\eta/16, \underline{g}/(2K_F^{\mathrm{an}})\}, \\ b_{\mathrm{an}} &= \underline{g}\, a_{\mathrm{an}}/4, & L_B &= P(4R_+ + 12(R_+/\tau_-)L_z), \\ K_{B,1} &= 6L_B, & K_{B,2} &= 128L_B/\eta, \\ K_{B,g} &= K_{B,2}/\underline{g}^2 + K_{B,1}K_F^{\mathrm{an}}/\underline{g}^3, & C_{\mathrm{zf}} &= \Lambda_\circ K_{B,g}/(R_- c_*). \end{aligned} \tag{60.1–60.2}$$

The constant $28/5$ is the enclosing true-flight bound in Theorem 2.2; it does not replace the intrinsic physical horizon or the physical clock.

Fix a limiting physical critical configuration Q_* of a designated m-flight word, with endpoint gradient zero and the original endpoint cap. It is the limit of actual regular trajectories of that word. Its endpoint incidences are one, and any number of interior incidences may be zero. Fix a nonincident marked disk on flight j whose perpendicular foot is interior and whose limiting signed impact is $\varsigma(R + u_0)$, $u_0 \in [0, 3/200]$. Let $q_* = (\alpha_0^*, \alpha_m^*)$ in fixed local lifts. The formal analytic reflection lift based at this configuration defines the formal action F and the signed line-clearance function

$$B(q) = \varsigma\,(z_{\mathrm{mark}} - x_j(q)) \cdot v_j(q)^\perp - R - u_0. \tag{60.3}$$

Thus $\nabla_q F(q_*) = 0$ and $B(q_*) = 0$ exactly. At every actual selected point the same expression is the true clearance from that marked disk minus u_0. Its use at intermediate formal points is an analytic calculation, not a visibility assertion.

Lemma 60.1 (Uniform formal gradient disk and signed-clearance tensors)**.** *On the real endpoint ball $|q - q_*| < a_{\mathrm{an}}$ the formal Hessian satisfies*

$$\underline{g} I \preceq G(q) \preceq 2\Lambda_\circ I, \qquad \|D_q G(q)[h]\|_{\mathrm{op}} \le K_F^{\mathrm{an}}|h|. \tag{60.4}$$

Every $|g| < b_{\mathrm{an}}$ has a unique formal inverse $q(g)$ in this ball, and it lies in $|q(g) - q_*| < a_{\mathrm{an}}/4$. Put $\mathcal{B}(g) = B(q(g))$. For all these gradients,

$$\|D_g^2 \mathcal{B}(g)\|_{\mathrm{op}} \le K_{B,g}\, \gamma^{d_j'}, \qquad d_j' = \min(j, m - 1 - j). \tag{60.5}$$

No positive lower bound on an interior incidence or on an unvisited clearance is required for this formal statement.

Proof. The limiting-reference version of Theorem 33.2 gives a holomorphic lift on $|h_0| + |h_m| < \eta$. Consider the polydisk $|h_0|, |h_m| < \eta/4$. Its weighted contact increment is at most $P\eta/2 = r_*/8$. All flight complexifications remain in the nonvanishing disks of Lemma 33.1. More explicitly, the two complex components X^+, X^- differ from their real reference values by at most a quarter of the reference flight length. Choose the square root of X^+X^- that is positive at the reference point. Its magnitude is at least three quarters of that reference length, and the holomorphic unit direction has complex Euclidean norm at most $5/3$. Circular normals and tangents have norm less than two on the contact ball. The endpoint first-variation identity therefore gives a holomorphic gradient with each component bounded by $4R_+$. This bound is independent of m: it is a bound for $R(-p_0, p_m)$, not for the total action.

For completeness, the first-variation identity continues to the formal lift because its reflection equations imply vanishing interior tangential derivatives of the same analytic action. It can also

be obtained by holomorphic continuation of the identity on the original regular branch. No intermediate formal contact is asserted physical.

Cauchy's formula on the smaller polydisk $|h_0|, |h_m| < \eta/8$ bounds every second partial derivative of each gradient component by $2(4R_+)(8/\eta)^2 = 512R_+/\eta^2$. For a two-by-two Hessian derivative, summing its two entries in each row and the two input coordinates bounds its Euclidean operator norm by four times this number times $|h|$. This is precisely K_F^{an} in (60.1).

At the critical physical limit $G_* \succeq RI \succeq 2\underline{g}I$, by the continuous reduced Hessian formula, and $G_* \preceq \Lambda_\circ I$. Integrating the preceding derivative bound along a real endpoint segment of length at most a_{an} changes the Hessian by at most $\underline{g}/2$. This proves (60.4), with room in both inequalities. For a prescribed $|g| < b_{\mathrm{an}}$, minimize $F(q_* + h) - g \cdot h$ on $|h| \le a_{\mathrm{an}}$. Strict convexity holds by (60.4). On the boundary its outward radial derivative is at least $\underline{g}a_{\mathrm{an}}^2 - |g|a_{\mathrm{an}} > 0$. The minimizer is therefore interior and is the unique inverse of the gradient in that ball. Strong monotonicity then gives $|q(g) - q_*| \le |g|/\underline{g} < a_{\mathrm{an}}/4$. The inverse function theorem makes it smooth on the whole gradient disk, and differentiation yields

$$\|D_g q\|_{\mathrm{op}} \le \underline{g}^{-1}, \qquad \|D_g^2 q\|_{\mathrm{op}} \le K_F^{\mathrm{an}} \underline{g}^{-3}. \tag{60.6}$$

We next check the depth factor for the individual marked clearance. This is a bound for its signed line coordinate, so no differentiability of a least-clearance minimizer is assumed. Let $s = |h_0| + |h_m|$ and $d = d_j'$. The weighted analytic localization gives

$$|\alpha_j(h) - \alpha_j^*| + |\alpha_{j+1}(h) - \alpha_{j+1}^*| \le Ps\gamma^d.$$

The corresponding complex position difference is at most $2R_+Ps\gamma^d$. Along the straight contact increment, all complex direction differentials have norm at most

$$\frac{1 + (5/3)^2}{3\tau_-/4}\,|dX| = \frac{136}{27\tau_-}|dX| < \frac{6}{\tau_-}|dX|.$$

Thus the flight-direction difference is at most $12(R_+/\tau_-)Ps\gamma^d$. The reference vector from departure contact to the marked center has length at most L_z. In (60.3), bound the position increment against the new direction, whose norm is less than two, and the direction increment against this fixed reference vector. The result is

$$|B(q_* + h) - B(q_*)| \le L_B s\gamma^d \le L_B\eta\gamma^d/2$$

on the larger polydisk. Cauchy's formula on the smaller polydisk therefore gives Euclidean derivative bounds

$$|D_q B| \le 6L_B\gamma^d = K_{B,1}\gamma^d, \qquad \|D_q^2 B\|_{\mathrm{op}} \le 128L_B\eta^{-1}\gamma^d = K_{B,2}\gamma^d.$$

The first constant bounds $4\sqrt{2}L_B$; for the second, each second partial is at most $64L_B/\eta$, and the two-by-two operator norm is at most twice that bound. Composing these inequalities with (60.6), keeping both second-order chain-rule terms, proves exactly (60.5). □

Call an actual one-sided boundary point *covered by a critical zero-offset chart* if its designated word and marked disk agree with one of the preceding limiting critical configurations, it has $B(q) = 0$, and its full contact configuration agrees with that formal inverse $q(g)$ at some $0 < |g| < b_{\mathrm{an}}$. Agreement here includes all $m + 1$ contacts, not only the two endpoint angles. The point must itself be an actual regular one-sided boundary record with all original physical cuts retained. The formal inverse supplies no new collision event. This definition contains the actual portions of every such critical boundary germ within the common formal gradient disk, irrespective of their possibly short or disconnected physical continuation. It does not include a branch whose corresponding formal critical clearance is nonzero.

Theorem 60.2 (Uniform full angular-source flux on the covered boundary)**.** *For every actual covered boundary point,*

$$\left|\left.\frac{\partial u}{\partial t}\right|_{\theta}\right| \le \Lambda_{\circ} K_{B,g} \gamma^{d'_j}. \tag{60.7}$$

Consequently the complete original positive marked endpoint charge on all covered boundary points at flight j, with both signs and any retained original exact labels, satisfies

$$\mathfrak{B}^{\mathrm{crit}}_{j,u_0} \le C_{\mathrm{zf}} \gamma^{d'_j}. \tag{60.8}$$

There is no deletion of small positive $|(p_0, p_m)|$ in this sum, no count of physical words in its constant, and no angular-kernel or whole-word pressure assumption. Summing all flights of depth at least D gives

$$\sum_{j:d'_j \ge D} \mathfrak{B}^{\mathrm{crit}}_{j,u_j} \le \frac{2C_{\mathrm{zf}}}{1-\gamma} \gamma^{D}, \qquad u_j \in [0, 3/200]. \tag{60.9}$$

For every fixed $0 < \vartheta_* < 1/2$, its product with m^2 tends to zero at $D = \lceil \vartheta_* m \rceil$, uniformly in the original radius and labels. These are bounds for the positive endpoint charge and for the total variation of its pushforward by the unchanged physical roof, not an assertion of a roof density for that boundary measure. The incoming critical fan at $g = 0$ itself is not this boundary-curve charge. Its original angular coefficient remains the separate quantity in Theorem 27.4; it is not assigned zero or included in (60.8) *by a bulk-null convention.*

Proof. Write $g = \nabla_q F$ at the selected actual point. Both $\mathcal{B}(0) = 0$ and $\mathcal{B}(g) = 0$ hold. The entire straight segment from 0 to g lies in the *formal* gradient disk of Lemma 60.1, where the proved analytic derivative bound is valid. The exact integral remainder is

$$D_g\mathcal{B}(g) \cdot g = \mathcal{B}(g) - \mathcal{B}(0) + \int_0^1 s D_g^2 \mathcal{B}(sg)[g,g]\, ds = \int_0^1 s D_g^2 \mathcal{B}(sg)[g,g]\, ds. \tag{60.10}$$

Its absolute value is at most $K_{B,g}\gamma^{d'_j}|g|^2/2$. At the actual point, the roof vector at fixed Gauss angle is $G^{-1}g/(g^{\mathsf{T}}G^{-1}g)$, so the normal speed is precisely

$$\left.\frac{\partial u}{\partial t}\right|_{\theta} = \frac{D_g\mathcal{B}(g) \cdot g}{g^{\mathsf{T}}G^{-1}g}.$$

Since the denominator is at least $|g|^2/(2\Lambda_{\circ})$, the quadratic factors cancel and prove (60.7). This calculation evaluates the physical normal speed at an actual point; it makes no physical-flow claim about the formal segment used in the integral identity.

The exact marked boundary formula (58.1) now bounds the charge by

$$\frac{\Lambda_{\circ} K_{B,g} \gamma^{d'_j}}{4\pi R_- c_*} \sum_{\varsigma=\pm 1} \int_0^{2\pi} d\vartheta = C_{\mathrm{zf}} \gamma^{d'_j}.$$

For each angular parameter and sign the limiting original finite record supplies only one marked physical source. Its possible assignment to a critical chart is made on this source before taking the upper bound. The bound is pointwise and identical for every such chart; it is not multiplied by their number.

Here is a precise disjoint assignment if multiple coordinate lifts were retained. At fixed R, m there are finitely many physical words and finitely many candidate marked disks per flight. A given word has at most one limiting normal-endpoint physical configuration. Indeed the support inequalities of Lemma 10.1 pass to its one-sided limits. If both endpoint gradients are zero, the endpoint normal incidences are one, so the summed inequality forces both endpoint contacts to agree. Both convex support gaps then vanish. Equality in the norm support inequality makes every pair of corresponding flight vectors positively collinear. Starting at the common first

contact, the incoming-root condition $c_i \geq 0$ chooses the same entry or tangent point of each next designated disk, successively. All contacts therefore agree, including possible limiting visited tangencies. Thus only finitely many reference configurations occur. Each of their formal inverse charts is continuous, and its equality with the actual finite record and all original inequalities is Borel. Assign an overlapping source point to the first such chart in a fixed ordering. Artificial angular seams are assigned once. The nonnegative source bound is unchanged. This proves the complete sum rather than a separate estimate per word.

There are at most two flight indices at each depth, and $\sum_{d\geq D}\gamma^d = \gamma^D/(1-\gamma)$. This proves (60.9). Multiplication by m^2 at the stated finite choice of D leaves a polynomial times an exponential with strictly negative exponent. No microscopic roof interval or outer limit has been changed. □

The exact remaining nonzero-offset term. This theorem treats a geometric class by a proved uniform normal-speed cancellation, not by assigning its positive trace zero. At an actual boundary point represented in a formal gradient chart with $\mathcal{B}(0) = a_w \neq 0$, the same exact calculation instead gives

$$\left.\frac{\partial u}{\partial t}\right|_{\theta} = \frac{-a_w + \int_0^1 sD_g^2\mathcal{B}(sg)[g,g]\,ds}{g^{\mathsf{T}}G^{-1}g}.$$

The offset is an actual signed clearance of that specified formal critical word, not a freely chosen clock or threshold. It cannot be dropped even if small. A complete weighted angular-source estimate for this offset term, for uncovered critical pieces, and for the original section/pairing boundary terms remains required to close the entire endpoint charge. The theorem neither interchanges a local critical-radius limit with $m \to \infty$ nor concludes the original essential-height or raw pointwise local limit assertion.

61. Radius-uniform cross-curve distortion on every real word

The transverse comparison required for a whole-cylinder pressure sum is stronger than distortion along one short curve. We prove that comparison here directly from the already established variational localization of module 308. In particular, the present argument does not assume that a physical word domain is convex, that two points can be joined through regular physical trajectories, or that a homogeneous planar cell has any unproved two-sided diameter property. The new conclusion is a uniform comparison of actual arclength Jacobians at different physical points and different cone directions. The localization input and its anchor-layer proof are inherited.

All trajectories below belong to the original unit-speed triangular Lorentz gas, with

$$R_- = \frac{9}{20},\quad R_+ = \frac{47}{100},\quad \tau_- = \frac{3}{50},\quad a = \frac{100}{47},\quad K = \frac{170}{9},\quad \varkappa_+ = \frac{20}{9},\quad \vartheta_s = \frac{47}{48}.$$

Let $\tau_+ < \infty$ be the uniform physical flight bound proved in module 282. As in module 313, tangent norms are the Euclidean norms in (s,φ), where $s = R\alpha$ and $p = \sin\varphi$. There is no measure normalization in a tangent Jacobian. The homogeneity partition is the original $q = 2$ partition with one fixed $k_0 \geq 2$:

$$H_k^+ = \{\pi/2 - k^{-2} < \varphi < \pi/2 - (k+1)^{-2}\},\qquad H_k^- = -H_k^+\quad (k \geq k_0),$$

and H_0 is the intervening central strip. Including strip boundaries by a fixed disjoint convention changes none of the inequalities below; true grazing points, at which the requested derivative is undefined, are excluded.

Put

$$N = 94,\qquad q_{\mathrm{a}} = \frac{3263680}{3263681},\qquad q = q_{\mathrm{a}}^{1/(4N)},\qquad C = 10^8, \tag{61.1}$$

where q_{a}, N, C are exactly the constants of module 308. In particular q is not substituted for the original guard constant $q_0 = 47/53$. Define the following finite, collision-independent constants:

$$C_Q = 2(2R_+C)^{1/4}q_{\mathrm{a}}^{-1/2}, \qquad C_\tau = (1+q^{-1})C_Q, \tag{61.2}$$

$$C_\varphi = \frac{\pi}{2}\left(\frac{C_Q}{R_-} + \frac{2C_\tau}{\tau_-}\right), \qquad D_c = \frac{2C_\varphi}{1-q}, \qquad D_\tau = \frac{2C_\tau}{1-q}, \tag{61.3}$$

$$D_v = \frac{K - a + (\tau_-^{-1} + (2a\tau_-^2)^{-1})D_c + \tau_-^{-2}D_\tau}{1-\vartheta_s}, \tag{61.4}$$

$$H_{\log} = \max\left\{2\pi 3^{2/3}, \frac{\pi^{5/3}}{2}k_0^2\right\}, \qquad D_{\log} = \frac{2H_{\log}C_\varphi^{1/3}}{1-q^{1/3}}, \tag{61.5}$$

$$D_A = \frac{(\varkappa_+ + K)D_\tau + \tau_+ D_v + D_c}{2a\tau_-}, \qquad D_{\mathrm{end}} = \log\frac{1+K^2}{1+a^2}, \tag{61.6}$$

$$D_\times = D_A + D_{\log} + D_{\mathrm{end}}, \qquad C_\times = \exp(D_\times). \tag{61.7}$$

These conservative constants may be very large. Their finiteness is uniform in the radius and every physical collision count; no assertion of a useful finite-sample numerical size is needed.

61.1. **A two-record consequence of the inherited anchor theorem.**

Lemma 61.1 (Paired real records, including disconnected word components)**.** *Let $m \geq 1$ and let $Q, \widehat{Q}$ be any two regular physical m-flight records at the same radius and with the same ordered lifted center word. Choose the common lift $O_0 = 0$ and write*

$$b_i = q^i + q^{m-i} \qquad (0 \leq i \leq m).$$

For the actual contacts, flight lengths, incidences, and collision angles one has

$$|Q_i - \widehat{Q}_i| \leq C_Q b_i, \qquad 0 \leq i \leq m, \tag{61.8}$$

$$|\tau_i - \widehat{\tau}_i| \leq C_\tau b_i, \qquad 0 \leq i < m, \tag{61.9}$$

$$|\varphi_i - \widehat{\varphi}_i| \leq C_\varphi b_i, \qquad 0 \leq i \leq m, \tag{61.10}$$

$$\sum_{i=0}^{m} |c_i - \widehat{c}_i| \leq D_c, \qquad \sum_{i=0}^{m-1} |\tau_i - \widehat{\tau}_i| \leq D_\tau. \tag{61.11}$$

In particular the actual two-sided phase distance is exponentially small toward the middle of a complete common physical word. The assertion requires no homogeneity or endpoint cap, and no minimum incidence or clearance. It compares two real records; the auxiliary minimum used in its proof is not a replacement trajectory.

Proof. Fix any one global minimum Y of the disk-product problem for this word and radius. It is a common comparison for both records; the proof does not require uniqueness of that minimum. In the notation of Corollary 48.4, let $d_i = \min(i, m-i)$ and $j(d) = \max\{0, \lfloor d/N \rfloor - 1\}$. The already proved quartic position budget, applied separately to the two real references and to this same Y, gives

$$|Q_i - Y_i|^4 \leq CE_R(Q)q_{\mathrm{a}}^{j(d_i)}, \qquad |\widehat{Q}_i - Y_i|^4 \leq CE_R(\widehat{Q})q_{\mathrm{a}}^{j(d_i)}.$$

Both original endpoint deficits are at most $2R \leq 2R_+$, including when i is an endpoint. Since $j(d) \geq d/N - 2$,

$$|Q_i - \widehat{Q}_i| \leq 2(2R_+C)^{1/4}q_{\mathrm{a}}^{j(d_i)/4} \leq C_Q q^{d_i} \leq C_Q b_i.$$

This is a consequence of the full inherited positive energy estimate, not an integration of a local endpoint derivative across a singular boundary. It proves (61.8).

The triangle inequality gives

$$|\tau_i - \widehat{\tau}_i| \le |Q_i - \widehat{Q}_i| + |Q_{i+1} - \widehat{Q}_{i+1}|.$$

Use $b_{i+1} \le q^{-1} b_i$ to prove (61.9). For nonzero vectors of lengths at least τ_-, their normalized directions differ by at most $2/\tau_-$ times their difference. Thus the unit direction of either flight adjacent to contact i differs from its paired direction by at most $2C_\tau b_i/\tau_-$; for the preceding flight use $b_{i-1} \le q^{-1} b_i$ directly in the sum of its endpoint bounds. The contact normals differ by at most $C_Q b_i/R_-$.

At $i < m$, the unit vector (c_i, p_i) is obtained by resolving the actual outgoing velocity in the normal and tangent frame of Q_i. The difference of the two resolved unit vectors is consequently at most the sum of the velocity and frame differences. At $i = m$, use the incoming resolution $(-v_{m-1} \cdot n_m, v_{m-1} \cdot t_m)$; it gives the same bound. Both resolved vectors lie on the right closed semicircle, with angles in $[-\pi/2, \pi/2]$. The angular difference on that semicircle is at most $\pi/2$ times their Euclidean chord distance. This proves (61.10), without an inverse-incidence differentiation of $\arcsin p$. Likewise the shorter arclength displacement of the contact on its radius-R circle is at most $\pi|Q_i - \widehat{Q}_i|/2$, so the same argument controls the actual phase metric modulo the circle seam.

Finally, $|c_i - \widehat{c}_i| \le |\varphi_i - \widehat{\varphi}_i|$ and $\sum_{i=0}^{m} b_i \le 2/(1-q)$. Summing the established inequalities proves (61.11). No connecting physical path or two-dimensional cell shape has appeared in the argument. □

Lemma 61.2 (The logarithmic incidence estimate on the actual strips)**.** *For two angles in the closure of the same member of the fixed $q = 2$ homogeneity partition,*

$$|\log\cos\varphi - \log\cos\widehat{\varphi}| \le H_{\log} |\varphi - \widehat{\varphi}|^{1/3}. \tag{61.12}$$

For any subset $J \subset \{0, \dots, m\}$ of matching homogeneous indices of the records in Lemma 61.1,

$$\sum_{i \in J} |\log c_i - \log \widehat{c}_i| \le D_{\log}. \tag{61.13}$$

Proof. In $\overline{H}_k^{\pm}$ put $\delta = |\varphi - \widehat{\varphi}|$. Then $\delta \le 3k^{-3}$ and $\cos\varphi, \cos\widehat{\varphi} \ge (2/\pi)(k+1)^{-2}$. The mean value bound for $\log\cos\varphi$ gives

$$|\log c - \log \widehat{c}| \le \frac{\pi}{2}(k+1)^2\delta \le 2\pi k^2 \delta \le 2\pi 3^{2/3}\delta^{1/3}.$$

For the central strip $c \ge \sin(k_0^{-2}) \ge (2/\pi)k_0^{-2}$ and $\delta \le \pi$. Hence the same mean value argument gives the other constant in (61.5). Both calculations hold on the closed strips. The original disjoint assignment of their boundary points is retained; no symmetry of that assignment is required. For paired records use (61.10), $(A+B)^{1/3} \le A^{1/3} + B^{1/3}$, and the two geometric sums:

$$\sum_{i \in J} |\log c_i - \log \widehat{c}_i| \le H_{\log} C_\varphi^{1/3} \sum_{i=0}^{m} (q^{i/3} + q^{(m-i)/3}) \le \frac{2H_{\log} C_\varphi^{1/3}}{1 - q^{1/3}}.$$

The common strip is used only where its logarithm is actually summed. □

61.2. **Cross-direction distortion and the exact endpoint factor.** For a nonzero vector ξ at x of slope in $[a, K]$, write

$$\mathcal{J}_m(x, \xi) = \frac{\|DT_R^m(x)\xi\|}{\|\xi\|}.$$

This is an actual forward arclength expansion in the unstable cone. When ξ is tangent to a smooth curve at x, it is precisely the Jacobian along that curve at that point. Curvature and a relation between two such curves are not needed for this pointwise comparison.

Theorem 61.3 (Uniform distortion across all physical components)**.** *Let $x, \widehat{x}$ generate two regular m-flight records with the same radius and the same full lifted physical word, and let $\xi, \widehat{\xi}$ be arbitrary nonzero vectors with slopes in $[a, K]$ at the two initial points. Suppose that the actual collision angles at each interior index $1 \le i < m$ lie in the closure of the same respective homogeneity strip. There is then the endpoint-exact bound*

$$\left|\log \frac{c_m(x)\mathcal{J}_m(x,\xi)}{c_m(\widehat{x})\mathcal{J}_m(\widehat{x},\widehat{\xi})}\right| \le D_{\times}. \tag{61.14}$$

Equivalently,

$$e^{-D_{\times}} \frac{c_m(\widehat{x})}{c_m(x)} \le \frac{\mathcal{J}_m(x,\xi)}{\mathcal{J}_m(\widehat{x},\widehat{\xi})} \le e^{D_{\times}} \frac{c_m(\widehat{x})}{c_m(x)}. \tag{61.15}$$

No homogeneity assumption is made on either endpoint in these two assertions. If the terminal angles also lie in one common closed strip, the same displayed constant gives the unweighted bound

$$\left|\log \frac{\mathcal{J}_m(x,\xi)}{\mathcal{J}_m(\widehat{x},\widehat{\xi})}\right| \le D_{\times}. \tag{61.16}$$

The constants are uniform in the radius, collision count, complete physical word, and the two cone directions. In particular the two points may belong to different connected regular components of that word, including components separated by true unvisited tangencies or any of the unchanged section and exact-label tests.

Proof. Let $v_i, \widehat{v}_i \in [a, K]$ be the slopes of the successive actual images of the two vectors, and put $\varkappa = 1/R$. The exact signed differential in Lemma 54.1 has a scalar minus sign; that sign changes neither slope nor Euclidean length. Its positive scalar numerator and slope map are

$$A_i = \tau_i(\varkappa + v_i) + c_i, \qquad v_{i+1} = f(v_i; c_i, c_{i+1}, \tau_i), \qquad f(v; c, d, \tau) = \varkappa + \frac{d(\varkappa + v)}{\tau(\varkappa + v) + c}. \tag{61.17}$$

Here $A_i \ge 2a\tau_- > 0$. The native contraction calculation gives $0 \le \partial_v f \le \vartheta_s = 47/48$. On the full parameter rectangle

$$v \in [a, K], \quad c, d \in [0, 1], \quad \tau \in [\tau_-, \tau_+]$$

the other exact derivatives satisfy

$$|\partial_c f| \le \frac{1}{2a\tau_-^2}, \qquad |\partial_d f| \le \frac{1}{\tau_-}, \qquad |\partial_\tau f| \le \frac{1}{\tau_-^2}. \tag{61.18}$$

For example $|\partial_c f| = d(\varkappa + v)/A^2$ and $|\partial_\tau f| = d(\varkappa + v)^2/A^2$; the bounds follow from $A \ge \tau(\varkappa + v)$ and $\varkappa + v \ge 2a$. The denominator stays positive throughout that algebraic parameter rectangle. A mean value estimate there interpolates numbers in an explicit rational function, not physical trajectories.

Writing $\delta v_i = |v_i - \widehat{v}_i|$, this proves

$$\delta v_{i+1} \le \vartheta_s \delta v_i + \frac{|c_i - \widehat{c}_i|}{2a\tau_-^2} + \frac{|c_{i+1} - \widehat{c}_{i+1}|}{\tau_-} + \frac{|\tau_i - \widehat{\tau}_i|}{\tau_-^2}.$$

The initial slope difference is at most $K - a$. Sum the recurrence, or sum its finite geometric expansion, and use (61.11). All terms are nonnegative and the initial term is paid once. Thus

$$\sum_{i=0}^{m-1} \delta v_i \le D_v. \tag{61.19}$$

The exact one-flight Euclidean Jacobian along the transported vector is

$$\frac{A_i}{c_{i+1}}\frac{\sqrt{1+v_{i+1}^2}}{\sqrt{1+v_i^2}}.$$

Multiplication therefore gives the actual identity

$$\log \mathcal{J}_m(x,\xi) = \sum_{i=0}^{m-1} \log A_i - \sum_{i=1}^{m} \log c_i + \frac{1}{2}\log\frac{1+v_m^2}{1+v_0^2}. \tag{61.20}$$

There is no incidence or Euclidean norm factor suppressed at the two ends. In particular multiplication by c_m cancels just the terminal logarithm in the second sum.

For the paired scalar numerators,

$$|A_i - \widehat{A}_i| \le (\varkappa_+ + K)|\tau_i - \widehat{\tau}_i| + \tau_+ \delta v_i + |c_i - \widehat{c}_i|.$$

Since both numerators are at least $2a\tau_-$, (61.11) and (61.19) imply

$$\sum_{i=0}^{m-1} |\log A_i - \log \widehat{A}_i| \le D_A. \tag{61.21}$$

After cancellation of the terminal incidence, the logarithmic incidences are precisely the indices $1,\dots,m-1$ covered by the assumption. Their total difference is at most $D_{\log}$ by Lemma 61.2. The last term of (61.20) telescopes before any bound is applied. The difference of its paired endpoint values is at most $\log((1+K^2)/(1+a^2)) = D_{\mathrm{end}}$. These three estimates prove (61.14) and its exact equivalent (61.15).

If the terminal index also has a common strip, apply (61.13) to all indices $1,\dots,m$ instead. Its constant was defined using the sum over $0,\dots,m$, so no additional endpoint constant is required. This proves (61.16). For $m = 1$ the interior sum is empty, the slope-sum bound includes only the initial slope, and the same proof gives both stated conclusions. Only actual finite words and their true reflection derivatives have been used. □

Lemma 61.4 (A physical terminal-incidence limit)**.** *In the original triangular lattice at $R = 9/20$, there is a family of regular one-flight records with the fixed lifted word $(O_0, O_1) = ((0,0),(1,0))$ for which c_0 stays strictly positive, $c_1 \downarrow 0$, and, for vectors ξ of fixed slope a,*

$$c_1 \mathcal{J}_1(x,\xi) \longrightarrow L_1 \in (0,\infty), \qquad \mathcal{J}_1(x,\xi) \longrightarrow \infty. \tag{61.22}$$

Every open flight segment in the family avoids every other obstacle. Thus the endpoint factor retained in (61.15) *is present on actual trajectories, including when the complete one-flight word is fixed.*

Proof. Start at $Q_0 = (R,0)$ with the outward unit velocity $v_\theta = (\cos\theta, \sin\theta)$, where

$$0 \le \theta < \theta_* := \arcsin(9/11).$$

The distance from Q_0 to O_1 is $11/20$. Direct intersection of this ray with the radius-R circle gives its first positive entry time and terminal incidence:

$$c_1(\theta) = \sqrt{1 - (11\sin\theta/9)^2}, \qquad \tau(\theta) = \frac{11}{20}\cos\theta - \frac{9}{20}c_1(\theta). \tag{61.23}$$

The time is positive: the difference between the squares of its two nonnegative terms is $(11/20)^2 - (9/20)^2 = 1/10$. The initial incidence is $c_0 = \cos\theta > 0$.

We check the entire physical segment, not just the entry point. For $0 \le t \le \tau(\theta)$, the bound $\tau \le (11/20)\cos\theta$ puts $Q_0 + tv_\theta$ in

$$[9/20, 1] \times [0, 11/40].$$

The nonzero rows of the triangular lattice have vertical coordinates of absolute value at least $\sqrt{3}/2$. Their distance from this rectangle is therefore at least $\sqrt{3}/2 - 11/40 > 9/20$; the strict inequality follows by squaring the positive quantities $\sqrt{3}/2$ and $29/40$, whose squared difference is $359/1600$. On the zero row, every center other than $(0,0), (1,0)$ is an integer horizontal translate with distance at least 1 from the rectangle. The initial obstacle is avoided for every $t > 0$ because

$$|Q_0 + tv_\theta|^2 = R^2 + 2Rt\cos\theta + t^2 > R^2.$$

The smaller positive root in (61.23) is the first contact with the target disk. These facts exclude every unvisited obstacle and every earlier contact. For $\theta < \theta_*$ the flight is regular and has exactly the stated physical word.

As $\theta \uparrow \theta_*$,

$$c_0 \longrightarrow \frac{2\sqrt{10}}{11}, \qquad \tau \longrightarrow \frac{1}{\sqrt{10}}, \qquad Q_1 \longrightarrow \left(\frac{139}{220}, \frac{9\sqrt{10}}{110}\right).$$

At each of these regular initial states take a nonzero vector of slope a; a local smooth curve with that tangent lies in the same regular one-flight domain after restriction to a sufficiently small neighborhood. Formula (61.17), with $\varkappa = 20/9$, gives

$$A_0 = \tau(\varkappa + a) + c_0, \qquad v_1 = \varkappa + \frac{c_1(\varkappa + a)}{A_0}, \qquad c_1\mathcal{J}_1 = A_0\frac{\sqrt{1+v_1^2}}{\sqrt{1+a^2}}.$$

Consequently (61.22) holds with

$$L_1 = \left(\frac{\varkappa + a}{\sqrt{10}} + \frac{2\sqrt{10}}{11}\right)\frac{\sqrt{1+\varkappa^2}}{\sqrt{1+a^2}} > 0.$$

No derivative at the limiting grazing state is used. Comparing this family with its regular member $\theta = 0$ shows directly why a uniform unweighted comparison without terminal homogeneity cannot be obtained by deleting the displayed terminal factor. This is a local geometric explanation of that factor; it is not an obstruction to the whole-pressure sum or to the original raw local-limit target. □

61.3. **The precise pressure and optical consequences.**

Corollary 61.5 (The required radius-uniform transverse curve comparison)**.** *Let $m \geq 1$ and let U, V be any two local smooth curves whose tangent slopes at $u \in U$ and $v \in V$ belong to $[-K, -a]$. Suppose the two actual backward records through times $0, -1, \ldots, -m$ are regular, have the same lifted physical word, and have matching homogeneity strips at times $-1, \ldots, -m$. Then*

$$C_\times^{-1} \leq \frac{J_U T_R^{-m}(u)}{J_V T_R^{-m}(v)} \leq C_\times. \tag{61.24}$$

In particular this holds on every one of the actual backward homogeneous cells specified in the preserved pressure candidate, with its literal time-index convention. It does not require the cell to meet a chosen transverse test. If homogeneity is specified only at $-1, \ldots, -m+1$, the exact endpoint-residual version is

$$C_\times^{-1}\frac{c(T_R^{-m}v)}{c(T_R^{-m}u)} \leq \frac{J_U T_R^{-m}(u)}{J_V T_R^{-m}(v)} \leq C_\times\frac{c(T_R^{-m}v)}{c(T_R^{-m}u)}.$$

Proof. The actual time reversal $I(s,\varphi) = (s, -\varphi)$ is a Euclidean arclength isometry and satisfies $T_R^{-1} = IT_RI$ on the regular domain. It turns both stable-cone directions into unstable-cone directions and the two backward records into two forward physical records with their order of centers reversed. The reflected angles lie in the closure of the same reflected strip, so the boundary-valid estimate of Lemma 61.2 applies. This argument does not assume that the original

disjoint assignment of strip boundaries commutes with reflection. Apply the appropriate form of Theorem 61.3. Norms before and after I are identical. The comparison includes every admissible curve tangent and needs neither common curvature nor a connection between the two curves. □

Corollary 61.6 (Whole-word normalized stable and optical cocycles)**.** *Let $x, \widehat{x} \in M'_R$ have the same actual m-flight lifted word and matching interior homogeneity strips $1, \dots, m-1$. No endpoint homogeneity, momentum cap, or positive incidence cutoff is imposed. For the true stable Euclidean arclength cocycle,*

$$C_\times^{-1} \le \frac{j_m(x)/c_0(x)}{j_m(\widehat{x})/c_0(\widehat{x})} \le C_\times. \tag{61.25}$$

For the original optical multiplier ω,

$$(30000^2 C_\times)^{-1} \le \frac{\omega(x)/(c_0(x)c_m(x))}{\omega(\widehat{x})/(c_0(\widehat{x})c_m(\widehat{x}))} \le 30000^2 C_\times. \tag{61.26}$$

These statements retain both physical endpoint factors explicitly. In particular an endpoint incidence is not declared harmless merely because a word or roof label is held fixed.

Proof. Start the backward comparison at $T_R^m x, T_R^m \widehat{x}$ and use the true stable tangent directions there. The backward Jacobians are exactly $1/j_m(x), 1/j_m(\widehat{x})$, by invariance of the stable line. The endpoint-weighted comparison of Theorem 61.3, after time reversal, compares $c_0(x)/j_m(x)$ to $c_0(\widehat{x})/j_m(\widehat{x})$. Taking reciprocals proves (61.25). The already proved optical comparison of module 313 is

$$\frac{c_m j_m}{30000} \le \omega \le 30000 c_m j_m.$$

Divide by the two actual endpoint incidences and compare the two points. One upper and one lower optical inequality cost 30000^2, not 30000 and not a factor raised to the number of collisions. This gives (61.26). □

Remark 61.7 (Completed geometric input and the remaining positive sum). Corollary 61.5 completes the radius-uniform cross-curve Jacobian comparison that remained conditional in the preserved whole-pressure candidate. Its proof is native and uses no radius-family interpretation of a fixed-table external theorem. It uses the established global minimum as a deterministic comparison only, and it includes all actual words, all positive incidences, and the exact homogeneity indices needed by its Jacobian formula.

The actual whole-cylinder pressure

$$\sup_{R,m} \sum_{w \in \mathcal{W}_m(R)} \sup_{x \in [w]_R \cap M'_R} j_m(x)$$

is not bounded merely by comparing two points in one homogeneous component. The positive sum still needs the complete physical barrier/continuation and planar refinement argument, or another proved payment for every cell supremum. In particular a component with arbitrarily small measure is not paid by module 315's bulk integrability theorem alone. The long-cell assignment and short-cell renewal in the retained candidate become valid conclusions only after their remaining actual planar source hypotheses are proved.

No original section, return, displacement, collision or physical roof test has been modified. Any such test, and any Borel selector taking values in $[0,1]$, may restrict the physical points compared here; they are not differentiated. This theorem itself is not a roof density estimate, a trace theorem, a uniform twisted spectral bound, or a proof of the original collision-normalized shortage limit. It changes neither the m^2 normalization nor the prescribed order of the original outer B limit and collision limsup.

62. Two-sided capacity of the actual marked angular source

The source in this section is the original one-copy marked flight, not a replacement collision process. A circle of directions is attached to each of the two sides of the marked disk. The disk is translated to the origin only to name its line coordinates. Following the line backwards through $j+1$ true collisions fixes the original initial disk; translating that disk to the fundamental cell recovers precisely one original marked copy. Following it forwards through $m-j$ collisions fixes the rest of the same record. In particular the two halves are not sampled independently.

Fix $R \in [R_-, R_+]$, $m \geq 1$, $0 \leq j < m$, and $u_0 \in [0, 3/200]$. At $u_0 = 0$ use the original positive-side unvisited trace: no collision is inserted at the marked foot. Every later first hit, including a later hit of the marked disk, remains a collision. At a positive level either prescribed incident side may be used; if two sides are counted their charges are added. The construction first uses all regular angular records. Original section, return, displacement, occupation and roof tests can subsequently be retained as selectors on this unchanged source.

Keep the exact constants of Sections 21, 25 and 57, and set

$$\sigma = \frac{47}{59}, \qquad E_* = \frac{1}{k_- a_\circ} = \frac{94}{\sqrt{1827}}, \qquad C_{\mathrm{foot}} = \frac{28/5 + R_+}{2R_-} = \frac{607}{90}, \qquad Q_{\mathrm{mb}} = C_{\mathrm{foot}} H_{\mathrm{nm}} E_*. \tag{62.1}$$

The enclosing horizon $28/5$ is a proved upper bound for the original physical flights. None of these constants depends on a word, collision count, clearance, radius, or exact record.

62.1. **Suprema on genuine one-sided source intervals.** For $l \geq 1$, let $P_l^\pm$ be the original momenta at the lth forward or backward collision from the foot, with the sign convention of Theorem 57.1. On a regular interval put $A_l^\pm = |\partial_\vartheta P_l^\pm|$. The following strengthens the pointwise root bound: it sums suprema on a complete physical partition before choosing any target.

Theorem 62.1 (One-sided physical branch capacity)**.** *For either time direction there is a countable partition $\mathcal{P}_l^\pm$ of the two angular circles, up to angular-null sets, into disjoint open source intervals. Each interval has a fixed actual lifted half-word. Its first $l-1$ collisions have the homogeneous and length cuts used in the native growth theorem; no terminal homogeneous subdivision is made. With*

$$a_I^\pm = \sup_{\vartheta \in I} (A_l^\pm(\vartheta))^{-1},$$

one has

$$\sum_{I \in \mathcal{P}_l^\pm} a_I^\pm \leq H_{\mathrm{nm}}, \qquad a_I^\pm \leq E_* \sigma^{l-1}. \tag{62.2}$$

These are suprema on actual intervals, not essential suprema of a chosen version of the terminal momentum density. Arbitrarily short intervals are included. The conclusion holds also for the specified zero-clearance trace.

Proof. Choose the common angular seam before constructing either time direction. A translated seam in the reversal $(\vartheta, \varsigma) \mapsto (\vartheta + \pi, -\varsigma)$ costs exactly the same one seam in Lemma 21.2; it does not introduce a second subdivision after the estimates.

For $l = 1$, the first-hit and coordinate-seam intervals in that lemma are the desired partition. There are at most M_* intervals for each side, and the exact first-hit derivative is $A_1 = \kappa a + c \geq k_- a_\circ$. Hence the sum is at most $2M_* E_* \leq H_{\mathrm{nm}}$.

For $l \geq 2$, use the source-labelled first-image family $\rho_C \, d\ell_C$ of Theorem 21.3. Its density suprema sum to at most A_*. For each C apply Theorem 25.1 through exactly $l-2$ further real collisions. Let V be one resulting short homogeneous curve and $J_V(T_R^{-(l-2)})$ its inverse arclength Jacobian. Cut V only at the finitely many next-hit branches required for the last actual collision.

There are at most J_f of them. On each such branch the exact terminal cancellation gives $|dp_{\mathrm{next}}/d\ell_V| \geq k_- t_*$. The inverse angular derivative on its original source interval is therefore bounded by

$$\frac{\|\rho_C\|_\infty}{k_- t_*} \sup_V J_V(T_R^{-(l-2)}).$$

Sum this bound over the last-hit cuts, the complete growth partition, and the labelled first-image curves. The result is $A_* C_g J_f/(k_- t_*) \leq H_{\mathrm{nm}}$. Every interval here is a pullback along injective regular curve branches. Homogeneity and length cuts preserve connectedness on those graphs. Thus these are disjoint intervals in the original angular source, not a multiset of alternative representations. The countable singular-null coverage is exactly the one proved in Theorem 25.1. No derivatives or component counts of later selectors occur.

For the second inequality use the actual positive optical product in (59.1). Its first free-flight matrix has $B = a \geq a_\circ$ and $D = 1$. Each additional interior collision and following flight replaces these entries by

$$B_{\mathrm{new}} = (1 + \ell k)B + \ell D, \qquad D_{\mathrm{new}} = kB + D, \qquad k = \frac{2}{Rc} > 0.$$

Here $\ell \geq t_* = 3/50$ and $c \leq 1$, so $1 + \ell k \geq 1 + 2t_*/R_+ = 59/47$. Positivity gives $B \geq a_\circ (59/47)^{l-1}$. The endpoint is incoming and has no extra kick in this product: $A_l = \kappa B + c_l D \geq k_- B$. This proves the pointwise bound and hence its interval supremum. It remains valid as any positive interior incidence decreases; no upper bound for a grazing kick was used.

At $u_0 = 0$ perform the same construction directly on the strict positive-side source of Theorem 57.4. The foot is outside every other disk by the strict standoff estimate. The first-hit intervals, their exact derivatives, the factor c in ρ_C, and all curve-class estimates of 279–283 hold at this level as well. Later genuine tangencies exclude only the countable angular set in the finite-time growth proof. The skipped tangency at the designated foot is fixed throughout and is never used as a reflection. This constructs the partition and proves its suprema at the trace itself; a weak convergence assertion is not used to transfer a sum of suprema. Reversal is the same arclength and angular isometry as in Theorem 57.1, with the original selectors transported through it. □

62.2. **An interval overlay estimate without a word-count factor.**

Lemma 62.2 (Weighted intersections of two interval families)**.** *Let $\mathcal{I}, \mathcal{J}$ be two countable families of disjoint open subintervals of one fixed bounded interval. For nonnegative weights a_I, b_J, with finite sums $A = \sum_I a_I$, $B = \sum_J b_J$, put $a_* = \sup_I a_I$ and $b_* = \sup_J b_J$, using zero for an empty family. Then*

$$\sum_{\substack{I \in \mathcal{I},\ J \in \mathcal{J} \\ I \cap J \neq \varnothing}} a_I b_J \leq b_* A + a_* B. \tag{62.3}$$

The same assertion holds on disjoint labelled copies of the interval, using the combined sums and suprema.

Proof. Write an interval's left endpoint as ℓ_I or ℓ_J. For each nonempty intersection charge its product to I if $\ell_I \geq \ell_J$, and to J otherwise. A fixed I receives at most one charge: two disjoint open J intervals starting no later than ℓ_I cannot both intersect I immediately to the right of ℓ_I. The same argument applies to J when its left endpoint is the larger one. Equal endpoints are assigned to I only. Thus charges to I sum to at most $b_* A$, and charges to J to at most $a_* B$. Apply this first to finite subfamilies. Every finite collection of intersections is contained in such subfamilies; nonnegative exhaustion proves the countable assertion. Summing over labelled

components proves the last claim. No lower interval length or connectedness of a union of intervals is required. □

For the original jth marked flight overlay $\mathcal{P}^{-}_{j+1}$ and $\mathcal{P}^{+}_{m-j}$ on each labelled angular circle. Denote the nonempty intersections by $\mathcal{O}_{m,j,R,u_0}$. An element has one complete genuine collision word and its indicated interior homogeneous cuts. Different intervals can carry the same word. They are not merged by assuming that its real physical domain is connected.

Theorem 62.3 (Complete two-sided marked inverse-weight capacity)**.** *Let $u_x = |\partial_{\alpha_0} p_m|^{-1}$ be the original inverse endpoint-momentum weight on the complete regular marked source. Then*

$$\sum_{O\in\mathcal{O}_{m,j,R,u_0}} \sup_{O} u_x \le Q_{\mathrm{mb}}(\sigma^j + \sigma^{m-j-1}). \tag{62.4}$$

This includes both mark signs and all actual half-words and their intersections, also at the one-sided zero-clearance level. Any original endpoint cap or exact-record selector can be retained inside the positive sum. Selectors are restricted within the existing intervals and do not create extra charges. Consequently, for arbitrary specified levels $u_j \in [0, 3/200]$,

$$\sum_{j:\,\min(j,m-1-j)\ge D} \sum_{O\in\mathcal{O}_{m,j,R,u_j}} \sup_{O} u_x \le \frac{2Q_{\mathrm{mb}}}{1-\sigma}\sigma^D. \tag{62.5}$$

An empty index set contributes zero.

Proof. On every actual overlay interval, Theorem 59.1 and Corollary 59.2 give

$$u_x = \frac{1}{RA^p_- A^p_+(h_- + h_+)}, \qquad h_- + h_+ \ge \frac{2}{\tau_+ + R_+}.$$

The two time directions fix the opposite clearance-column signs in this determinant. They yield a positive sum, not a difference that could vanish. Hence on $O = I \cap J$, $\sup_O u_x \le C_{\mathrm{foot}} a^-_I a^+_J$. Apply Lemma 62.2 and Theorem 62.1. The two sums of weights are at most H_{nm}; their respective maxima are at most $E_*\sigma^j$ and $E_*\sigma^{m-j-1}$. This gives (62.4) with exactly the constant in (62.1).

The physical uniqueness in the one-mark flux identifies each angular parameter and side with one actual marked copy at the fixed index j. Translating its starting disk does not add a lattice multiplicity. The overlay therefore sums the actual source before imposing any record, roof or section selector. Removing a test is used only to enlarge this positive upper bound. Finally sum $\sigma^j + \sigma^{m-j-1}$ over $D \le j \le m-1-D$. Each of the two geometric sums is at most $\sigma^D/(1-\sigma)$, proving (62.5). □

Scope of the capacity statement. Equation (62.4) is a genuine sum of physical suprema on a complete marked angular source. It does not follow by choosing one common endpoint momentum for all intervals. It supplies the comparison needed for the boundary theorem below, including short and disconnected real word domains. It is not the two-dimensional unrestricted pressure $\sum_w \sup_{[w]} j_m$ in Input 11.1, and is not yet a pointwise roof density. The original physical roof, its interval widths, all four recorded coordinates and every outer limit remain unchanged.

63. Uniform total turning of the physical marked boundary

The boundary measure in Theorem 58.1 cannot be bounded pointwise by deleting its nonzero critical offset. We instead control its complete angular variation on the actual marked source. The argument uses the physical one-sided interval partitions of Theorem 62.1 and their two-sided weighted overlay in Theorem 62.3. In particular its partition is not a hypothetical family of large comparison cells.

Keep the original endpoint cap of Lemma 4.1:

$$|p_0|, |p_m| \le 9/10, \qquad c_\circ = \sqrt{19}/10.$$

This cap follows from the original section tests. Restricting an upper-bound source to it adds no new condition to the observed event. Let $k_- = 100/47$, $k_+ = 20/9$, $U = 170/9$, $a_\circ = \sqrt{1827}/200$, $b_* = 97/200$, and $\varrho = 47/59$. Use exactly the fixed B_{cv}, k_0, ℓ_* and physical curve class of Theorem 25.1. Set

$$\begin{aligned}
B_2 &= B_{\mathrm{cv}}(1+U^2)^{3/2}, & \Gamma &= \sqrt{1+k_+^2}, \quad L_0 = \frac{\pi\sqrt{1+U^2}}{k_-},\\
L_h &= \frac{3\sqrt{1+U^2}}{k_-}, & c_H &= \sin(k_0^{-2}),\\
V_H &= \max\left\{-2\log c_H, \log(\pi/2) + 2\log(1+k_0^{-1})\right\}. & & \qquad (63.1)
\end{aligned}$$

Let D_B be the completed one-step derivative constant in (24.7), and K_0 the first-image derivative constant in (21.3). The following constants are finite and depend only on the original physical family and its already fixed curve class:

$$\begin{aligned}
D_1^{\mathrm{var}} &= \max\left\{(U/c_H + D_B)L_0^{2/3}, (2\pi U + D_B)L_h^{2/3}\right\},\\
V_J &= \frac{D_1^{\mathrm{var}}\Gamma^{1/3}L_0^{1/3}}{1-\varrho^{1/3}},\\
V_\rho &= V_H + \left(a_\circ^{-1} + b_* a_\circ^{-2} + K_0/2\right)\ell_*, \qquad (63.2)\\
V_t &= -2\log c_\circ + B_2 L_0/k_-,\\
V_1 &= \frac{L_0(k_+ b_* + 1)}{k_- a_\circ^2},\\
V_A &= \max\{V_1, V_\rho + V_J + V_t\}. \qquad (63.3)
\end{aligned}$$

No constant is tuned to the collision count, a clearance level, a roof window, an exact lattice label, or an outer cutoff.

Lemma 63.1 (Absolute homogeneous logarithmic variation)**.** *Let C' be a connected actual curve branch in the class of Theorem 25.1, and suppose its images through time l are homogeneous. If L_l is the arclength of its final image, then*

$$\mathrm{Var}_{C'} \log J_C(T_R^l) \le \frac{D_1^{\mathrm{var}}\Gamma^{1/3}}{1-\varrho^{1/3}} L_l^{1/3} \le V_J. \qquad (63.4)$$

The variation is along the full indicated connected branch, not just the difference between its endpoint values. No positive lower bound on the interior incidences is imposed.

Proof. Every unstable image graph has $k_- \le d\varphi/ds \le U$ and $-\pi/2 < \varphi < \pi/2$. Integration in φ therefore bounds its whole arclength by L_0. On a segment of length r in $H_k^{\pm}$ one has

$$r \le L_h k^{-3}, \qquad c \ge (2/\pi)(k+1)^{-2}.$$

Lemma 24.3 bounds the absolute logarithmic derivative per unit image arclength by $U/c + D_B$. Thus its absolute integral on this segment is at most

$$(2\pi U + D_B)k^2 r \le (2\pi U + D_B)L_h^{2/3} r^{1/3}.$$

On H_0 the analogous bound is $(U/c_H + D_B)r \le (U/c_H + D_B)L_0^{2/3} r^{1/3}$. These are estimates of the integral of the absolute derivative. They prove a one-step *total-variation* estimate with constant D_1^{var}; no conclusion about variation is inferred merely from a Hölder difference bound.

The exact adapted expansion of (24.2) gives $L_i \le \Gamma \varrho^{l-i} L_l$ for every intermediate image. Write the logarithm of the full physical Jacobian as the sum of its l actual one-step logarithms. Pullback by a monotone curve parametrization preserves their total variations. The triangle inequality for variation and the preceding absolute one-step bounds give

$$\operatorname{Var}_{C'} \log J_C(T_R^l) \le D_1^{\mathrm{var}} \sum_{i=1}^{l} L_i^{1/3} \le D_1^{\mathrm{var}} \Gamma^{1/3} L_l^{1/3} \sum_{r=0}^{l-1} \varrho^{r/3}.$$

Sum the convergent geometric series. The $l = 0$ Jacobian is one and has zero variation. Compact subarcs and monotone exhaustion give the same inequalities on open homogeneous intervals, including arbitrarily short intervals whose endpoints approach singularities. □

Lemma 63.2 (Uniform variation of a physical half-ray momentum derivative)**.** *On every one-sided angular branch of Theorem 62.1, let*

$$A_l^p(\vartheta) = |\partial_\vartheta p_l| > 0$$

be the actual momentum derivative after $l \ge 1$ true collisions from the marked foot. Restrict the terminal momentum to the original cap $|p_l| \le 9/10$. This restriction is an interval, possibly empty, and on it

$$\operatorname{Var} \log A_l^p \le V_A. \tag{63.5}$$

The same assertion holds in the reversed physical direction, with the transported original event and the same constants. Neither a terminal homogeneous strip count nor an extra collision is inserted.

Proof. The terminal curve is a genuine unstable graph in its direction of propagation. Its angle φ_l and momentum p_l are strictly monotone in its curve coordinate. The cap is consequently one interval. Since $k_0 \ge 2$ and $\arcsin(9/10) < \pi/2 - 1/4$, the cap is contained in the original central homogeneous strip H_0.

For $l \ge 2$, start from the exact first-image source $\rho_C \, d\ell_C$ of Theorem 21.3. Its first image C is already homogeneous and has length at most ℓ_*. The one-sided branch in Theorem 62.1 has homogeneous intermediate images; its last true collision, restricted to the cap, has its image in H_0. The arclength chain rule gives the exact identity

$$A_l^p = \frac{c_l V_l}{\sqrt{1 + V_l^2}} \frac{J_C(T_R^{l-1})}{\rho_C}. \tag{63.6}$$

Here ρ_C is the actual angular density before any Borel record selector is multiplied. The selector is not differentiated.

We verify each variation in (63.6). On a first-image homogeneous graph,

$$\log \rho_C = \log c_1 - \log a - \tfrac{1}{2} \log(1 + V_1^2).$$

The angle φ_1 is monotone. On H_0 the variation of $\log c_1$ is at most $-2 \log c_H$; on a signed H_k it is at most

$$\log \frac{\sin(k^{-2})}{\sin((k+1)^{-2})} \le \log(\pi/2) + 2 \log(1 + k_0^{-1}).$$

Both cases are bounded by V_H. The exact first-flight derivatives in (21.5) yield

$$|a_s| \le 1 + b_*/a_\circ, \qquad |V_{1,s}| \le K_0.$$

Using $a \ge a_\circ$ and $v/(1+v^2) \le 1/2$ for $v > 0$, their absolute integrals on the first piece show

$$\operatorname{Var} \log \rho_C \le V_\rho.$$

This retains the true grazing factor in ρ_C. It does not replace the density by a constant before summing the initial strips.

Lemma 63.1 gives variation at most V_J for the Jacobian in (63.6). At the terminal cap $c_l \ge c_\circ$ and φ_l is monotone, so

$$\operatorname{Var} \log c_l \le -2 \log c_\circ.$$

Geometric curvature gives $|V_{l,s}| \le B_2$. Moreover

$$\left| \partial_s \log \frac{V_l}{\sqrt{1+V_l^2}} \right| = \frac{|V_{l,s}|}{V_l(1+V_l^2)} \le B_2/k_-.$$

Its absolute integral is at most $B_2 L_0/k_-$. Thus the terminal factor has logarithmic variation at most V_t. Taking variation of (63.6) proves the second alternative in (63.3).

For $l = 1$, the branch partition uses the finite first-hit intervals, without an unnecessary infinite homogeneous cut. Put $b = \varsigma(R + u_0)$. The exact first-hit derivative is $A_1^p = \kappa a + c$, where $\kappa = 1/R$. Directly from (21.5),

$$\partial_s(\kappa a + c) = \kappa(p + bc/a) - p(\kappa + c/a) = c(\kappa b - p)/a.$$

Since $A_1^p \ge k_- a_\circ$, its logarithmic derivative has magnitude at most $(k_+ b_* + 1)/(k_- a_\circ^2)$. The first image has length at most L_0, proving the bound V_1.

Time reversal is the exact arclength isometry already used in Theorems 25.1 and 57.1. It reverses the momentum sign and transports the original selector; it preserves all the absolute derivative and variation bounds above. The same proof therefore covers the $j + 1$ backward collisions, including $j = 0$. All limiting statements use compact regular subarcs followed by exhaustion, not a fictitious reflection at the marked foot. □

Lemma 63.3 (Radial turning controlled by tangent turning)**.** *Let P be a C^2 regular plane curve on a connected interval. Suppose P is nonzero and its tangent has total turning V_ψ. Then*

$$\int |d \arg P| \le V_\psi + \pi. \tag{63.7}$$

If P passes through the origin at most once, the sum on its punctured intervals is at most $V_\psi + 2\pi$. The latter statement excludes the jump of the polar angle at the origin; it assigns no critical fan there a zero value.

Proof. On a compact nonzero subarc choose continuous angles $\alpha = \arg P$ and $\psi = \arg P'$. With $\delta = \psi - \alpha$, the exact derivative is

$$\alpha' = \frac{|P'|}{|P|} \sin \delta.$$

The 2π-periodic triangular function $h(\delta) = \arccos(\cos \delta)$ takes values in $[0, \pi]$ and has derivative $\operatorname{sgn}(\sin \delta)$ away from multiples of π. The absolutely continuous chain rule gives, almost everywhere,

$$|\alpha'| = \operatorname{sgn}(\sin \delta)\psi' - \frac{d}{ds} h(\delta).$$

At a level set $\delta \in \pi\mathbb{Z}$, its derivative is zero almost everywhere on that set; also $\alpha' = 0$ there, so the same identity holds with the sign set to zero. Integration gives

$$\int |\alpha'| \le \int |\psi'| + h(\delta_{\text{left}}) - h(\delta_{\text{right}}) \le V_\psi + \pi.$$

Exhausting the interval proves the first assertion. If there is one zero, apply it on both punctured sides and sum their tangent variations. This proves the second assertion and its explicit critical-point convention. □

Corollary 63.4 (Uniform turning on every actual two-sided angular piece)**.** *On any nonempty overlap piece from Theorem 62.3, restricted to the original endpoint cap, set* $P(\vartheta) = (-p_0(\vartheta), p_m(\vartheta))$*. Then*

$$\int_{\{P\neq 0\}} |\partial_\vartheta \arg P|\, d\vartheta \le 2\pi + V_A. \tag{63.8}$$

The bound is independent of the length of the overlap and of its distance from a critical record. All original Borel cuts may be retained as selectors without counting their connected components.

Proof. The signed derivative identity (59.4) makes both coordinate derivatives of P nonzero with fixed signs on the piece. Its tangent therefore stays in one open quadrant, and P can pass through the origin at most once. The cap restricts each monotone momentum to an interval, so their simultaneous restriction is still an interval.

The absolute slope ratio of the tangent is A^p_+/A^p_-. On its fixed quadrant the derivative of its tangent angle with respect to $\log(A^p_+/A^p_-)$ has absolute value

$$\frac{A^p_+/A^p_-}{1+(A^p_+/A^p_-)^2} \le \frac{1}{2}.$$

Lemma 63.2 bounds each of the two logarithmic variations by V_A, also after restriction to the overlap. Hence the tangent turning is at most V_A. Apply Lemma 63.3. A Borel selector bounded by one only decreases the nonnegative integral already bounded; it is neither differentiated nor used to create additional summation pieces. □

64. The complete regular clearance-boundary budget without an offset cutoff

Let $\mathfrak{B}^{\mathrm{reg}}_{j,u_0}$ be the original positive marked charge of (58.1), for one specified incident side of the clearance level $u_0 \in [0, 3/200]$, summed over both impact signs and all actual regular angular branches of the m-flight record. Here $0 \le j < m$, all visited collisions on an integrated branch are regular, and all original endpoint and exact-record tests are retained. The regular boundary is punctured only at $g = 0$, where the original incoming critical fan remains a different measure. There is no positive lower bound on $|g|$ or on an interior incidence.

For $u_0 = 0$ this is the actual $u > 0$ one-sided unvisited trace of Theorem 57.4; no collision is added at the mark. An isolated singular angular parameter is not used to define a new trace. In particular the notation does not absorb a distinct visited-grazing, section or pairing boundary, or a simultaneous critical corner fan. Every regular part of the prescribed marked boundary, including nonzero-offset and arbitrarily short components, is included.

Put

$$C_{\mathrm{ab}} = \frac{\Lambda_\circ(2\pi + V_A)}{4\pi c_* R_-^2}\,\frac{607}{90}\,H_{\mathrm{nm}}\,\frac{94}{\sqrt{1827}}. \tag{64.1}$$

The last three factors are exactly the complete two-sided root-weight constant in (62.4); in particular H_{nm} already includes both impact signs.

Theorem 64.1 (Full regular marked boundary, all critical offsets)**.** *For the unchanged Lorentz family and the complete indicated physical source,*

$$\mathfrak{B}^{\mathrm{reg}}_{j,u_0} \le C_{\mathrm{ab}}\big(\varrho^j + \varrho^{m-1-j}\big). \tag{64.2}$$

This bound is uniform in R, m, the level u_0, and every original exact record and roof selector between zero and one. It requires no critical zero-offset chart, no critical momentum cutoff, and no whole-word pressure or comparison-pool abundance premise.

Consequently, for any levels $u_j \in [0, 3/200]$ and any integer $D \geq 0$,

$$\sum_{\min(j,m-1-j)\geq D} \mathfrak{B}^{\mathrm{reg}}_{j,u_j} \leq \frac{2C_{\mathrm{ab}}}{1-\varrho}\,\varrho^D. \tag{64.3}$$

At $D = \lceil \vartheta_* m \rceil$, with fixed $0 < \vartheta_* < 1/2$, the product of this complete positive charge with m^2 converges to zero, uniformly in the original radius and exact labels and in the prescribed levels. Its pushforward by the unchanged physical roof has at most the same total variation.

Proof. On the original endpoint cap, the identities in Corollary 59.2 give

$$f^\varepsilon = \frac{W^\varepsilon u_x}{4\pi c_* h_x}, \qquad h_x = \frac{R^2\chi}{\det G} = R^2 e^{\mathsf{T}} G^{-1} e, \qquad e = g/|g|.$$

The last equality is the two-dimensional cofactor identity $e^{\perp\mathsf{T}} G e^\perp / \det G = e^{\mathsf{T}} G^{-1} e$. Since $G \preceq \Lambda_\circ I$,

$$f^\varepsilon \leq \frac{\Lambda_\circ}{4\pi c_* R_-^2}\, u_x. \tag{64.4}$$

The original bad-source weight satisfies $0 \leq W^\varepsilon \leq 1$; it is not set to zero on a zero-clearance boundary. There is exactly one original conditioning factor in (64.4).

Represent the fixed-level two-sign angular source by the actual one-sided interval partitions and their overlaps of Theorem 62.3. On each overlap take the supremum of the genuine root weight u_x, not a supremum on a formal nonphysical extension. Corollary 63.4 bounds the integral of $|\partial_\vartheta \theta|$ on its punctured cap by $2\pi + V_A$. Therefore

$$\mathfrak{B}^{\mathrm{reg}}_{j,u_0} \leq \frac{\Lambda_\circ(2\pi + V_A)}{4\pi c_* R_-^2} \sum_{\text{actual overlaps}} \sup u_x.$$

Insert the complete interval-overlay sum (62.4). It gives exactly (64.2) and (64.1). Countable partitions are handled first on finite compact regular subfamilies, followed by nonnegative exhaustion. Artificial seams are assigned once on the angular source. The overlap argument has already paid its possible count; it is not paid again by a collision word multiplier. Original labels and valid one-sided Borel trace selectors only restrict the positive measures at this stage.

This proof never divides the source into zero and nonzero formal critical offsets. In particular the term identified after Theorem 60.2 is retained through the exact turning integral, even when it concentrates sharply. The estimate bounds that integral and does not assert a uniform pointwise normal speed. Arbitrarily short genuine overlaps and components approaching a critical record are included with the same constant.

Summing (64.2) over $D \leq j \leq m-1-D$ gives twice a subseries of $\sum_{j\geq D} \varrho^j$, proving (64.3). An empty index range contributes zero. After the stated finite choice of D, multiplication by m^2 leaves a constant times $m^2 \varrho^{\vartheta_* m}$, tending to zero. All outer widths and the original central target remain fixed. The supremum over radius, labels and levels is permitted by the same displayed finite-count bound. Pushforward cannot increase the total mass of a positive measure, proving the roof-TV statement. □

The exact improvement and the remaining target. Theorem 64.1 closes the unrestricted *regular marked-boundary* angular summation, including the nonzero-offset term left in Section 60. The mechanism is the complete two-sided physical inverse-weight sum and uniform total turning, rather than smallness of a local offset. It preserves every actual marked copy and every original physical clock and record label. No claim that a small total variation is a small roof essential height is made.

Section 66 additionally pays the leading incoming zero-clearance marked critical fan when all visited incidences remain positive, including collapsed slices and shallow marks. The nonleading

critical remainders, distinct visited-grazing and section/pairing boundary measures, shallow low-load same-roof shortage and whole physical comparison capacity retain their stated obligations. In particular (64.3) does not imply the original m^2L^∞ local limit simply by multiplying a fixed shallow-source mass bound by m^2. The raw target (1.5), its exact arithmetic reference, conditioning and original order of limits remain unchanged.

65. A complete outgoing-angle capacity bound for physical classes

The cross-curve estimate of Section 61 can be summed over all physical classes that meet one prescribed transversal. Here we construct a genuine family of transversals which covers the entire collision cylinder, including its coordinate seam, and perform that sum. Short classes and disconnected classes are included. The result is an unconditional bound for a geometric capacity moment and for the unweighted sum over classes of a prescribed positive capacity. The capacity is the angular size of the set of actual outgoing directions present in a class; it is not a physical roof interval or a replacement for the original same-roof comparison pool.

Throughout this section $R \in [R_-, R_+] = [9/20, 47/100]$, the speed is one, and T_R is the original physical collision map. Put

$$\kappa_R = R^{-1}, \qquad a = \frac{100}{47}, \qquad U = \frac{170}{9}, \qquad \Gamma = \sqrt{1 + (20/9)^2}.$$

All curve lengths and tangent Jacobians are Euclidean arclength in the physical coordinates (s, φ), with $s = R\alpha$ and $p = \sin\varphi$. Fix once and for all the original $q = 2$ homogeneity partition, its k_0, the curvature allowance B_{cv}, the short length ℓ_* and the constants $\theta_* < 1, C_g$ of Theorems 24.2 and 25.1. Further fixed shortening of ℓ_*, if needed to apply their preliminary short-graph arguments, is absorbed in that same curve-growth theorem. None of these choices depends on the collision count, a record label, a roof interval, or an outer parameter of the local-limit problem.

65.1. **The initially uncut affine family.** The actual outgoing direction is the circle-valued function

$$\pi_R(s, \varphi) = s/R + \varphi \pmod{2\pi} \quad \text{on } M_R = (\mathbb{R}/2\pi R\mathbb{Z}) \times (-\pi/2, \pi/2). \tag{65.1}$$

In particular π_R is globally defined at the collision-circle seam. For $\vartheta \in \mathbb{R}/2\pi\mathbb{Z}$, its fiber $\Gamma_{R,\vartheta}$ is parametrized by

$$-\pi/2 < \varphi < \pi/2, \qquad s = R(\vartheta - \varphi) \pmod{2\pi R}. \tag{65.2}$$

It is an injective physical stable-cone arc: in a lift its slope is $d\varphi/ds = -\kappa_R$, its curvature is zero, and

$$|\Gamma_{R,\vartheta}| = \pi\sqrt{1 + R^2} \le S_{\mathrm{aff}} := \pi\sqrt{1 + R_+^2}. \tag{65.3}$$

These are the collision states of parallel outgoing rays; no orbit has been changed by introducing the foliation. A fiber generally crosses infinitely many current homogeneity strips. We do not partition it at those current strips before applying the next lemma.

Set

$$S_{\mathrm{im}} := \pi\sqrt{1 + a^{-2}}, \qquad N_{\mathrm{im}} := 1 + \left\lceil \frac{2S_{\mathrm{im}}}{\ell_*} \right\rceil, \qquad C_{\mathrm{aff}} := \max\{1, N_{\mathrm{im}}\Gamma\theta_* C_g\}. \tag{65.4}$$

Lemma 65.1 (Growth from a nonhomogeneous affine stable source)**.** *Let W be any open subarc of a fiber* (65.2)*, with $|W| \le \ell_*$. For every $N \ge 1$, let $\mathcal{G}_N^{\mathrm{aff}}(W)$ be obtained by N actual backward collisions, cutting at the actual regularity boundaries and at the homogeneity boundaries at times $-1, \dots, -N$, and making the fixed short-length subdivisions after each collision. Then*

$$\sum_{V \in \mathcal{G}_N^{\mathrm{aff}}(W)} \sup_V J_V T_R^N \le C_{\mathrm{aff}}. \tag{65.5}$$

The pullbacks to W of these open pieces are disjoint, and cover arclength-almost all of W. At every point of W with an actual regular backward record through time $-N$, the excluded cuts are locally finite. Such a point is either in one of the open pullbacks, or is an endpoint of at least one of them. In the latter case the one-sided curve Jacobian has the value of the actual finite regular derivative at that point.

Proof. Time reversal $I(s,\varphi) = (s,-\varphi)$ is an arclength isometry, satisfies $IT_RI = T_R^{-1}$, and turns W into a positive-cone affine graph of slope κ_R and curvature zero. We first check that the native one-step proof applies before this graph has been made homogeneous. The relative-angle and positive-foot estimates in Section 24 require only the cone bounds and the common short length. They give at most J_f actual first-hit pieces, each meeting each target strip at most once. The limiting first-hit argument of Lemma 24.1, including its initial-grazing case, also uses no current-strip restriction. There is consequently at most one central target piece. The exact adapted expansion (24.2) is valid for every positive initial incidence, and its target-strip sum is therefore precisely (24.4). Neither the initial incidence nor the current strip index is divided out or summed in this argument.

The curvature formula (23.3) and Theorem 23.2 also use only the initial cone and curvature bounds. Every regular homogeneous first image hence belongs to the inherited class with allowance B_{cv}, even though the source was not homogeneous. Reversing the resulting statement gives

$$\sum_{V\in\mathcal{H}^-(W)} \sup_V J_V T_R \le \Gamma\theta_*. \tag{65.6}$$

Here $\mathcal{H}^-(W)$ consists of the maximal actual homogeneous pieces of $T_R^{-1}W$, before their length subdivision. The factor Γ is the one endpoint conversion from the adapted norm to Euclidean arclength; it is not a factor at every subsequent collision.

On each of these graph pieces, $|d\varphi/ds| \ge a$ and φ is strictly monotone. In a local lift,

$$d\ell = \sqrt{1+|d\varphi/ds|^{-2}}\,|d\varphi| \le \sqrt{1+a^{-2}}\,|d\varphi|.$$

Its total length is at most S_{im}. The graph cannot acquire an extra turn by passing a coordinate seam: its lifted horizontal length is at most $\pi/a < 2\pi R_-$, while its collision angle stays in $(-\pi/2,\pi/2)$. A piece of length at most S_{im} can be subdivided into at most N_{im} pieces of length at most ℓ_*, with the standard lower length $\ell_*/2$ when a long piece is subdivided. For example, a piece of length $L > \ell_*$ can be divided into $\lceil L/\ell_*\rceil$ equal parts. Thus the sum in (65.6), after this fixed subdivision, is at most $N_{\mathrm{im}}\Gamma\theta_*$.

Each new piece is now an actual homogeneous stable graph in $I\mathcal{W}^u_R(B_{\mathrm{cv}})$. The stable inverse form of Theorem 25.1 applies to its remaining $N-1$ backward collisions. The chain rule, with the first-step supremum paid only once per first piece, gives

$$\sum_{V\in\mathcal{G}_N^{\mathrm{aff}}(W)} \sup_V J_V T_R^N \le C_g \sum_{V_1\in\mathcal{G}_1^{\mathrm{aff}}(W)} \sup_{V_1} J_{V_1} T_R \le C_g N_{\mathrm{im}}\Gamma\theta_*.$$

This includes $N=1$, since $C_g \ge 1$. At $N=0$ the analogous sum is one.

For coverage, the physical first-hit cuts are finite on each short graph, the target homogeneity cuts are countable, and the additional length cuts are finite on each homogeneous image. Finite iteration of a countable collection remains countable. Each regular branch is injective, so the points removed on the original W are countable. This is the actual curve-null argument; it is not an inference from the bulk measure of a singular set.

We need a little more than this null statement for endpoints. Fix a point $p \in W$ with a regular backward record of length N. On a neighborhood of p the finitely many actual iterates T_R^{-j} are smooth and have the same physical word. Each of their collision angles stays in a

compact subinterval of $(-\pi/2, \pi/2)$. Only finitely many homogeneity levels can therefore occur in a smaller neighborhood. On the corresponding stable image the derivative of its angle is nonzero, by cone invariance. A fixed such level cuts a one-sided regular graph at most once. The length subdivisions are locally finite there as well: their interior cut points are separated by the fixed positive image length $\ell_*/2$, and the finitely many image maps on this neighborhood have finite derivatives. Induction through the finitely many times gives finitely many cut points near p. Consequently, if p was removed as a cut, a one-sided adjacent open piece reaches it. The original finite regular map and the initial affine tangent are smooth at p, so their curve Jacobian extends to that endpoint with the actual derivative value. No derivative at a true grazing state is used. □

65.2. **All actual words, strip faces, and unique lifts.** Let $D^-_{N,R}$ be the open set of collision states having an actual regular backward record through $0, -1, \dots, -N$. All first collisions on these records are the physical first collisions; an earlier unvisited tangency is part of the excluded regularity boundary. For $y \in D^-_{N,R}$, lift its current scatterer to $O_0 = 0$. Its past then determines the unique tuple

$$((O_0, O_{-1}, \dots, O_{-N}); h_{-1}, \dots, h_{-N}), \qquad T_R^{-j} y \in H_{h_{-j}}. \tag{65.7}$$

The indices h_{-j} use the fixed original disjoint assignment of horizontal homogeneity boundaries. In particular the terminal past strip at $-N$ is included, and the current strip at 0 is not. Denote by $\mathfrak{C}_N(R)$ the nonempty sets of states with a fixed tuple (65.7). A class is the union of all its physical components; it is not counted again for each disconnected piece, section cut, or periodic lattice translate. Boundary-only faces allowed by the disjoint strip convention are also classes.

There are countably many such classes. Indeed the uniform finite flight bound gives a finite alphabet of physical center increments, and a finite tuple of homogeneity indices is countable. The classes are disjoint and cover $D^-_{N,R}$ exactly. Each is a countable union of compact sets in the collision cylinder: a fixed regular physical word has an open domain, the strip intervals with their assigned endpoints are countable unions of closed sets, and the finitely many relevant iterate maps are continuous on that domain. Intersect with a compact exhaustion of the domain to obtain the assertion. This observation also applies to classes consisting only of artificial-boundary points.

Write M'_R for the original set of points with regular two-sided orbits, and let $j_N(x) = J^s T_R^N(x)$ be its true stable arclength cocycle. For $C \in \mathfrak{C}_N(R)$ define

$$d_N(C) = \sup_{y \in C \cap M'_R} j_N(T_R^{-N} y), \qquad E_R(C) = \pi_R(C), \qquad \mathrm{cap}_R(C) = |E_R(C)|. \tag{65.8}$$

The supremum of an empty set is defined to be zero. The measure in the capacity is ordinary angle Lebesgue measure on $\mathbb{R}/2\pi\mathbb{Z}$, of total mass 2π. Because the projection is continuous and the classes are countable unions of compact sets, $E_R(C)$ is Borel. True stable invariance gives the exact identity

$$j_N(T_R^{-N} y) = \frac{1}{J_{E^s(y)} T_R^{-N}(y)} \qquad (y \in M'_R). \tag{65.9}$$

In particular the weights are finite, with the elementary bound $d_N(C) \le \Gamma \Lambda_*^{-N}$. There is no collision-flux, section-conditioning, or radius normalization in this tangent identity.

Lemma 65.2 (One short test pays every class it meets)**.** *For an affine stable source W as in Lemma 65.1,*

$$\sum_{\substack{C \in \mathfrak{C}_N(R) \\ C \cap W \neq \varnothing}} d_N(C) \le 3 C_\times C_{\mathrm{aff}} \qquad (N \ge 1). \tag{65.10}$$

This includes every disconnected class and every regular strip face which meets W.

Proof. Fix a class with $d_N(C) > 0$ which meets W, and choose a point $p_C \in C \cap W$. It need not have a regular future. It does have the required finite regular past. For every $y \in C \cap M'_R$ the two backward words agree, and their angles lie in the closures of the same strips at exactly $-1, \dots, -N$. The true stable direction at y and the affine tangent at p_C are both in $[-U, -a]$. Corollary 61.5 and (65.9) therefore give

$$d_N(C) \le \frac{C_\times}{J_W T_R^{-N}(p_C)}. \tag{65.11}$$

Taking the class supremum has cost only this common constant. No connecting path inside the class and no continuity of the true stable field on a test curve have been assumed.

Let $\mathcal{I}_N(W)$ be the open pullbacks to W of $\mathcal{G}_N^{\mathrm{aff}}(W)$. If p_C is in an interval $I \in \mathcal{I}_N(W)$, charge C to that interval. All interior points of I have the same actual word and the same strip indices, so at most one distinct class can be charged to its interior. If p_C is a removed cut, the endpoint part of Lemma 65.1 allows it to be charged to an adjacent interval. An interval has at most two endpoints, and each endpoint belongs to at most one class under the fixed disjoint convention. It can therefore receive at most two additional distinct endpoint classes. This gives multiplicity at most three, regardless of the number of components or intersections belonging to one class. One representative is chosen per class, not per intersection.

If $V = T_R^{-N} I$ is its charged image piece, continuity at a regular endpoint, when needed, implies

$$\frac{1}{J_W T_R^{-N}(p_C)} \le \sup_V J_V T_R^N.$$

Sum (65.11), use multiplicity three, and apply (65.5). The argument applies first to finite sets of classes and then to their increasing countable union. Classes of zero weight do not affect the sum. □

65.3. **Uniform capacity and the long-class consequence.** Put

$$N_{\mathrm{aff}} = 2 + \left\lceil \frac{2S_{\mathrm{aff}}}{\ell_*} \right\rceil, \qquad K_{\mathrm{tc}} = 3N_{\mathrm{aff}} C_\times C_{\mathrm{aff}}, \qquad \beta_{\mathrm{tc}} = \frac{2a}{\sqrt{1+U^2}} > 0. \tag{65.12}$$

Theorem 65.3 (All-class physical outgoing-angle capacity)**.** *For every radius in the original family and every $N \ge 1$,*

$$\sum_{C \in \mathfrak{C}_N(R)} d_N(C) \mathbf{1}_{E_R(C)}(\vartheta) \le K_{\mathrm{tc}} \qquad \text{for every } \vartheta \in \mathbb{R}/2\pi\mathbb{Z}, \tag{65.13}$$

$$\sum_{C \in \mathfrak{C}_N(R)} d_N(C) \operatorname{cap}_R(C) \le 2\pi K_{\mathrm{tc}}. \tag{65.14}$$

Consequently, for every $\delta > 0$,

$$\sum_{\substack{C \in \mathfrak{C}_N(R) \\ \operatorname{cap}_R(C) \ge \delta}} d_N(C) \le \frac{2\pi K_{\mathrm{tc}}}{\delta}. \tag{65.15}$$

If $\mathfrak{L}_{N,R}(\delta)$ denotes the classes which contain a connected physical unstable-cone arc of arclength at least δ, with slopes in $[\kappa_R, U]$, then

$$\sum_{C \in \mathfrak{L}_{N,R}(\delta)} d_N(C) \le \frac{2\pi K_{\mathrm{tc}}}{\beta_{\mathrm{tc}} \delta}. \tag{65.16}$$

The constants are uniform before any supremum in radius or collision count. No connectedness, minimum area, or positive lower bound for the capacity of an arbitrary class is required.

Proof. Parametrize $\Gamma_{R,\vartheta}$ by arclength on its open interval of length $L \le S_{\mathrm{aff}}$. For $n = \lceil 2L/\ell_* \rceil$, the open windows with centers at jL/n, $0 \le j \le n$, and half-width L/n, restricted to that interval, cover it and each have length at most ℓ_*. There are at most N_{aff} windows. They are the short affine sources of Lemma 65.2. They also cover a point at the collision-circle seam, since the original fiber is the smooth arc (65.2), not two arcs stopped at that seam.

Every class with $\vartheta \in E_R(C)$ meets at least one window. Sum (65.10) over the windows; counting a class more than once only increases this positive upper bound. This proves the pointwise assertion (65.13). Apply Tonelli's theorem to its countable nonnegative sum on the outgoing angle circle to obtain (65.14). Restricting to classes of capacity at least δ proves (65.15).

For the last assertion orient a connected unstable-cone arc so that s and φ increase in a lift, and write $v = d\varphi/ds \in [\kappa_R, U]$. Its outgoing-angle lift satisfies

$$\frac{d\vartheta_{\mathrm{lift}}}{d\ell} = \frac{v + \kappa_R}{\sqrt{1+v^2}} \ge \frac{2a}{\sqrt{1+U^2}} = \beta_{\mathrm{tc}}. \tag{65.17}$$

Moreover,

$$0 < d\vartheta_{\mathrm{lift}} = \left(1 + \frac{\kappa_R}{v}\right) d\varphi \le 2\, d\varphi.$$

Since the collision angle is in $(-\pi/2, \pi/2)$, the total outgoing-angle variation is at most 2π. Thus passage to the angle circle causes no loss of its one-dimensional image measure. The projection of an arc of length at least δ has measure at least $\beta_{\mathrm{tc}}\delta$ and is contained in $E_R(C)$. Apply (65.15) with this threshold. For an arc specified through a supremum of lengths, first take shorter arcs and then pass monotonically to the stated length. This proof uses the actual monotone cone arc, and does not infer a long arc from a large Euclidean diameter or a cell-shape assertion. □

65.4. **Exact pressure, optical, and observation scope.** Let $\mathcal{P}_N(R)$ denote the unchanged physical-word pressure of Input 11.1. Under $y = T_R^N x$, translation of the current lifted center to zero maps the forward word $(0, O_1, \dots, O_N)$ bijectively to the backward word

$$(0, O_{N-1} - O_N, \dots, O_1 - O_N, -O_N).$$

This is reversal of the ordered record followed by one lattice translation. It preserves the actual collision states and tangent norms. Distinct periodic lifts of this same record do not make new classes. Since the homogeneity convention assigns every finite regular state exactly once, including artificial strip faces,

$$\mathcal{P}_N(R) = \sum_{w \in \mathcal{W}_N(R)} \sup_{\substack{C \in \mathfrak{C}_N(R) \\ C \text{ has reversed word } w}} d_N(C) \le \sum_{C \in \mathfrak{C}_N(R)} d_N(C). \tag{65.18}$$

There is no deletion of boundary values in this comparison of suprema. All true grazing points remain outside M_R', as in the original definition of the stable derivative. The finite regular classes cover a full collision-measure set for each fixed radius and finite time; a common null set for uncountably many radii is neither used nor claimed.

Theorem 65.3 pays every term after multiplication by its actual outgoing-angle capacity. It also pays the entire unweighted long-class sum (65.16), without a planar-cell intersection or continuation assumption. It does not bound the unweighted sum on the right of (65.18) merely because small-capacity classes are short. More precisely, for any fixed $\delta > 0$,

$$\mathcal{P}_N(R) \le \frac{2\pi K_{\mathrm{tc}}}{\delta} + \sum_{w \in \mathcal{W}_N(R)} \sup_{\substack{C \in \mathfrak{C}_N(R) \\ C \text{ has reversed word } w \\ \mathrm{cap}_R(C) < \delta}} d_N(C). \tag{65.19}$$

This states exactly the still-needed short-capacity payment along this route; it is not an assertion that the last positive sum vanishes or is uniformly bounded. A complete physical renewal or

another genuine geometric payment must provide that additional bound. The geometric capacity in this section must in turn be connected to the original conditioned physical-roof source before it can supply a same-roof probability estimate.

The optical comparison already proved in module 313 remains endpoint-exact:

$$\frac{c(T_R^N x)j_N(x)}{30000} \le \omega_N(x) \le 30000c(T_R^N x)j_N(x).$$

For example the class weight

$$o_N(C) := \sup_{y\in C\cap M_R'} \frac{\omega_N(T_R^{-N}y)}{c(y)}$$

satisfies $30000^{-1}d_N(C) \le o_N(C) \le 30000d_N(C)$; the corresponding capacity moment is at most $30000 \cdot 2\pi K_{\mathrm{tc}}$. This comparison retains the actual terminal incidence, and introduces neither an incidence cutoff nor a fictitious measure normalization.

For a fixed Borel observation event A of the actual finite record, one may replace $d_N(C)$ by the supremum over $C \cap A \cap M_R'$ in every positive upper bound above. Such an event may specify the original section and return event, exact displacement and collision labels, and any interval of the original physical roof. The upper bound is uniform in this restriction, but gives no factor equal to the roof-interval width and does not authorize summing separately repeated class suprema over arbitrary additional labels. In particular $1/(4\pi R)$ and $1/c_*$ remain the inherited collision-flux and section-conditioning factors at their original measure interfaces; they are absent from the tangent and angle-capacity identities here. Neither the original m^2 normalization nor the prescribed outer-B, collision-lim sup order is changed. The final raw pointwise local-limit and four-coordinate inversion target consequently has precisely its previous physical and probabilistic scope.

66. Positive critical marked sources with collapsing clearance slices

A physical corner can have a positive incoming critical fan although its angular section at zero clearance has collapsed. Such a fan is not an angular-null error. We now pay its actual leading coefficient by transporting finitely many positive physical sectors to a common nearby positive clearance, and using the branch-supremum theorem there. This argument concerns the original unvisited level $u = 0$ and permits simultaneous unvisited tangencies and original section cuts. All visited incidences at a limiting germ remain positive. A trace at an actual visited grazing collision is a different measure.

Fix $R \in [9/20, 47/100]$, $m \ge 1$ and $0 \le j < m$. A marked critical germ has the original m-flight strict visited word, normal endpoints $p_0 = p_m = 0$, and an unvisited marked disk with a perpendicular interior foot on its jth flight at clearance zero. Its local physical side consists of actual regular records with $u > 0$, all the original other-obstacle inequalities, and the specified original section and finite record tests. Original physical roof intervals may also be imposed. Germs with the same limiting marked state and strict visited word have their physical side sets united and disjointized, not counted a second time. All actual marked disks remain separate marked copies.

Use the native normalized coordinates $y = G_x^{1/2}(q - q_x)$ and the physical angular fraction $\beta_y(x) \in [0, 1]$ from Section 27. The incoming leading coefficient is

$$\mathfrak{J}(x)\beta_y(x), \qquad \mathfrak{J}(x) = \frac{\sqrt{\det G_x}}{2R^2c_*}\,u_x, \qquad u_x = \frac{R|G_{0m}(x)|}{\det G_x}. \tag{66.1}$$

These are the actual auxiliary-action boundary values of (27.1). No derivative of the singular physical collision map is assigned at an unvisited tangency. The physical map equals this auxiliary map on the actual regular side.

66.1. **A physical representative at every small positive clearance.**

Lemma 66.1 (Analytic sectors and common one-sided levels)**.** *At a fixed marked critical germ as above, its finite physical-side conditions can be written locally as a finite Boolean combination of real analytic inequalities, including $u > 0$. The limiting angular indicator exists outside finitely many rays. If $\beta_y(x) > 0$, there are actual regular points $x(u)$ on this physical side for every sufficiently small $u > 0$, with their designated marked clearance exactly u and $x(u) \to x$. Representatives may additionally avoid the cuts and exceptional angular sets of both partitions in Theorem 62.1. For any finite collection of such germs they may be chosen at every common sufficiently small $u > 0$.*

Proof. Positive visited incidences give an invertible internal optical Hessian. The local strict-word reflection equations and all contact positions are therefore real analytic functions of the endpoint coordinates by the implicit-function theorem. The flight lengths are bounded away from zero. Fixing the finite word and a sufficiently small neighborhood, only finitely many lattice centers can meet one of its flights or its clearance neighborhood. Other centers have a fixed positive distance and impose no active local cut. The first-hit and unvisited-obstacle requirements are finite analytic sign conditions: the perpendicular-foot tests and squared line-distance tests may be used on their respective finite cases. Endpoint distance tests are analytic as well. The designated marked foot is strictly interior and separated from the visited contacts by the original standoff estimate.

Each of the original finitely many section rectangles is given by coordinate inequalities pulled back through finite regular maps. Occupation is a finite Boolean combination of those tests; the lifted displacement is fixed on the strict word. The physical roof is the same analytic length sum, so a fixed roof interval adds two analytic inequalities, including at a critical roof value. No replacement section or time is used. This reasoning concerns these original finite analytic tests; it makes no ray-persistence assertion for an arbitrary additional Borel selector.

In any analytic local coordinate z centered at the germ, remove constant truth values and identically zero functions from the finite Boolean formula. For each remaining analytic function f with $f(0) = 0$, let P_f be its first nonzero homogeneous Taylor term. The nonzero polynomial P_f vanishes on only finitely many rays of the unit circle. On each complementary open angular interval, the signs of all $f(re)$ are fixed for sufficiently small $r > 0$, uniformly on a compact smaller interval. The formula thus has a limiting indicator there. Boundary rays have zero angular measure. An invertible linear coordinate change preserves positivity of the angular measure, so $\beta_y > 0$ supplies at least one such open interval of actual physical directions. In the inherited local radial coarea formula the optical coefficient is continuous and bounded at this fixed positive-visited germ. Dominated convergence of the angular indicators therefore gives exactly $\mathfrak{J}(x)\beta_y(x)$ as its leading one-sided coefficient. In particular zero opening contributes zero to this leading coefficient; this is not an assertion about bulk-null traces or about a collision-uniform coarea remainder.

Take $z = (p_0, p_m)$. Its local change from q is invertible by the positive Hessian and nonzero optical corner in Lemma 27.1. The exact half-ray determinant of Theorem 59.1, continued on the same strict word to the limit, makes (ϑ, u) another local analytic coordinate system:

$$\left|\det D_{(\vartheta,u)}(p_0, p_m)\right| = A_-^p A_+^p (h_- + h_+) > 0.$$

Consequently $D_z u(0) \neq 0$. Since the physical side includes $u > 0$, a positive physical angular interval contains a direction e with $D_z u(0)e > 0$, after omitting the two zero directions of this linear form. The ray $z = re$ satisfies all selected physical sign conditions strictly for small $r > 0$, after constant truth values are accounted for. Along it

$$u(re) = rD_z u(0)e + O(r^2), \qquad \frac{d}{dr}u(re) = D_z u(0)e + O(r) > 0.$$

The one-variable inverse-function theorem supplies a representative at every small positive u, converging to the germ.

At each such representative the same sign conditions persist on an open neighborhood. Intersecting with its fixed-u coordinate line leaves a nonempty open angular interval. The two one-sided partitions in 319 omit only angular-null sets at that fixed level. Choose a point in this interval outside their union, at angular distance tending to zero from the original representative as $u \downarrow 0$. It still satisfies the same physical tests and still converges to x. For a finite family, take the minimum of the finitely many positive level ranges and perform these choices for every common u in that range. No common neighborhood size over all words, radii or collision counts is asserted. □

66.2. **The critical branch-supremum transfer.**

Lemma 66.2 (Both half-ray sums on positive physical critical germs)**.** *Let $\mathcal{F}_{m,j}(R)$ be the complete one-copy family of the specified zero-clearance marked critical germs with $\beta_y(x) > 0$. At their limiting strict-word optical values,*

$$\sum_{x \in \mathcal{F}_{m,j}(R)} (A^p_-(x))^{-1} \leq H_{\mathrm{nm}}, \qquad \sum_{x \in \mathcal{F}_{m,j}(R)} (A^p_+(x))^{-1} \leq H_{\mathrm{nm}}. \tag{66.2}$$

There is no assumption that a germ has a nonempty regular angular interval at $u = 0$.

Proof. First fix any finite set $\mathcal{F}$ of distinct merged germs. Translate each designated marked disk to the origin and retain its impact sign. A limiting angular direction and that sign determine one marked tangent line. Following its two halves, the next strictly met disk is unique, even when the limiting line is tangent to other earlier disks. Continuing through its positive-incidence reflections determines the complete strict visited word. Using an earlier tangency as a visited collision would introduce a zero incidence and would belong to a different, here unasserted stratum. This is the strict-hit uniqueness argument of Theorem 27.4, applied also backwards to recover the original initial disk at index $j + 1$.

Thus distinct merged germs with the same impact sign have distinct limiting angular points. If they give the same marked point, they are the same strict-word germ and their side selectors have already been united. The original initial-disk translation is unique and does not add a lattice multiplicity. Choose an angular seam avoiding the finitely many limiting points. The constants in 319 are uniform in this initial seam choice.

For every common sufficiently small $u > 0$, take representatives from Lemma 66.1 inside intervals of both partitions, away from their cuts and exceptional angular sets. For small enough u, no two representatives with the same sign can lie in the same backward interval $I \in \mathcal{P}^-_{j+1}$. Otherwise monotonicity and the pointwise lower bound in 319 give

$$|p_0(x(u)) - p_0(\widehat{x}(u))| \geq k_- a_\circ |\vartheta(x(u)) - \vartheta(\widehat{x}(u))|.$$

The left side tends to zero, whereas the right side stays bounded below by the separation of two distinct limiting angles. A finite family has a common positive minimum separation for each sign, so this contradiction excludes every such pair for all sufficiently small u. The same argument uses p_m and $\mathcal{P}^+_{m-j}$ for the forward halves. Both momenta tend to zero because the limiting endpoints are normal. No derivative of the selectors is involved.

The representatives therefore occupy distinct backward intervals, and also distinct forward intervals. Each reciprocal half derivative is bounded by the supremum of its actual interval. Applying 319 at this common positive u gives

$$\sum_{x\in\mathcal{F}}(A_{\pm}^{p}(x(u)))^{-1}\le H_{\mathrm{nm}}.$$

Every visited incidence at each fixed limiting germ is positive, so its finite strict-word optical matrix depends continuously on these representatives. Passing $u\downarrow 0$ in the finite sum proves the two inequalities for $\mathcal{F}$. Taking the supremum over finite subsets proves (66.2). Equivalently, this defines the full nonnegative sum without choosing an order; finiteness implies at most countably many positive terms.

The limit was taken at fixed R,m and finite $\mathcal{F}$ before that exhaustion. Uniformity in R,m comes from the same final constant, not from an interchange of local and long-time limits. In particular a collapsed angular slice at level zero never had to be populated by actual regular points. □

Theorem 66.3 (Complete positive marked critical-fan payment)**.** *Put* $\sigma=47/59$ *and retain* Q_{mb} *from* (62.1)*. For the original zero-clearance marked physical critical family with positive visited incidences,*

$$\sum_{x:\,\beta_y(x)>0}u_x\le Q_{\mathrm{mb}}\,\sigma^{\max(j,m-1-j)},\tag{66.3}$$

$$\mathfrak{F}_{m,j}(R):=\sum_{x}\mathfrak{J}(x)\beta_y(x)\le C_{\mathrm{pc}}\,\sigma^{\max(j,m-1-j)},\qquad C_{\mathrm{pc}}=\frac{170}{18c_*}\,Q_{\mathrm{mb}}.\tag{66.4}$$

Both impact signs, all actual marked disks and strict words, and all simultaneous finite original physical cuts are included. The bounds are uniform in radius and collision count and may be restricted to any original exact record or to critical values in any interval of the unchanged physical roof. Any further weights between zero and one can decrease this already bounded positive sum.

In particular the complete sum over all *marked flight indices, including shallow indices, satisfies*

$$\sum_{j=0}^{m-1}\mathfrak{F}_{m,j}(R)\le C_{\mathrm{pc}}m\,\sigma^{(m-1)/2},\qquad \sup_{R}m^{2}\sum_{j=0}^{m-1}\mathfrak{F}_{m,j}(R)\le C_{\mathrm{pc}}m^{3}\sigma^{(m-1)/2}\longrightarrow 0.\tag{66.5}$$

The same conclusion holds after the original exact-label suprema. No outer width or guard is changed.

Proof. The two actual half counts are $j+1$ and $m-j$. At the limiting strict-word germ, continuity and the positive optical recurrence of 319 give

$$(A_{-}^{p})^{-1}\le E_{*}\sigma^{j},\qquad (A_{+}^{p})^{-1}\le E_{*}\sigma^{m-1-j}.$$

The same exact optical determinant gives

$$u_x\le C_{\mathrm{foot}}(A_{-}^{p})^{-1}(A_{+}^{p})^{-1}.$$

Bound either factor individually and sum the other by Lemma 66.2. The two resulting bounds are $Q_{\mathrm{mb}}\sigma^{j}$ and $Q_{\mathrm{mb}}\sigma^{m-1-j}$. Their minimum is (66.3), since $0<\sigma<1$.

Use the actual coefficient identity (66.1) and the sharp inherited critical bound $\sqrt{\det G_x}/R^{2}\le 170/9$ from Lemma 27.1. The united one-copy fraction satisfies $0\le\beta_y\le 1$. Zero-fraction germs contribute zero by their defining angular coefficient, not by a bulk-null claim. Thus

$$\sum_{x}\mathfrak{J}(x)\beta_y(x)\le\frac{170}{18c_*}\sum_{\beta_y(x)>0}u_x,$$

proving (66.4). One conditioning factor and the original normalized fan fraction are retained.

For every $0 \le j < m$, $\max(j, m-1-j) \ge (m-1)/2$. Sum the nonnegative bounds and multiply by m^2 to obtain (66.5). Its scalar right-hand side tends to zero by geometric decay and is independent of radius, labels and chosen roof-value restrictions. Full positive sums precede every such supremum. The local representatives were used only to prove the finite-count inequalities, and no original outer limit is exchanged. □

What has been closed and what remains. Theorem 66.3 pays the complete *leading* incoming zero-clearance critical-fan coefficient in this actual physical stratum, including positive corner fans with collapsed angular slices and every shallow marked index. It extends the inherited unscaled coefficient budget by an explicit collision-normalized vanishing bound. It does not assert a uniform size for all local critical coordinate neighborhoods, a collision-uniform sum of the local coarea remainders, or a trace theorem at visited grazing. An arbitrary Borel pairing selector need not have the analytic sector property proved above; it may only restrict the already bounded family when its own leading coefficient exists with weight between zero and one.

The full regular angular marked charge is controlled by 320, and the leading physical marked critical fan by this section. Distinct visited-grazing, section/pairing boundary measures, additional multipliers on those boundary measures, nonleading critical remainders and the original same-roof positive-source comparison retain their exact remaining obligations. In particular (0.1) and the four-coordinate raw inverse theorem are not asserted complete.

67. Complete physical roof variation without a margin cutoff

The full-source weak roof estimates of Section 55 leave open a pointwise concentration at a common physical roof. We now bound that common-roof density by summing its actual interior variation and all geometric entrance and exit charges. This is a directional variation argument in the physical roof variable. No variation in the Gauss angle is asserted. The local critical remainders are included in the exact interior calculation; they are not summed after a word-dependent Taylor approximation.

Set $a = 5/6$, and retain the original regular endpoint cap

$$\mathcal{C}_{m,R} = \{x : |p_0(x)|, |p_m(x)| < a, \quad x \text{ is an actual regular } m\text{-flight record}\}.$$

Its boundary has collision area zero, so the strict convention has the same bulk measure as the closed convention in 314. Every original exact event lies in the corresponding closed cap. Define the positive majorant, on the unchanged unit-speed roof line, by

$$\overline{\mu}_{m,R} = (L_{m,R})_*(\mathbf{1}_{\mathcal{C}_{m,R}}\, d\nu/c_*). \tag{67.1}$$

This is not a new probability or a new conditioning event: its mass is at most c_*^{-1}, and the original initial-section indicator is restored by positive restriction. In particular, for every original exact record and every Borel $0 \le H \le 1$,

$$(L_{m,R})_*(\mathbf{1}_{\Pi_{n,k,m,R}} H\, d\nu_R^*) \le \overline{\mu}_{m,R}. \tag{67.2}$$

Neither side changes the collision count, physical clock or labels.

Keep the already fixed constants $\lambda_\circ, \Lambda_\circ, C_\omega, C_F, J_\times, C_{\mathrm{mom}}, C_{\mathrm{ab}}$ and $\sigma = 47/59$. Define

$$C_{\mathrm{int}} = \frac{\Lambda_\circ}{\lambda_\circ}\left(C_\omega + \frac{2C_F}{\lambda_\circ}\right), \tag{67.3}$$

$$D_{\mathrm{roof}} = \frac{C_{\mathrm{int}} J_\times}{c_*} + \frac{2C_{\mathrm{ab}}}{1-\sigma} + \frac{\Lambda_\circ C_{\mathrm{mom}}}{2c_* R_-^2} + \frac{U_2 C_{\mathrm{mom}}}{2c_*}, \qquad U_2 = 320/9, \qquad C_{\mathrm{roof}} = D_{\mathrm{roof}}/2.$$

These constants are fixed by the original physical family and the native homogeneous curve class, before every collision, radius, label or roof supremum. They contain no whole-word pressure or comparison-capacity assumption.

67.1. **The unweighted interior and the actual cap boundaries.**

Lemma 67.1 (The full regular interior has finite roof variation)**.** *On a regular fixed-Gauss-angle physical segment put $t = F$, $g = \nabla_q F$, $z = |g| > 0$ and $A = \omega/(4\pi c_* \chi)$. Then*

$$|\partial_t A| \le C_{\mathrm{int}} A I_m / z. \tag{67.4}$$

On the complete cap source, with all actual words summed,

$$\int_0^{2\pi} \sum_w \int_{\mathcal{D}_w(\theta)} |\partial_t A_w(t,\theta)|\, dt\, d\theta \le C_{\mathrm{int}} J_\times / c_*. \tag{67.5}$$

Here $\mathcal{D}_w(\theta)$ is the open set of actual cap records of the indicated word on that Gauss ray. Missing physical intervals are not filled in.

Proof. The proof of Theorem 30.4, before differentiating its guard, gives

$$|\partial_t \log A| \le \frac{\Lambda_\circ}{\lambda_\circ z} \left(C_\omega I_m + \frac{2C_F}{\lambda_\circ} \right).$$

The endpoint bounds used there depend only on $|p_0|, |p_m| \le 9/10$ and hence hold on this cap. Since $I_m \ge 1$, this proves (67.4), with no guard derivative and no factor ε^{-1}.

The exact endpoint-coordinate coarea formula in Theorem 10.2 and Proposition 10.4 applies to the complete regular cap just as to its original exact-event subsets: the uniqueness proof uses the physical word, and not the section test. The critical set has zero collision area. Consequently

$$\int_0^{2\pi} \sum_w \int_{\mathcal{D}_w(\theta)} A_w I_m / z\, dt\, d\theta = c_*^{-1} \int_{\mathcal{C}_{m,R}} Z_m\, d\nu \le J_\times / c_*.$$

The last inequality is the complete physical multiplier theorem 56.3; no independent product law is used. All equalities follow first on compact regular subsegments, then by nonnegative exhaustion. This proves the asserted full sum. □

Lemma 67.2 (The four endpoint-cap charges)**.** *The complete positive roof–angle entrance/exit charge on the four regular sides $p_0 = \pm a$, $p_m = \pm a$, with the other endpoint momentum in $[-a, a]$, is at most*

$$B_{\mathrm{cap}} = \frac{\Lambda_\circ C_{\mathrm{mom}}}{2 c_* R_-^2}. \tag{67.6}$$

All actual physical words are included before any original record restriction. Visited singular intersections have the separate zero trace stated below.

Proof. Consider $p_0 = a$ and write $v = p_m$. Since $a > 0$, the Gauss angle of $(-a, v)$ has absolute differential $a(a^2 + v^2)^{-1} dv$. The exact optical identity of Corollary 59.2 is

$$A = \frac{u_x}{4\pi c_* h_x}, \qquad h_x = R^2 e^{\mathsf{T}} G^{-1} e, \qquad A \le \frac{\Lambda_\circ}{4\pi c_* R_-^2} u_x.$$

The complete physical momentum-root bound of Theorem 26.4 applies at the common pair (a, v). Thus this side has charge at most

$$\frac{\Lambda_\circ C_{\mathrm{mom}}}{4\pi c_* R_-^2} \int_{-a}^{a} \frac{a}{a^2 + v^2}\, dv = \frac{\Lambda_\circ C_{\mathrm{mom}}}{8 c_* R_-^2}.$$

The remaining sides have exactly the same integral, either by changing the sign or interchanging the two endpoint coordinates. Summing four positive bounds proves (67.6). Cap corners are

assigned once; they cannot add a charge by duplicating a coordinate chart. This argument evaluates the proved pointwise root sum on its actual regular roots. It does not restrict an arbitrary version of a two-dimensional density to a null line. □

67.2. **Exhausting all physical entrances and exits.**

Lemma 67.3 (Complete fixed-count physical endpoint accounting)**.** *Fix R, m. Up to a null set of Gauss angles, every endpoint of a maximal regular fixed-angle interval in the cap has one of the following actual sources: a cap side; an unvisited-disk tangency; an interior visited grazing collision; or $g = 0$. The complete positive charge of the first two types is bounded by*

$$B_{\mathrm{cap}} + \frac{2C_{\mathrm{ab}}}{1-\sigma}. \tag{67.7}$$

The third type has zero coarea endpoint trace. The complete incoming charge at $g = 0$ is at most

$$B_{\mathrm{crit}} = \frac{U_2 C_{\mathrm{mom}}}{2c_*}. \tag{67.8}$$

There is no additional charge at a homogeneity cut, angular seam, nearest-obstacle switch or auxiliary coordinate boundary.

Proof. At fixed R, m the lifted word family is finite, by the actual finite horizon and lattice bound. The global uniqueness theorem 10.2 makes the regular roof–angle image for one word an open set, even when its physical domain is disconnected. At a fixed angle its maximal components are disjoint open roof intervals. The endpoint-limit argument in Theorem 5.3 applies without a section selector: both endpoint angles, every designated contact, flight, incidence and third-disk clearance have limits. Flight lengths stay at least τ_-. The cap keeps both endpoint incidences positive. If the limit has $g \neq 0$, no visited grazing, positive third-disk clearance and a strict cap inequality, the local physical map and roof–angle inverse extend through it. It therefore is not an endpoint of a maximal component. This proves the list. For a disjoint charge assignment, assign visited-grazing limits first, then the remaining critical limits. Among noncritical limits, assign an active unvisited constraint before a cap constraint. Thus a residual cap arc consists of actual regular roots of the momentum map, as required in Lemma 67.2.

Here is a precise way to account for simultaneous boundaries and coordinate limits. The closure of the finite-word configurations in the cap is compact. The formal analytic endpoint charts of Theorem 33.2, including their visited-grazing limits, give a finite local cover for this fixed R, m. On each such chart, actual admissibility is a finite Boolean combination of analytic sign conditions: the signed visited incidences and the distances and foot tests for finitely many candidate disks. Endpoint-distance cases are retained as well. This finite list follows from the horizon, not a sampled list of encountered disks. The cap adds four analytic inequalities.

Where $g \neq 0$, the continuous positive reduced Hessian in 260 makes (t, θ) local analytic coordinates, also for the formal calculation at a one-sided limit. On a fixed-angle slice a nonconstant analytic boundary function has isolated zeros on every compact subinterval. If it is identically zero on that slice it imposes a constant truth value there; exceptional angular levels of this type may be discarded when the function is not identically zero in a neighborhood. Indeed an accumulating sequence of such angular levels would make the two-variable analytic function identically zero. Identical constraints are removed from the local Boolean description. Thus Fubini reduces the calculation to ordinary one-dimensional intervals. A smooth partition of unity on the finite chart cover introduces artificial terms whose sum is zero before absolute values are taken. Equivalently one may use the global one-copy word image and cancel its artificial subinterval endpoints first. No new physical component is charged.

At a nongrazing unvisited boundary, its closest foot is strictly inside the indicated flight: a third disk is separated from either incident contact by the positive disk gap. The marked-foot standoff therefore applies. The signed line-clearance differential is nonzero by the actual determinant in Theorem 59.1. A regular boundary arc contributes its trace times $|d\theta|$, exactly as in Theorem 58.1. An arc on which the angle derivative vanishes has zero angular image, by the one-dimensional argument in that theorem. Intersections or endpoints with zero angular image introduce no extra interval-endpoint integral. If several disk constraints are active, assign the physical boundary germ to the first of them in a fixed ordering; the full positive marked sum is an upper bound. Coincident boundary arcs are assigned once, and isolated corners are treated by the same fixed-angle slicing. All actual marks, words and both impact signs are already present in Theorem 64.1 at $u_0 = 0$.

The proof of that theorem starts from all regular angular records and uses only the endpoint cap $|p_0|, |p_m| \le 9/10$; its later exact-record tests are optional positive selectors. It is therefore valid for the present smaller cap, without imposing an initial section. Also its proof bounds A itself: on this zero-clearance boundary the original W^{ε} equals one. Summing its finite-count estimate gives

$$\sum_{j=0}^{m-1} \mathfrak{B}_{j,0}^{\mathrm{reg}} \le C_{\mathrm{ab}} \sum_{j=0}^{m-1} (\sigma^{j} + \sigma^{m-1-j}) \le \frac{2C_{\mathrm{ab}}}{1-\sigma}.$$

Together with Lemma 67.2, this proves (67.7).

At any actually visited interior grazing limit the marked optical bound in 260 gives $\omega \to 0$, whereas $\chi \ge \lambda_{\circ}$. Hence $A \to 0$, even if $g \to 0$ at the same endpoint. This is the inherited zero *trace* proof, not an appeal to collision-area nullness. It is applied before an integral over endpoint angles. At fixed count the critical limiting configuration of a word, if it exists, is unique. The support inequalities of Lemma 10.1 pass to the limit; normal endpoints force identical endpoint contacts, and equality in the length support inequalities then forces the same successive entry or tangent contacts. Thus there are only finitely many such critical configurations at this stage.

For a critical germ with positive visited incidences, the local physical side is analytic. The cap and all visited-incidence tests are strict there. Every active physical boundary is therefore an unvisited clearance with the nonzero differential already proved above, so the specific nonzero-differential hypothesis of Theorem 27.4 is satisfied. In the gradient coordinates, the first nonzero homogeneous Taylor term of each active nonconstant inequality has only finitely many zero rays. Off those rays its sign is constant along every sufficiently short ray. This proves the angular sector limit, including simultaneous tangencies and collapsed slices, by exactly the analytic argument in Lemma 66.1. There is no section or arbitrary Borel selector in the majorant being differentiated. Its incoming angular trace is exactly $\mathfrak{J}(x)\beta_y(x)$, with the normalized angular change of variables of Lemma 27.3. The complete one-copy sector bound of Theorem 27.4 gives (67.8); it already includes regular critical points. At a visited-grazing critical germ the incoming trace is zero by the preceding optical argument. One may verify this by excising $|g| < \delta$ on its fixed local chart: $A \to 0$ uniformly there, while the angular range is at most 2π. A fixed finite family permits the local limit first, without a collision-uniform chart radius.

The positive Hessian makes every critical endpoint incoming on its fixed ray; no second outgoing critical charge is added. All local limits in this proof occur at fixed R, m, after which the displayed uniform bounds are applied. The complete regular interior and regular boundary are kept in their exact measures, so no uniform Taylor remainder or exchange of long-time and local limits is used. □

Theorem 67.4 (Uniform full-source variation and original-roof height)**.** *For every $m \geq 1$ and $R \in [9/20, 47/100]$, the measure* (67.1) *has a nonnegative density $\overline{p}_{m,R} \in BV(\mathbb{R})$ with*

$$\|D\overline{p}_{m,R}\|_{\mathrm{TV}} \leq D_{\mathrm{roof}}, \qquad \|\overline{p}_{m,R}\|_{\infty} \leq C_{\mathrm{roof}}, \qquad \|\overline{p}_{m,R}\|_{1} \leq c_{*}^{-1}. \tag{67.9}$$

In particular, for every original exact return, displacement and collision label, every Borel $0 \leq H \leq 1$, and every Borel physical roof set S, the unchanged source satisfies

$$\nu_{R}^{*}\big(\mathbf{1}_{\Pi_{n,k,m,R}} H \mathbf{1}_{\{L_{m,R} \in S\}}\big) \leq C_{\mathrm{roof}} |S|. \tag{67.10}$$

The corresponding density is at most C_{roof} almost everywhere. The complete physical sum precedes $\sup_{R,m,n,k}$ and the roof essential supremum.

Proof. The exact coarea formula gives the density

$$\overline{p}(t) = \int_{0}^{2\pi} \sum_{w} \mathbf{1}_{\mathcal{D}_w}(t,\theta) A_w(t,\theta)\, d\theta.$$

It is integrable with mass at most c_{*}^{-1}. On each maximal regular interval integrate $A_w \psi'$ by parts, for $\psi \in C_c^1(\mathbb{R})$. The endpoint trace and absolute continuity follow from the proof in 260 and 289 with weight one. First perform the calculation on a finite interval exhaustion. The absolute sums of interior derivatives and endpoint charges are finite after angular integration by the two preceding lemmas. The integral of $A_w|\psi'|$ is finite by the source mass. Consequently the exhaustion passes through the integrated identity and gives

$$\left| \int \overline{p}(t) \psi'(t)\, dt \right| \leq D_{\mathrm{roof}} \|\psi\|_{\infty}.$$

More explicitly, the derivative is the signed pushforward of the integrable interior derivatives and the finite entrance and exit trace measures just constructed. Thus it is a finite signed measure, and the inequality is the first bound in (67.9).

The density is supported in $[m\tau_{-}, m\tau_{+}]$. Its standard one-dimensional bounded-variation representative is zero outside that interval. To reach any nonnegative value v from zero and return to zero requires total variation at least $2v$. Approximation by Lebesgue points therefore gives $2\|\overline{p}\|_{\infty} \leq \|D\overline{p}\|_{\mathrm{TV}}$. This proves the remaining bounds. Finally (67.2) and positivity give (67.10) and its density version. At each fixed radius the null sets can be chosen for the countable original labels; no common null set across all real radii is asserted. □

Corollary 67.5 (The unchanged bad guard and exact roof Fourier payment)**.** *For the full cap majorant weighted by the original $W^{\varepsilon} = 1 - H^{\varepsilon}$, $\varepsilon > 0$, the same result holds with*

$$D_{\mathrm{roof}}^{\varepsilon} = \frac{C_{\mathrm{reg}}(1 + \varepsilon^{-1}) J_{\times}}{c_{*}} + \frac{2C_{\mathrm{ab}}}{1 - \sigma} + B_{\mathrm{cap}} + B_{\mathrm{crit}}. \tag{67.11}$$

Every original exact-event submeasure of this guarded majorant has roof density at most $D_{\mathrm{roof}}^{\varepsilon}/2$. For either of the two full cap majorants, if its variation bound is denoted by D and its density by p, then

$$\left| \int e^{ist} p(t)\, dt \right| \leq \min\{c_{*}^{-1}, D/|s|\} \quad (s \neq 0), \qquad \|p(\cdot + h) - p\|_{1} \leq D|h|. \tag{67.12}$$

The Fourier assertion in this corollary is for the specified full majorants. An arbitrary measurable restriction does not inherit their variation or Fourier bound from positive domination alone.

Proof. Use the exact regular-variation inequality of Theorem 30.4 in the interior. The unchanged guard is continuous on each one-sided finite-word closure and between zero and one. Every boundary charge is therefore bounded by the same unweighted charge; visited-grazing traces

remain zero. The preceding integration by parts proves (67.11). There is no change in the guard depth schedule or its physical collision and clearance factors.

For the Fourier bound, integrate by parts against the finite measure Dp, using compact support. For $s \neq 0$ the resulting factor is $1/(is)$; its absolute value gives $D/|s|$. The mass gives the other bound. For translations, the bounded-variation representative satisfies $|p(t+h) - p(t)| \leq |Dp|$ on the interval between t and $t+h$, up to endpoint conventions on null sets. Tonelli integration over t gives $D|h|$. □

The exact gain and the remaining normalized question. Theorem 67.4 is an unscaled, collision-uniform essential-height estimate on the complete physical source. It includes arbitrarily small positive incidence and clearance, all relevant words, the branch-dependent radii selected by one common physical roof, and the exact nonleading coarea contributions. It strengthens the full-source weak norm in 314 by a different geometric mechanism. The inherited results 260–322 remain intact.

It does not give an $o(m^{-2})$ height for a small selected source: a bounded density of small mass can still have a narrow positive spike. Nor does differentiation of the cap majorant differentiate the original section or a pairing selector, establish abundance of the same-roof comparison pool, or provide a uniform further conditioning denominator. At the original application width $\varepsilon = \varepsilon(B)$, the bound in (67.11) is fixed before the collision limit. No exchange of that limit with $B \to \infty$ is made. The exact shortage in (0.1), the joint twisted spectral interfaces, and the full raw local limit and four-coordinate inverse theorem retain their original unresolved status.

68. Complete canonical collision-boundary budgets

We now control the actual coordinate sides of the original section at all collision indices. The source is a genuine collision state, and all propagation uses its actual past and future flights. In particular, the quantity $\tau + R\cos\varphi$ below is an optical coefficient; it is not an extra segment of the physical roof.

Write $t_* = \tau_- = 3/50$, $k_- = 100/47$, $k_+ = 20/9$, $U = 170/9$, and $\sigma = 47/59$. Throughout this section retain the endpoint cap $|p_0|, |p_m| \leq 9/10$ whenever a Gauss-angle charge is estimated. Put $c_\circ = \sqrt{19}/10$. The source and root-supremum estimates themselves do not require this endpoint cap. Use precisely the fixed enlarged class

$$\widehat{B}_{\mathrm{cv}} \geq \max\{B_{\mathrm{cv}}, K_{\mathrm{mom}}, 2C_*\}, \qquad \widehat{k}_0, \quad \widehat{\ell}_*, \quad \widehat{C}_g$$

of (26.5). These are its own growth and subdivision constants; a constant for the smaller marked-foot class is not reused. Keep $J_f, J_{\mathrm{pre}}, J_{\mathrm{len}}, L_0, A_{\mathrm{mom}}$ from (24.1) and (26.7)–(26.9). Define

$$A_{\mathrm{fan}} = \frac{J_f J_{\mathrm{len}}}{t_*}\left(1 + \frac{2}{\widehat{k}_0 - 1}\right), \qquad E_{\mathrm{cs}} = \frac{1}{k_- t_*} = \frac{47}{6}, \tag{68.1}$$

$$H_{\mathrm{hor}} = \max\{J_{\mathrm{pre}} J_f E_{\mathrm{cs}}, A_{\mathrm{mom}} \widehat{C}_g J_f E_{\mathrm{cs}}\}, \qquad H_{\mathrm{ver}} = \max\{J_f E_{\mathrm{cs}}, A_{\mathrm{fan}} \widehat{C}_g J_f E_{\mathrm{cs}}\}.$$

The symbols "horizontal" and "vertical" refer to the unchanged (α, p) collision coordinates: $p = b$ and $\alpha = a$, respectively. The vertical source is parametrized by $\varphi = \arcsin p$, without changing its physical states.

68.1. **Actual first images and one-sided supremum sums.**

Lemma 68.1 (The two canonical sources in the qualified curve class)**.** *Fix a collision disk and translate its center to the origin. For $b \in (-1, 1)$ the horizontal source $p = b$, parametrized by α, has the complete first-image family of Lemma 26.2. For every fixed a the vertical source $\alpha = a$,*

parametrized by $\varphi \in (-\pi/2, \pi/2)$, has a complete countable family of short homogeneous first images in $\mathcal{W}^u_R(\widehat{B}_{\mathrm{cv}})$. Its exact pushed densities satisfy

$$\sum_C \|\rho_C\|_\infty \le A_{\mathrm{fan}}. \tag{68.2}$$

Both families are source-disjoint and cover their source up to its actual singular points and a countable set of subdivision points. Their total source lengths are 2π and π, respectively; neither source is normalized to a probability.

Proof. The horizontal statement, including its real first-hit test and uniformity as b tends to either grazing value, is exactly Lemmas 26.1 and 26.2. Here is also a common first-image calculation that fixes the optical conventions needed below. Put $c = \sqrt{1-b^2}$ on the horizontal source. Its outgoing direction is $\vartheta = \alpha + \arcsin b$, and its current contact is

$$q_0 = -Rb\, v_\vartheta^\perp + Rc\, v_\vartheta.$$

Consequently the next actual contact can be written $q_1 = -Rb\, v_\vartheta^\perp + a_1 v_\vartheta$, where $a_1 = \tau + Rc \ge t_*$. The line foot lies inside the current disk in general. The flight starts at the actual exit q_0; one must not apply a first-hit test starting from that foot. For a vertical source q_0 is fixed, and $q_1 = q_0 + a_1 v_\vartheta$ with $a_1 = \tau \ge t_*$. In the following identities take $b_1 = Rb$ in the horizontal case and $b_1 = 0$ in the vertical case, and let $d = \cos\varphi_1$:

$$\left|\frac{ds_1}{d\vartheta}\right| = a_1/d, \qquad V_1 = \kappa + d/a_1, \qquad \rho_1 = \frac{d}{a_1\sqrt{1+V_1^2}}, \tag{68.3}$$

$$(a_1)_{s_1} = p_1 + b_1 d/a_1, \qquad (V_1)_{s_1} = -\frac{p_1\kappa}{a_1} - \frac{2p_1 d}{a_1^2} - \frac{b_1 d^2}{a_1^3}.$$

The horizontal identities agree with (26.1)– (26.3), since $a_1 = R(c + \tau\kappa)$. For the vertical source they follow by differentiating $q_1 = q_0 + \tau v_\vartheta$ and projecting onto the target tangent and the flight direction. Thus both have $k_- \le V_1 \le U$ and

$$|(V_1)_{s_1}| \le \frac{k_+ + 2/t_*}{t_*} + \frac{R_+}{t_*^3} = K_{\mathrm{mom}}.$$

The vertical bound is smaller, since its last term is zero. This verifies membership in the already enlarged class before invoking any growth estimate.

For completeness the vertical source also has a uniform finite physical first-hit partition. Its current point is exterior to every other disk, with $|z - q_0| \ge 1 - R_+ > R_+$. The directions that enter one other disk form its open angular shadow of width less than π. Intersecting it with the outgoing semicircle gives an interval, with at most two tangency endpoints. The actual first contact has $0 < |z| \le D_f$, so at most $2N_f$ such endpoints must be cut. Distinct entry times cannot exchange order without a common point of two disjoint disks. Hence there are at most $J_f = 2N_f + 1$ first-hit intervals. Outward directions cannot reenter the current convex disk. Every flight being partitioned has the true lower bound $\tau \ge t_*$.

On each first image $V_1 > 0$, so φ_1 is strictly monotone and each homogeneous strip is met at most once. Its total length is at most L_0. Apply the fixed length subdivision with at most J_{len} pieces per strip. In the central strip $\rho_1 \le t_*^{-1}$; in each signed strip of index k it is at most $(t_* k^2)^{-1}$. Summing the actual densities, rather than counting infinitely many strips without their weights, proves (68.2). All collision angles on a connected image are represented by a continuous unwrapped lift, so crossing the numerical value 0 creates no new physical interval. Such a lift exists on its source interval; the monotone target φ_1 and the bound L_0 concern that same lift. The horizontal source itself is opened at one common fixed seam before its preliminary arcs are chosen; the retained J_{pre} includes that division. The vertical source is already an interval. Thus

J_f has only its proved physical tangency cuts. If a short-curve implementation additionally splits an image at a coordinate seam, the one possible split is included in $J_{\mathrm{len}} = 2 + \lceil 2L_0/\widehat{\ell}_* \rceil$, exactly the constant of (26.8), not silently charged to J_f. There is at most one such seam crossing because the lifted horizontal span is at most $\pi/k_- < 2\pi R_-$; after the fixed length subdivision, adding one cut fits within that displayed conservative count. In the one-flight case and at the uncut terminal image we use the continuous lift and do not insert that extra split. All removed tangencies are genuine singular endpoints, and the remaining subdivision cuts are countable. They have zero source length. No true grazing collision is deleted from an admissible physical record. □

For $l \geq 1$, propagate a source variation across exactly l actual flights, including the $l-1$ intermediate reflection kicks and neither an initial nor a terminal kick. In unfolded transverse coordinates let M_l be this positive optical matrix and put

$$(X_l, Y_l) = (\kappa_l, c_l)M_l, \qquad A_l^{\mathrm{hor}} = RcX_l + Y_l, \qquad A_l^{\mathrm{ver}} = Y_l. \tag{68.4}$$

These are the absolute derivatives of the actual terminal momentum with respect to α on a horizontal source, or φ on a vertical source. A common orientation sign is suppressed, not an absolute value inside either positive sum.

Theorem 68.2 (Complete branch-supremum bounds for canonical sources)**.** *For every R, $l \geq 1$, and either canonical source there is a countable partition into disjoint open source intervals $\mathcal{J}_l$. It retains every actual regular collision history, up to the null cuts just described. The first $l-1$ images are homogeneous and short according to the growth partition; the terminal image is not artificially divided into infinitely many strips. On these intervals,*

$$\sum_{I \in \mathcal{J}_l^{\mathrm{hor}}} \sup_I (A_l^{\mathrm{hor}})^{-1} \leq H_{\mathrm{hor}}, \qquad \sum_{I \in \mathcal{J}_l^{\mathrm{ver}}} \sup_I (A_l^{\mathrm{ver}})^{-1} \leq H_{\mathrm{ver}}, \tag{68.5}$$

$$\sup_I (A_l^{\mathrm{hor}})^{-1}, \quad \sup_I (A_l^{\mathrm{ver}})^{-1} \leq E_{\mathrm{cs}}\sigma^{l-1}.$$

These are suprema on whole intervals, not a root sum at a common target momentum. They hold without a terminal incidence cap and in the reversed physical direction with the same constants.

Proof. The initial transverse variation is $(Rc, 1)$ for a horizontal source and $(0, 1)$ for a vertical source. These conventions also follow directly from the physical differential of Lemma 54.1. If

$$B(c) = \begin{pmatrix} c & 0 \\ \kappa & 1 \end{pmatrix}, \quad P(\tau) = \begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix}, \quad K(c') = \begin{pmatrix} 1 & 0 \\ 2\kappa/c' & 1 \end{pmatrix},$$

then its positive one-flight matrix D satisfies $B(c')\mathsf{D} = K(c')P(\tau)B(c)$. At the terminal point $c_l(0,1)B(c_l)^{-1}K(c_l) = (\kappa_l, c_l)$. Thus multiplication leaves exactly the l real flights and $l-1$ intermediate kicks in M_l, with no terminal kick left over. The initial canonical vectors $(R, 0)$ and $(0, 1)$ are sent by $B(c)$ to $(Rc, 1)$ and $(0, 1)$, respectively. After the first true flight its positive displacement is $B_1 = Rc + \tau$ or $B_1 = \tau$, and its angular component is $D_1 = 1$. Each subsequent actual intermediate kick and flight acts by

$$B_{r+1} = (1 + \tau_r k_r)B_r + \tau_r D_r, \qquad D_{r+1} = k_r B_r + D_r, \qquad k_r = \frac{2}{Rc_r}.$$

Therefore $B_l \geq t_*(59/47)^{l-1}$. The terminal incoming momentum row is $\kappa_l B_l + c_l D_l \geq k_- B_l$. This proves the pointwise and hence the branch-supremum optical bounds, with their stated index $l-1$. It introduces no virtual physical flight.

For $l = 1$, use just the finite first-hit partitions from Lemma 68.1. Their sizes are at most $J_{\mathrm{pre}}J_f$ and J_f, giving the first alternatives in (68.1). For $l \geq 2$, push the exact first-image densities ρ_C through $l-2$ actual iterates. On every resulting short homogeneous graph V, the

final true collision has at most J_f physical branches and obeys the exact last-collision cancellation in the proof of Theorem 26.4:

$$\left|\frac{dp_l}{d\ell_{l-1}}\right| \geq k_- t_*.$$

For a source interval leading to one such final branch the chain rule therefore bounds the inverse source derivative by

$$\|\rho_C\|_\infty \sup_V J_V(T_R^{-(l-2)})\, E_{\mathrm{cs}}.$$

The growth theorem for the qualified enlarged class bounds the sum of the middle factor by $\widehat{C}_g$. Sum first over the at most J_f terminal physical branches and then over the full weighted first-image family. This proves the second alternatives in (68.1). No supremum is transferred from a pointwise endpoint kernel to a collection of branches.

The physical and homogeneous partitions and their monotone pullbacks are countable and source-disjoint. The terminal momentum is strictly monotone on each one. A terminal interval restriction therefore leaves at most one interval and does not increase any of the sums. A common fixed source seam is used before superposing two families. Time reversal $\mathcal{I}(\alpha,\varphi) = (\alpha,-\varphi)$ is an isometry on both source parameters; it preserves every bound and transports the actual source tests. The physical map is invertible on each regular history. Translating the selected collision disk to the origin consequently creates no multiplicity of initial points, lattice labels, or collision words. □

68.2. **Exact two-sided determinants, including the incidence factor.**

Lemma 68.3 (Canonical two-sided root weights)**.** *At an actual interior collision $i \in \{1,\dots,m-1\}$ write $c_i = \sqrt{1-p_i^2}$ and let (X_-,Y_-) and (X_+,Y_+) be (68.4) for the i backward and $m-i$ forward flights. Put $A_\pm^{\mathrm{hor}} = Rc_iX_\pm + Y_\pm$, $A_\pm^{\mathrm{ver}} = Y_\pm$, and*

$$d_\pm = \frac{Y_\pm}{Rc_iX_\pm + Y_\pm}, \qquad h_\pm = 1 + \frac{Rc_iX_\pm}{Y_\pm}, \qquad C_0 = \frac{t_* + R_+}{2t_*} = \frac{53}{12}.$$

The unchanged endpoint root weight $u_x = |\partial_{\alpha_0} p_m|^{-1}$ satisfies

$$u_x = \frac{c_i}{A_-^{\mathrm{hor}}A_+^{\mathrm{hor}}(d_- + d_+)} \leq \frac{C_0c_i}{A_-^{\mathrm{hor}}A_+^{\mathrm{hor}}}, \tag{68.6}$$

$$u_x = \frac{c_i}{A_-^{\mathrm{ver}}A_+^{\mathrm{ver}}(h_- + h_+)} \leq \frac{c_i}{2A_-^{\mathrm{ver}}A_+^{\mathrm{ver}}} \leq \frac{1}{2A_-^{\mathrm{ver}}A_+^{\mathrm{ver}}}. \tag{68.7}$$

In particular the horizontal bound retains the factor $c_i = \sqrt{1-b^2}$ uniformly as its level approaches grazing.

Proof. Up to a common sign in each endpoint row, the derivative columns in the central variables (α_i,φ_i) are

$$\begin{pmatrix} A_-^{\mathrm{hor}} & -Y_- \\ A_+^{\mathrm{hor}} & Y_+ \end{pmatrix}.$$

Indeed varying α_i with φ_i fixed gives the initial transverse field $(Rc_i,1)$ in both physical directions. Varying φ_i at fixed contact gives $(0,1)$ forward and $(0,-1)$ backward. Time reversal may reverse the entire backward momentum row but cannot change this relative sign between columns. The determinant thus has absolute value $A_-^{\mathrm{hor}}Y_+ + Y_-A_+^{\mathrm{hor}}$.

Collision-area invariance and the exact endpoint coarea identity give

$$d\nu = \frac{d\alpha_i\, dp_i}{4\pi} = \frac{c_i\, d\alpha_i\, d\varphi_i}{4\pi} = \frac{u_x}{4\pi}\, dp_0\, dp_m.$$

This proves both displayed identities; unlike a marked line foot, a canonical collision source has no additional factor R^{-1}. For the inequalities, write $M_l = \widetilde{M}P(\tau_1)$, with $P(\tau) = \left(\begin{smallmatrix} 1 & \tau \\ 0 & 1 \end{smallmatrix}\right)$. If $(x, y) = (\kappa_l, c_l)\widetilde{M}$, positivity implies $X_l = x$, $Y_l = \tau_1 x + y$, and hence

$$0 < X_l/Y_l \le 1/t_*.$$

Therefore $d_\pm \ge t_*/(t_* + R_+)$ and $h_\pm \ge 1$. The coefficient c_i arises from the original collision area and has not been replaced by one in (68.6). □

Corollary 68.4 (All-word two-sided supremum budgets)**.** *Intersect the two capped or uncapped one-sided partitions at the same canonical source. Let $\mathcal{O}_i^{\mathrm{hor}}$ or $\mathcal{O}_i^{\mathrm{ver}}$ denote the nonempty interval overlaps. Then*

$$\sum_{O \in \mathcal{O}_i^{\mathrm{hor}}} \sup_O u_x \le C_0 c_i H_{\mathrm{hor}} E_{\mathrm{cs}}(\sigma^{i-1} + \sigma^{m-i-1}), \tag{68.8}$$

$$\sum_{O \in \mathcal{O}_i^{\mathrm{ver}}} \sup_O u_x \le \tfrac{1}{2} H_{\mathrm{ver}} E_{\mathrm{cs}}(\sigma^{i-1} + \sigma^{m-i-1}). \tag{68.9}$$

The first assertion is uniform over all $b \in (-1, 1)$, and the second over all fixed contact angles. No exact-record selector is used to form new interval components.

Proof. Apply the weighted countable interval-overlay lemma 62.2 to the two actual source-disjoint families. For clarity its bound is $\sum_{I \cap J \ne \varnothing} a_I b_J \le (\sup_J b_J) \sum_I a_I + (\sup_I a_I) \sum_J b_J$. It follows by assigning an overlap to the interval whose left endpoint is later: the other disjoint family has at most one interval containing that endpoint. Equal endpoints are assigned once by a fixed rule; open intervals and finite exhaustion handle all countable cases. Use (68.5) for the full sums and the optical bounds with half lengths i and $m - i$ for their maxima, then apply Lemma 68.3. On a horizontal level c_i is a constant. On a vertical side it is at most one. The partitions cover all actual regular histories; their artificial endpoints have zero angular-image length and are not new physical records. □

68.3. **Uniform total turning on the actual source intervals.** Here are explicit variation constants for the same enlarged class. In the constants of (63.1)– (63.3), replace $B_{\mathrm{cv}}, k_0, \ell_*, D_B$ by $\widehat{B}_{\mathrm{cv}}, \widehat{k}_0, \widehat{\ell}_*, \widehat{D}_B$, where $\widehat{D}_B$ is (24.7) for that allowance. Equivalently, put

$$\begin{aligned}
\widehat{B}_2 &= \widehat{B}_{\mathrm{cv}}(1 + U^2)^{3/2}, \\
\Gamma &= \sqrt{1 + k_+^2}, \qquad L_h = 3\sqrt{1 + U^2}/k_-, \\
\widehat{c}_H &= \sin(\widehat{k}_0^{-2}), \\
\widehat{V}_H &= \max\{-2 \log \widehat{c}_H, \log(\pi/2) + 2 \log(1 + \widehat{k}_0^{-1})\}, \\
\widehat{D}_1 &= \max\{(U/\widehat{c}_H + \widehat{D}_B) L_0^{2/3}, (2\pi U + \widehat{D}_B) L_h^{2/3}\}, \\
\widehat{V}_J &= \frac{\widehat{D}_1 \Gamma^{1/3} L_0^{1/3}}{1 - \sigma^{1/3}}, \\
V_{\mathrm{cap}} &= -2 \log c_\circ, \\
V_{\rho,\mathrm{cs}} &= \widehat{V}_H + (t_*^{-1} + R_+ t_*^{-2} + K_{\mathrm{mom}}/2)\widehat{\ell}_*, \\
V_{t,\mathrm{cs}} &= V_{\mathrm{cap}} + \widehat{B}_2 L_0/k_-, \\
V_{1,\mathrm{cs}} &= \frac{L_0(k_+ R_+ + 1)}{k_- t_*^2}, \\
V_{A,\mathrm{cs}} &= \max\{V_{1,\mathrm{cs}}, V_{\rho,\mathrm{cs}} + \widehat{V}_J + V_{t,\mathrm{cs}}\}.
\end{aligned} \tag{68.10}$$

These finite constants precede every radius, collision count, source level, exact label, and roof supremum.

Lemma 68.5 (Logarithmic variation and regular angular turning)**.** *On every one-sided canonical source interval, after restricting its terminal momentum to* $[-9/10, 9/10]$*,*

$$\operatorname{Var}\log A_l^{\mathrm{hor}} \le V_{A,\mathrm{cs}}, \qquad \operatorname{Var}\log A_l^{\mathrm{ver}} \le V_{A,\mathrm{cs}}.$$

On every interior two-sided overlap, with $P = (-p_0, p_m)$*, one has*

$$\int_{\{P\neq 0\}} |d\arg P| \le 2\pi + V_{A,\mathrm{cs}}. \tag{68.11}$$

On a fixed initial-angle side, including both endpoint caps, the corresponding bound is

$$\int_{\{P\neq 0\}} |d\arg P| \le V_{\mathrm{end}} := 2\pi + \tfrac{1}{2}(V_{A,\mathrm{cs}} + V_{\mathrm{cap}}). \tag{68.12}$$

No critical incoming fan at $P = 0$ *is assigned zero mass by these regular, punctured-angle statements.*

Proof. All the following estimates concern absolute integrals of derivatives, not merely differences of their endpoint values. The proof of Lemma 63.1 applies to the enlarged class: on a homogeneous image of length r the absolute one-step log-Jacobian variation is at most $\widehat{D}_1 r^{1/3}$, and the adapted expansion gives $L_j \le \Gamma\sigma^{l-j}L_l$. Summation therefore bounds the full log-Jacobian variation by $\widehat{V}_J$.

For $l \ge 2$ the actual first-image density in (68.3) gives exactly

$$A_l^{\mathrm{hor}} \text{ or } A_l^{\mathrm{ver}} = \frac{c_l V_l}{\sqrt{1+V_l^2}} \frac{J_C(T_R^{l-1})}{\rho_C}.$$

The terminal cap is one interval because p_l is strictly monotone, and it lies in the central homogeneous strip. On the first homogeneous piece the variation of $\log c_1$ is at most $\widehat{V}_H$. The exact identities (68.3) give $|(a_1)_{s_1}| \le 1 + R_+/t_*$ and $|(V_1)_{s_1}| \le K_{\mathrm{mom}}$. Their integrals, with $a_1 \ge t_*$, bound $\operatorname{Var}\log\rho_C$ by $V_{\rho,\mathrm{cs}}$. The terminal factor has variation at most $V_{t,\mathrm{cs}}$, since φ_l is monotone, $c_l \ge c_\circ$, and $|(V_l)_s| \le \widehat{B}_2$. This proves the claimed bound for $l \ge 2$, preserving the exact initial density and all its grazing factors throughout the summation.

For $l = 1$ no initial homogeneous cut is needed. In either source $A_1 = \kappa a_1 + d$, and

$$\partial_{s_1} A_1 = d(\kappa b_1 - p_1)/a_1.$$

Its log-derivative is bounded by $(k_+R_+ + 1)/(k_-t_*^2)$. Integrate over a first-image graph of length at most L_0. This proves the remaining alternative. Time reversal preserves these absolute variation estimates.

On a two-sided interior overlap the two components of P' have fixed signs and nonzero magnitudes $A_-^{\mathrm{hor}}, A_+^{\mathrm{hor}}$, or $A_-^{\mathrm{ver}}, A_+^{\mathrm{ver}}$. Thus its tangent stays in one quadrant and has total turning at most

$$\tfrac{1}{2}\{\operatorname{Var}\log A_- + \operatorname{Var}\log A_+\} \le V_{A,\mathrm{cs}}.$$

One component of P is strictly monotone, so P crosses the origin at most once. Lemma 63.3 proves (68.11), including both punctured sides of a crossing. At a fixed initial angle use the source variable φ_0. Then $P' = (-c_0, \pm Y_m)$, its tangent remains in one quadrant, and $\operatorname{Var}\log c_0 \le V_{\mathrm{cap}}$ on the initial cap. The same lemma proves (68.12). A terminal-angle side is covered by time reversal, which preserves absolute turning. □

68.4. **Complete section-side charges, including the endpoint indices.** For one actual regular canonical side let its charge mean the sum, over its unique physical source points and all words, of $A|d\theta|$, where

$$A = \frac{\omega}{4\pi c_* \chi}, \qquad \theta = \arg(-p_0, p_m), \qquad P = (-p_0, p_m) \neq 0.$$

An additional existing selector of absolute value at most one can restrict this positive integral. This definition counts the physical side once. When a piecewise record weight is differentiated, its jump multiplies this one-copy charge; the two incident traces are combined into that jump before absolute values. If two incident traces are intentionally counted separately, twice this bound is available. It is not legitimate to count them twice and then omit the factor two.

Set

$$\begin{aligned} F_{\mathrm{cs}} &= \frac{\Lambda_\circ}{4\pi c_* R_-^2}, & C_{\mathrm{hor}} &= F_{\mathrm{cs}}(2\pi + V_{A,\mathrm{cs}}) C_0 H_{\mathrm{hor}} E_{\mathrm{cs}}, \\ C_{\mathrm{ver}} &= \tfrac{1}{2} F_{\mathrm{cs}}(2\pi + V_{A,\mathrm{cs}}) H_{\mathrm{ver}} E_{\mathrm{cs}}, & E_{\mathrm{hor}} &= \pi F_{\mathrm{cs}} C_{\mathrm{mom}}, \\ E_{\mathrm{ver}} &= F_{\mathrm{cs}} V_{\mathrm{end}} H_{\mathrm{ver}}. \end{aligned} \tag{68.13}$$

Theorem 68.6 (Uniform canonical side charges on the physical source)**.** *For every $R \in [R_-, R_+]$, $m \geq 2$, and $1 \leq i \leq m-1$, the complete regular charge at $p_i = b \in (-1,1)$ or at $\alpha_i = a \pmod{2\pi}$ satisfies respectively*

$$\mathfrak{S}_{i,b}^{\mathrm{hor}} \leq C_{\mathrm{hor}} \sqrt{1-b^2} (\sigma^{i-1} + \sigma^{m-i-1}), \tag{68.14}$$

$$\mathfrak{S}_{i,a}^{\mathrm{ver}} \leq C_{\mathrm{ver}} (\sigma^{i-1} + \sigma^{m-i-1}). \tag{68.15}$$

The vertical source may be restricted to the actual momentum interval of an original rectangle, or to any positive Borel subsource, without an added component count. At either endpoint index 0 or m a horizontal level in the endpoint cap has charge at most E_{hor}, and a vertical level has charge at most E_{ver}. These endpoint assertions hold also for $m = 1$.

In particular each fixed type and level of interior side has total charge at most $2C/(1-\sigma)$ after all its collision indices are summed. The indices satisfying $\min(i-1, m-i-1) \geq D$ have total charge at most $2C\sigma^D/(1-\sigma)$. These are sums over complete physical histories before any count, radius, label or source-level supremum.

Proof. The exact optical identity of Corollary 59.2 gives

$$A = \frac{u_x}{4\pi c_* h_x}, \qquad h_x = R^2 e^{\mathsf{T}} G^{-1} e \geq R_-^2/\Lambda_\circ.$$

It follows that $A \leq F_{\mathrm{cs}} u_x$. On each interior overlap use Lemma 68.5 and then Corollary 68.4. Nonnegative countable summation gives (68.14)–(68.15). A Borel restriction is made after this positive summation; it is not assumed to have a finite number of components or a differentiable boundary. The positive source density, the conditional factor c_*^{-1} and the original collision-area normalization have all been retained.

For an initial horizontal endpoint side $p_0 = b \neq 0$, parametrize by the actual terminal momentum $v = p_m$. Its absolute angular differential is $|b|(b^2+v^2)^{-1} dv$. The pointwise physical root sum of Theorem 26.4, evaluated at the actual pair (b, v), gives the complete charge bound

$$F_{\mathrm{cs}} C_{\mathrm{mom}} \int_{-9/10}^{9/10} \frac{|b|}{b^2+v^2}\, dv \leq \pi F_{\mathrm{cs}} C_{\mathrm{mom}} = E_{\mathrm{hor}}.$$

For $b = 0$ its regular polar angle is constant on either side of $v = 0$, so the regular angular charge is zero. Its critical incoming source is a separate endpoint source and is not discarded. Time reversal proves the terminal horizontal assertion.

For a fixed initial angle, on each forward fan interval

$$u_x = \frac{1}{Rc_0X_m + Y_m} \le \frac{1}{Y_m}.$$

The branch-supremum sum in Theorem 68.2 is therefore at most H_{ver}. Multiply by F_{cs} and the complete turning bound (68.12). This proves the stated vertical endpoint bound, and time reversal proves the final one. The original cap and any rectangle-side momentum interval merely restrict these positive charges. A coordinate seam, subdivision endpoint or section corner has zero angular-image length and creates no additional charge. All visited grazing limits and critical incoming fans retain their separately proved physical traces.

Finally sum the two geometric sequences in the interior bounds. The deep-index restriction starts each surviving exponent at least D, giving the asserted exponential tail. For every fixed R, m all local source limits and countable exhaustions precede these uniform estimates. No long-time or outer-cutoff limit was interchanged. □

A useful grazing-level consequence. The factor in (68.6) and (68.14) is essential rather than cosmetic. For the original signed homogeneous levels $\varphi = \pm(\pi/2 - k^{-2})$ it is $\cos\varphi = \sin(k^{-2}) \le k^{-2}$. Thus their complete horizontal charges are summable over k, even though there are infinitely many such levels. This consequence does not assert that every whole-word cell has already been assigned to such a boundary; any use requiring that assignment must prove it separately.

69. Exact marked records and a uniform physical-roof Fourier bound

The original section Y_R^* is the disjoint union of its four specified squares and the specified rectangle U. In particular it has $N_{\mathrm{hor}} = 10$ horizontal sides and $N_{\mathrm{ver}} = 10$ vertical sides, counted with their actual coordinate levels and finite side intervals. A square crossing the circular coordinate seam is glued there; the seam is not another side. We use precisely these rectangles, the original occupation convention and the original conditional measure. No differentiability is assigned to an arbitrary measurable selector.

The constants of Theorem 68.6 give

$$\begin{aligned} B_{\mathrm{sec,int}} &= \frac{2(N_{\mathrm{hor}}C_{\mathrm{hor}} + N_{\mathrm{ver}}C_{\mathrm{ver}})}{1-\sigma}, \\ B_{\mathrm{sec,end}} &= 2(N_{\mathrm{hor}}E_{\mathrm{hor}} + N_{\mathrm{ver}}E_{\mathrm{ver}}). \end{aligned} \tag{69.1}$$

All collision indices $1, \dots, m-1$ are included in the first constant; both actual endpoint indices $0, m$ are included in the second. When $m = 1$ the interior sum is empty and its displayed bound remains valid. Retaining the individual factors $\sqrt{1-b^2}$ on horizontal sides can only improve this coarse common constant.

Write $\eta_R = \mathbf{1}_{Y_R^*}$. At a fixed actual collision count $m \ge 1$, let

$$K = K^{\mathrm{c}}_{m,R} \in \mathbb{Z}^2, \qquad A = A_{m,R} = \sum_{j=0}^{m-1} \eta_R \circ T_R^j, \qquad L = L_{m,R} = \sum_{j=0}^{m-1} \tau_R \circ T_R^j.$$

The initial visit is counted in A and the terminal visit is tested separately. The roof L is the original unit-speed physical time. For n, k use the unchanged event $\Pi_{n,k,m,R}$ of (1.1). For real frequencies $\xi \in \mathbb{R}^2$, $\zeta, \upsilon \in \mathbb{R}$, define

$$\begin{aligned} Q_{n,k,m}(x) &= \eta_R(x)\eta_R(T_R^m x)\mathbf{1}_{\{K=k, A=n\}}, \\ Q_m^{\xi,\zeta,\upsilon}(x) &= \eta_R(x)\eta_R(T_R^m x)\exp\{i(\xi\cdot K + \zeta A + \upsilon m)\}. \end{aligned} \tag{69.2}$$

The harmless collision-count phase $e^{i\upsilon m}$ makes the fixed collision mark explicit. The other three discrete frequency coordinates retain the two lattice labels and the exact occupation count.

The fourth nontrivial collision-observable frequency below is the real roof frequency s. Thus (ξ_1, ξ_2, ζ, s) are exactly the original four transform coordinates at fixed m; vm records only that fixed collision tag and introduces no fifth target coordinate or new inversion problem.

Let w be either 1 or the original W^{ε}, with $\varepsilon > 0$ fixed before any collision limit. Define the positive or complex physical-roof measures

$$\mu^{w}_{n,k,m,R} = L_*\big(wQ_{n,k,m}\, d\nu/c_*\big) = L_*\big(w\mathbf{1}_{\Pi_{n,k,m,R}}\, d\nu^*_R\big), \tag{69.3}$$
$$\mu^{w;\xi,\zeta,\upsilon}_{m,R} = L_*\big(wQ^{\xi,\zeta,\upsilon}_{m}\, d\nu/c_*\big).$$

Set $D^1_{\mathrm{base}} = D_{\mathrm{roof}}$ and $D^{W^{\varepsilon}}_{\mathrm{base}} = D^{\varepsilon}_{\mathrm{roof}}$ from (67.3) and (67.11), and put

$$\begin{aligned} D^{w}_{\mathrm{exact}} &= D^{w}_{\mathrm{base}} + B_{\mathrm{sec,end}} + B_{\mathrm{sec,int}}, \\ D^{w}_{\mathrm{twist}}(\zeta) &= D^{w}_{\mathrm{base}} + B_{\mathrm{sec,end}} + 2|\sin(\zeta/2)|B_{\mathrm{sec,int}}, \\ D^{w}_{\mathrm{twist}} &= D^{w}_{\mathrm{base}} + B_{\mathrm{sec,end}} + 2B_{\mathrm{sec,int}}. \end{aligned} \tag{69.4}$$

Here and below a superscript w names one of these two physical weights, not an additional source assumption.

Lemma 69.1 (Exact jump accounting for the original finite records)**.** *On each regular physical word the weights* (69.2) *are locally constant away from the preimages of the twenty original section sides. Differentiation in the roof coordinate at fixed Gauss angle adds total section-boundary charge at most $B_{\mathrm{sec,end}} + B_{\mathrm{sec,int}}$ for an exact record, and at most $B_{\mathrm{sec,end}} + 2|\sin(\zeta/2)|B_{\mathrm{sec,int}}$ for the unit-modulus marked twist. These bounds include all actual words and all possible occupation labels. They do not depend on the number of components selected by one exact label.*

Proof. A fixed lifted physical word fixes K, and all collision states vary analytically on its regular endpoint chart. Every membership test in Y^*_R is a finite Boolean combination of its original coordinate inequalities. Consequently an exact-record indicator or its marked phase is constant in each complementary chamber. On a compact regular fixed-angle interval, a nonconstant analytic side function has only finitely many zeros. A slice on which one such function is identically zero is exceptional for angular integration unless the function vanishes on a neighborhood; the latter is impossible for a genuine canonical coordinate side since T^i_R is a local diffeomorphism. To verify the exceptional-set assertion, use a countable local analytic chart cover: an accumulating sequence of constant-zero angular slices forces every coefficient of the local power series to vanish. Thus the exceptional angular levels form a null set. Exhaustion treats endpoints and any accumulation at a physical singular boundary.

On a regular interior side where only the visit at index i changes, A changes by 1 or -1. An exact indicator jumps by at most one; the marked phase jumps in modulus by $|e^{i\zeta} - 1| = 2|\sin(\zeta/2)|$. If several visit boundaries coincide, order the affected visits and telescope their changes. The same upper bound is paid once for each changed visit, all of which are present in (69.1). For an exact indicator the bound one per such surface is still valid. The endpoint factors $\eta_R(x)$ and $\eta_R(T^m_R x)$ have jump at most one in modulus, even for the twisted weight. The initial occupation contribution is part of its endpoint value, and the terminal test creates no fictitious occupation at index m.

The exact canonical flux identity in Theorem 58.1 expresses the contribution of a section side as its one-copy charge $\mathscr{A}|d\theta|$ times the absolute jump, where $\mathscr{A} = \omega/(4\pi c_* \chi)$ is the coarea coefficient (and not the occupation count A). The incident regular pieces are combined into this jump before absolute values. Hence an artificial subdivision does not pay twice for the same unchanged weight. Apply Theorem 68.6 to every actual horizontal or vertical side at each indicated collision index. Restricting to its finite rectangle-side interval is a positive restriction of that complete source. Summing the geometric depth bounds over $1 \le i \le m-1$ and then the

two endpoint bounds proves exactly (69.1) and the asserted jump charges. Section corners or transverse intersections have zero angular-image length. Coincident regular boundary arcs are treated by the jump or the telescoping assignment just given, not duplicated as new trajectories. This proof uses the geometry of the original finite rectangles, not a variation claim for an arbitrary Borel set. □

Theorem 69.2 (Uniform bounded variation for exact physical records)**.** *For every $R \in [R_-, R_+]$, every $m \geq 1$, all original labels n, k, and either specified weight w, the first measure in* (69.3) *has a nonnegative density $p^w_{n,k,m,R} \in BV(\mathbb{R})$ satisfying*

$$\|Dp^w_{n,k,m,R}\|_{\mathrm{TV}} \leq D^w_{\mathrm{exact}}, \qquad \|p^w_{n,k,m,R}\|_1 \leq 1, \qquad \|p^w_{n,k,m,R}\|_\infty \leq C_{\mathrm{roof}}. \tag{69.5}$$

For $w = 1$ this is the original density $p_{n,R}(k, m, t)$. The twisted measure has a complex density $p^{w;\xi,\zeta,\upsilon}_{m,R} \in BV(\mathbb{R}; \mathbb{C})$ with

$$\|Dp^{w;\xi,\zeta,\upsilon}_{m,R}\|_{\mathrm{TV}} \leq D^w_{\mathrm{twist}}(\zeta) \leq D^w_{\mathrm{twist}}, \quad \|p^{w;\xi,\zeta,\upsilon}_{m,R}\|_1 \leq 1, \quad |p^{w;\xi,\zeta,\upsilon}_{m,R}| \leq \overline{p}_{m,R} \quad a.e. \tag{69.6}$$

All densities have support in $[m\tau_-, m\tau_+]$. The variation bounds precede the supremum over R, m, n, k and all real discrete-mark frequencies. As throughout the original problem, exceptional density sets may depend on the fixed radius.

Proof. Every original section state has $|p| < 5/6$, as in the endpoint cap of Section 67. Therefore both weights in (69.2) vanish outside the source of its full positive majorant. Their absolute values are at most one. The exact coarea formula consequently gives integrable densities dominated in modulus by $\overline{p}_{m,R}$, and hence their stated height bounds. Their L^1 norms are at most one, rather than only c_*^{-1}, because the initial factor has been restored and $c_*^{-1} \int \eta_R \, d\nu = 1$. Their roof and support are the actual ones.

We verify the derivative, since domination alone would not prove it. Use the maximal regular fixed-angle physical intervals of Lemma 67.3. On each chamber cut by the original section sides, multiply the exact coarea coefficient by the constant record weight. Its interior derivative has absolute value bounded by the unweighted derivative of (67.4), or by the unchanged guarded derivative in Corollary 67.5. Hence the sum of interior derivatives is bounded by precisely the interior part of D^w_{base}. All artificial seams and length subdivisions cancel before taking absolute values.

The physical interval endpoints have the complete accounting already proved in Lemma 67.3. At a cap or an unvisited-disk boundary, multiplying a one-sided trace by a number of modulus at most one cannot increase its positive charge. All actual words are retained, so no cancellation across different words is needed for this upper bound. At a visited grazing limit the inherited optical trace is zero, still after the bounded multiplier. At a critical endpoint the finite original section inequalities have one-sided angular sector limits: in a fixed local analytic chart, the first nonzero homogeneous term of each active inequality has only finitely many zero rays. Off those rays its sign is constant along a sufficiently short ray. The finite Boolean record and its phase therefore have a limit of modulus at most one on almost every incoming ray. Their incoming charge is bounded by the complete positive critical sector charge of Lemma 67.3. No critical sector has been replaced by zero merely because its source has area zero.

All these physical endpoint contributions, including the full critical fan, are therefore bounded by the boundary part of D^w_{base}. The remaining section jumps have exactly the bounds of Lemma 69.1. Integrate by parts against a compactly supported C^1 test function on a finite exhaustion of the regular chambers. The positive absolute sums of interior, physical endpoint and section-jump measures are finite after angular integration. The source integral against the derivative of the test function is integrable as well. Exhaustion thus produces a finite signed,

or complex, distributional derivative, with total variation at most the constants in (69.4). This proves (69.5)–(69.6).

This construction treats the entire finite set of original section tests before the collision-uniform bound. It does not count the components of a selected event, assume independent visits, discard a collision, or fill a physically missing roof interval. The local limits are taken at each fixed R, m before applying the displayed uniform estimates. □

Corollary 69.3 (Uniform roof Fourier tail with exact marks)**.** *For either density in Theorem 69.2, let D denote its corresponding variation constant. For every $s \neq 0$ and $h \in \mathbb{R}$,*

$$\left| \int_{\mathbb{R}} e^{ist} p(t)\, dt \right| \leq \min\{1, D/|s|\}, \qquad \|p(\cdot + h) - p\|_1 \leq D|h|. \tag{69.7}$$

In particular, uniformly in $m, R, \xi, \zeta, \upsilon$,

$$\left| \int w\, \eta_R(T_R^m x) e^{i\{\xi \cdot K^{\mathrm{c}}_{m,R}(x) + \zeta A_{m,R}(x) + \upsilon m + sL_{m,R}(x)\}}\, d\nu_R^*(x) \right| \leq \min\{1, D^w_{\mathrm{twist}}/|s|\}. \tag{69.8}$$

No lattice or occupation frequency, including an arithmetic zero class or a nonzero spectral peak, has been omitted. If desired, the same uniform estimate gives the quantitative squared tail

$$\int_{|s|>S} |\widehat{p}(s)|^2\, ds \leq 2D^2/S \qquad (S > 0).$$

Proof. The densities have compact physical-roof support. Integration by parts in the distributional identity for their finite derivative measure gives $|\widehat{p}(s)| \leq \|Dp\|_{\mathrm{TV}}/|s|$. Their L^1 bounds give the other term. Integration of the absolute variation measure between t and $t+h$, followed by Tonelli, gives the translation estimate also for the complex density. The characteristic expression is exactly the Fourier transform of the specified twisted roof measure, with the original conditional source and terminal event. Integrating $D^2 s^{-2}$ on the two tails proves the last assertion. □

Scope relative to the raw local limit. The new estimate differentiates every original finite section and occupation test and retains the complete physical displacement and collision marks. It supplies a uniform high-roof-frequency bound for those exact measures and their unit-modulus joint twists. An arbitrary Borel selector H still has only the height domination of Theorem 67.4; no Fourier or variation estimate for such an H follows from this result. Nor is the $|s|^{-1}$ majorant integrable over the whole high-frequency line, or small enough by itself to prove the original collision-normalized m^2 pointwise local limit. The remaining joint twisted spectral estimates, normalization payments and original-order limiting obligations must be supplied at their stated strengths. No return event, microscopic window, probability quantifier, physical clock or arithmetic factor has been changed to obtain the present bound.

70. A physical boundary stopping proof of the full word-pressure bound

The capacity moment in Section 65 does not, by itself, pay the small-capacity part of the original physical-word pressure. We now bound that physical-word sum by a different geometric mechanism, choosing one nearly maximizing past-strip class per physical word. We do not assert a uniform sum over every homogeneous class. We stop an actual affine transversal inside the same complete physical word and the same past homogeneity labels, and transport its word weight to a genuine one-dimensional source at a small positive interior level. The possible exits are exhaustively classified below. No lower bound on a cell's area, length, capacity, or angle is required.

All collisions, their lifted centers, the radius $R \in [R_-, R_+] = [9/20, 47/100]$, and the physical unit-speed flow are unchanged. The stable cocycle is Euclidean arclength in (s, φ), where $s = R\alpha$.

Put

$$\kappa_R = R^{-1}, \qquad \Gamma = \sqrt{1 + (20/9)^2}, \qquad \sigma = 47/59.$$

Use the original $q = 2$ homogeneity strips and their fixed k_0. Their interior boundary levels form the set

$$\mathcal{H} = \{\, \pm(\pi/2 - k^{-2}) : k \geq k_0 \,\}, \qquad S_{\mathcal{H}} := \sum_{h \in \mathcal{H}} \cos h = 2 \sum_{k \geq k_0} \sin(k^{-2}) \leq \frac{2}{k_0 - 1}.$$

This is a sum of actual endpoint incidence factors, not a sum of strip widths.

We use the horizontal collision-source constants of Theorem 68.2, writing $H_{\mathrm{H}} = H_{\mathrm{hor}}$ and $E_{\mathrm{H}} = E_{\mathrm{cs}} = 47/6$: for every prescribed $\varphi_i = h \in (-\pi/2, \pi/2)$ and either time direction, the source parameter is α_i on its one physical circle. For $l \geq 1$ the genuine branch partition has inverse terminal momentum-derivative weights a_I with

$$\sum_I a_I \leq H_{\mathrm{H}}, \qquad a_I \leq E_{\mathrm{H}} \sigma^{l-1}. \tag{70.1}$$

The initial source need not be homogeneous. There is no terminal homogeneity subdivision. These are the unconditional horizontal-source conclusions of that theorem, before any endpoint cap or angular-turning estimate. Its auxiliary growth cuts use the qualified enlarged class and its constants from module 284. The stopped word classes below still use the original k_0 and its original strips; these two partitions need not coincide. Both collections of source cuts will be avoided inside open physical source intervals. Lemma 68.3 gives the exact two-half determinant inequality

$$u_x := |\partial_{\alpha_0} p_N|^{-1} \leq C_{\mathrm{H}} \cos\varphi_i \, a_I^- \, a_J^+, \qquad C_{\mathrm{H}} = \frac{\tau_- + R_+}{2\tau_-} = \frac{53}{12}, \quad 1 \leq i < N, \tag{70.2}$$

at a point of the actual overlap $I \cap J$. Here the two half-lengths are i and $N - i$. At $i = 0$ and $i = N$, respectively, the exact identities are $u_x = |\partial_{\alpha_0} p_N|^{-1}$ on the forward horizontal source and $u_x = |\partial_{\alpha_N} p_0|^{-1}$ on the backward horizontal source. The latter follows from preservation of $d\alpha\, dp$, and is not a time-reversal substitution of a roof.

Corollary 68.4, using the interval overlay lemma of Section 62, therefore gives at every common horizontal source level

$$\sum_O \sup_O u_x \leq \cos h \, C_{\mathrm{H}} H_{\mathrm{H}} E_{\mathrm{H}} (\sigma^{i-1} + \sigma^{N-i-1}), \qquad 1 \leq i < N. \tag{70.3}$$

For either endpoint the corresponding one-sided sum is at most H_{H}. All source intervals fix actual lifted half-words. Their overlaps consequently fix the complete physical word after the initial disk is translated to the original fundamental copy. They include arbitrarily short source intervals.

The marked-flight bound from Theorem 62.3 is used only at positive clearance levels:

$$\sum_{O \in \mathcal{O}_{N,j,R,u}} \sup_O u_x \leq Q_{\mathrm{mb}} (\sigma^j + \sigma^{N-j-1}), \quad 0 \leq j < N, \quad 0 < u \leq 3/200. \tag{70.4}$$

Both mark signs and every actual source copy are already included in this formula.

70.1. **An endpoint-exact comparison with the affine inverse weight.** Let $D^-_{N,R}$ be the open set of states with regular physical past through times $0, -1, \ldots, -N$. For $y \in D^-_{N,R}$ put

$$e_R = \frac{(1, -\kappa_R)}{\sqrt{1 + \kappa_R^2}}, \qquad w_N(y) = \frac{1}{|DT_R^{-N}(y) e_R|}. \tag{70.5}$$

The vector is the unit tangent of the intrinsic affine fiber $\Gamma_{R,\pi_R(y)}$ from Section 65. The function w_N is positive and continuous on every finite regular word domain. It does not require an infinite future or a choice of an invariant stable line.

Lemma 70.1 (The affine weight and the original endpoint momentum)**.** *For every regular physical N-flight record $x, T_R x, \dots, T_R^N x$, $N \geq 1$, one has*

$$w_N(T_R^N x) \leq R_+ \Gamma\, c_0(x)\, |\partial_{\alpha_0} p_N(x)|^{-1}. \tag{70.6}$$

If $x \in M_R'$, then

$$j_N(x) \leq C_\times w_N(T_R^N x). \tag{70.7}$$

Neither estimate has a terminal incidence cutoff.

Proof. Write the positive matrix product of Lemma 54.1 as

$$\mathsf{D}_N = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \qquad DT_R^N = (-1)^N \mathsf{D}_N, \qquad \det \mathsf{D}_N = c_0/c_N.$$

Every entry is positive. Since a variation of α_0 at fixed p_0 has $\delta s_0 = R\,\delta\alpha_0$ and $\delta\varphi_0 = 0$, while $\delta p_N = c_N \delta\varphi_N$, the original derivative is exactly

$$u_x = \frac{1}{R c_N C}. \tag{70.8}$$

There is one radius factor and one terminal incidence in this identity.

The inverse matrix sends $(1, -\kappa_R)$ to

$$\frac{c_N}{c_0}(D + \kappa_R B,\ -C - \kappa_R A).$$

Consequently

$$w_N(T_R^N x) = \frac{c_0 \sqrt{1 + \kappa_R^2}}{c_N \sqrt{(D + \kappa_R B)^2 + (C + \kappa_R A)^2}}. \tag{70.9}$$

The square root in the denominator is at least C. Dividing (70.9) by $c_0 u_x$ proves (70.6). In particular no factor $1/c_N$ has been dropped: the identical factor in (70.8) cancels before taking the uniform bound.

For (70.7), true invariance gives $j_N(x) = 1/J_{E^s(T_R^N x)} T_R^{-N}$. At the same current point, both this stable direction and e_R have slopes in $[-K, -a]$. Apply Corollary 61.5 with these two vectors at the same point. The physical word and all its past strip labels agree automatically. Inverting its two positive Jacobians gives (70.7). No continuity assertion for an invariant line near a singular point is involved. □

70.2. **The exhaustive physical exit of a fixed affine interval.** In this subsection a word is normalized by putting its current scatterer at the origin. Reversing its ordered centers and translating its initial scatterer to the origin recovers one original forward physical word, bijectively. No periodic translate is counted twice.

Fix $N \geq 1$. Let $\mathcal{U}$ be the open subset of $D_{N,R}^-$ having a specified complete lifted past word and specified *open* homogeneity strips at exactly the times $-1, \dots, -N$. There is no current homogeneity condition. For any $y \in \mathcal{U}$ let I_y be the connected open interval of $\Gamma_{R,\pi_R(y)} \cap \mathcal{U}$ containing y.

Lemma 70.2 (Physical boundary stopping)**.** *At either endpoint of I_y, at least one of the following events occurs in the limiting prescribed record, written in forward order $0, \dots, N$:*

(1) *the current angle φ_N is $+\pi/2$ or $-\pi/2$;*
(2) *for some $0 \leq i < N$, φ_i reaches a level $h \in \mathcal{H}$ bounding its fixed open strip;*

(3) *a flight $0 \le j < N$ is tangent in its interior to an unvisited lattice disk.*

All incidences $c_0, \dots, c_{N-1}$ in this limiting record are positive. In the third alternative the tangent foot is strictly inside the flight; no additional collision is inserted there.

For a chosen event, a tail of I_y reaches arbitrarily small positive interior values of its defining level distance while remaining in the same complete word and all the same open strips. Its image under that distance contains $(0, \eta)$ for some $\eta > 0$. The number η may depend on the word, the chosen interval, and R.

Proof. Parametrize the intrinsic affine fiber by $-\pi/2 < \varphi_N < \pi/2$ as in (65.2). An endpoint exists in the closed interval $[-\pi/2, \pi/2]$, and the interval itself supplies an approaching continuous path of actual regular records. All listed centers are fixed along it. Their contact points lie on the corresponding compact circles. Every physical flight has length in $[\tau_-, \tau_+]$.

Every one of the finitely many prescribed past strips has closure strictly inside $(-\pi/2, \pi/2)$. Thus $c_0, \dots, c_{N-1}$ have positive lower bounds along this interval; these bounds are not asserted uniform in its strip labels. Compactness gives a limiting prescribed record. The limit is unique: going backwards, the intersection with each prescribed previous circle is a simple first-entry root because that circle's limiting incidence is positive. The implicit circle-intersection formula consequently continues this prescribed branch uniquely and continuously at each of the finitely many steps. An unvisited tangency does not obstruct this *formal prescribed-word* continuation. No claim is made that the continued branch beyond the endpoint remains a physical first-hit branch.

Limits of the actual closed flight segments avoid every open disk. Only finitely many lattice disks can meet the uniformly bounded neighborhood of any one of these finitely many segments. Hence, if the current angle is nongrazing, no past strip boundary is reached, and no segment is tangent to an unvisited disk, every listed collision remains a strict physical first collision on a neighborhood of the endpoint. All unvisited clearances there are strict, and the finitely many prescribed strip conditions remain open. The endpoint would then belong to $\mathcal{U}$, and the affine interval would extend through it. This contradicts its maximality and proves the exhaustive list.

An unvisited contact cannot be an endpoint of the flight, since distinct radius-R disks are disjoint with a positive gap. A line segment avoiding an open disk can touch its boundary at an interior point only tangentially. This proves the stated nature of the third event. The geometric standoff of the original marked source applies to this foot.

For a terminal grazing event take the interior distance $\pi/2 - \varepsilon\varphi_N$, with $\varepsilon \in \{-1, 1\}$ the selected terminal sign. For a homogeneity event take $\varepsilon(\varphi_i - h)$, where ε is its side inside the selected open strip. For an unvisited tangency choose a sufficiently short tail on which the foot remains inside the same flight and the signed normal side of the marked disk is fixed; take the physical clearance $u > 0$ of this line from that disk. In each case the distance is continuous and strictly positive on the tail and tends to zero at the endpoint. The continuous image of an interval is an interval. Since it contains one positive value and values arbitrarily close to zero, it contains $(0, \eta)$ for some $\eta > 0$. This is the intermediate-value theorem on actual records, with no extension across the stopping boundary and no alteration of an event test. □

Lemma 70.3 (Common interior levels with actual source intervals)**.** *Select finitely many distinct physical words, one open past-strip class and one stopped affine interval for each. Assign each selected word to one of its stopping events. Group terminal events by their sign; homogeneity events by (i, h, ε); and unvisited events by their marked flight index j, with both mark signs kept on the original two source circles.*

Within any such group, every sufficiently small common interior level has a representative of each assigned word in its chosen open class. Each representative may be chosen away from all

cuts and exceptional sets of the one-sided source partitions. Different assigned words then belong to different complete overlay intervals, or different one-sided intervals in an endpoint group.

Proof. For a finite group take the minimum of the positive level ranges in Lemma 70.2; for marked flights also keep the common level at most 3/200. Its intermediate-value conclusion gives the required physical representatives. No uniform level neighborhood over an infinite word family is used.

The relevant source slice is open in its one-dimensional source parameter. For an internal horizontal source this follows by applying the regular physical map T_R^i to an open neighborhood of the selected record: the exact level $\varphi_i = h'$ cuts this open set in an open α_i interval. The same argument applies at $i = 0, N$.

For completeness, the marked source has the same property by an explicit physical determinant. At the departure state of flight j, put $\vartheta = s_j/R + \varphi_j$, let v be the outgoing unit vector, and put $b = (O - q_j) \cdot v^{\perp}$, $t = (O - q_j) \cdot v$ for the chosen marked center O. Direct differentiation gives

$$\partial_{s_j}\vartheta = \kappa_R, \quad \partial_{\varphi_j}\vartheta = 1, \quad \partial_{s_j} b = -c_j - \kappa_R t, \quad \partial_{\varphi_j} b = -t, \qquad \det \frac{\partial(\vartheta, b)}{\partial(s_j, \varphi_j)} = c_j > 0.$$

The signed clearance u differs from b by a fixed sign and the constant R on this tail. Thus (ϑ, u) is a local coordinate system there. A fixed positive u gives an open angular interval inside the same class, with the same interior foot and the same mark sign.

The excluded cuts and exceptional sets of either actual source partition have one-dimensional Lebesgue measure zero. Two such null sets cannot exhaust a nonempty open source interval. Choose the representative outside them. It belongs to one genuine one-sided interval in each required time direction. Every such interval fixes its complete physical half-word; an overlap fixes the full word. The original one-copy normalization follows the unique actual backward collisions to the initial disk and translates that disk to the fundamental copy. Consequently two different assigned physical words cannot occupy the same overlay interval. The same argument with one time direction proves the endpoint assertion. □

70.3. **The uniform unweighted sum.** Set

$$\mathcal{B}_* = 2H_{\mathrm{H}} + 2S_{\mathcal{H}}H_{\mathrm{H}} + \frac{4S_{\mathcal{H}}C_{\mathrm{H}}H_{\mathrm{H}}E_{\mathrm{H}}}{1-\sigma} + \frac{2Q_{\mathrm{mb}}}{1-\sigma}, \qquad Q_{\mathrm{stop}} := \max\{1, R_+\Gamma C_\times^2 \mathcal{B}_*\}. \tag{70.10}$$

All constants have been fixed by the physical geometry, original homogeneity family, native source growth, and cross-curve theorem. In particular Q_{stop} is independent of N, R, physical word, exact record label, roof interval, and the outer parameters of the raw local-limit problem.

Theorem 70.4 (Full original physical-word pressure)**.** *For the original true stable Euclidean arclength cocycle,*

$$P_N(R) := \sum_{w \in \mathcal{W}_N(R)} \sup_{x \in [w]_R \cap M'_R} j_N(x) \le Q_{\mathrm{stop}} \qquad (R \in [R_-, R_+],\ N \ge 0). \tag{70.11}$$

Every physical collision word is included. No capacity abundance premise, connectedness premise for a word domain, exclusion of short cells, positive endpoint incidence cap, or pressure theorem for a different model is used.

Proof. The case $N = 0$ is the original convention $P_0 = 1$. For $N \ge 1$ let $\mathcal{D}_w$ be the finite regular past domain of the current-normalized version of the forward word w. Put

$$S_w = \sup_{y \in \mathcal{D}_w} w_N(y).$$

A word with empty such domain has zero contribution. Equation (70.7) gives

$$P_N(R) \le C_\times \sum_w S_w. \tag{70.12}$$

The finite cone expansion bounds these affine inverse Jacobians, so each S_w is finite.

Fix $0 < \epsilon < 1$ and any finite set of contributing physical words. For each choose y_w off all past homogeneity boundaries with

$$w_N(y_w) \ge (1-\epsilon)S_w.$$

This choice is legitimate: on the finite regular domain w_N is continuous, and the pullback of every fixed homogeneity level under a regular finite iterate is a smooth coordinate level with empty interior. The countably many such levels have locally finite intersection with a neighborhood on which the finitely many collision angles remain away from grazing. Their complement is therefore dense.

Choose the open past-strip class containing y_w, stop its affine interval by Lemma 70.2, and assign one stopping event. For every actual representative z in that same class, the complete-word estimate of Corollary 61.5 gives

$$(1-\epsilon)S_w \le w_N(y_w) \le C_\times w_N(z) \le C_\times R_+ \Gamma\, c_0(z) u_x(z). \tag{70.13}$$

Only the matching past strips $-1, \dots, -N$ are used here. There is no current strip and no future invariant-line regularity requirement on z. In particular the transfer does not follow a physical continuation from y_w to z.

Sum (70.13) by the finite source groups supplied by Lemma 70.3. It remains to bound the sum of $c_0 u_x$ of their selected representatives.

For a terminal grazing group, choose the common nongrazing level $\varphi_N = \varepsilon(\pi/2 - \delta)$. The endpoint one-sided source bound, the uniqueness of the assigned interval, and $c_0 \le 1$ pay the group by H_{H}. The two terminal signs cost $2H_{\mathrm{H}}$ in total.

For a homogeneity group (i, h, ε) take the common level $\varphi_i = h + \varepsilon\delta$. If $i = 0$, then $c_0 = \cos(h + \varepsilon\delta)$ exactly, so the one-sided bound pays its sum of $c_0 u_x$ by $\cos(h + \varepsilon\delta) H_{\mathrm{H}}$. The selected S_w on the left of (70.13) do not depend on δ. Letting this common positive parameter decrease to zero gives the budget $\cos h\, H_{\mathrm{H}}$. Both sides of every level and every level together cost $2S_{\mathcal{H}} H_{\mathrm{H}}$.

For $1 \le i < N$, use $c_0 \le 1$ and (70.3). The same common-level argument and the limit of its explicit cosine give the budget

$$\cos h\, C_{\mathrm{H}} H_{\mathrm{H}} E_{\mathrm{H}} (\sigma^{i-1} + \sigma^{N-i-1})$$

for that group. The sum over the two sides, all levels, and all internal indices is at most

$$2S_{\mathcal{H}} C_{\mathrm{H}} H_{\mathrm{H}} E_{\mathrm{H}} \sum_{i=1}^{N-1} (\sigma^{i-1} + \sigma^{N-i-1}) \le \frac{4S_{\mathcal{H}} C_{\mathrm{H}} H_{\mathrm{H}} E_{\mathrm{H}}}{1-\sigma}.$$

This is where the actual collision incidence $\cos h$ is essential: a uniform bound without it would not pay the infinitely many original homogeneity boundaries.

For an unvisited group at index j, choose a common positive clearance u and use (70.4). The original two marked signs and the correct physical copy are already included. Its budget is $Q_{\mathrm{mb}}(\sigma^j + \sigma^{N-j-1})$. Thus all flight indices cost at most

$$Q_{\mathrm{mb}} \sum_{j=0}^{N-1} (\sigma^j + \sigma^{N-j-1}) \le \frac{2Q_{\mathrm{mb}}}{1-\sigma}.$$

No zero-clearance trace, grazing derivative, or factor counting unvisited disk labels is substituted into this step.

Combining the four positive budgets proves, for the chosen finite physical word family,

$$\sum_{w} S_w \le \frac{C_\times R_+ \Gamma}{1-\epsilon}\mathcal{B}_*.$$

Every selected word was assigned once. Source partitions, levels, and representatives may depend on this finite family, but the displayed constant does not. Nonnegative exhaustion therefore proves the same inequality for the full word set; in the present finite-horizon geometry there are already only finitely many N-flight words. Let $\epsilon \downarrow 0$ and apply (70.12). This gives (70.11) with exactly (70.10). The common-level limits occurred at fixed R, N and a fixed finite selected family. No physical collision-time limit, roof essential supremum, or outer limit has been interchanged. □

Corollary 70.5 (Completion of the original geometric pressure input)**.** *Input 11.1 holds on the original radius family with*

$$C_0 = 30000, \qquad C_1 = 6400, \qquad Q_* = Q_{\mathrm{stop}}.$$

Consequently the formerly pressure-dependent constants in module 266 may be taken unconditionally as

$$C_\star = \frac{30000 Q_{\mathrm{stop}}}{2c_*\lambda_\circ}, \qquad K_\star = 6400 D Q_{\mathrm{stop}}.$$

Proof. The two optical inequalities are Theorem 54.3; the remaining full physical-word sum is Theorem 70.4. All parameters, measures, clock conventions, and exact-record selectors in that input have been retained. Thus Theorem 11.3 and Corollary 11.4 in module 266 apply with these constants. Their explicit descendants in modules 267–269, namely Theorems 12.2, 13.1, 14.2, and 15.1, likewise retain their stated conclusions with $Q_* = Q_{\mathrm{stop}}$. These are inherited consequences activated by the newly proved geometric input; their proof bodies and all their other hypotheses remain unchanged. □

Scope of the closure. This theorem pays the original unweighted physical-word pressure, including the small-capacity word suprema left by Section 65; it does not replace that capacity theorem. The proof is independent of a planar-cell abundance hypothesis and of a radius-uniform interpretation of a fixed-table external pressure theorem. The remaining spectral, arithmetic, conditional microscopic-return, and raw pointwise local-limit obligations retain their original quantifiers and order of limits. A uniform word-pressure bound is a completed geometric input to those obligations, not a claim that their full conjunction has already been proved. In particular the normalized positive payments in modules 267–269 concern precisely the incidence events specified there. They do not supply an m^{-2} same-roof estimate for a single unvisited-clearance witness, nor prove that the retained low-load shortage in (15.8) vanishes in its original ordered limits.

71. Length-sensitive transport of actual clearance sources

We keep the physical model, radius interval, original section and unit-speed clock fixed. In this section a marked closest point is a coordinate on a free line, never a reflection. The purpose is to retain the length of a small source in a complete branch-supremum estimate. An unscaled source mass or a bound at one common endpoint momentum would not supply that estimate. The full-word pressure theorem of Section 70 is not used below.

Use the enlarged physical curve class of (26.5), with its own curvature allowance $\widehat{B}_{\mathrm{cv}}$. Set

$$t_* = 3/50, \quad k_- = 100/47, \quad k_+ = 20/9, \quad U = 170/9, \quad \sigma = 47/59, \quad \Gamma = \sqrt{1+k_+^2}.$$

Choose its own homogeneous cutoff $\widehat{k}_0$ so that

$$\theta_{\mathrm{r}} := \sigma + \frac{J_f}{k_- t_* (\widehat{k}_0 - 1)} < \frac{1+\sigma}{2} < 1. \tag{71.1}$$

The one-step and distortion proofs in Theorems 24.2 and 24.4 apply with this fixed allowance and cutoff. Take a common short length $\ell > 0$ small enough for those proofs, the first-hit geometry, and the relative-angle condition below. The choices are independent of radius, collision count, mark level, original exact labels and outer parameter B. Let D be the resulting Euclidean distortion constant and put

$$D_\ell = e^{D\ell^{1/3}}, \qquad C_\ell = \frac{3\Gamma D_\ell}{\ell(1-\theta_{\mathrm{r}})}, \qquad G_\ell = \Gamma + C_\ell \ell. \tag{71.2}$$

71.1. **The complete renewal, including the last generation.** At each step take every maximal regular physical homogeneous image. Only images longer than ℓ are subdivided, into pieces of lengths in $[\ell/2, \ell]$. Images of length at most ℓ are not artificially subdivided. Write $\mathcal{G}_n(C)$ for the complete resulting partition of the nth image of a qualified curve C, retaining each source label. Call a piece long when its length is at least $\ell/3$.

Theorem 71.1 (Length-sensitive physical renewal)**.** *For every qualified homogeneous curve C of length at most ℓ, every $n \geq 0$ and every original radius,*

$$\sum_{V \in \mathcal{G}_n(C)} \sup_V J_V(T_R^{-n}) \leq \Gamma\theta_{\mathrm{r}}^n + C_\ell|C| \leq G_\ell. \tag{71.3}$$

The sum contains all regular physical descendants; the omitted singular source points have arclength zero. The same statement holds after physical reversal. It is an estimate of actual inverse Jacobian suprema, not an average or an independent-branch model.

Proof. First consider descendants having no long piece at the positive generations $1, \dots, n$. They cannot pass through a length subdivision, since every child of such a subdivision has length at least $\ell/2$. At each generation their inverse adapted-Jacobian sum is therefore bounded by the full maximal homogeneous sum in (24.4). Products and nonnegative summation give θ_{r}^n. The endpoint equivalence of the adapted and Euclidean curve norms costs Γ once for the entire n-step Jacobian, and not once per step. Their total contribution is at most $\Gamma\theta_{\mathrm{r}}^n$. This also covers $n = 0$.

Every other terminal piece has a unique last long ancestor W at a time $k \in \{1, \dots, n\}$. Denote its original preimage by C_W. Its images are regular and homogeneous, and $\ell/3 \leq |W| \leq \ell$. The Euclidean distortion theorem yields

$$\sup_W J_W(T_R^{-k}) \leq D_\ell \frac{|C_W|}{|W|} \leq \frac{3D_\ell}{\ell}|C_W|.$$

The descendants after this last long ancestor have no later long piece. The preceding adapted-norm argument bounds the complete sum of their inverse suffix Jacobians by $\Gamma\theta_{\mathrm{r}}^{n-k}$. For every fixed k, the genuine preimage intervals C_W of the long pieces are disjoint in C; therefore $\sum_W |C_W| \leq |C|$. The contribution of all last-long ancestors is consequently at most

$$\frac{3\Gamma D_\ell}{\ell}|C| \sum_{k=1}^{n} \theta_{\mathrm{r}}^{n-k} \leq C_\ell|C|.$$

The endpoint $k = n$ is included and has exponent zero. No terminal long piece is discarded. All calculations can first be made for a finite subfamily; nonnegative exhaustion gives the complete sum. The physical one-step singularities and homogeneous boundaries have countably many preimages on the qualified curves, with zero arclength as in the native growth proof. Reversal is the exact arclength isometry conjugating T_R to T_R^{-1}; it transports the original observations and does not replace the section. This proves (71.3). □

71.2. **A short interval cover of the true clearance event.** Write N_* for the complete candidate-center bound of (21.6), using the proved physical horizon, and put

$$\delta_* = 3/200, \qquad a_{\mathrm{c}} = 9/200, \qquad K_{\mathrm{c}} = 2N_*, \qquad L_{\mathrm{c}} = \frac{2N_*\sqrt{1+U^2}}{k_- a_{\mathrm{c}}}. \tag{71.4}$$

Let $E_{\delta,R} = \{N_{\delta,R} \geq 1\}$ be the original event that the actual outgoing flight has an unvisited clearance witness of width δ. All physical visibility and interior-foot conditions in that definition are retained.

Lemma 71.2 (Actual-flight clearance cover)**.** *For $0 < \delta \leq \delta_*$, on any qualified short graph C the set $C \cap E_{\delta,R}$ is covered, apart from endpoints, by at most K_{c} disjoint source intervals whose combined arclength is at most $L_{\mathrm{c}}\delta$. The same bound holds on a short horizontal momentum graph, using $0 \leq V \leq U$ in the proof. The cover consists of true outgoing states. It can be larger than the selected event, but every subsequent evolution always takes its actual first hit.*

Proof. Use boundary arclength s, outgoing angle φ, physical velocity v and $V = d\varphi/ds$. For a candidate center z put

$$t_z = (z-q)\cdot v, \qquad b_z = (z-q)\cdot v^{\perp}, \qquad \eta_z = \arg(z-q) - \arg v.$$

Differentiating the actual circular geometry gives

$$b_z' = -\cos\varphi - t_z(\kappa + V), \qquad t_z' = -\sin\varphi + b_z(\kappa+V), \qquad \kappa = 1/R. \tag{71.5}$$

Since $|z-q| \geq 1 - R_+$, we have $\eta_z' \leq (1-R_+)^{-1} - k_- < 0$ even when $V = 0$. The short length was chosen so that the variation of each η_z is at most $\pi/2$. Thus its positive-foot set is an interval. On that interval b_z is strictly decreasing.

The two bands $R < |b_z| < R + \delta$, with $t_z > 0$, each give at most one interval. Throughout either band,

$$t_z^2 = |z-q|^2 - b_z^2 \geq (1-R_+)^2 - (R_+ + \delta_*)^2 = \frac{1827}{40000} > a_{\mathrm{c}}^2.$$

Equation (71.5) therefore gives $|b_z'| \geq k_- a_{\mathrm{c}}$. The length in s of each band is at most $\delta/(k_- a_{\mathrm{c}})$. Conversion to curve arclength costs at most $\sqrt{1+U^2}$. Sum over both signs and the at most N_* possible centers, and merge overlapping intervals. Merging decreases both the number of components and their combined length.

Every genuine witness belongs to this cover. We did not impose the upper foot restriction $t_z < \tau$ when forming the cover. A covered state whose candidate foot is beyond an earlier collision is still evolved to that earlier first collision; its inclusion only enlarges a positive source estimate. Endpoints of the weak bands do not alter the length bound. On a regular graph each such endpoint is isolated because of the strict derivative just proved. The same reasoning, with $V = 0$, proves the horizontal assertion. □

71.3. **The two physical source classes and their exact weights.** We use (i) the two marked angular circles of Theorem 21.3, with source coordinate $\xi = \vartheta$ and measure $\sum_{\varsigma=\pm1} d\vartheta$, and (ii) the horizontal momentum fibre F_a, $a \in (-1,1)$, with $\xi = \alpha_0$ and measure $d\alpha_0$. Their respective masses are 4π and 2π. For (i), $u_0 \in [0, \delta_*]$; at $u_0 = 0$ use the original strict positive-side unvisited trace, with no reflection at the foot. Both first-image families are now formed with the same enlarged class, cutoff $\widehat{k}_0$ and length ℓ fixed above.

Here are fully specified uniform source budgets. Set

$$M_* = 8N_* + 1, \qquad L_0 = \frac{\pi\sqrt{1+U^2}}{k_-},$$

$$J_0 = 2 + \left\lceil \frac{2\pi R_+}{\ell} \right\rceil, \qquad J_{\mathrm{len}} = 2 + \left\lceil \frac{2\max\{L_0, 2\pi R_+\sqrt{1+U^2}\}}{\ell} \right\rceil, \tag{71.6}$$

$$A_{\mathrm{ang}} = \frac{2M_* J_{\mathrm{len}}}{a_\circ}\left(1 + \frac{2}{\widehat{k}_0 - 1}\right), \qquad A_{\mathrm{hor}} = \frac{J_0 J_f J_{\mathrm{len}}}{t_*}\left(1 + \frac{2}{\widehat{k}_0 - 1}\right),$$

$$A_{\mathrm{src}} = \max\{A_{\mathrm{ang}}, A_{\mathrm{hor}}\}, \qquad E_{\mathrm{lc}} = \frac{1}{k_- t_*} = \frac{47}{6},$$

$$H_{\mathrm{lc}} = \max\left\{2M_* E_{\mathrm{lc}}, J_0 J_f E_{\mathrm{lc}}, \frac{A_{\mathrm{src}} G_\ell J_f}{k_- t_*}\right\}. \tag{71.7}$$

Shorten ℓ also below ℓ_{pre} in (26.7). The exact first-image formulas (21.2) and (26.3) then show that the sum of density suprema in either family is at most A_{src}. These bounds retain the initial grazing factor c before summing every homogeneous strip. They are valid at $u_0 = 0$ by the direct trace construction of Theorem 57.4. The bounds are not conditional probabilities on the source fibres.

Theorem 71.3 (Localized half-source capacity)**.** *Start from either of the preceding physical sources and take its first true collision Y. Fix $h, n \geq 0$ and select $T_R^h Y \in E_{\delta,R}$. Propagate n further true collisions from that selected outgoing state, and then one last true collision. Thus the half-count from the original source is exactly*

$$l = h + n + 2. \tag{71.8}$$

Let $A_l = |\partial_\xi p_l|$ be the actual final momentum derivative. There is a countable disjoint family of regular source intervals, with fixed physical half-words, covering the selected event up to source-null sets, such that

$$\sum_I \sup_I A_l^{-1} \leq K_{\mathrm{loc}}(\delta + \theta_{\mathrm{r}}^n), \qquad K_{\mathrm{loc}} := \frac{A_{\mathrm{src}} G_\ell J_f}{k_- t_*}(\Gamma K_{\mathrm{c}} + C_\ell L_{\mathrm{c}}). \tag{71.9}$$

Every individual interval has the bound

$$\sup_I A_l^{-1} \leq E_{\mathrm{lc}} \sigma^{l-1}. \tag{71.10}$$

The complete unselected half-source, at every $l \geq 1$, has a corresponding interval partition with sum at most H_{lc} and the same individual bound. All assertions hold in the reversed physical direction with the original observations transported.

Proof. Push the first-image family through h true collisions. By Theorem 71.1, the sum of the suprema of all exact transported densities on the resulting short homogeneous curves is at most $A_{\mathrm{src}} G_\ell$. On each such curve use the disjoint clearance cover of Lemma 71.2. Apply the renewal inequality separately to its at most K_{c} intervals through n further true collisions. Their combined inverse-Jacobian sum is at most

$$\Gamma K_{\mathrm{c}} \theta_{\mathrm{r}}^n + C_\ell L_{\mathrm{c}} \delta.$$

This is where both component count and actual selected length are paid. A small mass is not substituted for an inverse-Jacobian supremum.

On each resulting preterminal curve use the exact last-collision cancellation (26.18). The derivative of final momentum per incoming arclength is at least $k_- t_*$, and there are at most J_f genuine next-hit pieces. Do not make an infinite terminal homogeneous subdivision. Chain

the exact source density, inverse transport Jacobian and terminal derivative; then sum over the last-hit cuts, selected covers, transported curves and first-image family. This gives the first inequality in (71.9). All intervals are pullbacks through injective physical branches, so disjoint covers remain disjoint in the original source. Intermediate physical and homogeneous cuts have the native zero-arclength preimages; a bounded Borel observation can be retained inside the resulting positive integrals without differentiating it or counting its components.

For the individual inequality use the actual positive transverse Jacobi product, not the averaged density. On a marked angular source its first incoming-side entries are $B = a^{\mathrm{foot}} \geq a_\circ > t_*$ and $D = 1$. On a horizontal momentum source they are $B = R\cos\varphi_0 + \tau \geq t_*$ and $D = 1$; its exact first endpoint derivative is $\cos\varphi_0 + \tau/R + c_1$. Each intervening kick and following physical flight replaces them by

$$B_{\mathrm{new}} = (1 + \tau k)B + \tau D, \qquad D_{\mathrm{new}} = kB + D, \qquad k = 2/(Rc) > 0.$$

Since $\tau \geq t_*$ and $c \leq 1$, the factor of B is at least $1 + 2t_*/R_+ = 59/47$. The terminal endpoint is incoming: its momentum derivative is $\kappa B + c_l D$, with no additional kick. It is therefore at least $k_- t_* \sigma^{-(l-1)}$. This proves (71.10) pointwise and on every subinterval, with no positive lower incidence cutoff.

For the complete unselected source with $l \geq 2$, use the first family, renewal through $l - 2$ collisions and the same final cancellation. The sum is at most $A_{\mathrm{src}} G_\ell J_f/(k_- t_*)$. At $l = 1$ the marked circles have at most $2M_*$ first-hit/seam intervals; the horizontal fibre has at most $J_0 J_f$. Apply the individual bound. These are exactly the three terms in H_{lc}. Direct construction at the strict zero-clearance trace uses the same first-image density, standoff and class estimates, as in Theorem 62.1; no weak-limit transfer of a supremum is invoked. Physical reversal preserves arclength and source-angle measure and reverses momentum signs. It proves the backward assertions without changing the conditioning event. □

Corollary 71.4 (The selected angular source mass)**.** *For either source after its first true collision, and every $h \geq 0$,*

$$\lambda\{T_R^h Y \in E_{\delta,R}\} \leq A_{\mathrm{src}} G_\ell L_{\mathrm{c}} \delta. \tag{71.11}$$

For the horizontal source selected before its first collision,

$$\int_{F_a} \mathbf{1}_{E_{\delta,R}}\, d\alpha_0 \leq J_0 L_{\mathrm{c}} \delta / R_-. \tag{71.12}$$

These are one-dimensional physical source estimates. They will become cap-boundary charges only after the exact Jacobian cancellation proved in the next section.

Proof. The total transported density supremum is at most $A_{\mathrm{src}} G_\ell$. Integrate it on the clearance covers of combined length at most $L_{\mathrm{c}}\delta$, and sum the exact labelled sources. For the initial horizontal fibre divide the circle into the J_0 preliminary arcs and apply the horizontal cover directly. Since $ds = R\, d\alpha_0$, the conversion costs at most R_-^{-1}. Every retained original observation is restored by positive restriction. No selected event has been replaced by a normalized fibre law. □

72. A selected physical roof height for an actual clearance witness

We now use the length-sensitive source estimate at the same physical roof as the observation. Keep the complete cap $\mathcal{C}_{m,R}$, its strict regular convention, and the original normalization of (67.1). Thus $a = 5/6$, every original exact event is a positive restriction of this cap source, and the unconditioned collision measure is divided by c_* exactly once. For $m \geq 1$, $0 \leq j < m$ and $0 < \delta \leq \delta_* = 3/200$, define

$$E_{j,\delta,R} = \{x : N_{\delta,R}(T_R^j x) \geq 1\}, \qquad \mu_{m,R;j,\delta} = (L_{m,R})_*(\mathbf{1}_{\mathcal{C}_{m,R}} \mathbf{1}_{E_{j,\delta,R}}\, d\nu/c_*). \tag{72.1}$$

The index j denotes an actual flight between collisions j and $j+1$. No collision is inserted at a closest point. The measure is a positive majorant on the original roof line; it is not a new conditional law.

72.1. **Turning constants for the enlarged physical source class.** The enlarged class, cutoff $\widehat{k}_0$, short length ℓ, and constants in Section 71 are used throughout this section. To avoid reusing a constant for a smaller curve class, set

$$\begin{aligned}
B_2 &= \widehat{B}_{\mathrm{cv}}(1+U^2)^{3/2},\\
a_0 &= 2k_-t_*, \qquad L_h = 3\sqrt{1+U^2}/k_-,\\
c_H &= \sin(\widehat{k}_0^{-2}), \qquad c_a = \sqrt{1-a^2} = \sqrt{11}/6,\\
D_B &= t_*^{-1} + 2/(t_*a_0) + (B_2/a_0)(1+(2k_-)^{-1}) + B_2,\\
V_H &= \max\{-2\log c_H, \log(\pi/2) + 2\log(1+\widehat{k}_0^{-1})\},
\end{aligned} \tag{72.2}$$

$$\begin{aligned}
D_{\mathrm{var}} &= \max\{(U/c_H + D_B)L_0^{2/3}, (2\pi U + D_B)L_h^{2/3}\},\\
V_J &= D_{\mathrm{var}}\Gamma^{1/3}L_0^{1/3}/(1-\sigma^{1/3}),\\
V_\rho &= V_H + (a_\circ^{-1} + b_*a_\circ^{-2} + K_0/2)\ell,\\
V_t &= -2\log c_a + B_2L_0/k_-,\\
V_1 &= L_0(k_+b_*+1)/(k_-a_\circ^2),\\
V_A^{\mathrm{lc}} &= \max\{V_1, V_\rho + V_J + V_t\},\\
T_{\mathrm{lc}} &= 2\pi + V_A^{\mathrm{lc}}.
\end{aligned} \tag{72.3}$$

Here L_0, Γ, U and σ have the meanings fixed above, $a_\circ = \sqrt{1827}/200$, $b_* = 97/200$, and K_0 is the exact first-image derivative bound in (21.3). All these constants are fixed before a collision, radius, label, level or roof supremum.

Lemma 72.1 (Turning on the actual selected interval covers)**.** *On each overlap of the two physical half-source interval partitions of Theorem 71.3, use the original cap and set* $P(\vartheta) = (-p_0(\vartheta), p_m(\vartheta))$*. The complete punctured turning integral on that overlap obeys*

$$\int_{\{P\neq 0\}} |d\arg P| \le T_{\mathrm{lc}}. \tag{72.4}$$

This includes the selected covers of that theorem and the strict unvisited trace at $u_0 = 0$*. Every retained Borel observation may be inserted as a selector in this positive integral.*

Proof. We reproduce the variation argument for the enlarged class. On an actual homogeneous image segment of arclength r in a signed strip H_k one has

$$r \le L_h k^{-3}, \qquad c \ge (2/\pi)(k+1)^{-2}.$$

The exact one-step logarithmic derivative (24.6), with (72.2), has absolute value at most $U/c + D_B$ per unit image arclength. Its absolute integral is at most $(2\pi U + D_B)k^2 r \le (2\pi U + D_B)L_h^{2/3}r^{1/3}$. On the central strip the same integral is at most $(U/c_H + D_B)L_0^{2/3}r^{1/3}$. Each unstable graph has total arclength at most L_0, because $V \ge k_-$ and its angle ranges in $(-\pi/2, \pi/2)$. Adapted expansion gives the intermediate length bound $L_i \le \Gamma\sigma^{l-i}L_l$. Summing the absolute integrals of the actual one-step logarithms therefore bounds the variation of a full homogeneous arclength Jacobian by V_J. This is an estimate of total variation; it is not inferred merely from a bound on differences of endpoint values.

For a half-ray with $l \geq 2$ true collisions, its exact angular first-image density is $\rho_C = c_1/(a^{\mathrm{foot}}\sqrt{1+V_1^2})$. The monotonicity of φ_1 and the strip bounds give variation of $\log c_1$ at most V_H. The exact first-flight identities (21.5) give $|(a^{\mathrm{foot}})_s| \leq 1 + b_*/a_\circ$ and $|V_{1,s}| \leq K_0$. Thus $\operatorname{Var} \log \rho_C \leq V_\rho$. This step retains the first grazing factor. The arclength chain rule is exactly

$$A_l^p = |\partial_\vartheta p_l| = \frac{c_l V_l}{\sqrt{1+V_l^2}} \frac{J_C(T_R^{l-1})}{\rho_C}.$$

The terminal cap is one interval, since the physical endpoint momentum is strictly monotone. It lies inside the central strip for every $\widehat{k}_0 \geq 2$. On it $c_l \geq c_a$, and geometric curvature gives $|V_{l,s}| \leq B_2$. Consequently the terminal factor has logarithmic variation at most V_t. The preceding three bounds give $\operatorname{Var} \log A_l^p \leq V_\rho + V_J + V_t$.

For $l = 1$ the actual incoming endpoint derivative is $A_1^p = \kappa a^{\mathrm{foot}} + c$. With signed impact $b = \varsigma(R + u_0)$, direct differentiation gives

$$\partial_s(\kappa a^{\mathrm{foot}} + c) = c(\kappa b - p)/a^{\mathrm{foot}}.$$

Its logarithmic derivative has magnitude at most $(k_+ b_* + 1)/(k_- a_\circ^2)$, hence variation at most V_1. Physical reversal proves the same statement for the backward half. Refining a connected branch by any of the interval covers in Section 71 only decreases these variations.

On an overlap, both coordinate derivatives of P have nonzero fixed signs by (59.4). Its tangent remains in one open quadrant, and P has at most one zero. Its tangent angle, as a function of the logarithm of the absolute slope ratio, has derivative bounded by $1/2$. Since each half logarithmic variation is at most V_A^{lc}, the tangent turning is at most V_A^{lc}. Lemma 63.3 gives (72.4), with 2π allowing a single zero. This punctured estimate omits a polar-angle jump at zero; it does not assign the separate incoming critical fan a zero value. Borel selectors multiply the bounded positive integral and are never differentiated. Compact regular exhaustion proves all open-interval and one-sided trace statements. □

72.2. **All outgoing boundary charges of the selected source.** For a regular physical root write u_x for the absolute inverse endpoint-momentum Jacobian, and $h_x = R^2 e^{\mathsf{T}} G^{-1} e$. The exact identity of Corollary 59.2 gives

$$A = \frac{u_x}{4\pi c_* h_x} \leq C_A u_x, \qquad C_A = \frac{\Lambda_\circ}{4\pi c_* R_-^2}. \tag{72.5}$$

Set

$$\begin{aligned} C_{\mathrm{foot}} &= 607/90, & K_{\mathrm{bd}} &= C_A T_{\mathrm{lc}} C_{\mathrm{foot}} E_{\mathrm{lc}}, \\ C_{\mathrm{side}} &= \frac{K_{\mathrm{bd}}(2K_{\mathrm{loc}} + H_{\mathrm{lc}})}{1-\sigma}, & C_\delta &= 2K_{\mathrm{bd}} H_{\mathrm{lc}}, \\ M_{\mathrm{cap}} &= \max\{A_{\mathrm{src}} G_\ell L_{\mathrm{c}}, J_0 L_{\mathrm{c}}/R_-\}, & C_{\mathrm{cap,loc}} &= 4C_A M_{\mathrm{cap}}/a. \end{aligned} \tag{72.6}$$

All marked sums in these constants include both impact signs.

Lemma 72.2 (Selected old clearance boundaries and new level boundaries)**.** *The total regular boundary charge of the selected source* (72.1) *on all original unvisited tangencies is at most*

$$C_{\mathrm{side}}\{\delta + \theta_{\mathrm{r}}^j + \theta_{\mathrm{r}}^{m-1-j}\}. \tag{72.7}$$

The additional regular boundary at clearance level δ on flight j has charge at most

$$C_\delta\{\theta_{\mathrm{r}}^j + \theta_{\mathrm{r}}^{m-1-j}\}. \tag{72.8}$$

These are complete positive boundary sums before any original exact-label or roof restriction.

Proof. Consider an old unvisited tangency on flight k. The two actual half-lengths from its unreflected foot are $k+1$ and $m-k$. If $k<j$, the selected flight is in the forward half. In (71.8) the precise indices are

$$h=j-k-1, \qquad n=m-j-1, \qquad l=m-k.$$

Both h and n are nonnegative. The backward half is unselected. The interval-overlay inequality (62.3), the selected capacity (71.9), and the individual bound (71.10) give

$$\sum_{\text{actual selected overlaps}} \sup u_x \le C_{\mathrm{foot}}E_{\mathrm{lc}}\{K_{\mathrm{loc}}\sigma^k(\delta+\theta_{\mathrm{r}}^{m-j-1})+H_{\mathrm{lc}}\sigma^{m-k-1}\}.$$

Indeed the exact determinant bounds each root by $C_{\mathrm{foot}}(A_-^pA_+^p)^{-1}$. The overlay charges a product intersection to the interval with the later left endpoint; each interval is charged at most once to an interval whose left endpoint precedes its own. Its two sums are the selected total times the opposite individual maximum and the unselected total times the selected individual maximum. This proves the displayed bound even for arbitrarily short intervals. It is not a product of physical probabilities. Counts of words or components are not multiplied onto it afterwards.

For $k>j$ use exact reversal. The selected backward half now has

$$h=k-j-1, \qquad n=j, \qquad l=k+1,$$

and its bound is

$$C_{\mathrm{foot}}E_{\mathrm{lc}}\{K_{\mathrm{loc}}\sigma^{m-k-1}(\delta+\theta_{\mathrm{r}}^j)+H_{\mathrm{lc}}\sigma^k\}.$$

At $k=j$ no separate selected-half assertion is needed: pay the whole original boundary by $C_{\mathrm{foot}}E_{\mathrm{lc}}H_{\mathrm{lc}}(\sigma^j+\sigma^{m-j-1})$. Multiply these three bounds by C_AT_{lc}, using Lemma 72.1. For $k\neq j$ the selected trace is covered by the weak clearance intervals of Lemma 71.2. Their newly included endpoints have zero source-angle measure by its strict impact derivative; no separate positive trace is removed. At $k=j$ the whole positive boundary was already retained.

Summing over every k gives the sharper upper bound

$$\frac{K_{\mathrm{bd}}}{1-\sigma}\{2K_{\mathrm{loc}}\delta+(K_{\mathrm{loc}}+H_{\mathrm{lc}})(\theta_{\mathrm{r}}^j+\theta_{\mathrm{r}}^{m-1-j})\}.$$

Here, for example, the unselected tail for $k<j$, together with its $k=j$ endpoint, is bounded by $H_{\mathrm{lc}}\sigma^{m-j-1}/(1-\sigma)$; the other tail is its reversal. Since $\sigma\le\theta_{\mathrm{r}}$, this proves (72.7). The marked angular construction is a one-copy parametrization after fixing the incident flight index and mark; the complete two-sign source already includes every admissible marked displacement and physical word. Thus no unvisited disk, collision word or shallow index has been omitted.

The new boundary is at $u_0=\delta$ on flight j. The unselected half-capacity bound and the same turning argument give $K_{\mathrm{bd}}H_{\mathrm{lc}}(\sigma^j+\sigma^{m-j-1})$ for one incident side. We allow both sides explicitly, obtaining (72.8). For an event with several qualifying marks, assign each physical boundary germ to its first active mark in a fixed ordering; discarding this assignment only enlarges the complete positive marked sum. A boundary where a selected foot reaches a flight endpoint is impossible in this range: at an incident contact its distance to a third center is at least $1-R$, whereas $R+\delta_*<1-R$ uniformly. The two incident disks themselves are never counted as unvisited marks. Thus the displayed list of new physical boundaries is complete. □

Lemma 72.3 (Selected endpoint-cap payment by the physical Jacobian)**.** *The complete residual regular cap-side charge of* (72.1) *is at most* $C_{\mathrm{cap,loc}}\delta$.

Proof. On $p_0=a$, write $v=p_m$. On every genuine regular root branch of the endpoint map its inverse Jacobian obeys the exact identity

$$u_x\,|dv|=|d\alpha_0|.$$

The other component of the endpoint map is the unchanged coordinate p_0, so this is simply its triangular determinant on the actual fibre, with no probability normalization. Moreover $|d\arg(-a,v)| = a(a^2+v^2)^{-1}|dv| \le |dv|/a$. By (72.5) the selected side charge is at most C_A/a times the selected $d\alpha_0$ mass. For $j=0$ use (71.12); for $j \ge 1$ use (71.11) with $h = j-1$. Each is at most $M_{\mathrm{cap}}\delta$.

The side $p_0 = -a$ has the same estimate. For $p_m = \pm a$ apply physical reversal, which changes the selected index to $m-1-j$; the preceding initial and later-index cases cover it without an additional factor. Adding the four sides gives exactly the claimed constant. At simultaneous boundaries assign visited grazing first, then remaining critical endpoints, then original unvisited constraints, then the new level constraint, and finally the cap. The residual cap roots are therefore genuine regular roots where the displayed Jacobian identity applies. Corners and coordinate seams are assigned once. In particular this proof does not restrict an arbitrary almost-everywhere version of a two-dimensional density to a null line. □

72.3. **The selected interior and the incoming critical endpoints.**

Theorem 72.4 (Complete physical selected roof height)**.** *The measure* (72.1) *is absolutely continuous. Its density $p_{m,R;j,\delta}$ satisfies, for every allowed m, R, j, δ,*

$$\|p_{m,R;j,\delta}\|_{L^\infty(\mathbb{R})} \le C_{\mathrm{clr,roof}}\{\delta^{1/4} + \theta_{\mathrm{r}}^{j} + \theta_{\mathrm{r}}^{m-1-j}\}, \tag{72.9}$$

where a fixed admissible choice is

$$C_{\mathrm{clr,roof}} = \frac{C_{\mathrm{int}} J_\times}{c_* R_-^{1/4}} + C_{\mathrm{side}} + C_\delta + C_{\mathrm{cap,loc}}. \tag{72.10}$$

The complete actual-word, physical mark and angular sums precede every essential roof supremum and every supremum over the original radius and exact labels. All original joint return, displacement and collision tests, the original section, and every Borel selector $0 \le H \le 1$ may be retained by positive restriction. The theorem is for their unchanged physical time and makes no assertion about an independent momentum process or a modified conditional event.

Proof. Absolute continuity follows by positive domination by the complete cap measure in (67.1) and its physical coarea. For the localized interior variation use the exact pointwise bound (67.4). The full multiplier theorem (56.5) and the exact clearance flux give

$$\begin{aligned}\int \sum_w \int_{\mathcal{D}_w(\theta)\cap E_{j,\delta,R}} |\partial_t A_w|\, dt\, d\theta &\le \frac{C_{\mathrm{int}}}{c_*} \int \mathbf{1}_{E_{j,\delta,R}} Z_m \, d\nu \\ &\le \frac{C_{\mathrm{int}} J_\times}{c_*} \nu(E_{j,\delta,R})^{1/4} \\ &\le \frac{C_{\mathrm{int}} J_\times}{c_* R_-^{1/4}} \delta^{1/4}.\end{aligned}$$

Here $Z_m = I_m/|\nabla_q F|$ is the entire original singular multiplier, and invariance together with (B.3) gives $\nu(E_{j,\delta,R}) \le \int N_\delta \, d\nu = \delta/R$. This calculation intersects actual observables on the same orbit; it uses neither independence nor a positive incidence margin. The factor c_*^{-1} enters once, outside the unconditioned localized multiplier estimate.

We give the endpoint-to-height step explicitly, since a small bulk mass alone does not prove it. Fix m, R. In the complete analytic physical endpoint charts of Lemma 67.3, add the finitely many signed impact and interior-foot conditions defining $E_{j,\delta,R}$. For almost every Gauss angle these give ordinary open selected subintervals of the original physical roof intervals. The same finite Boolean analytic description, compact chart cover and one-dimensional slicing argument used there applies. Equalities have the explicit weak-boundary traces already counted above;

identically satisfied constraints are removed from the local Boolean description. Artificial seams and homogeneous cuts cancel before absolute values are taken. Distinct actual word sheets and all genuine physical intervals remain in the sum.

The possible interval endpoints are thus visited grazing, a critical point, an original unvisited tangency, a new clearance-δ level, or a cap side. At a visited grazing the actual optical factor ω tends to zero while $\chi \geq \lambda_\circ$; hence $A = \omega/(4\pi c_* \chi)$ has zero physical coarea trace, including a simultaneous critical limit. This is the actual optical trace assertion of Lemma 67.3, not an inference from the zero collision-area measure of the singular set.

Every remaining critical endpoint has positive visited incidences. It is an incoming endpoint for the roof. Indeed put $g = re$ on its fixed Gauss ray. The positive reduced Hessian gives

$$\frac{dq}{dr} = G^{-1}e, \qquad \frac{dF}{dr} = re^{\mathsf{T}}G^{-1}e > 0 \quad (r > 0).$$

At $r = 0$ the roof is its lower one-sided limit. Such a critical point can therefore be a left endpoint of a selected interval, or support no interval at all; it cannot be its right endpoint. All positive incoming critical fan values are retained in this statement. No bound sets those values to zero, including a critical point on an active clearance boundary. Local charts and these conclusions are used at fixed m, R before nonnegative exhaustion; no uniform Taylor radius is presumed.

On each selected fixed-angle interval (a_q, b_q), use the actual unweighted smooth density and its physical right trace. Absolute continuity on compact subintervals, followed by the proved finite interior and outgoing boundary budgets, yields

$$A_q(t) \leq A_q(b_q^-) + \int_t^{b_q} |\partial_u A_q(u)|\, du, \qquad a_q < t < b_q. \tag{72.11}$$

The whole physical roof lies in $[m\tau_-, m\tau_+]$, so there is no unaccounted endpoint at infinity. The right trace is either zero visited grazing or a regular charge covered by Lemmas 72.2 and 72.3; a remaining critical point was just proved to be incoming. For finitely many compact regular intervals sum (72.11); integrate the Gauss angle and then exhaust all actual intervals. Tonelli and nonnegative convergence apply to both upper budgets. Their total bounds prove

$$\|p_{m,R;j,\delta}\|_\infty \leq \frac{C_{\mathrm{int}} J_\times}{c_* R_-^{1/4}} \delta^{1/4} + C_{\mathrm{side}}(\delta + \theta_{\mathrm{r}}^j + \theta_{\mathrm{r}}^{m-1-j}) + C_\delta(\theta_{\mathrm{r}}^j + \theta_{\mathrm{r}}^{m-1-j}) + C_{\mathrm{cap,loc}}\delta.$$

Since $0 < \delta \leq \delta_* < 1$, this implies (72.9). An arbitrary original Borel selector is restored only after this positive bound and cannot increase its density. No variation of an indicator-cut density is asserted. For each radius the conclusion is an essential supremum; uniform constants then permit a supremum over radii. A single exceptional null set common to uncountably many radii is not required. □

The exponent $1/4$ comes from the actual localized singular multiplier, whereas the two geometric tails come from the number of real collisions between the marked flight and either endpoint. The result controls the selected density at a common physical roof. It does not yet pay the endpoint layers after multiplication by m^2, and it does not assert abundance of the original nongrazing comparison pool.

73. Deep clearance at the original normalization and the same comparison pool

We apply the physical selected roof height, rather than an integrated clearance mass, to the original persistent positive source. The original radius family, collision map, section and return tests, unit-speed clock, all joint labels, and conditioning are retained. At the outer stage fix B and all its original parameters $\varepsilon(B), h_B, r_B, \gamma_B, K_B$; in particular $\gamma_B \geq 1$. No parameter is chosen from a spectral estimate at a collision-dependent window.

Let

$$q_0 = 47/53, \qquad d'_j = \min(j, m-1-j), \qquad \delta_{j,B} = 2\varepsilon(B) q_0^{d'_j/4}, \qquad 0 \le j < m. \tag{73.1}$$

These are exactly the clearance thresholds of the original guard. For an integer $D \ge 0$ define the physical union

$$\mathcal{E}^{\mathrm{clr}}_{B,m}(D) = \bigcup_{\substack{0 \le j < m \\ d'_j \ge D}} \{x : N_{\delta_{j,B},R}(T_R^j x) \ge 1\}. \tag{73.2}$$

If the index range is empty, the union is empty. At a regular target point this is the event that at least one actual original clearance witness occurs in the indicated interior flight range. Different witnesses may occur at different roofs of the same paired segment; no common witness on the whole segment is required.

Theorem 73.1 (Complete deep-clearance height and its collision payment)**.** *For every finite m, B, D satisfying $2\varepsilon(B) q_0^{D/4} \le \delta_*$, every original exact event $\Pi_{n,k,m,R}$, and every original Borel selector $0 \le H \le 1$, the pushforward*

$$(L_{m,R})_*(\mathbf{1}_{\Pi_{n,k,m,R}} H \mathbf{1}_{\mathcal{E}^{\mathrm{clr}}_{B,m}(D)}\, d\nu_R^*)$$

has a density of essential height at most

$$Q_{\varepsilon(B)}(D) = 2C_{\mathrm{clr,roof}} \left\{ \frac{(2\varepsilon(B))^{1/4} q_0^{D/16}}{1 - q_0^{1/16}} + \frac{\theta_{\mathrm{r}}^D}{1 - \theta_{\mathrm{r}}} \right\}. \tag{73.3}$$

The same bound holds for the unchanged persistent source restricted at the target roof to $\mathcal{E}^{\mathrm{clr}}_{B,m}(D)$, and for any positive subsource of that persistent source.

Fix $\eta > 0$, and put

$$q_{\mathrm{c}} = \max\{q_0^{1/16}, \theta_{\mathrm{r}}\} < 1, \qquad D_m = \left\lceil \frac{(2+\eta)\log(m+1)}{|\log q_{\mathrm{c}}|} \right\rceil. \tag{73.4}$$

For every fixed outer B, the size condition preceding (73.3) *holds at all sufficiently large m, and*

$$\sup_{R,n,k} m^2 \| \mathsf{U}^{\mathrm{per}}_{B,m}[\mathcal{E}^{\mathrm{clr}}_{B,m}(D_m)] \|_\infty \le A_B (m+1)^{-\eta}, \tag{73.5}$$

where

$$A_B = 2C_{\mathrm{clr,roof}} \left\{ \frac{(2\varepsilon(B))^{1/4}}{1 - q_0^{1/16}} + \frac{1}{1 - \theta_{\mathrm{r}}} \right\}. \tag{73.6}$$

All norms are essential suprema on the complete original roof line. The asserted inner suprema follow after summing all physical words, all marks, both impact signs and every relevant flight index.

Proof. By positivity, the selected union measure is at most the sum of the single-flight measures in (72.1). Every width in the index range is at most $2\varepsilon(B) q_0^{D/4} \le \delta_*$, so Theorem 72.4 applies to each term. There are at most two flight indices of each nonnegative depth. Thus

$$\sum_{d'_j \ge D} \delta_{j,B}^{1/4} \le \frac{2(2\varepsilon(B))^{1/4} q_0^{D/16}}{1 - q_0^{1/16}}.$$

Also the two endpoint tails are separate subseries:

$$\sum_{D \le j \le m-1-D} (\theta_{\mathrm{r}}^j + \theta_{\mathrm{r}}^{m-1-j}) \le \frac{2\theta_{\mathrm{r}}^D}{1 - \theta_{\mathrm{r}}}.$$

These exact finite-index bounds prove (73.3). Double witnesses enlarge the positive upper sum but are counted once in the union itself. The original section and full joint label are restored through the positive domination (67.2); no new conditioning factor is introduced.

For the persistent source use only the elementary coefficient inequality (12.1):

$$0 \le c_{\mathrm{per}}^{\varepsilon,\gamma} \le f^{\varepsilon}(t,\theta) \le A(t,\theta).$$

Multiply this inequality by the unchanged target selector and $\mathbf{1}_{\mathcal{E}_{B,m}^{\mathrm{clr}}(D)}$, then perform the existing one-copy roof–angle integration. This proves domination by the same physical selected majorant at the target roof. A minimum at unrelated segment endpoints is not estimated by a small bulk set. This use of the elementary coefficient inequality does not invoke the whole-word incidence pressure theorem or assume a uniform comparison-capacity abundance statement.

The definition (73.4) gives

$$q_0^{D_m/16} \le q_{\mathrm{c}}^{D_m} \le (m+1)^{-(2+\eta)}, \qquad \theta_{\mathrm{r}}^{D_m} \le (m+1)^{-(2+\eta)}.$$

In particular $2\varepsilon(B) q_0^{D_m/4} \le 2\varepsilon(B)(m+1)^{-4(2+\eta)}$. At each fixed B this is eventually at most δ_*, without any restriction on its original positive width beyond finiteness. Insert these inequalities into (73.3) and use $m^2 \le (m+1)^2$ to obtain (73.5). The finitely many earlier collision counts do not affect a limsup.

The selected depth is only a substitution in a finite-count bound valid for every D. It is not a shrinking physical roof, section, residence or comparison window. Moreover $D_m = O(\log m) = o(m)$, so the interior range contains $m - O(\log m)$ actual flights and is nonempty for every sufficiently large count. The statement is therefore not an empty-event payment. At fixed B its rate is uniform in the original radius and exact labels, since all constants were fixed before these suprema. This proves the theorem in the prescribed order. □

73.1. **An exact shortage equivalence with the original full pool.** Keep the existing source $\mathsf{U}_{B,m}^{\mathrm{low}}$ in (15.2), with precisely its old incidence-load, section and pairing selectors. Define two positive restrictions, with no renormalization, by

$$\begin{aligned} \mathsf{U}_{B,m}^{\mathrm{low,deep}} &= \mathsf{U}_{B,m}^{\mathrm{low}}[\mathcal{E}_{B,m}^{\mathrm{clr}}(D_m)], \\ \mathsf{U}_{B,m}^{\mathrm{low,end}} &= \mathsf{U}_{B,m}^{\mathrm{low}}[\mathcal{E}_{B,m}^{\mathrm{clr}}(D_m)^c]. \end{aligned} \tag{73.7}$$

The notation end describes the location of possible clearance witnesses on this complement: every such witness lies in one of the two layers $d_j' < D_m$. It makes no assertion that this remaining positive source is small or has enough comparison capacity. Define

$$\mathsf{R}_{B,m}^{\mathrm{end}} = (\mathsf{U}_{B,m}^{\mathrm{low,end}} - K_B P_{r_B,m}^{\mathrm{ng}})_+. \tag{73.8}$$

The comparison pool $P_{r_B,m}^{\mathrm{ng}}$ is the same full original nongrazing pool as in (15.2). It is not restricted to the new source class, and its multiplier and its microscopic width are unchanged.

Theorem 73.2 (Original-height equivalence of the two shortages)**.** *At every finite count there is the exact positive identity and bound*

$$\mathsf{U}_{B,m}^{\mathrm{low}} = \mathsf{U}_{B,m}^{\mathrm{low,deep}} + \mathsf{U}_{B,m}^{\mathrm{low,end}}, \tag{73.9}$$

$$0 \le \mathsf{R}_{B,m}^{\mathrm{low}} - \mathsf{R}_{B,m}^{\mathrm{end}} \le \mathsf{U}_{B,m}^{\mathrm{low,deep}}. \tag{73.10}$$

For every fixed B, M, the difference has zero original ordered height:

$$\limsup_{m\to\infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} (\mathsf{R}_{B,m}^{\mathrm{low}}(t) - \mathsf{R}_{B,m}^{\mathrm{end}}(t)) = 0. \tag{73.11}$$

Consequently the old shortage obligation (15.8) *is equivalent, with exactly the same order of limits, to*

$$\lim_{B\to\infty}\limsup_{m\to\infty}\sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} (\mathsf{U}^{\mathrm{low,end}}_{B,m}(t) - K_B P^{\mathrm{ng}}_{r_B,m}(t))_+ = 0 \qquad \textit{for every fixed } M. \tag{73.12}$$

This equivalence itself uses the new physical height theorem and the elementary persistent domination, without invoking the inherited pressure theorem or assuming comparison-pool abundance.

Proof. The event in (73.2) and its complement partition the same original target source. This proves (73.9) before and after pushforward. For $u, v, c \geq 0$,

$$0 \leq (u + v - c)_+ - (u - c)_+ \leq v.$$

Take $u = \mathsf{U}^{\mathrm{low,end}}$, $v = \mathsf{U}^{\mathrm{low,deep}}$ and $c = K_B P^{\mathrm{ng}}_{r_B}$. The same value of c is essential for this inequality to compare the specified shortages. It proves (73.10) pointwise at almost every roof. The low-load deep subsource is a positive subsource of the full persistent deep source. Theorem 73.1 therefore bounds its complete-roof normalized height by $A_B(m+1)^{-\eta}$ for all sufficiently large m. Restriction to the original central set only decreases the essential supremum. This proves (73.11).

At every fixed B the two nonnegative normalized shortage heights thus have the same collision limsup, including when that value is infinite: each is bounded above by the other plus a nonnegative sequence tending uniformly to zero. Sending B to infinity last proves the exact equivalence of the two displayed obligations. The original $\varepsilon(B), h_B, r_B, \gamma_B, K_B$ stay fixed through the collision limsup, and all label and radius suprema remain inside it. No roof supremum is interchanged with a limit or replaced by an averaged norm. □

Proved payment and the remaining minimal closure condition. The new conclusion pays a complete class of actual clearance witnesses in the original m^2L^∞ roof topology. Its estimate covers every physical word and every mark in the middle $m - O(\log m)$ flights, including zero-incidence-load source, and retains its original persistent coefficient and all conditioning factors. The logarithmic layers are the result of a proved uniform geometric tail and singular multiplier bound; the mass outside them was not declared negligible by a source partition alone.

Equation (73.12) is still a substantive mathematical obligation. The present proof supplies no same-roof lower bound for $P^{\mathrm{ng}}_{r_B}$ on the remaining positive source. A guard witnessed only by shallow clearance can still have zero incidence load and a positive coefficient. Shallow incidences, nongrazing critical entrances, original section and pairing cuts, and short or capacity-deficient segments are not removed by the new decomposition. Establishing (73.12) requires a physical comparison or decay mechanism strong enough for its full suprema and original ordered limits; a fixed unscaled endpoint-layer bound does not suffice.

With the inherited analytic scalar input and the retained native incidence/load payment theorems in their stated scopes, the implication (15.6) makes (73.12) the remaining positive-source condition for the unchanged scalar raw pointwise assertion. The complete native pressure theorem of Section 70, with its full proof and original scope, remains part of the author proof chain. The new clearance payment does not require invoking it and does not alter its mathematical status. Independent expert reconstruction of that chain remains an evidence task, distinct from the unresolved mathematical closing conditions. The full joint spectral and arithmetic estimates, microscopic conditional denominator, return/path transfer and four-coordinate local inverse retain their existing separate dependencies and scope. No new unconditional raw local limit or completed inverse theorem is asserted by this section.

74. One-copy variation of the complete exact-label vector

All collision states, section rectangles, labels and physical roofs in this section are those of Section 69. Fix $m \geq 1$ and $R \in [R_-, R_+]$. The initial section is counted in $A_{m,R}$ and the terminal membership is tested separately, exactly as in (1.1). Write $\ell = (n, k) \in \mathbb{Z} \times \mathbb{Z}^2$; components outside the actual label range are zero. For $w = 1$, or for the original $w = W^\varepsilon$ with $\varepsilon > 0$ fixed, retain $p^w_{\ell,m,R} = p^w_{n,k,m,R}$ from (69.3) and put

$$D^w_{\mathrm{all}} = D^w_{\mathrm{base}} + B_{\mathrm{sec,end}} + 2B_{\mathrm{sec,int}}. \tag{74.1}$$

Every constant on the right has already been defined on the original physical family in (69.1) and (69.4). In particular there is only the original conditioning factor c_*^{-1}. The complete physical word sum is part of those constants.

Theorem 74.1 (Summed variation over every exact physical label)**.** *For every R, m and either specified weight,*

$$\sum_\ell \|Dp^w_{\ell,m,R}\|_{\mathrm{TV}} \leq D^w_{\mathrm{all}}, \tag{74.2}$$

$$\sum_\ell \|p^w_{\ell,m,R}\|_1 \leq 1, \qquad \sum_\ell \|p^w_{\ell,m,R}\|_\infty \leq \tfrac{1}{2} D^w_{\mathrm{all}}, \tag{74.3}$$

$$\sum_\ell \|p^w_{\ell,m,R}(\cdot + h) - p^w_{\ell,m,R}\|_1 \leq D^w_{\mathrm{all}}|h| \qquad (h \in \mathbb{R}). \tag{74.4}$$

Every component has support in the actual interval $[m\tau_-, m\tau_+]$. The sums precede the suprema over the radius and collision count. No factor equal to the number of occupation or lattice labels is present.

Proof. For clarity, use the unit vectors e_ℓ of the label space with its ℓ^1 norm. On a regular physical word define

$$v(x) = \eta_R(x)\eta_R(T_R^m x) e_{(A_{m,R}(x), K^{\mathrm{c}}_{m,R}(x))}.$$

This vector has norm zero or one. Cut each maximal fixed-angle physical interval by all the actual preimages of the original finite section sides. The analytic slicing and exhaustion in Lemma 69.1 give complementary regular chambers on which v is constant. The physical displacement is fixed on a word. Artificial chart seams with the same vector on both sides cancel before absolute values.

Let $\mathscr{A}$ be the physical coarea coefficient in Lemma 69.1. Inside a chamber, the derivative of $w\mathscr{A}v$ has ℓ^1 norm at most the absolute derivative of $w\mathscr{A}$. Thus the complete interior charge is bounded by the interior part of D^w_{base}. At every genuine physical endpoint, the vector has one-sided norm at most one. The cap and unvisited-disk charges, the incoming positive critical sectors and the proved zero visited-grazing traces are exactly those in Lemma 67.3. Multiplication by this vector does not increase any positive one-sided charge. Charges on different physical words are kept in their already complete positive sum; no cancellation between different words is assumed. The boundary part of D^w_{base} therefore pays all those endpoints.

At a regular interior visit crossing, the vector changes from one unit vector to another if both endpoint tests hold, and remains zero otherwise. Its ℓ^1 jump is at most two. At the initial or terminal section boundary it changes between zero and a single unit vector, at cost at most one. A change in the initial occupation is included in that initial transition; no visit at index m is added. For coincident visit boundaries, order the affected visits and telescope the vector changes. The total is bounded by two for each changed interior visit and one for each changed endpoint test. All these sides, with their original finite side intervals, are already included in (69.1). Their complete charge is at most $B_{\mathrm{sec,end}} + 2B_{\mathrm{sec,int}}$.

One can express this calculation without any vector-valued differentiation theorem. Choose any finite collection of labels and compactly supported C^1 scalar tests ψ_ℓ, each with $\|\psi_\ell\|_\infty \le 1$. Integrate $\sum_\ell p^w_{\ell,m,R}\psi'_\ell$ by parts first on a finite regular chamber exhaustion. Each interior, physical endpoint and section jump is bounded by precisely the positive charges above. The source against the test derivatives is integrable. The finite total positive charge permits passage to the exhaustion, just as in Theorem 69.2. Taking independent tests for each component gives the sum of its distributional variation norms. Increasing the finite label set proves (74.2). Equivalently, at fixed count only finitely many labels are attainable by the finite physical horizon, but the bound does not use their number.

The exact label events partition the original source with both endpoint memberships. Positivity and $0 \le w \le 1$ give

$$\sum_\ell \int p^w_{\ell,m,R}(t)\,dt = \int w\eta_R(T_R^m x)\,d\nu^*_R(x) \le 1.$$

Each nonnegative compactly supported BV component starts and ends at zero. Reaching a value a at a Lebesgue point and returning to zero costs variation at least $2a$. Consequently $2\|p^w_{\ell,m,R}\|_\infty \le \|Dp^w_{\ell,m,R}\|_{\mathrm{TV}}$. Summing proves the height estimate. Finally, the BV representative satisfies

$$|p(t+h)-p(t)| \le |Dp|([\min(t,t+h),\max(t,t+h)])$$

outside irrelevant endpoint conventions. Tonelli integration in t, followed by the positive label sum, proves (74.4). The support follows from the true flight bounds. Null sets are understood at the fixed radius, as in the original target. □

Corollary 74.2 (Bounded tests of the exact labels)**.** *For any complex numbers a_ℓ with $|a_\ell| \le 1$, the density $\sum_\ell a_\ell p^w_{\ell,m,R}$ has L^1 norm at most one and variation at most D^w_{all}. Its translations have L^1 error at most $D^w_{\mathrm{all}}|h|$, and its roof Fourier transform is bounded by $\min\{1, D^w_{\mathrm{all}}/|s|\}$ for $s \ne 0$.*

Proof. The summed L^1 and derivative norms give convergence and the asserted bounds, first on finite label sums and then by their absolute norm convergence. Integrating the finite derivative measure by parts on the compact physical roof support gives the Fourier assertion. The translation assertion follows from (74.4). □

Information retained. This result adds a summed label budget to the separate bounds in Theorem 69.2; it does not sum those separate bounds by counting labels. The coefficients in Corollary 74.2 are independent of the roof and trajectory within one label. An arbitrary Borel trajectory selector has no asserted BV derivative. Such selectors will instead be handled by a positive trajectory-level error in the next section. All physical boundaries, the original guard when present, and all original conditional and arithmetic conventions remain unchanged.

75. The actual return tower and positive finite-band likelihoods

We retain the probability $\nu^*_R = \nu|_{Y^*_R}/c_*$ and the unit-speed physical clock. All maps in this section act on the full-measure invariant set of trajectories whose forward and backward collisions are regular. This removes no positive collision mass and assigns no value to a separate singular boundary trace.

Lemma 75.1 (The conservative physical return tower)**.** *Let $r_R(x) = \min\{j \ge 1 : T^j_R x \in Y^*_R\}$ be the true first return in collisions and $F^*_R x = T_R^{r_R(x)}x$. The return is finite almost everywhere on the original section, F^*_R preserves ν^*_R, and*

$$\int_{Y^*_R} r_R\,d\nu \le 1, \qquad \mathbb{E}_{\nu^*_R} N_{n,R} \le \frac{n}{c_*}, \qquad \nu^*_R\{N_{n,R} > L\} \le \frac{n}{c_* L} \quad (L>0), \tag{75.1}$$

where

$$N_{n,R} = \sum_{a=0}^{n-1} r_R \circ (F_R^*)^a$$

is the actual collision count of the nth return. Its physical time $S_{n,R} = L_{N_{n,R},R}$ *satisfies* $\mathbb{E}_{\nu_R^*} S_{n,R} \le \tau_+ n/c_*$.

Proof. Suppress R and write $Y = Y_R^*$. The actual invertible collision map preserves ν. Let $E \subset Y$ consist of points with no future visit to Y. The sets $T^j E$, $j \ge 0$, are disjoint: an intersection at times $i < j$ would give a future visit to Y at time $j - i$ from a point of E. Their equal measures have sum at most one, so $\nu(E) = 0$. Points of Y with only finitely many future visits belong to $\bigcup_{j\ge 0} T^{-j}E$, also null. The same argument for T^{-1} gives infinitely many past visits almost everywhere on Y.

Put $Y_a = \{x \in Y : r(x) = a\}$. The sets $T^a Y_a$, $a \ge 1$, partition Y up to a null set: for almost every terminal point, its nearest strictly earlier visit gives the unique starting point and the unique positive return time. Thus for every measurable $B \subset Y$, invariance and disjointness give

$$\nu((F^*)^{-1}B) = \sum_{a\ge 1} \nu(Y_a \cap T^{-a}B) = \sum_{a\ge 1} \nu(T^a Y_a \cap B) = \nu(B).$$

Division by the original c_* proves induced invariance.

The true tower levels $T^j\{x \in Y : r(x) > j\}$, $j \ge 0$, are disjoint. Indeed, an intersection at levels $i < j$ implies that a point with return time exceeding j has a visit to Y at time $j - i \le j$, a contradiction. Consequently

$$\int_Y r\, d\nu = \sum_{j\ge 0} \nu\{x \in Y : r(x) > j\} = \nu\left(\bigsqcup_{j\ge 0} T^j\{x \in Y : r(x) > j\}\right) \le 1.$$

Equality is not needed and is not asserted without saturation of the collision space. Induced invariance gives the expectation of the sum defining N_n and hence the second bound in (75.1). Integrating $\mathbf{1}_{\{N_n > L\}} \le N_n/L$ proves its last bound. Finally $S_n \le \tau_+ N_n$ on each true trajectory. No mixing, independence, spectral estimate or ergodicity assertion enters this argument. □

Let $K_{n,R} = K^{\mathrm{c}}_{N_{n,R},R}$ be the actual two-dimensional lattice displacement at that return. The full probability law is

$$\mu_{n,R} = \mathrm{Law}_{\nu_R^*}(K_{n,R}, N_{n,R}, S_{n,R}) \quad \text{on } \mathbb{Z}^2 \times \mathbb{N} \times \mathbb{R}. \tag{75.2}$$

Fixing n in this law retains the two displacement coordinates, true collision count and physical time. Fixing m instead gives the original coordinates (k_1, k_2, t, n) in (1.5). These are two slices of the same joint records, not different conditioning events or a replacement inverse problem.

Lemma 75.2 (Exact components and a positive proof restriction)**.** *The* (k, m) *component of* (75.2) *is exactly* $p_{n,R}(k, m, t)\, dt$. *For an integer* $L \ge 1$ *put*

$$Q_{n,R}^L = \mathbf{1}_{\{m\le L\}}\mu_{n,R}, \qquad H_{n,R}^L = \mu_{n,R} - Q_{n,R}^L, \qquad \delta_{n,R}(L) = \|H_{n,R}^L\|_{\mathrm{TV}}.$$

Both are positive measures, and

$$\delta_{n,R}(L) \le \frac{n}{c_* L}, \qquad \sum_{m\le L}\sum_{k} \|Dp_{n,R}(k, m, \cdot)\|_{\mathrm{TV}} \le D^1_{\mathrm{all}} L. \tag{75.3}$$

Neither measure is renormalized. This restriction is a device of the proof and is removed from the conclusions about $\mu_{n,R}$.

Proof. Starting in Y_R^*, the event $N_{n,R} = m$ says that the time-m state is in that section and precisely n visits, including the initial one, occur at indices $0, \dots, m-1$. It is therefore exactly the occupation and endpoint event in (1.1). Adding the true displacement label gives its stated density. The tail bound is (75.1). At each m, fixing one occupation label in Theorem 74.1 decreases its positive variation sum. Summing over at most L actual collision counts proves (75.3). Almost-sure successive recurrence from Lemma 75.1 shows that all components together have mass one. □

75.1. **A positive kernel with an exact compact frequency support.**

Lemma 75.3 (The kernel and its first absolute moment)**.** *With its continuous value at zero, put*

$$k(t) = \frac{3}{8\pi}\left(\frac{\sin(t/4)}{t/4}\right)^4, \qquad k_{\mathfrak{b}}(t) = \mathfrak{b}k(\mathfrak{b}t) \quad (\mathfrak{b} > 0). \tag{75.4}$$

Then $k \geq 0$, $\int k = 1$, $\int t^2 k(t)\,dt = 12$, *and* $\kappa_{\mathrm{ker}} := \int |t| k(t)\,dt \leq \sqrt{12}$. *For the convention* $\widehat{f}(s) = \int e^{ist} f(t)\,dt$,

$$\widehat{k}(s) = \eta(s) = \begin{cases} 1 - 6s^2 + 6|s|^3, & |s| \leq 1/2, \\ 2(1-|s|)^3, & 1/2 \leq |s| \leq 1, \\ 0, & |s| \geq 1. \end{cases} \tag{75.5}$$

The bandwidth $\mathfrak{b}$ *is independent of the outer geometric parameter* B *in the original shortage limit.*

Proof. The displayed η is compactly supported and C^2. Its third derivative has jumps $12, -48, 72, -48, 12$ at respectively $-1, -1/2, 0, 1/2, 1$. Four integrations by parts, or the same distributional identity tested against e^{-its}, give for $t \neq 0$

$$t^4 \int \eta(s) e^{-its}\,ds = 24\cos t - 96\cos(t/2) + 72 = 192\sin^4(t/4).$$

Thus $(2\pi)^{-1}\int \eta(s)e^{-its}\,ds$ equals the k in (75.4); at zero both equal $3/(8\pi)$ since $\int \eta = 3/4$. This also proves nonnegativity, and the displayed formula gives an integrable $O(|t|^{-4})$ tail. Fourier inversion for this integrable transform pair gives $\int k = \eta(0) = 1$. Since $t^2 k$ is integrable, differentiation of its transform is justified by domination, and $\int t^2 k(t)\,dt = -\eta''(0) = 12$. Cauchy–Schwarz against the probability density k gives the asserted first absolute moment. Scaling gives transform $\eta(s/\mathfrak{b})$ and first absolute moment $\kappa_{\mathrm{ker}}/\mathfrak{b}$ for $k_{\mathfrak{b}}$. □

Convolution below acts only on the real roof coordinate and preserves every discrete label. The norm $\|\cdot\|_{\mathrm{TV}}$ of a signed measure is its full variation, without a factor one half.

Theorem 75.4 (Positive likelihoods on the original trajectories)**.** *Fix* $n \geq 1$, $L \geq 1$, $\mathfrak{b} > 0$, $R \in [R_-, R_+]$ *and an integer* $J \geq 1$. *For every label* $\ell = (k, m)$ *let* I_ℓ *be empty or a union of at most* J *bounded physical-roof intervals. These intervals and labels specify the unchanged event*

$$A = \{x : S_{n,R}(x) \in I_{(K_{n,R}(x), N_{n,R}(x))}\}.$$

On the very same trajectories define the analytical likelihood

$$\lambda_{A,\mathfrak{b}}(x) = (k_{\mathfrak{b}} * \mathbf{1}_{I_{(K_{n,R}(x), N_{n,R}(x))}})(S_{n,R}(x)). \tag{75.6}$$

Then $0 \le \lambda_{A,\mathfrak{b}} \le 1$. *If* $\mathcal{V}_L = D^1_{\mathrm{all}}L$ *and* $\delta = \delta_{n,R}(L)$, *then*

$$\|\mu_{n,R} - k_{\mathfrak{b}} * \mu_{n,R}\|_{\mathrm{TV}} \le 2\delta + \frac{\kappa_{\mathrm{ker}}\mathcal{V}_L}{\mathfrak{b}}, \tag{75.7}$$

$$\mathbb{E}_{\nu^*_R}|\mathbf{1}_A - \lambda_{A,\mathfrak{b}}| \le \delta + \frac{\kappa_{\mathrm{ker}}J\mathcal{V}_L}{\mathfrak{b}} =: \mathcal{E}_n(L,\mathfrak{b},J), \tag{75.8}$$

$$\left|\mathbb{E}_{\nu^*_R}V(\mathbf{1}_A - \lambda_{A,\mathfrak{b}})\right| \le \|V\|_\infty \mathcal{E}_n(L,\mathfrak{b},J) \tag{75.9}$$

for every bounded Borel observation V *of the original trajectories. All actual collision counts remain in* $\mu_{n,R}$, A *and* $\lambda_{A,\mathfrak{b}}$. *No positive lower bound on any interval length, incidence or clearance is required.*

Proof. For the positive proof restriction Q^L, integrate the all-component translation estimate in (75.3) against $k_{\mathfrak{b}}$. Tonelli and the triangle inequality give

$$\|Q^L - k_{\mathfrak{b}} * Q^L\|_{\mathrm{TV}} \le \mathcal{V}_L \int |h| k_{\mathfrak{b}}(h)\, dh = \kappa_{\mathrm{ker}}\mathcal{V}_L/\mathfrak{b}.$$

The positive measures H^L and $k_{\mathfrak{b}} * H^L$ both have mass δ; their difference has variation at most 2δ. This proves (75.7) for the full law.

For a union I of at most J intervals, translation of its indicator satisfies

$$|I \triangle (I+h)| \le 2J|h|.$$

It follows first for one interval by its two endpoints and then for a finite union by subadditivity; it remains true for overlapping interval presentations. For a nonnegative BV component density p, its compact physical support and Theorem 74.1 give $\|p\|_\infty \le \|Dp\|_{\mathrm{TV}}/2$. Therefore

$$\int p(t)|\mathbf{1}_I(t) - \mathbf{1}_I(t-h)|\, dt \le J|h|\|Dp\|_{\mathrm{TV}}.$$

Positivity of the kernel permits integration of this inequality in h and summation over every component of Q^L. The resulting bound is $\kappa_{\mathrm{ker}}J\mathcal{V}_L/\mathfrak{b}$. On the removed positive tail, both the indicator and likelihood belong to $[0,1]$, so the absolute difference costs at most its mass δ. This proves (75.8) on all the original trajectories. Multiplication by $|V| \le \|V\|_\infty$ proves (75.9). It uses no regularity or independence of the observation and no BV assertion for V. Every exchange in this proof is of nonnegative integrals or an absolutely bounded integral at the stated finite parameters. □

75.2. **An exact four-coordinate frequency formula.** Put

$$\widehat{\mu}_{n,R}(\xi,\upsilon,s) = \mathbb{E}_{\nu^*_R} e^{i(\xi\cdot K_{n,R} + \upsilon N_{n,R} + sS_{n,R})}.$$

For one exact label (k,m) and one bounded interval union I, put

$$\lambda_{k,m,I,\mathfrak{b}} = \mathbf{1}_{\{K_{n,R}=k, N_{n,R}=m\}}(k_{\mathfrak{b}} * \mathbf{1}_I)(S_{n,R}).$$

Corollary 75.5 (The full finite-band transform uses the true law)**.** *The exact identity is*

$$\mathbb{E}_{\nu^*_R}\lambda_{k,m,I,\mathfrak{b}} = (2\pi)^{-4}\int_{[-\pi,\pi]^3}\int_{-\mathfrak{b}}^{\mathfrak{b}} e^{-i(\xi\cdot k + \upsilon m)}\eta(s/\mathfrak{b})\widehat{\mathbf{1}_I}(-s)\widehat{\mu}_{n,R}(\xi,\upsilon,s)\, ds\, d\xi_1\, d\xi_2\, d\upsilon. \tag{75.10}$$

No torus frequency or arithmetic class is omitted. A general event in Theorem 75.4 is the positive sum of these one-label likelihoods. No interchange of an infinite complex label series with the displayed frequency integral is required.

Proof. Character orthogonality on the full three-dimensional torus gives the exact label indicator. Since η is even,

$$(k_{\mathfrak{b}} * \mathbf{1}_I)(t) = (2\pi)^{-1} \int_{-\mathfrak{b}}^{\mathfrak{b}} e^{ist} \eta(s/\mathfrak{b}) \widehat{\mathbf{1}_I}(-s)\, ds.$$

Multiply the two identities and integrate on the actual probability space. The frequency domain is bounded and $|\widehat{\mathbf{1}_I}(s)| \le |I|$, so Fubini is justified without a collision cutoff. This proves (75.10) with its factor $(2\pi)^{-4}$. The last assertion follows by monotone summation of nonnegative trajectory likelihoods. A discrete label having zero physical mass contributes zero. This does not assert that smoothing has zero mass on every zero-mass roof window within an attainable discrete class. □

Corollary 75.6 (Arbitrary polynomial normalized accuracy)**.** *Fix $P > 0$, $\kappa \ge 0$ and $J_0 \ge 1$. If $J_n \le J_0 n^{\kappa}$, choose*

$$L_n = \lceil n^{P+3} \rceil, \qquad \mathfrak{b}_n = n^{2P+\kappa+5}. \tag{75.11}$$

Uniformly in the original radius, all labels, all interval events as above and all $\|V\|_\infty \le 1$, the errors in (75.7)–(75.9), multiplied by n^2, are $O(n^{-P})$. The constants depend only on the displayed fixed parameters and the original physical family.

Proof. For $n \ge 1$, $n^{P+3} \le L_n \le 2n^{P+3}$. Consequently

$$n^2 \delta_{n,R}(L_n) \le c_*^{-1} n^{-P}, \qquad \frac{n^2 \kappa_{\mathrm{ker}} J_n D^1_{\mathrm{all}} L_n}{\mathfrak{b}_n} \le 2\kappa_{\mathrm{ker}} J_0 D^1_{\mathrm{all}} n^{-P}.$$

For the law error omit J_n and use $n^{-\kappa} \le 1$. These are substitutions into inequalities already valid for every finite L and every positive bandwidth. They exchange no geometric outer limit and impose no lower microscopic roof width. □

Corollary 75.7 (Fixed collision count and the original m^2 scale)**.** *At any fixed true collision count m, consider the entire positive exact-label subprobability with components $p^w_{n,k,m,R}(t)\, dt$, where $w = 1$ or the original fixed W^ε. The conclusions (75.7)–(75.9) hold with $\delta = 0$ and $\mathcal{V}_L = D^w_{\mathrm{all}}$, using the original physical source weight $w\eta_R(T_R^m x) d\nu_R^*$. If $J_m \le J_0 m^{\kappa}$ and $\mathfrak{b}_m = m^{P+\kappa+2}$, the absolute event and observation errors multiplied by m^2 are $O(m^{-P})$. There is no collision restriction or renormalization in this assertion.*

Proof. Apply Theorem 74.1 directly to the full fixed-count label vector and repeat the positive kernel argument. The source is a subprobability and every test remains on that source; its mass need not be replaced by one in any equality. Multiplying its event bound by m^2 gives $\kappa_{\mathrm{ker}} J_0 D^w_{\mathrm{all}} m^{-P}$. □

Corollary 75.8 (The actual conditional denominator)**.** *Let H be any Borel information event on the original trajectories, $a = \mathbf{1}_H \mathbf{1}_A$, $b = \mathbf{1}_H \lambda_{A,\mathfrak{b}}$, and $p = \mathbb{E}a$, $p_{\mathfrak{b}} = \mathbb{E}b$. If the applicable absolute error bound is E, then*

$$|p - p_{\mathfrak{b}}| \le E.$$

If the actual selected mass satisfies $p > E$, both normalized measures exist and their full variation distance obeys

$$\left\| \frac{a\, d\nu_R^*}{p} - \frac{b\, d\nu_R^*}{p_{\mathfrak{b}}} \right\|_{\mathrm{TV}} \le \frac{2E}{p}. \tag{75.12}$$

If instead $p_{\mathfrak{b}} > E$ is verified, the bound $2E/(p_{\mathfrak{b}} - E)$ is valid. The same statements hold with the fixed-collision source of Corollary 75.7.

Proof. The positive trajectory estimate gives $\mathbb{E}|a-b| \le E$, hence the mass bound. If $p > E$, then $p_{\mathfrak{b}} \ge p - E > 0$. Add and subtract $b\, d\nu_R^*/p$. The first variation cost is at most E/p; the second is $p_{\mathfrak{b}}|p^{-1} - p_{\mathfrak{b}}^{-1}| = |p - p_{\mathfrak{b}}|/p \le E/p$. This proves (75.12). If $p_{\mathfrak{b}} > E$, use $p \ge p_{\mathfrak{b}} - E > 0$ in the same calculation. The proof requires only the stated positive base measure, so applies also to the fixed-collision case. □

Relation to the original raw target. The interval event, each collision and its unit-speed physical time remain exact throughout. Convolution defines an analytical likelihood on those trajectories; it is not a new physical clock or a changed return condition. The absolute estimates are uniform even when the original roof intervals shrink arbitrarily. Dividing by their lengths, or by a small selected-event probability, requires the corresponding additional estimate; neither is implicit here. In particular each information restriction has its own actual denominator. A lower bound for a larger event cannot be inherited by restriction, and a zero arithmetic event cannot be conditioned upon. These conclusions concern ν_R^*; they do not automatically transfer to a stationary entrance law with an unbounded return-roof density. The full joint spectral evaluation of (75.10), every relevant arithmetic peak, the original endpoint-layer shortage and the four-coordinate raw local inverse remain separate requirements. No averaged limit or finite frequency calculation is being substituted for their proofs.

76. Endpoint-tested marked roof operators on the actual collision chain

We pass from scalar label tests to smooth observations of the physical collision states. The estimate below is a weak operator estimate with specified test norms. Its proof uses the actual two-endpoint stationary action, including the full inverse-gradient moment; a derivative of the expanding forward collision map is never replaced by a bounded derivative. Throughout, $R \in [R_-, R_+]$, $m \ge 1$, and $w = 1$ or the original W^ε with $\varepsilon > 0$ fixed. The clock, section, labels, and endpoint memberships are those of (1.1).

Put $\mathcal{X} = (\mathbb{R}/2\pi\mathbb{Z}) \times [-1,1]$. For a complex C^1 function a on this compact cylinder, including continuous one-sided derivatives at its boundary, set

$$a_0 = \|a\|_\infty, \qquad a_1 = \max\{\|\partial_\alpha a\|_\infty, \|\partial_p a\|_\infty\}, \qquad \|a\|_{C^1} = a_0 + a_1. \tag{76.1}$$

Use the same convention for b. Let $\varphi_0, \dots, \varphi_m$ be real $C^1(\mathcal{X})$ functions and write, on the actual regular m-collision chain,

$$\Psi_m(x) = \sum_{i=0}^{m} \varphi_i(T_R^i x), \qquad l_i = \max\{\|\partial_\alpha \varphi_i\|_\infty, \|\partial_p \varphi_i\|_\infty\}, \qquad \mathcal{L}_m = \sum_{i=0}^{m} l_i(\sigma^i + \sigma^{m-i}). \tag{76.2}$$

These functions are observations of the real orbit. They impose no independence assumption and remove no collisions. Define constants using precisely the inherited physical Hessian and Jacobi bounds:

$$C_{\mathrm{test,end}} = \Lambda_\circ(\lambda_\circ^{-1} + R_-^{-1}), \qquad C_{\mathrm{test,phase}} = \frac{\Lambda_\circ}{\lambda_\circ}(J_* + C_c), \qquad J_{\mathrm{test}} = J_\times/c_*. \tag{76.3}$$

Here J_* and C_c are defined in (29.1) and (29.5), and $J_\times$ in (56.1). The constants contain exactly one original conditioning factor. They have no dependence on m, a positive incidence cutoff, a label, a roof value, or a new observation event.

Lemma 76.1 (Actual Gauss derivatives of the endpoint tests and orbit phase)**.** *On every regular fixed-Gauss-angle segment, let $t = F(q)$, $q = (\alpha_0, \alpha_m)$, $g = \nabla_q F$, $z = |g| > 0$, $e = g/z$ and $H = e^{\mathsf{T}} G^{-1} e$. Then*

$$q_t = \frac{G^{-1}e}{zH}, \qquad z_t = \frac{1}{zH}, \qquad \|q_t\|_\infty \le \frac{\Lambda_\circ}{\lambda_\circ z}. \tag{76.4}$$

With the collision states and momenta of this segment,

$$z\big|\partial_t a(\alpha_0,p_0)\big|\le C_{\mathrm{test,end}}a_1,\qquad z\big|\partial_t b(\alpha_m,p_m)\big|\le C_{\mathrm{test,end}}b_1, \tag{76.5}$$

$$z|\partial_t\Psi_m|\le C_{\mathrm{test,phase}}\mathcal{L}_m. \tag{76.6}$$

In particular, if $l_i\le L$ for every i, then $\mathcal{L}_m\le 2L/(1-\sigma)$, uniformly in the entire collision count. There is no inverse incidence in the interior observable derivatives in this assertion.

Proof. At fixed e, differentiation of $\nabla F(q)=ze$ gives $dq/dz=G^{-1}e$. Differentiation of the physical action therefore gives $dt/dz=g\cdot(dq/dz)=zH$. This proves both identities in (76.4); in particular the factor H must not be omitted from q_t. The Hessian bounds (4.3) imply $\Lambda_\circ^{-1}\le H\le\lambda_\circ^{-1}$ and $\|G^{-1}e\|_\infty\le\lambda_\circ^{-1}$, proving the displayed inequality. By the exact first variation $g=R(-p_0,p_m)$, we have $p_0=-ze_1/R$ and $p_m=ze_2/R$. Consequently each endpoint momentum derivative is at most $\Lambda_\circ/(R_-z)$ in absolute value. The chain rule and the two actual partial-derivative norms in (76.1) give (76.5).

For completeness, the momentum derivative at an interior collision is obtained geometrically, without differentiating a square root of the incidence. Let $u_i=n'(\alpha_i)$ be the unit tangent and $v_i=(x_{i+1}-x_i)/\tau_i$ the unit outgoing flight. For $i<m$, the true tangential momentum is $p_i=v_i\cdot u_i$; at $i=m$ use $p_m=v_{m-1}\cdot u_m$, since specular reflection preserves the tangential component. For an endpoint direction η,

$$|D_qp_i[\eta]|\le\|D_qv_i[\eta]\|_2+|D_q\alpha_i[\eta]|\quad(i<m),$$

with v_{m-1} in the terminal formula. The unweighted Jacobi localization (29.3), the flight direction bound (29.7), and $s_{i+1},s_{i-1}\le\sigma^{-1}s_i$ wherever the indicated indices exist yield

$$|D_qp_i[\eta]|\le C_cs_i\|\eta\|_\infty,\qquad|D_q\alpha_i[\eta]|\le J_*s_i\|\eta\|_\infty\quad(0\le i\le m). \tag{76.7}$$

This proof also includes $m=1$. Every flight denominator is the true length, bounded below by τ_-; no denominator c_i has been introduced. Applying the chain rule to each $\varphi_i(\alpha_i,p_i)$ and then inserting $\eta=q_t$ gives (76.6). Finally

$$\sum_{i=0}^{m}s_i=\frac{2(1-\sigma^{m+1})}{1-\sigma}\le\frac{2}{1-\sigma}$$

proves the uniform phase assertion. Every differentiation stays inside the same actual stationary physical branch. It is not a bound on DT_R^i with initial momentum held fixed. □

For $\ell=(n,k)$ define the complex physical roof measure by

$$f_{\ell,m,R}^{w;a,b,\Psi}(t)\,dt=(L_{m,R})_*\left(w\,\eta_R(x)\eta_R(T_R^mx)\mathbf{1}_{\{A_{m,R}=n,\,K_{m,R}^{\mathrm c}=k\}}a(x)b(T_R^mx)e^{i\Psi_m(x)}\,\frac{d\nu(x)}{c_*}\right). \tag{76.8}$$

The existence of an L^1 density follows from the same exact physical coarea formula in Theorem 69.2, by domination with the complete capped source; the zero-measure critical bulk is exhausted as in Section 67. Each density is supported in the unchanged physical interval $[m\tau_-,m\tau_+]$.

Theorem 76.2 (Summed variation with specified smooth trajectory tests)**.** *For the complete source and all its exact labels,*

$$\sum_\ell\|f_{\ell,m,R}^{w;a,b,\Psi}\|_1\le a_0b_0, \tag{76.9}$$

$$\sum_\ell\|Df_{\ell,m,R}^{w;a,b,\Psi}\|_{\mathrm{TV}}\le D_{\mathrm{all}}^wa_0b_0+J_{\mathrm{test}}\left[C_{\mathrm{test,end}}(a_1b_0+a_0b_1)+C_{\mathrm{test,phase}}\mathcal{L}_ma_0b_0\right]=:\mathcal{V}_m^w(a,b,\Psi). \tag{76.10}$$

The label sum is taken before every supremum in R and m. In particular the right side is uniform in those variables for bounded C^1 endpoint tests and a uniform bound on the partial derivatives of the phase observations. All original exact section events, unvisited constraints, critical endpoints and visited singular limiting configurations are included with their proved physical charges.

Proof. On a regular physical word and one true section chamber let $M = a(x)b(T_R^m x)e^{i\Psi_m(x)}$, and use the label vector v from the proof of Theorem 74.1. Its coefficient is $wAMv$, where $A = \omega/(4\pi c_* \chi)$ is the unchanged physical coarea coefficient. Since $|e^{i\Psi_m}| = 1$, $|M| \le a_0 b_0$, and Lemma 76.1 gives

$$|\partial_t M| \le \frac{1}{z}\left[C_{\text{test,end}}(a_1 b_0 + a_0 b_1) + C_{\text{test,phase}}\mathcal{L}_m a_0 b_0\right]. \tag{76.11}$$

In differentiating wAM, the term $M\partial_t(wA)$ is paid by $a_0 b_0$ times the inherited complete base variation. For the additional term $wA\partial_t M$ use $0 \le w \le 1$ and the full inverse-gradient moment, not a pointwise incidence cutoff:

$$\int_0^{2\pi} \sum_{\text{actual words } v} \int_{\mathcal{D}_v(\theta)} \frac{A_v(t,\theta)}{z}\, dt\, d\theta = \frac{1}{c_*}\int_{\mathcal{C}_{m,R}} \frac{1}{|\nabla_q F|}\, d\nu \le \frac{1}{c_*}\int \frac{I_m}{|\nabla_q F|}\, d\nu \le J_{\text{test}}. \tag{76.12}$$

The equality is the one-copy physical coarea formula on the original cap, first on compact regular pieces and then by nonnegative exhaustion. The inequality uses $I_m \ge 1$ and Theorem 56.3. It contains one division by c_*; the separately conditioned inequality from315 is not inserted and divided by c_* a second time. In particular all near-critical positive bulk is retained in this integral.

At a true physical endpoint, the inherited one-sided limits of every finite-chain collision state exist. The C^1 functions on $\mathcal{X}$ are continuous there and bounded, so their multiplier has a one-sided limit of modulus at most $a_0 b_0$. The complete cap and unvisited-disk charges therefore increase by at most this factor. The positive incoming critical sectors are charged in the same way; no critical value is declared zero. At an actual visited-grazing boundary the already proved limit $A \longrightarrow 0$ remains zero after bounded multiplication. This is a trace calculation on each finite chain, not a deduction from the collision measure of the singular set.

At a regular section cut, M is continuous along the same physical branch. The label vector jumps at cost two for an interior visit and at cost one for an endpoint membership. Multiplication by M pays at most $a_0 b_0$ times those costs. The coincidence, seam and all-word accounting is precisely the one-copy accounting in Theorem 74.1. A guard derivative has already been paid in D^w_{base}, and is not paid a second time in (76.12).

To conclude without assuming a vector-valued integration theorem, test a finite set of components against independent complex C^1_c functions of supremum norm at most one. Integration by parts on a finite regular chamber exhaustion gives the sum of the base, section and extra interior charges just computed. Their full positive total is finite by Lemma 67.3, Theorem 69.2, and (76.12), so the exhaustion passes to the distributional derivatives. Maximizing each component test independently, then increasing the finite label set, gives (76.10). The sum of the L^1 norms is at most the integral of the absolute multiplier against the exact partition of the original section source. That integral is at most $a_0 b_0 \int w\eta_R(T_R^m x) d\nu_R^* \le a_0 b_0$, which proves (76.9). □

Corollary 76.3 (The weak marked roof operator and its actual norms)**.** *For real $\xi \in \mathbb{R}^2$, $\zeta, \upsilon, s \in \mathbb{R}$, define a bilinear form on $C^1(\mathcal{X}) \times C^1(\mathcal{X})$ by*

$$\begin{aligned}\langle \mathcal{K}^{w;\xi,\zeta,\upsilon}_{m,R}(s)a, b\rangle = \int & w\eta_R(T_R^m x)a(x)b(T_R^m x) \\ & \cdot e^{i(\xi\cdot K^c_{m,R}(x) + \zeta A_{m,R}(x) + \upsilon m + sL_{m,R}(x))}\, d\nu_R^*(x).\end{aligned} \tag{76.13}$$

It defines a bounded linear map $C^1(\mathcal{X}) \to C^1(\mathcal{X})^*$, where the dual acts linearly on the second test, and for $s \neq 0$,

$$\|\mathcal{K}^{w;\xi,\zeta,\upsilon}_{m,R}(s)\|_{C^1\to(C^1)^*} \le \min\left\{1, \frac{D^w_{\mathrm{all}} + 2J_{\mathrm{test}}C_{\mathrm{test,end}}}{|s|}\right\}. \tag{76.14}$$

The bound is uniform in m, R and all displayed real mark frequencies. Insertion of $e^{i\Psi_m}$ adds at most $J_{\mathrm{test}}C_{\mathrm{test,phase}}\mathcal{L}_m$ to the numerator. For any bounded label coefficients (c_ℓ), the corresponding tested roof density has translation error at most $\sup_\ell |c_\ell|\, \mathcal{V}^w_m(a, b, \Psi)|h|$ in L^1.

Proof. Sum the densities in (76.8) with coefficients $e^{i(\xi\cdot k+\zeta n+\upsilon m)}$. The absolute L^1 and variation sums in Theorem 76.2 justify this operation and bound the resulting density and derivative. For a compactly supported BV function f and $s \neq 0$, distributional integration by parts gives $|\int e^{ist} f(t)dt| \le \|Df\|_{\mathrm{TV}}/|s|$. The L^1 bound gives the other part of the minimum. If both C^1 test norms are at most one, then $a_0b_0 \le 1$ and $a_1b_0 + a_0b_1 \le 2$; the displayed constant follows without discarding either partial-derivative norm. The phase addition follows from the same calculation. For general bounded label coefficients, the derivative norm is at most $\sup_\ell |c_\ell|$ times the summed bound in (76.10). Integrating that derivative measure along a roof displacement and taking total variation proves the translation assertion with its full coefficient norm. □

Relation to the original local problem. At fixed m the phase υm records the collision mark; it does not create an extra coordinate beyond the original four-coordinate slice (k_1, k_2, n, t). Formula (76.13) is evaluated on the actual physical source with the original entrance law and terminal membership. The full chain phase is an optional observation with the exact norm stated above, and no derivative of an arbitrary Borel trajectory selector is claimed. Such selectors are covered by the positive likelihood estimate of Theorem 75.4 instead.

Estimate (76.14) supplies a weak tested $1/|s|$ roof estimate. It is neither a strong anisotropic transfer-operator estimate nor a decay estimate in the number of returns. In particular $1/|s|$ is not integrable at infinity; absolute Fourier inversion on an expanding frequency range does not follow from this bound alone. The arithmetic peak expansion, uniform spectral remainder, microscopic pointwise local theorem and long-time four-coordinate inverse require their stated additional inputs. No supremum or ordered limit in the original target is exchanged here. The endpoint stationary-action derivative used in the proof does not replace the actual long-time forward-orbit Jacobian in that inverse problem.

77. Explicit reconstruction of the inherited section multipliers

The occupation observable is the indicator of the actual section Y^*_R. The historical native section-multiplier lemma and module 84 already prove its strong multiplier property, entire occupation phase and weak parameter continuity; modules 84–86 contain the inherited full fixed-band spectral and arithmetic arguments. Those results are retained, with their stated geometric and analytic hypotheses. This section supplies a fully explicit reconstruction of the section interface and its external hypotheses. It is a proof-completeness addition, not a claim that those multiplier or fixed-band theorems are new. No section side is rounded and no return event is changed.

Use the common coordinates (α, φ), with $p = \sin\varphi$, and the radius-dependent arclength $s = R\alpha$. The classical stable cones in arclength coordinates have slopes between $-K$ and $-a$, where the uniform choices

$$a = 100/47, \qquad K = 170/9 \tag{77.1}$$

follow from the actual curvature and flight bounds. In the common coordinates every such cone is contained in

$$-\overline{K} \le d\varphi/d\alpha \le -\underline{a}, \qquad \underline{a} = 45/47, \qquad \overline{K} = 799/90. \tag{77.2}$$

The common cone contains the physical stable directions. The test arcs used below are tangent to this cone, have the fixed curvature and short-length bounds, and are closed under restriction. Invariance of the enlarged test family under every T_R is a separate spectral requirement. The sign is reversed from the unstable curves in Section 71. The change $s = R\alpha$ and its inverse have uniformly bounded derivatives on the given radius interval.

We use precisely the stable-curve norm construction of Demers–Zhang [6, Sect. 3.3, Lemmas 5.2–5.3]. In that construction fix common admissible exponents $\alpha_{\mathrm{DZ}}, \beta_{\mathrm{DZ}}, p_{\mathrm{DZ}}, q_{\mathrm{DZ}}$ and curve length and curvature bounds, with $0 < q_{\mathrm{DZ}} < p_{\mathrm{DZ}} \le 1/3$, $0 < \alpha_{\mathrm{DZ}} < 1 - \varsigma_0$ for the fixed summability parameter $\varsigma_0 \in (0,1)$ of that construction, and $0 < \beta_{\mathrm{DZ}} \le \min\{\alpha_{\mathrm{DZ}}, p_{\mathrm{DZ}} - q_{\mathrm{DZ}}\}$. The norms are denoted $\|\cdot\|_{\mathcal{B}}$ and $|\cdot|_{\mathcal{B}_w}$ on the common cylinder. The strong norm dominates the stable norm and the weak norm tests homogeneous stable arcs against functions with $C^{p_{\mathrm{DZ}}}$ norm at most one. Here C^p has the cited little Hölder convention, the closure of smooth functions in the Hölder norm on each test arc. The stable part has the following defining consequence: on an admissible subarc V,

$$\left| \int_V h\psi \, dm_V \right| \le \|h\|_{\mathcal{B}} |V|^{\alpha_{\mathrm{DZ}}} |\psi|_{C^{q_{\mathrm{DZ}}}(V)}. \tag{77.3}$$

The strong multiplier proof also applies in each radius-specific Demers–Zhang arclength space, with common norm parameters and curve bounds. Its constants include the uniformly bounded coordinate scalings. The weak difference of two moving sections is asserted on the fixed common test-curve construction just specified; no identification of different dynamical spaces is implicit. In the measure embedding of this construction a function h represents $h\nu$; the constant function 1 therefore represents ν. The coordinate expressions are $d\nu = \cos\varphi \, d\alpha \, d\varphi/(4\pi) = \cos\varphi \, ds \, d\varphi/(4\pi R)$. These expressions describe the reference measure and are not a replacement of h by a density with respect to flat area. In particular no division by $\cos\varphi$ is used at grazing. Lemma 3.4 and the measure embedding of the cited construction make the mass functional $\ell(h) = \int h \, d\nu$ continuous on $\mathcal{B}_w$, by testing against the constant function 1. The assertions concern these norm constructions. They do not assert that the complete jointly twisted collision operators already satisfy the separate hypotheses (H1)–(H5), (C1)–(C4), or the spectral conclusions of that reference.

Lemma 77.1 (Finite transverse partition for the actual section)**.** *For every $R \in [R_-, R_+]$ there is a modulo-zero partition of the collision cylinder into at most* 121 *open simply connected cells on each of which $\eta_R = \mathbf{1}_{Y_R^*}$ is constant. Each cell has at most four smooth sides. Every side is uniformly transverse to* (77.2)*; a lower bound for the sine of the angle is*

$$\vartheta_{\mathrm{sec}} = \frac{\underline{a}}{\sqrt{1 + \overline{K}^2}} > 0. \tag{77.4}$$

Every homogeneity strip meets at most 121 *cells. The original section is contained in $|p| < 5/6$, and its four moving vertical sides have speed at most* 6 *in α, while its four moving horizontal sides have speed less than* 11 *in φ.*

Proof. The four squares and the rectangle U have at most ten distinct α side values and ten distinct p side values. Cut the angular circle at the common seam $\alpha = \pi$, which lies outside all these rectangles for the entire radius range, and extend the coordinate sides into full coordinate lines. At most eleven angular intervals and eleven p intervals result. Removing the lines gives the asserted partition. A cell remains a rectangle after $p = \sin\varphi$, so it is simply connected and has at most four sides. Empty or coincident cells reduce the count.

A unit tangent to a stable arc has components proportional to $(1, v)$ with $-\overline{K} \le v \le -\underline{a}$. Its angle with a vertical line has sine at least $(1 + \overline{K}^2)^{-1/2}$; its angle with a horizontal line has sine

at least $\underline{a}(1+\overline{K}^2)^{-1/2}$. This proves (77.4), also at the artificial seam. The homogeneity-strip count is bounded by the total cell count.

Here is an explicit check of the moving sides. Write $F(R,q)$ for the polynomial with coefficient $\sqrt{3}$ defining q_R in the physical model. For $49/100 \le q \le 53/100$,

$$F_q = 8R(1+3q^2) - 4 + 12q^2 + \sqrt{3}(-12q + 4q^3) < -2.$$

Indeed its first three terms are at most $8(47/100)(1+3(53/100)^2) - 4 + 12(53/100)^2$, and its last term is at most $(17/10)(-12(49/100)+4(53/100)^3)$: the expression in parentheses is negative and $\sqrt{3} > 17/10$. Their sum is -2.6842844. Also $0 < F_R \le 8(53/100)(1+(53/100)^2) < 6$. Implicit differentiation therefore gives $0 < q'_R < 3$ and $|\theta'_R| = 2q'_R/(1+q_R^2) < 6$. These estimates hold on the whole given root interval, including its closed endpoints used for the bounds; uniqueness of the physical root is already part of the retained geometry.

The two moving squares have angular center speed $|\theta'_R|$ and momentum center speed $|(\sin\theta_R)'|$, both at most 6. Furthermore

$$\sin\theta_R + h_0 \le \frac{2(53/100)}{1+(53/100)^2} + \frac{1}{200} < \frac{5}{6}.$$

All the other section momentum coordinates satisfy this same cap. Thus every moving horizontal side in φ coordinates has speed at most $6/\sqrt{1-(5/6)^2} = 36/\sqrt{11} < 11$. The initial normal square, the square centered at $(\pi/6, -1/2)$, and U are fixed. There are exactly four possibly moving sides of each orientation. No assertion of independence of their motion or of orbit visits has entered the calculation. □

Theorem 77.2 (Strong multiplication by the exact section and occupation phase)**.** *Let $\mathcal{B}$ have the Demers–Zhang norm construction just specified. There is a constant C_{sec} independent of R such that multiplication by η_R and by $f_{R,z} = \exp(z\eta_R)$ is bounded on $\mathcal{B}$, with*

$$\|M_{\eta_R}\|_{\mathcal{B}\to\mathcal{B}} \le C_{\mathrm{sec}}, \qquad \|M_{f_{R,z}}\|_{\mathcal{B}\to\mathcal{B}} \le C_{\mathrm{sec}} \max\{1, e^{\operatorname{Re} z}\}. \tag{77.5}$$

For every real occupation frequency ζ the second bound is C_{sec} at $z = i\zeta$. The map $z \mapsto M_{f_{R,z}}$ is entire in operator norm, and, for every integer $r \ge 1$,

$$M_{f_{R,z}} = I + (e^z - 1)M_{\eta_R}, \qquad \partial_z^r M_{f_{R,z}} = e^z M_{\eta_R}. \tag{77.6}$$

The constant depends on the fixed norm parameters, the common curve bounds, and the explicit transverse partition constants, but on no collision count or observation window.

Proof. Choose the Hölder exponent used in Lemma 5.3 of the cited author version as $\gamma = \max\{p_{\mathrm{DZ}}, 2\beta_{\mathrm{DZ}} + \epsilon_{\mathrm{DZ}}\}$ with one admissible $\epsilon_{\mathrm{DZ}} > 0$. On every cell of Lemma 77.1, both multipliers are constant. Their piecewise C^γ norms are therefore at most 1 and $\max\{1, e^{\operatorname{Re} z}\}$, respectively. The partition is modulo zero, open and simply connected; its boundary count, transversality and cells per homogeneity strip have all been checked explicitly. Lemmas 5.2–5.3 consequently give the two bounded multiplication operators on the strong space. All parameters entering their estimates are uniform here. In particular no factor $|\zeta|$ is incurred for the piecewise constant occupation phase.

On smooth densities the identity $e^{z\eta_R} = 1 + (e^z - 1)\eta_R$ holds almost everywhere. The bounded extensions and density of the original smooth class in $\mathcal{B}$ prove the first operator identity in (77.6). Its norm-convergent power series proves the asserted entire dependence and every derivative formula. This argument invokes the precise multiplier theorem, not its spectral corollaries. □

Theorem 77.3 (Weak perturbation estimate for the moving physical section)**.** *Use the common stable-curve realization above. There is a constant C_{move}, independent of $R, S \in [R_-, R_+]$, such*

that

$$\|M_{\eta_R} - M_{\eta_S}\|_{\mathcal{B}\to\mathcal{B}_w} \le C_{\mathrm{move}}|R-S|^{\alpha_{\mathrm{DZ}}}, \tag{77.7}$$

$$\|M_{f_{R,z}} - M_{f_{S,z}}\|_{\mathcal{B}\to\mathcal{B}_w} \le |e^z-1|C_{\mathrm{move}}|R-S|^{\alpha_{\mathrm{DZ}}}. \tag{77.8}$$

The measures and section normalizing constant are the original ones. These inequalities concern multiplication only; a moving collision map has its own additional perturbation estimates.

Proof. Write $d = |R-S|$. For $d=0$ both assertions are identities. The symmetric difference of the two sections is contained in the union of strips swept out by the eight moving sides. Their coordinate widths are at most $11d$, by Lemma 77.1. Along any stable arc satisfying (77.2), both coordinates are strictly monotone in arclength and their absolute speeds are at least ϑ_{sec}. The intersection with any one of these strips has length at most $11d/\vartheta_{\mathrm{sec}}$.

Cut an admissible short stable arc at the coordinate sides of both section grids and, if necessary, at the angular seam. There are at most twenty side values of each orientation. On each chart piece a monotone arc therefore has at most forty-one complementary intervals; allowing a seam split gives the safe bound eighty-two. The difference $\eta_R - \eta_S$ is constant on each such interval, with value 0, 1, or -1. Every nonzero interval is contained in a swept strip. One way to see the last assertion is to refine also by the strip endpoints, which are among the same two sets of side values: membership in each strip is then constant on a component. Thus all nonzero subarcs have length at most $11d/\vartheta_{\mathrm{sec}}$. Curvature, cone membership and homogeneity are preserved by restriction.

For a weak test ψ of $C^{p_{\mathrm{DZ}}}$ norm at most one, its restricted $C^{q_{\mathrm{DZ}}}$ norm is bounded by a uniform constant C_{pq}, since $p_{\mathrm{DZ}} \ge q_{\mathrm{DZ}}$ and the maximum curve length is fixed. Applying (77.3) on each nonzero interval gives

$$\left|\int_W h(\eta_R-\eta_S)\psi\, dm_W\right| \le 82C_{pq}\left(\frac{11}{\vartheta_{\mathrm{sec}}}\right)^{\alpha_{\mathrm{DZ}}} d^{\alpha_{\mathrm{DZ}}}\|h\|_{\mathcal{B}}.$$

This is first proved for smooth densities, for which restrictions and pointwise products are literal. Taking the weak norm and then using the bounded strong multipliers of Theorem 77.2 extends it to their strong-space completion. The continuous embedding into the weak space makes the extension the same operator. This proves (77.7); the first identity in (77.6) proves (77.8). □

Corollary 77.4 (Exact occupation and endpoint tests in a qualified transfer formula)**.** *Suppose, separately, that the physical operator*

$$\mathcal{A}_{R;\xi,s}h = \mathcal{L}_R(e^{i(\xi\cdot\kappa_R + s\tau_R)}h)$$

has a bounded realization on this strong space at the displayed parameters. Define $\mathcal{L}_{R;\xi,\zeta,s} = \mathcal{A}_{R;\xi,s}M_{f_{R,i\zeta}}$*. For every finite collision count* m *for which the realization is available, its physical moment is exactly*

$$\frac{1}{c_*}\ell\left(M_{\eta_R}\mathcal{L}^m_{R;\xi,\zeta,s}(\eta_R\nu)\right) = \int \eta_R(T_R^m x)e^{i(\xi\cdot K^c_{m,R}+\zeta A_{m,R}+sL_{m,R})}\, d\nu^*_R(x). \tag{77.9}$$

The section multipliers in this formula satisfy (77.5)*, and the dependence on* ζ *is entire.*

Proof. The physical transfer convention is $(\mathcal{L}_R h)(b) = h(b\circ T_R)$ on measures. An induction shows that m iterates multiply the trajectory by the displayed flight and displacement phases and by $\prod_{j=0}^{m-1} e^{i\zeta\eta_R(T_R^j x)}$. The latter product is exactly $e^{i\zeta A_{m,R}}$. The terminal multiplier tests $\eta_R(T_R^m x)$ and contributes no extra occupation count. The initial measure is $\eta_R\nu$; the single division by c_* gives the original entrance law. This proves the equality on the actual physical trajectories. When the bounded realization is used, the multiplier theorem justifies its section factors. The inherited fixed-band realization and power estimates have their own proof bodies and hypotheses;

this multiplication argument supplies no extension of their constants to a growing roof-frequency band. □

Reconstructed interface and retained spectral scope. The original section indicator, full occupation Fourier phase and moving section cuts have explicit strong multiplier and weak perturbation estimates with checked geometric hypotheses. This reconstruction does not reopen the existing fixed-band author proofs or replace them by a new assumption. Control on growing roof-frequency bands is a separate possible spectral estimate and is not supplied by this multiplier calculation. The retained fixed-window theorem will evaluate the scalar bridge in Section 78 without such an additional estimate. The original endpoint shortage (73.12), the normalized local derivative charge on that alternative route, and the actual microscopic denominator for conditional conclusions remain at their stated scopes.

78. Absolute Fourier inversion of the unchanged microscopic intervals

The weak roof estimate in Section 76 has a nonintegrable $|s|^{-1}$ tail when it is used alone. The transform of an *exact bounded roof interval* supplies another inverse power. We exploit this factor without replacing the interval by a likelihood or enlarging it. The resulting formula is an absolutely convergent inversion of the actual event. Its error after division by the true interval length is stated explicitly.

The earlier exact reconstruction arguments are retained. Historical native 64 already gives a sharp full-law interval integral with an extracted second-derivative budget; native 67 gives absolute sharp inversion for the stationary flight-age overlap; native 93 gives coefficient-first raw reconstruction after its exact germ subtraction. Theorem 75.4 already permits different windows at different labels, and Corollary 75.8 already uses their actual selected denominator for positive likelihoods. No priority is claimed for Fourier inversion, interval transforms, label summation or denominator certification as general methods. The addition here is a direct full-joint interval corollary of the complete first-order BV budget, including its precise sharp tail, the original $m^2/|I|$ payment and specified endpoint tests, with neither a remaining likelihood kernel nor a germ subtraction.

Fix $R \in [R_-, R_+]$, the actual collision count $m \geq 1$, and either $w = 1$ or the original $w = W^\varepsilon$ with $\varepsilon > 0$ fixed. Use the endpoint tests and physical orbit phase of (76.8), and abbreviate

$$V(x) = a(x)b(T_R^m x)e^{i\Psi_m(x)}, \qquad A_V = a_0 b_0, \qquad D_V = \mathcal{V}_m^w(a, b, \Psi).$$

The phase is optional. All tests have precisely the C^1 norms stated there. Write $\mathbf{j} = (k_1, k_2, n) \in \mathbb{Z}^3$ and $f_{\mathbf{j}} = f^{w;a,b,\Psi}_{(n,k),m,R}$; this is only an ordering of the unchanged discrete record. Define its genuine joint transform

$$\begin{aligned} \mathcal{F}^{w,V}_{m,R}(\vartheta, s) &= \sum_{\mathbf{j}} e^{i\vartheta\cdot\mathbf{j}} \int_{\mathbb{R}} e^{ist} f_{\mathbf{j}}(t)\, dt \\ &= \int w\eta_R(T_R^m x)V(x)e^{i\{\xi\cdot K^{\mathrm{c}}_{m,R}(x)+\zeta A_{m,R}(x)+sL_{m,R}(x)\}}\, d\nu^*_R(x). \end{aligned} \tag{78.1}$$

Here $\vartheta = (\xi_1, \xi_2, \zeta) \in [-\pi, \pi]^3$. Theorem 76.2 gives

$$\sum_{\mathbf{j}} \|f_{\mathbf{j}}\|_1 \leq A_V, \qquad \sum_{\mathbf{j}} \|Df_{\mathbf{j}}\|_{\mathrm{TV}} \leq D_V, \qquad |\mathcal{F}^{w,V}_{m,R}(\vartheta, s)| \leq \min\{A_V, D_V/|s|\} \quad (s \neq 0). \tag{78.2}$$

At fixed m only finitely many discrete records are attainable, by the physical finite horizon. None of the following constants uses their number.

Let I be a union of at most $J \geq 1$ bounded intervals, let $d = |I|$, and use the convention $\widehat{\mathbf{1}_I}(s) = \int_I e^{ist}\, dt$. Endpoints may be open or closed, because the true roof laws in question have

densities. Merging overlapping intervals gives disjoint intervals without increasing J, and direct integration gives

$$|\widehat{\mathbf{1}_I}(s)| \le \min\{d, 2J/|s|\} \quad (s \neq 0). \tag{78.3}$$

For $S > 0$ define

$$\mathfrak{I}_S(\mathbf{j}, I; V) = (2\pi)^{-4} \int_{[-\pi,\pi]^3} \int_{-S}^{S} e^{-i\vartheta\cdot\mathbf{j}} \widehat{\mathbf{1}_I}(-s) \mathcal{F}^{w,V}_{m,R}(\vartheta, s)\, ds\, d\vartheta. \tag{78.4}$$

This is a sharp frequency truncation, and is not asserted positive.

Theorem 78.1 (Exact interval inversion and its full tail)**.** *For every specified physical parameter, exact label and interval union,*

$$\int_I f_{\mathbf{j}}(t)\, dt = (2\pi)^{-4} \int_{[-\pi,\pi]^3} \int_{\mathbb{R}} e^{-i\vartheta\cdot\mathbf{j}} \widehat{\mathbf{1}_I}(-s) \mathcal{F}^{w,V}_{m,R}(\vartheta, s)\, ds\, d\vartheta. \tag{78.5}$$

The integral on the right is absolutely convergent. If $d > 0$, then

$$\left| \int_I f_{\mathbf{j}} - \mathfrak{I}_S(\mathbf{j}, I; V) \right| \le \frac{D_V}{\pi}\, \mathfrak{t}_J(d, S) \le \frac{2JD_V}{\pi S}, \tag{78.6}$$

$$\mathfrak{t}_J(d, S) = \begin{cases} d[1 + \log(2J/(dS))], & dS \le 2J, \\ 2J/S, & dS \ge 2J. \end{cases} \tag{78.7}$$

For $d = 0$ the event and every displayed integral vanish. The assertions hold for all original lattice and occupation labels, including zero arithmetic classes, and for the full physical roof support. There is exactly one conditioning factor c_*^{-1}, already present in $f_{\mathbf{j}}$ and D_V.

Proof. For bounded s the integrand in (78.5) is bounded by dA_V. At $|s| \ge 1$, its absolute value is bounded by $2JD_V/s^2$. The torus has finite volume; these two estimates prove absolute integrability.

For completeness we justify inversion without assuming pointwise Fourier inversion for a BV density. Let $k_{\mathfrak{b}}$ be the probability kernel of Lemma 75.3. The BV translation bound implies

$$\|k_{\mathfrak{b}} * f_{\mathbf{j}} - f_{\mathbf{j}}\|_1 \le \|Df_{\mathbf{j}}\|_{\mathrm{TV}} \kappa_{\mathrm{ker}}/\mathfrak{b} \longrightarrow 0.$$

For this intermediate convolution, the compact-frequency formula and orthogonality on the entire three-dimensional torus give the right side of (78.5) with the additional factor $\eta(s/\mathfrak{b})$. Every integral is then over a bounded frequency domain, so Fubini is valid. The left side tends to $\int_I f_{\mathbf{j}}$ by the displayed L^1 estimate. Since $|\eta| \le 1$ and $\eta(s/\mathfrak{b}) \to 1$, the already proved integrable bound permits dominated convergence on the right. The convolution has disappeared from the resulting identity. Its left side is the exact original interval event, with its original width and trajectory information.

The three torus integrals cancel three of the four factors 2π in an absolute tail bound. Equations (78.2) and (78.3) therefore yield

$$\left| \int_I f_{\mathbf{j}} - \mathfrak{I}_S(\mathbf{j}, I; V) \right| \le \frac{D_V}{\pi} \int_S^{\infty} \min\{d/u, 2J/u^2\}\, du.$$

Splitting this positive integral at $2J/d$ proves (78.7). Its integrand is always at most $2J/u^2$, proving the last bound in (78.6) for every $S > 0$, including $dS < 2J$. No roof supremum, label supremum, collision limit or outer geometric limit was exchanged. □

Corollary 78.2 (All labels and their different exact windows)**.** *For each attainable $\mathbf{j}$ choose an empty set or a union $I_{\mathbf{j}}$ of at most J bounded intervals, and choose coefficients $|c_{\mathbf{j}}| \le 1$. Summing*

the exact component identities gives

$$\left|\sum_{\mathbf{j}} c_{\mathbf{j}} \left(\int_{I_{\mathbf{j}}} f_{\mathbf{j}} - \mathfrak{I}_S(\mathbf{j}, I_{\mathbf{j}}; V) \right)\right| \le \frac{2JD_V}{\pi S}. \tag{78.8}$$

The bound has no factor counting records, selected windows or components of their physical preimages.

Proof. Integrate the full torus first in (78.4); orthogonality selects exactly $\widehat{f_{\mathbf{j}}}(s)$. For that component the same proof of (78.6) gives $2J\|Df_{\mathbf{j}}\|_{\mathrm{TV}}/(\pi S)$. Sum these positive error bounds and apply (78.2). A finite count has finitely many attainable labels, so no infinite complex-series interchange is needed. The norm sum explains why the estimate is independent of that finite number. □

Corollary 78.3 (The original density normalization)**.** *For a nonempty original interval union I, of its actual length d,*

$$\frac{m^2}{d} \left| \int_I f_{\mathbf{j}} - \mathfrak{I}_S(\mathbf{j}, I; V) \right| \le \frac{2JD_V m^2}{\pi d S}. \tag{78.9}$$

In particular, for any $P > 0$, the analytical choice $S = m^{P+2}/d$ makes this upper bound $2JD_V m^{-P}/\pi$. This holds for every positive d, however small. A claim at this bandwidth must evaluate the joint transform on that whole bandwidth; no available low-frequency theorem is silently extended to it.

Proof. Divide (78.6) by the true interval length and multiply by the original m^2. Substitute the stated finite S. There is no lower bound on d and no modification of the event. □

78.1. **Conditional observations use the actual event mass.** Let $\mathcal{A}$ be one exact record with its interval union, or an event formed by the different exact windows in Corollary 78.2, with coefficients all equal to one. Use the positive physical source $w\eta_R(T_R^m x)d\nu_R^*$ and write

$$p = \int \mathbf{1}_{\mathcal{A}} w\eta_R(T_R^m x)d\nu_R^*, \qquad N = \int V\mathbf{1}_{\mathcal{A}} w\eta_R(T_R^m x)d\nu_R^*.$$

Let D_S and N_S be the corresponding sharp-frequency expressions with test 1 and V, respectively. The former is real by conjugate symmetry. Put $E_0 = 2JD_{\mathrm{all}}^w/(\pi S)$ and $E_V = 2JD_V/(\pi S)$; the smaller single-window bounds in (78.6) may be substituted when appropriate.

Proposition 78.4 (Certified denominator and conditional test)**.** *One has $|p - D_S| \le E_0$ and $|N - N_S| \le E_V$. If $D_S > E_0$, then the actual selected probability satisfies $p \ge D_S - E_0 > 0$, and*

$$\left| \frac{N}{p} - \frac{N_S}{D_S} \right| \le \frac{E_V + A_V E_0}{D_S}. \tag{78.10}$$

No conditional law is asserted when the actual event has zero mass. This statement concerns exactly the displayed smooth endpoint and trajectory-phase observations. It does not assign their variation bound to an arbitrary additional Borel information selector.

Proof. The two absolute errors follow from Theorem 78.1 and Corollary 78.2. The positivity certificate follows from the first error. Since $|V| \le A_V$ on the original trajectories, positivity gives $|N| \le A_V p$. The exact identity

$$\frac{N}{p} - \frac{N_S}{D_S} = \frac{N - N_S}{D_S} + \frac{N(D_S - p)}{pD_S}$$

proves (78.10). This uses the mass of precisely $\mathcal{A}$, not that of a containing event. In particular, if $p = 0$ the positivity certificate cannot hold because $|D_S| \le E_0$. □

78.2. **A sufficient pointwise bridge, with the missing input exposed.** We state a sufficient implication to identify what the new interval formula does and does not complete. It retains the existing raw arithmetic reference, all its peaks, and the ordered outer limit. Here $w = 1$, $V = 1$, and $p_{\mathbf{j},m,R}$ is the original density. Every raw-target supremum below uses the original ranges $R \in [R_-, R_+]$, $1 \le n \le m$ and $k \in \mathbb{Z}^2$. Set

$$g_{\mathbf{j},m,R}(u) = m^{-2}\mathcal{L}_{m,R}(k_1, k_2, u, n).$$

Fix M. At each outer stage B choose a positive h_B before the collision limit, and for every central t put $I_{B,t} = (t - h_B, t + h_B)$. These intervals are analysis tests of the same physical roof, not a redefinition of any conditioning event. For any finite bandwidth $S_{m,B} > 0$ define

$$\Omega^p_{m,B} = \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} |Dp_{\mathbf{j},m,R}|([t - h_B, t + h_B]),$$

$$\Omega^g_{m,B} = \sup_{R,n,k} \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \frac{1}{2h_B}\int_{t-h_B}^{t+h_B} |\mathcal{L}_{m,R}(k_1, k_2, u, n) - \mathcal{L}_{m,R}(k_1, k_2, t, n)|\, du,$$

$$\Xi_{m,B} = \sup_{R,n,k} \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \frac{m^2}{2h_B}\left|\mathfrak{I}_{S_{m,B}}(\mathbf{j}, I_{B,t}; 1) - \int_{I_{B,t}} g_{\mathbf{j},m,R}(u)\, du\right|.$$

The derivative measure is that of the complete actual exact-label density, after every physical word and boundary has been included.

Proposition 78.5 (An ordered raw-density implication)**.** *At every finite m, B,*

$$\sup_{R,n,k} \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} |m^2 p_{\mathbf{j},m,R}(t) - \mathcal{L}_{m,R}(k_1, k_2, t, n)| \le \Omega^p_{m,B} + \Omega^g_{m,B} + \Xi_{m,B} + \frac{D^1_{\mathrm{all}} m^2}{\pi h_B S_{m,B}}. \quad (78.11)$$

Consequently the original raw target follows if each of the four terms on the right tends to zero under $\lim_{B\to\infty} \limsup_{m\to\infty}$. With $S_{m,B} = m^{P+2}/h_B$, $P > 0$, the last term already tends to zero for every fixed B. The local derivative-measure condition, the reference modulus and the scalar band comparison are displayed separately in this general inequality. The following subsection uses the retained fixed-window theorem to pay the latter two; they must not be relisted as new unproved requirements when that inherited theorem is invoked at its original scope.

Proof. For a BV representative at a Lebesgue point t, its difference from almost every value on $I_{B,t}$ is bounded by $|Dp|([t - h_B, t + h_B])$. Integrating gives

$$\left|p(t) - \frac{1}{2h_B}\int_{I_{B,t}} p(u)\, du\right| \le |Dp|([t - h_B, t + h_B]).$$

Insert the exact interval average between the raw density and the reference, then insert its sharp-frequency expression and the reference interval average. These four differences are bounded, respectively, by Ω^p, the normalized tail (78.9) with $J = 1, d = 2h_B$, Ξ, and Ω^g. This proves (78.11), first for each fixed radius and almost every roof, and then with the stated suprema. Take the collision limsup with h_B already fixed; take the outer limit last. No interchange is needed. The displayed bandwidth reduces its paid term to $D^1_{\mathrm{all}} m^{-P}/\pi$. □

78.3. **Evaluation of the bridge by the retained fixed-window theorem.** The preceding sufficient inequality must be read together with the written fixed-window theorem already present in historical native 87. Its Theorem v41-uniform-transition, equation v41-uniform-interval, concerns precisely the event (1.1) and the finite-peak reference (1.4). It includes the moving nonunit branches; it is not restricted to one fixed radius or to a unit arithmetic factor. We retain that theorem with its existing analytic hypotheses. The following implication uses its conclusion, without reasserting an independent reconstruction of its whole spectral proof.

For a fixed $h > 0$ set

$$\mathscr{E}_m(h) = \sup_{R,\, 1\le n\le m,\, k\in\mathbb{Z}^2,\, t\in\mathbb{R}} \left| \frac{m^2}{2h} \int_{t-h}^{t+h} p_{n,R}(k,m,u)\, du \right. \\ \left. -\mathcal{L}_{m,R}(k_1,k_2,t,n) \right|. \tag{78.12}$$

The retained fixed-window conclusion is $\mathscr{E}_m(h) \to 0$ for every fixed $h > 0$. Its proof fixes the spectral band before the collision limit, then removes that band by ordered interval envelopes. It supplies no rate uniform over collision-dependent choices of h.

Lemma 78.6 (Uniform roof modulus of the retained reference)**.** *Retain the finite peak data and their uniform properties in* (1.4) *and historical native 87. Let N_{pk} be the number of peak charts, B_{pk} a common upper bound for $|B_j(R)|$, and $a > 0$ the common lower bound for $\operatorname{Re} H_j$. For each chart use its compact parameter support K_j, on a neighborhood of which its scalar data are defined. Put*

$$b_j(R) = (a_j(R))_3, \qquad \kappa_j(R) = -\log|z_j(R)| \ge 0, \qquad \omega_m = \max_j \sup_{R\in K_j} |b_j(R)| e^{-m\kappa_j(R)},$$

where an empty supremum is zero. Then $\omega_m \to 0$. With

$$M_0 = (2\pi)^{-2} a^{-2}, \qquad M_1 = M_0\sqrt{2/(\pi a)}, \qquad D_m = \frac{N_{\mathrm{pk}} B_{\mathrm{pk}}}{c_*}\{M_0\omega_m + M_1 m^{-1/2}\},$$

one has $D_m \to 0$ and

$$\sup_{R,n,k,t} |\partial_t \mathcal{L}_{m,R}(k_1,k_2,t,n)| \le D_m. \tag{78.13}$$

In particular $\Omega^g_{m,B} \le h_B D_m/2$ for every finite outer stage B.

Proof. The retained peak theorem gives continuity on each K_j, bounded roof frequencies, and $\kappa_j(R) = 0 \Rightarrow b_j(R) = 0$. The last implication is the original no-roof-resonance assertion; it does not say that every continued peak has roof frequency zero. Let $b_* < \infty$ bound all $|b_j|$. For any $\epsilon > 0$, the finite union of compact sets $\{|b_j| \ge \epsilon\} \cap K_j$ either is empty or has a strictly positive minimum c_ϵ of κ_j. Consequently

$$\omega_m \le \epsilon + b_* e^{-mc_\epsilon}$$

in the nonempty case, and $\omega_m \le \epsilon$ otherwise. First let $m \to \infty$, then $\epsilon \downarrow 0$. No rate for c_ϵ is required.

The retained peaks are interior real maxima of $\log|\lambda_j|$, so $\operatorname{Re}\nabla \log\lambda_j(R, a_j) = 0$. Consequently $\mu_j = i^{-1}\nabla\log\lambda_j(R,a_j)$ is real, as asserted in historical native 87. All Gaussian arguments below are therefore real. For every real z and every displayed complex symmetric H, the defining Gaussian integral gives

$$|G_H(z)| \le (2\pi)^{-4}\int_{\mathbb{R}^4} e^{-a|v|^2/2}\, dv = M_0, \qquad |\partial_{z_3} G_H(z)| \le (2\pi)^{-4}\int_{\mathbb{R}^4} |v_3| e^{-a|v|^2/2}\, dv = M_1.$$

Differentiation is justified by the same integrable Gaussian majorant with the factor $|v_3|$. In each term of (1.4), differentiating the phase costs $|b_j|M_0$, while differentiating the Gaussian argument costs $m^{-1/2}M_1$. Its coefficient has modulus at most $c_*^{-1}B_{\mathrm{pk}}e^{-m\kappa_j}$. Sum the finitely many terms and take the real part to prove (78.13). This estimate is global in the real roof target, so averaging it over the analysis interval does not require that every point of that interval be central. Finally $(2h_B)^{-1}\int_{-h_B}^{h_B}|u|\, du = h_B/2$ proves the last assertion. □

Proposition 78.7 (The fixed-window input evaluates the scalar bridge)**.** *Assume the retained fixed-window conclusion* (78.12) *and its finite peak data at their original scope. At each finite*

outer stage choose $h_B > 0$ before the collision limit. For the quantities in Proposition 78.5, one has

$$\Xi_{m,B} \le \mathscr{E}_m(h_B) + \frac{h_B D_m}{2} + \frac{D^1_{\mathrm{all}} m^2}{\pi h_B S_{m,B}}. \tag{78.14}$$

Thus, with $S_{m,B} = m^{P+2}/h_B$ and fixed $P > 0$, both $\Xi_{m,B}$ and $\Omega^g_{m,B}$ tend to zero for each fixed B. More directly, writing $\mathcal{E}_M$ for the original raw-error collision limsup,

$$\mathcal{E}_M \le \limsup_{m\to\infty} \Omega^p_{m,B} \qquad \text{for every fixed } B. \tag{78.15}$$

In particular the sufficient remaining condition on this route is

$$\lim_{B\to\infty} \limsup_{m\to\infty} \Omega^p_{m,B} = 0.$$

This is a condition on the complete normalized local derivative charge of the original physical density. It is not proved by the global unscaled BV estimate and is not claimed necessary.

Proof. The label density here is exactly $p_{n,R}(k, m, \cdot)$, and its interval mass is the original nth-return event with collision count m, by (1.1). Apply the retained fixed-window theorem with $J = (-h_B, h_B)$. The interval length is positive and fixed at this stage, so division by $2h_B$ is legitimate and $\mathscr{E}_m(h_B) \to 0$.

Insert the actual interval mass between its sharp-frequency expression and the reference interval average in the definition of Ξ. The first error, multiplied by $m^2/(2h_B)$, is at most $D^1_{\mathrm{all}} m^2/(\pi h_B S_{m,B})$ by Corollary 78.3. The difference between the normalized actual mass and the reference at its center is bounded by $\mathscr{E}_m(h_B)$. The difference between that reference value and its interval average is at most $h_B D_m/2$, by Lemma 78.6. This proves (78.14). The stated bandwidth makes its last term $D^1_{\mathrm{all}} m^{-P}/\pi$. Every term therefore vanishes with B fixed.

For the sharper raw implication, no frequency truncation is needed. At each Lebesgue point of the exact-label BV density, insert its exact interval average directly between the density and the reference at the center. The local-variation estimate in the proof of Proposition 78.5 gives

$$\sup_{R,n,k} \operatorname*{ess\,sup}_{t\in \mathcal{C}_{m,R,n,k}(M)} |m^2 p_{n,R}(k, m, t) - \mathcal{L}_{m,R}(k_1, k_2, t, n)| \le \Omega^p_{m,B} + \mathscr{E}_m(h_B).$$

The inherited fixed-window theorem is uniform over all the displayed radii and labels. Take the collision limsup, keeping h_B fixed, to obtain (78.15); only then take the outer limit. Exceptional Lebesgue-point sets may depend on the radius, as allowed by the original $\sup_R \operatorname{ess\,sup}_t$ order. □

This argument does not claim a new strong-operator estimate on a growing roof band. It evaluates the particular scalar sharp interval expression through the retained fixed-window law and the proved tail. A direct growing-band spectral estimate is another possible method, not a prerequisite for this implication. The analysis intervals do not replace the original microscopic conditioning events. The conclusion in this subsection is for $w = 1, V = 1$; its use for additional observation classes requires their own stated fixed-window theorem.

Why the local variation condition cannot be omitted. The following analytical example tests the implication only; it is not substituted for a billiard source. Fix $\tau_0 \in (\tau_-, \tau_+)$ and, for large m, put

$$q_m(t) = m^{-2} \mathbf{1}_{[m\tau_0, m\tau_0 + e^{-m}]}(t), \qquad g_m = 0.$$

It is a nonnegative subprobability with the correct enclosing physical support, $\|Dq_m\|_{\mathrm{TV}} = 2m^{-2}$, and $\|q_m\|_1 = m^{-2}e^{-m}$. Every interval of a fixed positive length $2h$ satisfies $m^2(2h)^{-1} \int_I q_m \le e^{-m}/(2h)$, uniformly in its location. Nevertheless $m^2\|q_m\|_\infty = 1$. Moreover $|\widehat{q_m}(s)| \le m^{-2}e^{-m}$, so its sharp-frequency window integral satisfies

$$\frac{m^2}{2h} |\mathfrak{I}_S(I)| \le \frac{S e^{-m}}{\pi}.$$

It thus has vanishing normalized finite-band error at every polynomial bandwidth, as well as the common BV bound and vanishing normalized interval masses. Its local derivative charge, however, remains of order m^{-2} at the spike. This verifies that the new absolute inversion and a low- or polynomial-frequency comparison cannot alone be promoted to an essential-height assertion. A physical mechanism paying the actual local charge or the original same-roof shortage is still needed. The written fixed-band joint spectral, arithmetic and resonance arguments in historical native 84–87 are retained at their stated scope. The present bandwidth grows with the collision count and the reciprocal microscopic width; no new strong-operator estimate on that growing band has been proved. Nevertheless Proposition 78.7 evaluates the scalar analysis-window expression through the retained fixed-window law and leaves the complete normalized local charge as a sufficient closing condition on that route. The exact arithmetic reference, other observation classes, microscopic conditional denominators and long-time local inverse retain their own input-specific obligations.

79. Clearance-selected endpoint kernels and critical boundary tightness

The endpoint layers left by Theorem 73.1 cannot be discarded by their collision-area mass. We establish a different physical estimate: a clearance source at any flight has a small complete endpoint-momentum kernel after propagation along its longer side. This pays the first outgoing flight as well as the later flights. It subsequently gives a uniform tightness estimate for positive-clearance critical sectors. The norm of that conclusion and its distinction from the original roof-height norm are part of the statement.

For $1 \le j \le m-2$, the selected endpoint estimate below already follows from Theorem 71.3 in the two physical time directions. The additional first-image estimate retains a witness selected before the first true collision and thereby completes $j = 0$ and $j = m-1$. The longer-side argument is also present in the retained zero-clearance fan theorem 66.3; that theorem already pays its entire zero-level stratum at the original collision normalization. The new critical conclusion here concerns a positive clearance width, with the different norm explicitly displayed below.

All constants and curves are those of Section 71. In particular the original radius range, the actual first-hit map, the two impact signs and every unvisited disk are retained. Put

$$\begin{aligned} c_H &= \sin(\widehat{k}_0^{-2}), \qquad C_H = \max\{c_H^{-1}, (\pi/2)(1+\widehat{k}_0^{-1})^2\}, \\ D_{\mathrm{first}} &= C_H \,\frac{1+\tau_+/R_-}{t_*/R_+}\sqrt{1+U^2}, \qquad (79.1) \\ S_{\mathrm{first}} &= K_{\mathrm{c}} A_{\mathrm{hor}}, \qquad M_{\mathrm{first}} = D_{\mathrm{first}} J_0 L_{\mathrm{c}}/R_-, \\ C_{\mathrm{first}} &= \frac{J_f}{k_- t_*}\{\Gamma S_{\mathrm{first}}/\theta_{\mathrm{r}} + C_\ell M_{\mathrm{first}}\}, \\ C_{\mathrm{end}} &= \max\{K_{\mathrm{loc}}, C_{\mathrm{first}}, J_0 J_f E_{\mathrm{lc}}\}. \qquad (79.2) \end{aligned}$$

These constants are finite and independent of m, R, j, δ, a, v. The enclosing physical flight bound τ_+, rather than a sampled maximum, occurs in (79.1).

79.1. **Weighted first images of a selected horizontal source.**

Lemma 79.1 (The first selected flight retains its length)**.** *For a fixed horizontal source F_a, let its initial outgoing flight satisfy the original clearance event $E_{\delta,R}$, where $0 < \delta \le \delta_*$. A positive cover of its first physical image by homogeneous short labelled curves C carries the exact image densities ρ_C and satisfies*

$$\sum_C \|\rho_C\|_\infty \le S_{\mathrm{first}}, \qquad \sum_C \|\rho_C\|_\infty |C| \le M_{\mathrm{first}}\delta. \qquad (79.3)$$

The cover may include further initial states, but each of them follows its actual first collision. There is no reflection at a clearance foot.

Proof. Divide F_a into the J_0 short preliminary arcs of (71.6). On each arc, Lemma 71.2 covers its selected points by at most K_{c} disjoint initial intervals. The combined $d\alpha_0$ mass of all these intervals is at most $J_0 L_{\mathrm{c}}\delta/R_-$.

For each initial interval take its actual first-hit pieces and their first homogeneous images. Cut images into the same short lengths as in the proof of (71.7). An initial subinterval cannot increase the number of first-hit pieces on its preliminary arc, and on each first-hit piece the first-image momentum angle is strictly monotone. Consequently each target homogeneity strip is still met at most once. Applying the proof of the A_{hor} budget to at most K_{c} intervals per preliminary arc gives the first inequality in (79.3). Source labels distinguish any coincident target coordinate images.

For the second inequality use the exact density of Lemma 26.1,

$$\rho_C = \frac{d}{RA\sqrt{1+V_1^2}}, \qquad A = \sqrt{1-a^2} + \tau/R, \qquad d = \cos\varphi_1.$$

On a central homogeneous image, the ratio of the largest and smallest d is at most c_H^{-1}. On a signed strip of index $k \ge \widehat{k}_0$ it is at most

$$\frac{\sin(k^{-2})}{\sin((k+1)^{-2})} \le \frac{\pi}{2}\left(\frac{k+1}{k}\right)^2 \le \frac{\pi}{2}(1+\widehat{k}_0^{-1})^2.$$

Also $t_*/R_+ \le A \le 1 + \tau_+/R_-$ and $1 \le \sqrt{1+V_1^2} \le \sqrt{1+U^2}$. Thus every such image and each of its subcurves satisfies

$$\sup_C \rho_C \le D_{\mathrm{first}} \inf_C \rho_C, \qquad \|\rho_C\|_\infty |C| \le D_{\mathrm{first}} \int_C \rho_C \, d\ell_C.$$

The formulas hold on open regular pieces; taking the endpoint limits within one homogeneity strip gives the same inequalities. Their exact integrals sum to the $d\alpha_0$ mass of the initial cover. The latter intervals are disjoint after merging within each preliminary arc. The first-hit and homogeneous cuts have zero source measure, so summing proves the second inequality. The countable strip sum is legitimate by nonnegativity. No bounded distortion across different grazing strips is asserted. □

79.2. **The complete selected kernel.** Write $\mathcal{R}_{m,R}(a,v)$ for the unique regular physical roots in (26.16). At such a root put $u_x = |\partial_{\alpha_0} p_m(x)|^{-1}$ and define

$$D_{m,R;j,\delta}(a,v) = \sum_{\substack{x\in\mathcal{R}_{m,R}(a,v)\\ N_{\delta,R}(T_R^j x)\ge 1}} u_x, \qquad 0 \le j < m. \tag{79.4}$$

The root convention counts a physical state once, including states on artificial homogeneity or coordinate seams.

Theorem 79.2 (Clearance-selected endpoint-momentum kernel)**.** *For all $m \ge 1$, $R \in [R_-, R_+]$, $a, v \in (-1,1)$, $0 \le j < m$ and $0 < \delta \le \delta_*$,*

$$D_{m,R;j,\delta}(a,v) \le C_{\mathrm{end}}\{\delta + \theta_{\mathrm{r}}^{\max(j,m-1-j)}\}. \tag{79.5}$$

The corresponding selected pushforward of $d\alpha_0$ on F_a has this density bound. Therefore, with the original conditioning and any original Borel selector $0 \le H \le 1$,

$$(p_0,p_m)_{\#}(H\mathbf{1}_{E_{j,\delta,R}}\, d\nu_R^*) \le \frac{C_{\mathrm{end}}}{4\pi c_*}\{\delta + \theta_{\mathrm{r}}^{\max(j,m-1-j)}\}\, da\, dv. \tag{79.6}$$

All original roof, return, displacement, collision and observation tests may be retained by positive restriction. These inequalities are a coordinate disintegration of the real orbit law and do not make the momentum process Markov.

Proof. First prove the forward bound with exponent $m-1-j$. If $1 \le j < m$, apply Theorem 71.3 to the horizontal source with

$$h = j-1, \qquad n = m-j-1, \qquad l = h+n+2 = m.$$

Its inverse-derivative supremum sum is at most $K_{\mathrm{loc}}(\delta + \theta_{\mathrm{r}}^{m-j-1})$. On each source interval the terminal momentum is strictly monotone, so it contributes at most one root at a specified terminal value. The same sum bounds its selected density almost everywhere by one-dimensional coarea.

For $j = 0$ and $m \ge 2$, use Lemma 79.1. Propagate each first-image curve through exactly $m-2$ further actual collisions and then take the final collision without an infinite terminal homogeneity cut. The renewal inequality (71.3), weighted by $\|\rho_C\|_\infty$, gives a sum of incoming density suprema at most

$$\Gamma\theta_{\mathrm{r}}^{m-2}\sum_C \|\rho_C\|_\infty + C_\ell \sum_C \|\rho_C\|_\infty |C| \le \Gamma S_{\mathrm{first}}\theta_{\mathrm{r}}^{m-2} + C_\ell M_{\mathrm{first}}\delta.$$

The exact last-collision derivative cancellation (26.18) and at most J_f actual last-hit pieces multiply this bound by $J_f/(k_- t_*)$. Since $\theta_{\mathrm{r}}^{m-2} = \theta_{\mathrm{r}}^{-1}\theta_{\mathrm{r}}^{m-1}$, this is at most $C_{\mathrm{first}}(\delta + \theta_{\mathrm{r}}^{m-1})$. The generation count is $1 + (m-2) + 1 = m$. For $m = 1$, the unselected root sum is at most $J_0 J_f E_{\mathrm{lc}}$ by the first-hit count and its exact derivative lower bound. This proves the asserted forward density bound also in that case.

We justify the bound at every regular root value, including partition seams. Take any finite selected set of distinct regular roots at (a, v). A genuine width-δ witness has strict interior-foot and strict clearance inequalities; it persists in a sufficiently small regular neighborhood of each root. The inverse branches and positive weights are continuous. Their sum is bounded almost everywhere at nearby terminal values by the density estimate just proved, so it obeys the same bound at v. Exhausting finite root sets gives the complete series. A weak width-δ version, when needed, follows by using $\delta + \eta \le \delta_*$ and letting $\eta \downarrow 0$; at $\delta = \delta_*$ one can instead retain the strict convention of the theorem.

Now reverse each actual finite record. The collision reverser is $(\alpha, p) \mapsto (\alpha, -p)$, the selected flight has index $m-1-j$, and the two endpoint momenta become $(-v, -a)$. A flight of length τ is the same unoriented physical segment after reversal. For a designated unvisited disk its closest-foot coordinate changes from t_{foot} to $\tau - t_{\mathrm{foot}}$, so the strict interior-foot condition and its unsigned clearance are preserved. The two incident disks are exchanged and the same third disks remain unvisited on this flight. The original terminal disk has one definite lattice center; subtracting that center re-centers the reversed starting disk in the original fundamental cell uniquely. This bijection introduces no lattice or word multiplicity. Every original observation is transported on this same reversed orbit, rather than assumed invariant. Area preservation of the physical collision map gives equality of the original and reversed inverse root weights. Explicitly, on a regular branch write DT_R^m in the coordinates (α, p). Its determinant is one; the lower-left entry of its inverse has the opposite sign and the same magnitude as the original lower-left entry. The two momentum sign changes preserve that magnitude. Thus the reversed forward bound is $C_{\mathrm{end}}(\delta + \theta_{\mathrm{r}}^{j})$. Taking the smaller of the two bounds, using $0 < \theta_{\mathrm{r}} < 1$, proves (79.5). The two sides were evaluated on the same trajectory; no product of their probabilities was used.

Finally integrate the fibre inequality in a and restore the original positive selectors. The collision-coordinate measure is $d\alpha_0\, da/(4\pi)$ and the original section conditioning contributes exactly c_*^{-1}. This proves (79.6); it is not a rescaling by the selected source mass. □

79.3. **Uniform tightness of positive-clearance critical sectors.** Consider the physical one-copy critical sectors of Section 27, using the finite analytic side conditions justified in Lemma 66.1. All visited incidences at the limiting germ are positive; unvisited tangencies, finite original section cuts and corners are allowed. Keep the original fractions $\beta_y(x)$, $\beta_z(x)$ and actual critical coefficient $\mathfrak{J}(x)$. At a marked limiting flight count a witness of width δ when its strict auxiliary flight has an interior closest foot and clearance in $[0,\delta]$ from an unvisited disk. This definition is used only for its actual physical-side germ. It does not insert a collision at clearance zero.

Theorem 79.3 (Positive-clearance critical boundary tightness)**.** *For $0<\delta<\delta_*$, the complete one-copy family $\mathcal{F}_{m,j,\delta}(R)$ of those critical germs with at least one specified-flight witness obeys*

$$\sum_{x\in\mathcal{F}_{m,j,\delta}(R)} \mathfrak{J}(x)\beta_y(x) \le C_{\mathrm{crit}}\{\delta+\theta_{\mathrm{r}}^{\max(j,m-1-j)}\}, \qquad C_{\mathrm{crit}}=\frac{U_2C_{\mathrm{end}}}{2c_*}. \tag{79.7}$$

A germ with several witnesses is counted once in this sum. Original exact-record and critical-roof-value restrictions, and further weights in $[0,1]$, may be retained.

For the original clearance guard $\delta_{j,B}=2\varepsilon(B)q_0^{\min(j,m-1-j)/4}$ with $2\varepsilon(B)<\delta_*$, the leading critical coefficient of the union of all its clearance witnesses is at most

$$C_{\mathrm{crit}}\left\{\frac{4\varepsilon(B)}{1-q_0^{1/4}}+m\theta_{\mathrm{r}}^{(m-1)/2}\right\}. \tag{79.8}$$

Thus its unscaled complete critical coefficient is uniformly tight as the original guard is removed, after the collision limsup and the original radius and exact-label suprema.

Proof. Fix R,m,j and a finite set of distinct one-copy critical germs. Use the common small endpoint-momentum disk and their inverse branches from the proof of Theorem 27.4. On an actual regular side of a germ whose limiting witness has width at most δ, that witness has width less than $\delta+\eta$ for every fixed $\eta>0$ on a sufficiently small neighborhood. Its strict foot remains interior by the uniform standoff from the incident contacts. If the limiting clearance is zero, the actual unvisited side has positive clearance there. Choose $\eta>0$ with $\delta+\eta\le\delta_*$.

The physical-side inverse images are disjoint after the original one-copy merging. Use the unconditioned coordinate measure $d\alpha_0\,dp_0$ here. Its endpoint-momentum pushforward has density given by the regular fibre root sum (79.4), by one-dimensional coarea in α_0 followed by integration in p_0. Theorem 79.2 therefore gives almost everywhere in $z=(p_0,p_m)$

$$\sum_x u_x(z)\mathbf{1}_{E_x}(z)\le C_{\mathrm{end}}\{\delta+\eta+\theta_{\mathrm{r}}^{\max(j,m-1-j)}\}.$$

The finite analytic physical-side conditions have angular limits outside finitely many rays, by Lemma 66.1. Polar Fubini, continuity of the finite inverse weights and bounded convergence along good radii tending to zero therefore yield

$$\sum_x u_x\beta_z(x)\le C_{\mathrm{end}}\{\delta+\eta+\theta_{\mathrm{r}}^{\max(j,m-1-j)}\}.$$

Let $\eta\downarrow 0$, then exhaust finite germ families. These limits occur at the fixed physical R,m,j; their resulting bound has constants uniform in those parameters. Finally the exact angular comparison (27.6) and $\lambda_{\max}(G_x)/R^2\le U_2$ give (79.7). The only section-conditioning factor is the c_*^{-1} already present in that critical coarea comparison; none was used in the unconditioned kernel inequality. This passage controls actual one-sided coefficients, rather than assigning zero trace to a collision-area-null set.

For the original union, positivity bounds its one-copy coefficient by the sum of the m selected coefficients. There are at most two indices at each depth, so

$$\sum_{j=0}^{m-1}\delta_{j,B}\le\frac{4\varepsilon(B)}{1-q_0^{1/4}},\qquad\sum_{j=0}^{m-1}\theta_{\mathrm{r}}^{\max(j,m-1-j)}\le m\theta_{\mathrm{r}}^{(m-1)/2}.$$

Insert these finite-count bounds into (79.7). Complete positive sums precede all radius, label and roof-value restrictions. Taking the collision limsup first, and only then $B\to\infty$ along the original $\varepsilon(B)\to 0$, proves the stated unscaled tightness. □

The resulting gain and the exact limitation. Theorem 79.3 treats positive-width clearance witnesses at every flight, including both shallow endpoint layers. The previous zero-clearance coefficient payment is retained and is sharper on its own zero-level stratum. The new argument retains selected physical length through the first collision and then uses the longer genuine propagation direction. It supplies a uniform small-source estimate in the endpoint-momentum kernel and in the complete critical boundary coefficient norm.

It does not bound the original shortage in $m^2L^\infty(dt)$. In particular multiplying the first term of (79.8) by m^2 at a fixed B gives no vanishing bound. A leading critical coefficient is also not a collision-uniform estimate of the nonleading coarea remainder at a common original roof. Those remainders, the original same-pool endpoint-layer shortage, and the joint spectral, arithmetic and conditioning inputs remain as specified in the unchanged main theorem and dependency ledger. No neighborhood radius uniform over the full critical family has been inferred from finite-germ exhaustion.

80. The exact singular part of the physical roof derivative

The complete roof variation in Sections 74 and 76 is a finite measure. Finiteness alone does not describe its singular part. We now identify that part on the *entire* physical family, including every original section cut, unvisited boundary and critical entrance. The conclusion holds for the actual smooth-guard and C^1 trajectory tests of (76.8); those functions are not assumed subanalytic. The geometric decomposition is made before those weights are inserted.

The finite physical graph and its parameterized semialgebraicity are inherited from historical native164, not new assertions about the orbit model. Historical native40, native92 and native93 already extract and reconstruct finite-count singular germs for their specified subanalytic insertions. On that older constructible scalar subclass, their finite analytic preparation together with the retained BV bound already excludes a Cantor part; no new priority is claimed for that consequence. Historical native153 already proves the signed projected physical boundary current on its selected Morse charts, including the combination of coincident decision interfaces. That signed-current principle is also inherited. The new statement here is its full physical derivative decomposition for the current smooth observation class, with its exact physical coefficients and the complete collision-uniform variation budget. No definability of a new observation is silently added to the hypotheses.

Fix $m\ge 1$ and $R\in[9/20,47/100]$. Write

$$f_\ell=f_{\ell,m,R}^{w;a,b,\Psi},\qquad D=\mathcal{V}_m^w(a,b,\Psi),\qquad \ell=(n,k),$$

with precisely the choices and norms in Theorem 76.2. In particular $w=1$ or the original W^ε, with $\varepsilon>0$ fixed, and there is exactly one factor c_*^{-1}. There are finitely many attainable labels at this fixed collision count. They are not pooled below.

80.1. **Pushforward along an actual boundary side.**

Lemma 80.1 (A common finite family of physical roof-side maps)**.** *Fix the true collision count $m \geq 1$. There is a finite integer N_m and, for each original radius $R \in I$, a finite set $\mathcal{Z}_{m,R} \subset [m\tau_-, m\tau_+]$, with*

$$\#\mathcal{Z}_{m,R} \leq N_m,$$

having the following property. In the signed integration-by-parts formula for any original exact-record roof density in Theorems 69.2 and 74.1, the regular physical boundary terms can be written as a finite sum

$$(t_\gamma)_*(b_\gamma(r)\,dr).$$

Here r is a rational coordinate on a Gauss-direction arc, $b_\gamma \in L^1$ is the signed or complex actual *one-copy flux, and t_γ is the original physical roof on that side. Each t_γ is C^1 and either constant, with value in $\mathcal{Z}_{m,R}$, or has derivative nowhere zero on its open parameter interval. The finitely many discarded parameter points have zero flux. The positive-visited critical incoming contributions are also supported on $\mathcal{Z}_{m,R}$.*

The same set works simultaneously for every attainable exact label, the marked phases, either $w = 1$ or the original smooth guard $w = W^\varepsilon$, and the bounded C^1 endpoint tests for which Theorem 76.2 proves finite variation. Neither a positive incidence cutoff nor a replacement section is used. The set is geometric: it is independent of ε, of the particular smooth function Θ, and of these test functions or phases. No estimate uniform in m is asserted for N_m.

Proof. We spell out the physical construction because local analyticity alone would not give the asserted finite family uniformly in R. The finite semialgebraic first-hit graph already proved in the preserved native module 164 is the input here. We recall its defining variables, rather than add a symbolic replacement model.

The uniform physical horizon gives a finite set of possible next-center increments

$$\mathcal{A} = \{z \in \Lambda \setminus \{0\} : |z| \leq \tau_+ + 2R_+ + 1\}.$$

For a word $C_0 = 0, \ldots, C_m$ with increments in $\mathcal{A}$, use real contact points $Q_i = C_i + Rn_i$, unit normals n_i, unit velocities v_i, and real lengths τ_i. The equations are

$$Q_{i+1} = Q_i + \tau_i v_i, \qquad v_i = v_{i-1} - 2(v_{i-1} \cdot n_i)n_i, \qquad v_{i-1} \cdot n_i < 0$$

at every visited reflection, together with the original outgoing sign at the initial collision and $\tau_- \leq \tau_i \leq \tau_+$. To impose the first-hit rule, exclude an interior intersection of the *actual* flight with every candidate disk preceding its designated endpoint. A disk that can meet that segment lies within $\tau_+ + 2R_+$ of its starting center. Its squared distance to the segment is determined by the two endpoint cases and by the interior orthogonal-foot case. These are finite polynomial inequalities after the positive projection denominator is retained. The current disk is treated with its actual outgoing sign. No ray is restarted at a line foot.

The same finite enlargement gives every candidate for the original least third-disk clearance. A nearest-neighbor center other than the terminal center has distance at most $1 + R$ from the initial contact, so its distance from the entire segment is at most $1 + R$. The least third-disk *clearance* is therefore at most 1. A minimizing center is consequently within $\tau_+ + 2R_+ + 1$ of the starting center. Square roots are encoded by their nonnegative root and squared equation. Thus all physical incidences and original third-disk clearances are semialgebraic on this graph.

Use finitely many rational charts for each normal circle, with one-copy ownership and the finitely many seam sides retained until cancellation. For a fixed chart rotation the charts have the form

$$n(x) = \mathrm{Rot}_{\alpha_*}\left(\frac{1 - x^2}{1 + x^2}, \frac{2x}{1 + x^2}\right), \qquad |x| < 1.$$

The moving center normal of each original rectangle is semialgebraic in R, because its half-angle is the specified unique root of the original polynomial in q_R. An angular side test is $n \cdot n_* > \cos h$, with its actual fixed half-width h, and its companion momentum test is linear. Fixed real numbers such as $\cos h_0$ are allowed as coefficients. Retain $\eta_0 = \eta_m = 1$, $\sum_{i=0}^{m-1} \eta_i = n$, and $C_m = k$ exactly. There are finitely many words and labels at this fixed m. All these statements concern the deterministic physical graph and the original roof $L = \sum_i \tau_i$.

On a regular endpoint sheet, write $q = (\alpha_0, \alpha_m)$, $F = L$, and $g = \nabla_q F = R(-p_0, p_m)$. The exact endpoint and optical identities make F, g, and their derivatives semialgebraic in rational normal charts: differentiation in α is multiplication of the chart derivative by $(1 + x^2)/2$. Away from $g = 0$, adjoin the positive variable $|g|$ and use finitely many rational charts

$$e = \frac{g}{|g|} = \mathrm{Rot}_{\theta_*}\left(\frac{1 - r^2}{1 + r^2}, \frac{2r}{1 + r^2}\right)$$

for its true Gauss direction. The map $q \mapsto (F, e)$ has the nonzero Jacobian χ of the retained optical coarea proof. Consequently its regular fibers are discrete; since they are semialgebraic, they are finite. The same statement holds for the endpoint projection on a fixed regular word, by its nonzero optical twist. Finite semialgebraic cell decomposition therefore provides finitely many genuine inverse sheets, retaining their physical multiplicities. It does not assign a nonphysical orbit to a missing sheet.

Now take the complete one-sided graph of these sheets in variables (R, r, t), retaining the side identities before any observation weight is inserted. Every closure and boundary in this construction is taken *fiberwise at the fixed radius*, before forming the parameterized family. For example, membership of (r, t) in the closure of the R-fiber is expressed by

$$\forall a > 0\ \exists (r', t') \quad (R, r', t') \text{ belongs to that sheet}, \qquad (r - r')^2 + (t - t')^2 < a^2.$$

Quantifier elimination makes this fiberwise closure semialgebraic. The fiberwise relative boundary is obtained by the corresponding Boolean operations. Thus limits from another radius never create physical sides at an exceptional radius. At every fixed R the regular image is open in (r, t), so its boundary has dimension at most one. These boundaries include the original unvisited constraints, caps used in the positive majorant, every actual section side at every collision index, and their finite intersections. They are semialgebraic, because closure, projection and finite Boolean operations preserve semialgebraicity. At positive visited incidences the critical set has $g = 0$ and positive endpoint Hessian. Its points are isolated on each fixed-R sheet. This is a zero-dimensional semialgebraic fiber and hence finite. Over each such critical point the Gauss-direction closure records its actual incoming sectors at the single value $t = F(q)$. We keep those sectors, including unvisited-tangency corners; they are not declared null because the physical source point is null.

Apply C^1 cylindrical cell decomposition with R ordered before r. It is applied to the finite geometric side graphs, their relative closures, and their projection-critical sets. A side with nonconstant Gauss projection is then expressed on finitely many cells as $t = t_\gamma(R, r)$. For a fixed radius, further refine by the semialgebraic sets where $\partial_r t_\gamma$ is zero, positive, or negative. On an interval in the zero set the fundamental theorem of calculus makes t_γ constant. On each remaining interval the derivative never vanishes. A side whose Gauss projection is constant has zero $d\theta$-flux on its regular part and does not create an omitted charge. Its intersections and the finitely many point cells also have zero such flux. The constant values and the critical roofs form $\mathcal{Z}_{m,R}$.

This use of finite decomposition is precisely the finite definable-family assertion in the cell and smooth-cell decomposition theorems, for example [10, Theorems 2.9–2.10 and 6.6–6.7]. Their hypotheses apply to the semialgebraic graphs just constructed. The finite cylindrical

decomposition has finitely many cells across the entire original radius interval, including its exceptional parameter cells. Thus the number of its fixed-radius interval pieces, and hence the number of their constant values and critical roofs, has a finite common upper bound N_m. These theorems provide no upper bound independent of the formula size, which grows with m.

The inherited physical integration-by-parts formula on a roof interval is the interior derivative plus its signed left and right traces. For a regular side in the displayed chart, $d\theta = 2\,dr/(1+r^2)$ up to its chosen orientation. Accordingly $b_\gamma(r)$ is the signed trace, including the difference of the actual two incident record weights, times this coordinate factor. Coincident genuine arcs have their incident values combined before assigning the flux. The physical trace and section-charge theorems give its L^1 bound; they do not require a bounded trace at each parameter endpoint. Artificial angular charts, length cuts and homogeneity cuts cancel between their two incident copies before this representation. At a visited grazing limit the optical trace is zero by Lemma 67.3, including the simultaneous critical limit at its stated scope. Thus it creates no additional side measure. The positive visited critical fan is the separately retained incoming measure described above.

Finally none of the smooth observation weights changes a geometric sheet or introduces a jump side. The original W^ε is continuous on each physical sheet, including where two branches of a least clearance exchange; its interior derivative is the one already controlled in the guarded BV proof. A bounded C^1 endpoint test similarly adds only its proved interior derivative and multiplies existing traces. Exact labels and discrete twists are constant on the original section chambers. They therefore change b_γ and the critical sector coefficients but not the finite side maps or their constant values. This proves the claimed common set and its scope. An arbitrary discontinuous Borel trajectory selector does not have this conclusion. □

Lemma 80.2 (An integrable weight on a geometric roof side)**.** *Let $J \subset \mathbb{R}$ be an open interval, let $\gamma : J \to \mathbb{R}$ be C^1, and let $b \in L^1(J;\mathbb{C})$. If γ' is nowhere zero, then*

$$\gamma_*(b(r)\,dr) = h(t)\,dt, \qquad h(t) = \mathbf{1}_{\gamma(J)}(t)\frac{b(\gamma^{-1}(t))}{|\gamma'(\gamma^{-1}(t))|}, \qquad \|h\|_1 = \|b\|_1.$$

If $\gamma \equiv \tau$, the same pushforward is $(\int_J b(r)\,dr)\delta_\tau$. Consequently a finite collection of the geometric sides in Lemma 80.1, with arbitrary integrable weights, has a pushforward which is the sum of an L^1 density and a finite atomic measure. A zero value of γ' at an isolated subdivision point creates no atom.

Proof. On each compact subinterval of J, a continuous nowhere-vanishing derivative has constant sign and a positive lower bound in absolute value. The inverse-function theorem and the usual substitution formula therefore give the first identity there, including the displayed absolute denominator. Exhaust J by compact subintervals. For $|b|$ this passage is monotone; for b it follows by domination with $|b|$. This proves the full identity and norm equality without a lower bound for $|\gamma'|$ near an endpoint of J. For a constant map the pushforward identity is its definition. The geometric lemma provides a finite subdivision into these two cases and finitely many points. Points have zero $b(r)\,dr$ mass. Summing the absolutely integrable pieces proves the assertion. In particular, a nonconstant side may produce an unbounded but integrable density near a turning roof; it does not produce a singular-continuous measure there. □

80.2. **The signed physical coefficient, before absolute values.** Use the one-copy physical integration-by-parts construction of Theorem 76.2. On each open regular roof–angle chamber the actual coefficient is

$$w(t,\theta)A(t,\theta)V(t,\theta)v_\ell(t,\theta), \qquad V = a(x)b(T_R^m x)e^{i\Psi_m(x)},$$

where v_ℓ is the original section/record indicator and $A = \omega/(4\pi c_*\chi)$. This construction differentiates the complete physical density. It does not differentiate an arbitrary additional Borel trajectory selector.

For each regular geometric side γ in a rational Gauss-angle chart, let $r \in J_\gamma$ be its parameter, let $t_\gamma(r)$ be its actual roof, and let $b_{\ell,\gamma}(r)\,dr$ be its *signed* coefficient in that integration-by-parts formula. Explicitly, on a chamber interface with increasing-roof traces C_ℓ^- and C_ℓ^+,

$$b_{\ell,\gamma}(r) = (C_\ell^+(r) - C_\ell^-(r))|d\theta/dr|. \tag{80.1}$$

On the exterior side of a physical source the incident trace C_ℓ is zero; the resulting boundary coefficient need not vanish. At a regular section cut the continuous factor wAV multiplies the actual difference of the label indicators; an internal visit is not charged as two visits of the same label. At an entrance to a physical interval the coefficient is positive for $w = V = 1$; at an exit it is negative. For complex tests the signed formula still applies. Artificial chart seams are canceled before (80.1) is formed. Coincident real constraints describe one interface with its two actual traces, not extra copies of that interface. A side with constant Gauss angle has zero angular charge and is omitted.

Let $\mathcal{K}_{m,R}$ be the one-copy positive-visited critical germs of the retained physical action. Write t_x for a germ's critical roof and G_x for its endpoint Hessian. In the coordinates $y = G_x^{1/2}(q - q_x)$ let $\beta_{\ell,y}(x)$ be the angular fraction of its actual physical tangent sectors with the exact record ℓ and both original endpoint tests. It is zero when the record is absent. At that germ the smooth weights have the one-sided limits w_x, V_x along its designated physical word. The incoming critical coefficient is

$$c_{\ell,x} = w_x V_x\, \mathfrak{J}(x)\beta_{\ell,y}(x), \qquad \mathfrak{J}(x) = \frac{\omega(x)}{2c_*\sqrt{\det G_x}}. \tag{80.2}$$

Formula (80.2) uses the retained one-sided sector theorem. It does not declare an unvisited tangency to be a regular collision. A critical limit with a visited grazing contact has zero coarea trace by Lemma 67.3, even when several constraints meet. The values V_x are limits of the actual smooth observations on that same word. No time-reversal invariance of the section or of those observations is required.

Let $\mathcal{S}_{m,R} = \mathcal{Z}_{m,R}$ be the finite set supplied by Lemma 80.1: it contains the critical roofs and the roof values of all constant-roof regular sides. It depends only on the physical geometry and the fixed exact section at this collision count, not on a, b, Ψ, Θ, or ε. For $\tau \in \mathcal{S}_{m,R}$ set

$$a_\ell(\tau) = \sum_{\substack{x\in\mathcal{K}_{m,R}\\ t_x=\tau}} c_{\ell,x} + \sum_{\gamma:\ t_\gamma\equiv\tau} \int_{J_\gamma} b_{\ell,\gamma}(r)\,dr. \tag{80.3}$$

All terms at the *same physical roof and same exact record* are added with their signs before an absolute value is taken. Coincident roofs in distinct records are not combined. A zero value of $a_\ell(\tau)$ is allowed. In particular critical coefficients alone need not be the full atomic part.

Theorem 80.3 (No singular-continuous derivative for the full smooth class)**.** *For every specified physical parameter and test there are $d_\ell \in L^1(\mathbb{R};\mathbb{C})$ such that, exactly as finite distributional measures,*

$$Df_\ell = d_\ell(t)\,dt + \sum_{\tau\in\mathcal{S}_{m,R}} a_\ell(\tau)\delta_\tau. \tag{80.4}$$

There is no singular-continuous part. The density d_ℓ is the sum of the actual regular interior derivative and the one-dimensional coarea densities of all nonconstant physical roof sides. The

atomic coefficients are exactly (80.3). *Moreover*

$$\sum_{\ell}\left(\|d_\ell\|_1+\sum_{\tau\in\mathcal{S}_{m,R}}|a_\ell(\tau)|\right)=\sum_{\ell}\|Df_\ell\|_{\mathrm{TV}}\le D. \tag{80.5}$$

In particular the bound retains all words, all original labels and the original conditioning, with no label-count factor. The bound is uniform in m,R when the norms in Theorem 76.2 are uniformly bounded. Neither the number of sides and atoms nor their separation is claimed uniform as $m\to\infty$.

The canonical BV traces satisfy

$$f_\ell(\tau+)-f_\ell(\tau-)=a_\ell(\tau). \tag{80.6}$$

On every interval avoiding $\mathcal{S}_{m,R}$ the density has an absolutely continuous representative. These statements do not make the entire density differentiable at a jump, nor turn a finite-count geometric assertion into a collision-normalized pointwise estimate.

Proof. The proof of Theorem 76.2 constructs Df_ℓ by integrating the regular chamber derivative and all actual one-sided charges. Its full positive source bound is finite, including the inverse-gradient multiplier and every endpoint/section charge. Consequently it is legitimate to use that signed source identity, and then project it to the physical roof, before taking variation.

The regular part is absolutely continuous in t: at fixed Gauss angle its signed density is the actual t derivative of wAV on each chamber. Its full integral of the absolute value is finite by that same proof. Fubini therefore gives an L^1 roof density for its sum. This argument uses the actual regular derivative of the smooth observations and guard and needs no definability of those functions. In particular the minimum in the least-clearance definition is handled by its retained absolutely continuous one-dimensional representative; an artificial nearest-center switch is not a new physical endpoint.

The complete noncritical geometric side collection has the finite description of Lemma 80.1. Its signed weights satisfy $\sum_{\ell,\gamma}\int|b_{\ell,\gamma}|<\infty$ by the complete boundary accounting. On a nonconstant monotone side apply Lemma 80.2; its projected derivative is absolutely continuous, including the one-sided approach to a turning endpoint. On a constant-roof side the same lemma gives precisely the second sum in (80.3). Endpoints of those regular sides have zero angular integration measure. Their critical limits have instead already been assigned to the incoming critical construction; this avoids counting a critical atom twice.

At a positive-visited critical germ the inherited local sector coarea theorem and continuity of the weights give exactly (80.2). The positive endpoint Hessian makes the critical configurations isolated on each regular sheet; the finite semialgebraic description in Lemma 80.1 makes their fixed-count family finite. All visited-grazing traces are zero as actual one-sided traces; collision-area nullness is not substituted for this fact. The exhaustive classification in Lemma 67.3 and the exact section accounting in Theorem 76.2 leave no further source measure. In particular no singular-continuous measure can remain after these pushforwards. This proves (80.4) and its coefficient formula.

The absolutely continuous and atomic parts in (80.4) are mutually singular. The total variation of their sum is therefore the sum of their variation masses, also for complex measures. After all same-roof cancellations in (80.3), this gives the equality in (80.5). Its inequality is the existing complete variation theorem, not a separate sum of per-label constants.

Finally, for fixed ℓ define a representative by integrating the right side of (80.4) from any point to the left of its compact physical support. It has that same distributional derivative and equals f_ℓ almost everywhere: their difference has zero distributional derivative and vanishes on a left

exterior interval. The L^1 primitive is continuous, and an atomic primitive has the jump equal to its coefficient. This proves (80.6) and absolute continuity away from the finite atom set. □

80.3. **The normalized jump constraint for the original raw theorem.** Take $w = 1$, $V = 1$, and the original density $p_{n,R}(k, m, t)$. Let

$$E_m(M) = \sup_{R,n,k} \operatorname*{ess\,sup}_{t \in \mathcal{C}_{m,R,n,k}(M)} |m^2 p_{n,R}(k, m, t) - \mathcal{L}_{m,R}(k_1, k_2, t, n)|.$$

The full retained arithmetic reference is continuous in the physical roof at fixed m, R, n, k; every continued near-unit branch remains in it. Let

$$\mathcal{S}^{\mathrm{cen}}_{m,R,\ell}(M) = \mathcal{S}_{m,R} \cap \operatorname{int} \mathcal{C}_{m,R,n,k}(M).$$

Only this interior condition permits comparison with both traces inside the displayed central domain; an isolated endpoint of that domain is not silently treated as an interior roof.

Proposition 80.4 (Necessary full-label central jump payment)**.** *For every finite collision count,*

$$m^2 |a_\ell(\tau)| \le 2E_m(M) \quad (\tau \in \mathcal{S}^{\mathrm{cen}}_{m,R,\ell}(M)), \tag{80.7}$$

$$\sup_R m^2 \sum_\ell \sum_{\tau \in \mathcal{S}^{\mathrm{cen}}_{m,R,\ell}(M)} |a_\ell(\tau)|^2 \le 2D^1_{\mathrm{all}} E_m(M). \tag{80.8}$$

Hence the original raw theorem implies that both left sides tend to zero as $m \to \infty$. These conclusions are necessary tests of that theorem. Neither one is asserted sufficient for it.

Proof. Fix a radius, label and interior central roof τ. The definition of the essential supremum bounds the normalized error at almost every roof on each side of τ in a small central interval. Choose Lebesgue points approaching from those two sides. The BV traces exist, and the retained reference has the same continuous limit from each side. Passing to the two limits gives

$$|m^2 p(\tau+) - \mathcal{L}(\tau)| \le E_m(M), \qquad |m^2 p(\tau-) - \mathcal{L}(\tau)| \le E_m(M).$$

Subtract and use (80.6) to prove (80.7). This operation is performed at each fixed radius and uses no common exceptional roof set across all radii.

Multiply the pointwise coefficient bound by $|a_\ell(\tau)|$, then sum over all the specified physical central atoms and labels. By (80.5) with $w = V = 1$ this gives

$$m^2 \sum_{\ell,\tau:\,\mathrm{central}} |a_\ell(\tau)|^2 \le 2E_m(M) \sum_{\ell,\tau} |a_\ell(\tau)| \le 2D^1_{\mathrm{all}} E_m(M).$$

The same constant holds at every radius, proving the asserted supremum. There is no multiplication by the number of labels or physical words. If $E_m(M)$ tends to zero, the two conclusions follow in the original collision order. □

Scope. The theorem gives a common signed physical formula on the full current smooth observation class, whose weights need not be tame, and ties all its singular derivative mass to actual constant-roof sides and critical entrances. It does not suppress its positive critical trace, negative exit charge or sharp integrable turning density. In particular the complete normalized local derivative charge in (78.15) is still not proved small. The original same-roof shortage in (73.12) also retains its exact source, comparison pool and ordered limits. All previous directed-current, paired-flux and critical-orbit theorems remain available with their original stronger or different conclusions. The present structural result is not presented as a rediscovery of their mechanisms.

81. The exact jump spectrum of the full marked physical roof

The absence of a singular-continuous derivative in Section 80 identifies the entire nondecaying mean-square part of the differentiated physical roof transform. The result keeps the two displacement frequencies and the occupation frequency; it does not average over collision counts or change a conditioning event. It is an exact identity at every finite count, with a collision-uniform bound and a necessary consequence of the original raw limit. Its high-frequency limit is not interchanged with the collision limit.

Use the source and tests of (76.8). Write $\ell = (k_1, k_2, n)$, $f_\ell = f^{w;a,b,\Psi}_{(n,k),m,R}$, $D = \mathcal{V}^w_m(a, b, \Psi)$ and $A_V = a_0 b_0$. At fixed m, R only finitely many labels are attainable. By Theorem 80.3,

$$Df_\ell = d_\ell(t)\,dt + \sum_{u\in\mathcal{S}_{m,R}} a_\ell(u)\delta_u, \qquad a_\ell(u) = f_\ell(u+) - f_\ell(u-), \tag{81.1}$$

where $d_\ell \in L^1$ and $\mathcal{S}_{m,R}$ is finite. Zero coefficients are permitted. The geometric support is not asserted uniformly separated. All coincident contributions to a given (ℓ, u) have already been combined with their signs. In particular

$$\sum_\ell \left(\|d_\ell\|_1 + \sum_u |a_\ell(u)| \right) \le D. \tag{81.2}$$

This is the existing complete all-label variation budget applied to the mutually singular parts of its exact derivative.

Use the full marked transform $\mathcal{F}(\vartheta, s) = \mathcal{F}^{w,V}_{m,R}(\vartheta, s)$ of (78.1), with $\vartheta \in [-\pi, \pi]^3$. For $S > 0$ define

$$\mathcal{A}(\vartheta, s) = \sum_{\ell,u} e^{i\vartheta\cdot\ell + isu} a_\ell(u), \qquad \mathcal{G}(\vartheta, s) = \sum_\ell e^{i\vartheta\cdot\ell}\widehat{d}_\ell(s), \tag{81.3}$$

$$\mathcal{J}_{m,R}(w, V) = \sum_{\ell,u} |a_\ell(u)|^2, \qquad \mathcal{W}_S = \frac{1}{2S(2\pi)^3}\int_{-S}^{S}\int_{[-\pi,\pi]^3} s^2 |\mathcal{F}(\vartheta, s)|^2\,d\vartheta\,ds. \tag{81.4}$$

The convention is $\widehat{f}(s) = \int e^{ist} f(t)\,dt$.

Theorem 81.1 (All-label physical jump spectrum)**.** *For every actual $m \ge 1$, $R \in [R_-, R_+]$, original specified weight and tests,*

$$-is\mathcal{F}(\vartheta, s) = \mathcal{A}(\vartheta, s) + \mathcal{G}(\vartheta, s), \qquad \lim_{S\to\infty} \mathcal{W}_S = \mathcal{J}_{m,R}(w, V) \le D^2. \tag{81.5}$$

The first identity holds for every real s and full-torus frequency. At fixed m, R,

$$\sup_\vartheta |s\mathcal{F}(\vartheta, s) - i\mathcal{A}(\vartheta, s)| \longrightarrow 0 \qquad (|s| \to \infty). \tag{81.6}$$

For a precise finite-frequency error, put

$$\mathcal{Q}_S = \frac{1}{2S}\int_{-S}^{S}\sum_\ell |\widehat{d}_\ell(s)|^2\,ds, \tag{81.7}$$

$$\mathcal{C}_S = \sum_\ell \sum_{\substack{u,v\in\mathcal{S}_{m,R}\\ u\ne v}} |a_\ell(u)a_\ell(v)| \min\left\{1, \frac{1}{S|u - v|}\right\}. \tag{81.8}$$

Then

$$|\mathcal{W}_S - \mathcal{J}_{m,R}(w, V)| \le \mathcal{C}_S + 2D\sqrt{\mathcal{Q}_S} + \mathcal{Q}_S. \tag{81.9}$$

Both $\mathcal{Q}_S$ and $\mathcal{C}_S$ tend to zero for each fixed m, R. No positive uniform lower bound for distinct roof gaps, or rate uniform in m, R, is asserted.

Proof. Compact support and distributional integration by parts give

$$-is\widehat{f}_\ell(s) = \widehat{d}_\ell(s) + \sum_u a_\ell(u)e^{isu}.$$

At $s = 0$ this is the vanishing total mass of a compactly supported derivative. Summing the finite label set gives the first identity, with the displayed sign.

For $g \in L^1(\mathbb{R})$, approximate g in L^1 by finite linear combinations of bounded interval indicators. Their transforms tend to zero by direct integration, and the transform error is at most the L^1 error, uniformly in frequency. Thus $\widehat{g}(s) \to 0$. Apply this to every d_ℓ. The finite label sum yields $\sup_\vartheta |\mathcal{G}(\vartheta, s)| \le \sum_\ell |\widehat{d}_\ell(s)| \to 0$, which proves (81.6).

Full-torus orthogonality, before the roof-frequency average, gives exactly

$$\mathcal{W}_S = \frac{1}{2S}\int_{-S}^{S} \sum_\ell \left| \widehat{d}_\ell(s) + \sum_u a_\ell(u)e^{isu} \right|^2 ds. \tag{81.10}$$

No factor counting labels is introduced. The purely atomic part has mean

$$\mathcal{J}_{m,R}(w, V) + \sum_\ell \sum_{u \ne v} a_\ell(u)\overline{a_\ell(v)}\frac{\sin(S(u-v))}{S(u-v)}. \tag{81.11}$$

The second term has modulus at most $\mathcal{C}_S$. The diagonal is the square of the combined coefficient at each coincident roof, not the sum of squares of an artificial partition.

At every s,

$$\sum_\ell \left| \sum_u a_\ell(u)e^{isu} \right|^2 \le \left(\sum_{\ell,u} |a_\ell(u)| \right)^2 \le D^2.$$

Cauchy–Schwarz in the frequency average bounds the mixed term in (81.10) by $2D\sqrt{\mathcal{Q}_S}$; the remaining term is $\mathcal{Q}_S$. This proves (81.9). The finite cross sum tends to zero. The function $\sum_\ell |\widehat{d}_\ell(s)|^2$ is bounded by D^2 and tends to zero at both ends of the line, so its symmetric averages tend to zero. The claimed limit and upper bound follow. All operations occur at the specified finite count and radius, after including the entire physical source. □

Remark 81.2 (Coalescing roofs and the limit order). The cross term (81.11) retains coalescing roof values. For example, the derivative $\delta_0 - \delta_\delta$ of $\mathbf{1}_{(0,\delta)}$ has atomic energy 2 for every $\delta > 0$, but finite-frequency energy $2 - 2\sin(S\delta)/(S\delta)$, tending to zero as $\delta \downarrow 0$ with S fixed. This analytical check is not a substitute billiard model. A uniform variation bound alone does not justify moving the frequency limit through a radius supremum or collision limit. Nor does the $o(1/|s|)$ remainder assert absolute integrability.

81.1. **A necessary central spectrum at the original normalization.** Return to $p_\ell = p_{n,R}(k, m, \cdot)$ and $D_0 = D^1_{\mathrm{all}}$. Fix $M_1 > M > 0$ and a C^1 function $0 \le \psi \le 1$ on $[0, \infty)$ equal to one on $[0, M^2]$ and zero on $[M_1^2, \infty)$. Define the analytical roof test

$$\chi_{\ell,m,R}(t) = \psi\left(\left| \frac{(k_1, k_2, t, n) - m\mu_R}{\sqrt{m}} \right|^2 \right), \qquad h_\ell(t) = \chi_{\ell,m,R}(t)p_\ell(t). \tag{81.12}$$

This is a bounded observation of the original physical record; it does not replace its entrance law, terminal test or microscopic interval. Its choice is fixed before the collision limit. With the exact jumps $a_\ell(u)$ of p_ℓ, put

$$\mathcal{J}^{\chi}_{m,R} = \sum_{\ell,u} |\chi_{\ell,m,R}(u)a_\ell(u)|^2.$$

The same Fourier calculation as in the proof of Theorem 81.1 gives the energy limit $\mathcal{J}^{\chi}_{m,R}$ for the marked transform of (h_ℓ). This is justified by its following product decomposition, rather than by classifying the label-dependent cutoff as an endpoint test in (76.8). The BV product rule gives

$$Dh_\ell = (\chi_{\ell,m,R} d_\ell + \chi'_{\ell,m,R} p_\ell)\, dt + \sum_u \chi_{\ell,m,R}(u) a_\ell(u) \delta_u.$$

Its summed variation is at most $D_0 + 2M_1\|\psi'\|_\infty/\sqrt{m}$, since $|\chi'| \le 2M_1\|\psi'\|_\infty/\sqrt{m}$ and $\sum_\ell \|p_\ell\|_1 \le 1$.

Theorem 81.3 (The raw target forces a complete normalized jump spectrum)**.** *Let $E_m(M_1)$ be the raw error in* (1.5) *at central parameter M_1, before taking its limit. For every m,*

$$\sup_R m^2 \mathcal{J}^{\chi}_{m,R} \le 2D_0 E_m(M_1). \tag{81.13}$$

Consequently the original raw target implies

$$\lim_{m\to\infty} \sup_R m^2 \mathcal{J}^{\chi}_{m,R} = 0. \tag{81.14}$$

This is a necessary condition, not a sufficient one.

Proof. If $\chi_{\ell,m,R}(u) \neq 0$, u is interior to the larger central roof interval. The retained finite-peak reference is continuous there, including all near-unit branches. Approach u from both sides by Lebesgue points in that interval. The definition of $E_m(M_1)$ and the BV traces give

$$|m^2 p_\ell(u\pm) - \mathcal{L}_{m,R}(k_1, k_2, u, n)| \le E_m(M_1).$$

Hence $m^2|a_\ell(u)| \le 2E_m(M_1)$, equivalently the physical trace argument of Proposition 80.4. Multiply by $|a_\ell(u)|\chi_{\ell,m,R}(u)^2$ and sum all labels and roofs. The complete budget $\sum_{\ell,u} |a_\ell(u)| \le D_0$ proves (81.13). The target at M_1 proves the limit. The allowable null set may depend on R. The frequency limit identifying $\mathcal{J}^{\chi}$ was taken at fixed m, R; it has not been passed through a supremum or collision limit. □

81.2. **The inherited zero-clearance payment in this signed spectrum.** Take the one-copy union of positive-visited critical germs with at least one zero-clearance unvisited-disk witness covered by Theorem 66.3. Unite actual incident sectors first. When a germ has several witnesses, assign it once by a fixed ordering of its finite flight indices, nearby centers and signs; then split by its exact-label sectors. These sectors partition each physical fan, so the sum over labels introduces no extra multiplicity. This is an enumeration of the actual source, not a change of the source or event. At each germ include its actual sector and continuous bounded test value. Let $b_\ell(u)$ be this part of the atomic coefficient in (81.1), grouped at the same exact label and roof. For the positive unweighted source these incoming coefficients are nonnegative; tested coefficients can be complex. They need not be the entire atom at that roof.

Corollary 81.4 (Paid fans with every roof coincidence retained)**.** *With $\sigma = 47/59$ and the inherited C_{pc}, put $B_m = A_V C_{\mathrm{pc}} m \sigma^{(m-1)/2}$. Then*

$$\begin{aligned} \sum_{\ell,u} |b_\ell(u)| &\le B_m, \\ \left| \sum_{\ell,u} |a_\ell(u)|^2 - \sum_{\ell,u} |a_\ell(u) - b_\ell(u)|^2 \right| &\le 2DB_m + B_m^2. \end{aligned} \tag{81.15}$$

For fixed original guard width and fixed test norms, the last bound vanishes after multiplication by any prescribed power of m, uniformly in R. The same conclusion holds with the central tests (81.12)*. This transfers the already proved zero-clearance fan payment to the new signed spectrum.*

It makes no claim about the positive-clearance critical sources, other constant-roof side charges, or the absolutely continuous derivative.

Proof. The positive fan bound (66.4) includes both impact signs, every actual marked disk and visited word, and the original finite section, record and roof restrictions. The once-only union is bounded by the positive sum over all witnesses. Sum over $j = 0, \dots, m-1$ and use $\max(j, m-1-j) \geq (m-1)/2$. This gives $C_{\mathrm{pc}} m \sigma^{(m-1)/2}$. The guard is at most one, and the continuous test multiplier has modulus at most A_V at the fixed positive-visited germ. Combining true labels and coincident roofs before applying triangle inequalities gives the first bound.

For arbitrary complex vectors,

$$\bigl|\|a\|_2^2 - \|a-b\|_2^2\bigr| \leq 2\|a\|_2\|b\|_2 + \|b\|_2^2.$$

Use $\|a\|_2 \leq \|a\|_1 \leq D$ and $\|b\|_2 \leq \|b\|_1 \leq B_m$. Multiplication of coefficients by $0 \leq \chi \leq 1$ decreases these norm bounds. Exponential decay dominates every fixed power of m. The conditioning is already in D and C_{pc}; there is no second division by c_*. The proof invokes the established physical fan theorem and does not claim that theorem as new. □

Relation to the unchanged raw objective. The special-BV theorem excludes a Cantor derivative for the specified physical observations. The exact all-label spectrum identifies its atomic part with the full signed cancellations and arithmetic labels. These results do not establish vanishing of the remaining normalized jumps or continuous local derivative charges. Smooth narrow spikes can have no jumps at all. The original same-roof shortage (73.12) remains the closing condition on its existing route. The alternative sufficient bridge of Proposition 78.7 still requires its complete normalized local derivative charge, in the original order of limits. Fixed-count structure and the new frequency identity are not uniform dynamical estimates of that charge. Microscopic conditioning and the four-coordinate inverse retain their actual denominator and Jacobian hypotheses.

Appendix A. Reconstruction of the retained physical anchor input

This appendix reconstructs the specific consequences of native modules 287 and 294 needed in the new sublevel proof. Its statements are inherited results, not newly claimed contributions. The exact remote body of 287 is retained under `provenance/sublevel-continuation-freeze/provenance/287-original.tex`. The full module 294 is loaded natively in the present main; its predecessor copy is also preserved.

Lemma A.1 (Buffered all-direction line covering, inherited)**.** *Every straight segment of length $H = 28/5$ has a lattice center at perpendicular distance at most $\omega = 179/400$ from its line, with perpendicular foot strictly inside the segment.*

Proof. For a primitive triangular lattice vector of length l, the parallel lattice rows have spacing $h = \sqrt{3}/(2l)$. At a segment midpoint q choose a nearest row and then a nearest lattice point on it. The two coordinates of $z - q$ parallel and perpendicular to this primitive direction obey $|x| \leq l/2$, $|y| \leq h/2$. For a direction at angle δ to that vector, its transverse projection is at most $W = (h|\cos\delta| + l|\sin\delta|)/2$, and its longitudinal projection at most $\sqrt{l^2+h^2}/2$.

By lattice symmetries reduce to an unoriented angle $0 \leq \theta \leq \pi/6$ and put $t = \tan\theta/\sqrt{3}$. The primitive lattice vectors and the bounds obtained from their row rectangles are

t interval	v	l^2	$\sup W^2$
$[0, 1/60]$	$(1, 0)$	1	961/4804
$[1/60, 1/12]$	$(11/2, \sqrt{3}/2)$	31	13697401/73866304
$[1/12, 1/5]$	$(5/2, \sqrt{3}/2)$	7	529/3136
$[1/5, 1/3]$	$(3/2, \sqrt{3}/2)$	3	75/448

Here is an explicit check that these are bounds for whole intervals. For the first row, $W^2 = 3(1+2t)^2/(16(1+3t^2))$, increasing since $2-3t > 0$. For each other row write $v = ((2a+1)/2, \sqrt{3}/2)$ with $a = 5, 2, 1$, $s = 2a+1$, $N = a^2+a+1$. Then

$$W = \frac{\sqrt{3}}{4\sqrt{1+3t^2}} \left(\frac{s+3t}{2N} + |st-1| \right).$$

On the listed interval $t \le 1/s$, the positive numerator decreases with derivative $3/(2N) - s < 0$, and the denominator increases. Evaluate its left endpoint, and the right endpoint for the first row, to obtain the table. Every listed rational is strictly below $(179/400)^2$. The largest longitudinal bound is $\sqrt{31+3/124}/2 = \sqrt{3847/124}/2 < H/2$, since $96175 < 97216$. Therefore the selected foot lies strictly inside the original segment. This argument neither changes the disk radii nor discards a direction. □

Lemma A.2 (Physical-time turning and the 94-flight anchor, inherited)**.** *Every actual regular trajectory segment of physical duration H has total interior incidence cosine at least $\gamma = 1/4480$. Every actual free flight has length at most H, and every complete 94-flight block has an interior collision with $c_i \ge 1/416640$.*

Proof. A collision changes the unit velocity by a vector of length $2c_i$. Hence for a physical segment $q(s)$, $0 \le s \le H$, with initial outgoing velocity v, direct integration gives

$$|q(s) - q(0) - sv| \le 2H \sum_{0<t_i<H} c_i.$$

Endpoint collisions do not affect this estimate: an initial collision is accounted for by the outgoing v, and the velocity change at H does not affect positions up to H. If the sum were less than $1/4480$, the deviation would be less than $1/400$. Apply Lemma A.1 to the initial straight reference segment and evaluate at its interior perpendicular foot. The actual position would then have distance less than $179/400 + 1/400 = 9/20 \le R$ from a lattice center, contradicting exterior nonpenetration. This proves the incidence sum.

A straight free flight of length exceeding H would contain an H segment intersecting a disk interior by the same covering lemma, a contradiction. Consecutive distinct disks are separated by $\tau_- = 3/50$, so 94 flights take at least $94\tau_- = 141/25 > H$. Apply the physical-time bound to their first H units. The relevant collisions are among the 93 interior collisions of the block. Thus their total cosine is at least $1/4480$, and one is at least $1/(4480 \cdot 93) = 1/416640$. All collisions are actual; no probabilistic independence, symbolic model or number of words is used. □

Appendix B. An exact flux identity for unvisited near misses

The incidence load does not see a trajectory whose original guard is witnessed only by clearance. We introduce a counting observable on the *same* physical flights, not a new collision map. Throughout this section let

$$g_* = 1 - 2R_+ = 3/50, \qquad \delta_* = g_*/4 = 3/200, \qquad 0 < \delta \le \delta_*.$$

The symbol δ_* here is a one-flight geometric constant, not the outer microscopic width $\delta(B)$. All disks still have radius R.

For a regular outgoing collision x, write its flight as $q(x) + tv(x)$, $0 \le t \le \tau(x)$. Let $\mathcal{Z}_\delta(x)$ be the set of lattice centers other than the two incident centers for which

$$0 < \min_{0 \le t \le \tau(x)} |q(x) + tv(x) - z| - R < \delta, \qquad N_\delta(x) = |\mathcal{Z}_\delta(x)|.$$

This counts near misses, not collisions. In particular neither the collision index, the physical roof, nor the section-visit count is incremented by N_δ. A center is counted once on a flight.

Tangencies may be assigned value zero on their collision-area null set; no assertion about a positive pairing-endpoint trace follows from this convention.

Lemma B.1 (An interior foot and uniform endpoint separation)**.** *Every center counted by N_δ has its perpendicular foot in the open physical flight. If t_z is the time of that foot, then*

$$t_z \geq g_* - \delta, \qquad \tau - t_z \geq g_* - \delta. \tag{B.1}$$

The candidate centers, after translating the initial incident center to zero, belong to the finite set

$$\{z \in \Lambda \setminus \{0\} : |z| \leq \tau_+ + 2R_+ + \delta_*\}.$$

Write N_* for its cardinality. This constant is independent of the itinerary length and of all exact record labels.

Proof. At either incident point, the distance to a third center is at least $1-R$, since distinct lattice centers have distance at least one. Its distance to the third disk is therefore at least $1 - 2R \geq g_*$. A minimum below $\delta \leq g_*/4$ cannot occur at an endpoint. The nearest point on a line segment is then its perpendicular foot, whose distance from z is at most $R + \delta$. By the reverse triangle inequality its distance along the line from either endpoint is at least $(1 - R) - (R + \delta) \geq g_* - \delta$. Finally a foot is at distance at most $R + \tau_+$ from the initial center. Adding its distance from z gives the finite-set bound. This proof uses the actual segment, including both of its endpoints, rather than an infinite-line substitute for the clearance event. □

Put $v_\vartheta = (\cos\vartheta, \sin\vartheta)$ and $v_\vartheta^\perp = (-\sin\vartheta, \cos\vartheta)$. For a mark z define the signed line coordinate $b = (z - q) \cdot v_\vartheta^\perp$. Translate the marked center to zero. For $\sigma \in \{-1, 1\}$ and $0 < u < \delta_*$ the oriented line with $b = \sigma(R + u)$ has closest point $-\sigma(R + u)v_\vartheta^\perp$. Follow this line backwards to its last actual collision. Reduce that collision by a lattice translation to the initial fundamental disk, and denote the resulting outgoing state and marked-center displacement by

$$X_R(\vartheta, \sigma, u), \qquad Z_R(\vartheta, \sigma, u).$$

These maps use straight free flight until an *actual* disk is met. There is no reflection at the closest point. The finite-horizon assumption ensures actual collisions in both directions, apart from null singular parameters on which these maps can be defined arbitrarily. The interval from the preceding collision to the following one remains the original physical flight.

Theorem B.2 (One-copy clearance flux and exact multiplicity)**.** *For every nonnegative Borel function a of an outgoing state and a marked-center displacement,*

$$\int \sum_{z \in \mathcal{Z}_\delta(x)} a(x, z)\, d\nu(x) = \frac{1}{4\pi R} \sum_{\sigma = \pm 1} \int_0^{2\pi} \int_0^\delta a(X_R(\vartheta, \sigma, u), Z_R(\vartheta, \sigma, u))\, du\, d\vartheta. \tag{B.2}$$

In particular,

$$\int N_\delta\, d\nu = \delta / R. \tag{B.3}$$

Both statements hold for every $R \in [R_-, R_+]$. The multiplicity on the left is the number of physically distinct unvisited disks, not an upper bound for the number of collision words.

Proof. We first verify the local measure, including its radius factor. Write $q = R(\cos\alpha, \sin\alpha)$ and $\vartheta = \alpha + \arcsin p$. At fixed ϑ,

$$\left.\frac{\partial b}{\partial \alpha}\right|_\vartheta = -R\cos(\vartheta - \alpha), \qquad \frac{\partial \vartheta}{\partial p} = (1 - p^2)^{-1/2}.$$

Consequently

$$\left|\det D_{(\alpha,p)}(\vartheta, b)\right| = R, \qquad d\nu = \frac{d\vartheta\, db}{4\pi R}. \tag{B.4}$$

The two signs $b = \sigma(R + u)$ both have $|db| = du$.

There is also a global multiplicity assertion to prove. A marked physical flight maps to its unique oriented line relative to its marked center. Conversely, the closest point of every line just described is in the free region: its distance to any other lattice center is at least $1 - R - u > R$. In fact the disk of radius $R + \delta_*$ around the marked center does not intersect any other actual disk, since $2R + \delta_* < 1$. The central disk is missed because the line has impact magnitude greater than R. Following the line in both directions therefore gives a physical flight whose incident disks are different from the marked disk. Its foot is interior and its clearance from that disk is u. This is inverse to the marked-flight map, modulo the prescribed lattice translation. Multiple near misses on one flight give different marked copies; they are neither lost nor identified.

For measure-theoretic precision, partition the line parameters by the actual backward incident disk and the forward incident disk, and exclude their tangent boundaries. These partitions are Borel; on each regular piece the disk-intersection formulas and their inverses are smooth. Tangency boundaries satisfy the line equation $|b_z| = R$. For any fixed alternative center their equation in (ϑ, u) is a graph in u (with one of the two fixed signs), hence has planar Lebesgue measure zero. There are countably many centers. The corresponding singular collision sets are also null by (B.4) on regular charts. Exhaust these charts by relatively compact subcharts and disjointize them. Applying the local Jacobian formula to bounded functions on finite unions, then using monotone convergence for both chart unions and $a \wedge n$, proves (B.2) without a prior integrability assumption. Setting $a = 1$ gives $2(2\pi)\delta/(4\pi R) = \delta/R$. □

Corollary B.3 (All physical depths with the original normalization)**.** *For arbitrary $0 < \delta_j \le \delta_*$, $0 \le j < m$,*

$$\int \sum_{j=0}^{m-1} N_{\delta_j}(T_R^j x)\, d\nu(x) = \frac{1}{R} \sum_{j=0}^{m-1} \delta_j.$$

For every original exact event Π the same integral with $\mathbf{1}_\Pi\, d\nu_R^*$ is at most $(Rc_*)^{-1} \sum_j \delta_j$. In particular, if $2\varepsilon \le \delta_*$ and $r = q_0^{1/4}$, then

$$\nu_R^*\Big(\Pi \cap \{\exists j : \Delta_j < 2\varepsilon r^{d'_j}\}\Big) \le \frac{4\varepsilon}{R_- c_*(1 - r)}. \tag{B.5}$$

Proof. The collision measure ν is invariant under the original elastic collision map. This also follows from the oriented-line Jacobian (B.4), whose magnitude is the same at both incident circles, and specular reflection, which preserves boundary flux. Thus apply (B.3) at each j. The original initial restriction has density $\mathbf{1}_{Y_R^*}/c_* \le 1/c_*$ with respect to ν; further exact labels only decrease the positive integral. Outside the null exact-tangency set, existence of a clearance witness implies $N_{2\varepsilon r^{d'_j}} \ge 1$. There are at most two flight indices of each nonnegative depth, so $\sum_j 2\varepsilon r^{d'_j} \le 4\varepsilon/(1 - r)$. No independence or return-conditioned invariant measure is introduced. □

The estimate (B.5) is an integrated mass estimate. Multiplication of it by m^2 at fixed $\varepsilon(B)$ is not a proof about a target-roof density.

Appendix C. Clearance strips transverse to physical test curves

A measure bound alone gives no short-curve control. The positive separation between an unvisited disk and the incident disks supplies a transverse derivative which remains nonzero even as an incident angle approaches grazing. We specify the time orientation explicitly: forward near misses are tested on increasing-slope curves, and *past-flight* near misses on decreasing-slope curves.

Use physical arclength $s = R\alpha$ increasing counterclockwise and $p = \sin\varphi$. Fix a finite $U > 0$ and put

$$\begin{aligned} \kappa_- &= 1/R_+, \qquad \kappa_+ = 1/R_-, \\ D_0 &= \kappa_+ + U + (1 - R_+)^{-1}, \\ \ell_0 &= \pi/(2D_0), \qquad a_0 = \kappa_-(g_* - \delta_*), \\ C_{\mathrm{geo}} &= 2N_*\sqrt{1+U^2}/a_0. \end{aligned} \tag{C.1}$$

All lengths below are Euclidean arclength in (s, φ), not configuration-space arclength. A test is a connected graph over an s interval, of length at most ℓ_0. Let E_δ^+ be the forward physical near-miss event $\{N_\delta \geq 1\}$, and let E_δ^- be the analogous event on the preceding actual flight.

Theorem C.1 (A grazing-uniform, one-flight strip bound)**.** *For every C^1 test graph W^+ with $0 \leq d\varphi/ds \leq U$, and every C^1 test graph W^- with $-U \leq d\varphi/ds \leq 0$,*

$$|W^+ \cap E_\delta^+| \leq C_{\mathrm{geo}}\delta, \qquad |W^- \cap E_\delta^-| \leq C_{\mathrm{geo}}\delta. \tag{C.2}$$

The bound is uniform in R, in the homogeneity strip of the test, and in its distance to the grazing boundary. Arbitrary Borel restrictions, including exact record restrictions, can be added to the sets on the left. Regular pieces are understood with their physical multiplicity one.

Proof. On W^+ the direction angle is $\vartheta = s/R + \varphi(s)$. For a fixed candidate center $z \neq 0$ set

$$t_z = (z - q)\cdot v_\vartheta, \qquad b_z = (z - q)\cdot v_\vartheta^\perp.$$

Differentiation in the increasing s coordinate gives the signed identity

$$b_z' = -\cos\varphi - t_z\vartheta', \qquad \kappa_- \leq \vartheta' \leq \kappa_+ + U. \tag{C.3}$$

At a physical near miss, Lemma B.1 yields $t_z \geq g_* - \delta_*$. Hence $-b_z' \geq a_0$ there. It is the $t_z\vartheta'$ term, not a lower bound on $\cos\varphi$, which prevents degeneration.

We must not multiply this estimate by an uncontrolled number of components. Choose a continuous argument $\beta(s) = \arg(z - q(s))$ on the graph. Since $|z - q| \geq 1 - R_+$, one has $|\beta'| \leq (1 - R_+)^{-1}$. Moreover

$$(\beta - \vartheta)' \leq (1 - R_+)^{-1} - \kappa_- < 0,$$

because $R_+ < 1/2$. Its total variation is at most $D_0|W^+| \leq \pi/2$. Thus the set $t_z > 0$, equivalently $\cos(\beta - \vartheta) > 0$, is at most one interval: two distinct positive-cosine intervals are separated by a negative interval of angular length π. On the interval $t_z > 0$, formula (C.3) says that b_z strictly decreases. Each of the two impact bands $(R, R + \delta)$ and $(-R - \delta, -R)$ therefore has multiplicity at most one in this coordinate. Restricting to the portions which are actual near misses can break them into pieces, but cannot increase that multiplicity. By the one-dimensional change of variables on this monotone map,

$$\int_{\{s:\ z \text{ is a physical near miss}\}} ds \leq 2\delta/a_0.$$

Since $d\ell \leq \sqrt{1+U^2}\,ds$, summing over the finite candidate set from Lemma B.1 proves the first bound. This sum counts only centers in one physical flight. It does not count iterated collision words or homogeneous continuity components.

For the second bound, the collision time-reversal involution is $\mathcal{I}(s, \varphi) = (s, -\varphi)$. The outgoing velocity at $\mathcal{I}x$ reverses the incoming velocity at x. Consequently its next straight flight is the reversal of the previous actual flight at x. This proves

$$E_\delta^- = \mathcal{I}E_\delta^+ = T_R E_\delta^+ \tag{C.4}$$

on the regular collision space. The involution is an isometry in (s, φ) and sends W^- to an increasing-slope graph. Apply the first bound. Borel restrictions decrease lengths. Taking monotone limits of truncated regular test pieces gives the same bound at grazing limits; an exactly tangent parameter contributes zero transverse length by the strict derivative above. This last statement does not delete any paired-source boundary trace in the older coarea construction. □

Corollary C.2 (Bounded multipliers on stable tests)**.** *If $|h| \le H$ pointwise on regular collision states, then for every W^- as above and every bounded Borel test ψ,*

$$\left| \int_{W^-} \mathbf{1}_{E_\delta^-} h\psi \, d\ell \right| \le H\|\psi\|_\infty \min\{|W^-|, C_{\mathrm{geo}}\delta\}.$$

This is a bound for bounded functions with specified representatives. It is not an assertion that multiplication by the indicator is bounded on an entire space of anisotropic distributions.

Proof. Bound the integrand by $H\|\psi\|_\infty \mathbf{1}_{E_\delta^-}$ and use Theorem C.1 together with the trivial length bound. Representatives on singular sets are fixed to zero. □

Appendix D. Clearance gains after arbitrarily many actual collisions

We next give a consequence which is uniform in the full collision count, rather than only a one-flight estimate. Its additional input is the retained stable-curve growth bound, stated explicitly here. It is not the stronger target-roof local limit estimate.

Input statement D.1 (Retained homogeneous curve growth)**.** Use a family $\mathcal{W}_R^s$ of short homogeneous admissible stable graphs in the above physical coordinates, with slopes in $[-U, 0]$, length at most ℓ_0, and the inherited common curvature bound. For every $W \in \mathcal{W}_R^s$ and $l \ge 0$, its regular inverse image has a countable partition $\mathcal{G}_l(W)$ into such graphs, up to arclength-null sets, with one-copy inverse branches and

$$\sum_{V \in \mathcal{G}_l(W)} \sup_V J_V T_R^l \le C_g. \tag{D.1}$$

Here $J_V T_R^l$ is the Euclidean arclength derivative along V; C_g is independent of R, l, W. At $l = 0$ the partition is $\{W\}$.

This is the curve-growth interface of historical module 241, not its whole-cell pressure conclusion. Length thresholds can be decreased: subdividing a pre-existing short component into lengths at most ℓ_0 multiplies its Jacobian upper sum by at most a fixed number, independent of l. The precise external interface is discussed in Appendix H. We retain this input separately from the unconditional one-flight geometry of Theorems B.2 and C.1.

For fixed representatives of bounded Borel functions put

$$\mathcal{S}_a(f) = \sup_{W \in \mathcal{W}_R^s} |W|^{-a} \int_W |f| \, d\ell, \qquad 0 \le a < 1.$$

This absolute stable-test seminorm is specified here to avoid identifying it with a full anisotropic distribution norm. With respect to the invariant measure ν, the ordinary density transfer is $\mathcal{L}_R h = h \circ T_R^{-1}$. For the original record observable $\mathbf{g}_R = (\kappa_1, \kappa_2, \tau, \mathbf{1}_{Y_R^*})$ and real $\xi = (\xi_1, \xi_2, \xi_t, \xi_n)$, define

$$\mathcal{L}_{R,\xi} h(x) = e^{i\xi \cdot \mathbf{g}_R(T_R^{-1}x)} h(T_R^{-1}x).$$

The three discrete frequencies may be taken modulo 2π; the physical-roof frequency is real and unbounded. This definition does not assert boundedness of this operator on an unstated Banach space. On bounded functions its modulus is simply $|h| \circ T_R^{-1}$.

Theorem D.2 (A uniform future-propagated clearance gain)**.** *Under Input D.1, let $|h| \le H$ pointwise on the regular collision space, with $h = 0$ off E_δ^-. Then for every $l \ge 0$, real ξ, and $0 \le a < 1$,*

$$\int_W |\mathcal{L}_{R,\xi}^l h|\, d\ell \le H \min\{|W|, C_g C_{\mathrm{geo}} \delta\}, \tag{D.2}$$

$$\mathcal{S}_a(\mathcal{L}_{R,\xi}^l h) \le H (C_g C_{\mathrm{geo}} \delta)^{1-a}. \tag{D.3}$$

The constants do not depend on the entire future collision word or on any of the four real frequencies. The bounds also hold for any bounded multiplier of modulus at most one applied at arbitrary intermediate collisions, with the same support at the marked time.

Proof. On regular branches positivity and $|e^{i\xi\cdot \mathbf{g}_R}| = 1$ give $|\mathcal{L}_{R,\xi}^l h(x)| \le H \mathbf{1}_{E_\delta^-}(T_R^{-l} x)$. Change variables separately on the inverse branches of W. The full nonnegative sum, completed by monotone convergence, satisfies

$$\int_W |\mathcal{L}_{R,\xi}^l h|\, d\ell \le H \sum_{V \in \mathcal{G}_l(W)} (\sup_V J_V T_R^l)\, |V \cap E_\delta^-| \le H C_{\mathrm{geo}} \delta C_g.$$

The pointwise bound by H supplies the other term in the minimum. For $x > 0$ and $D > 0$, splitting into $x \le D$ and $x \ge D$ gives $x^{-a} \min(x, D) \le D^{1-a}$. Apply this with $x = |W|$ and $D = C_g C_{\mathrm{geo}} \delta$ to obtain (D.3). The modulus of a product of additional multipliers is at most one, so precisely the same positive upper bound applies. No stochastic independence, wordwise supremum in place of a physical integral, or Fourier inversion has been used. □

Corollary D.3 (The complete depth schedule in stable tests)**.** *Let $f_0 = \mathbf{1}_{Y_R^*}/c_*$. For $0 \le j < m$ set*

$$h_j = \mathbf{1}_{E_{\delta_j}^-} \mathcal{L}_{R,\xi}^{j+1} f_0, \qquad f_{m,j} = \mathcal{L}_{R,\xi}^{m-j-1} h_j.$$

Then

$$\mathcal{S}_a(f_{m,j}) \le c_*^{-1} (C_g C_{\mathrm{geo}} \delta_j)^{1-a}. \tag{D.4}$$

If $2\varepsilon \le \delta_*$ and $\delta_j = 2\varepsilon r^{d'_j}$, the terminal collision density restricted to the union of all original clearance witnesses has absolute stable-test seminorm at most

$$\frac{2 (C_g C_{\mathrm{geo}})^{1-a} (2\varepsilon)^{1-a}}{c_* (1 - r^{1-a})}, \qquad 0 \le a < 1. \tag{D.5}$$

This holds uniformly over m, R and real record frequencies. Further original exact labels and terminal-section restrictions may be retained on the left by positive domination.

Proof. The initial density and all of its twisted iterates have modulus at most $1/c_*$. By (C.4), the past-flight event at collision $j+1$ is exactly the forward clearance mark on flight j. Thus the exponents $j+1$ and $m-j-1$ insert the mark at the actual collision time, including $j = 0$ and $j = m-1$ with zero remaining future steps. Theorem D.2 proves (D.4). The union indicator is bounded by the sum of its physical one-flight indicators. Subadditivity of $\mathcal{S}_a$ and at most two indices at each depth give

$$\sum_{j=0}^{m-1} \delta_j^{1-a} \le \frac{2 (2\varepsilon)^{1-a}}{1 - r^{1-a}}.$$

This proves (D.5). Arbitrary exact-record selectors are bounded by one on the same trajectories, so they cannot increase the positive majorant. They are not replaced by independent weights, and no probability measure is renormalized. □

The estimate is genuinely uniform in the full number of physical collisions and in the full scheduled set of clearance marks. Its norm is nevertheless a stable-test norm on terminal collision space, not the essential height of a roof density. In particular, the absence of frequency growth in this absolute estimate is not decay as $|\xi_t| \to \infty$ and does not justify integration over that unbounded frequency. Such decay, or another rigorously justified pointwise bridge, is still required for the original raw target.

Appendix E. Exact-record disintegration at an unvisited mark

The flux coordinates also give a direct representation of the remaining roof problem which does not discard the exact return, displacement or collision information. We state it as an equality of measures first, because a conditional angular roof kernel need not have a density without proof.

For $0 \le j < m$ define the initial state

$$X_{R,j}(\vartheta, \sigma, u) = T_R^{-j} X_R(\vartheta, \sigma, u)$$

on its regular set, with a fixed null-set convention otherwise. The complete actual record from this state is used. Displacements are computed in the original lattice basis, relative to its initial lift; translating the marked disk to zero changes none of these relative displacements. For almost every u, define the positive Borel measure on the original roof line

$$\mathcal{K}_{m,R,n,k,j,u}(A) = \sum_{\sigma=\pm 1} \int_0^{2\pi} \mathbf{1}_{\Pi_{n,k,m,R}}(X_{R,j}) \mathbf{1}_{\{L_{m,R}(X_{R,j}) \in A\}} \, d\vartheta. \tag{E.1}$$

It has mass at most 4π. No density or angular derivative bound is included in this definition.

Theorem E.1 (Disintegration of a physically marked exact record)**.** *For every Borel set $A \subset \mathbb{R}$,*

$$\int \mathbf{1}_\Pi N_\delta(T_R^j x) \mathbf{1}_{\{L_{m,R} \in A\}} \, d\nu_R^* = \frac{1}{4\pi R c_*} \int_0^\delta \mathcal{K}_{m,R,n,k,j,u}(A) \, du. \tag{E.2}$$

The identity also holds against every nonnegative Borel roof test. All singular boundary sets in these integrals use the original collision-area null-set convention, not deletion of paired boundary coefficients.

Proof. Replace $d\nu_R^*$ by $c_*^{-1} d\nu$ inside the integral, since $\Pi \subset Y_R^*$. Change the initial collision coordinate to $y = T_R^j x$ using invariance of ν. Apply Theorem B.2 with

$$a(y, z) = \mathbf{1}_\Pi(T_R^{-j} y) \mathbf{1}_{\{L_{m,R}(T_R^{-j} y) \in A\}}.$$

This is a bounded nonnegative function on the original physical states, independent of which near-miss mark is selected once y is fixed. The resulting integral is exactly (E.2). Countable regular partitions and Tonelli's theorem justify the integration over u; their exceptional parameters have two-dimensional measure zero. For arbitrary nonnegative roof tests use monotone simple-function approximation. In particular the factor is $1/(4\pi R c_*)$, with no second section normalization and no new conditioning on the near miss. □

E.1. **A sufficient pointwise bridge, with its status explicit.** Fix M. The following measure domination is one concrete, stronger sufficient condition for paying all original clearance witnesses: there exists C_M such that, for all sufficiently large m, all R, n, k, j and almost every $0 < u < \delta_*$,

$$\mathcal{K}_{m,R,n,k,j,u}\big|_{\mathcal{C}_{m,R,n,k}(M)} \le C_M m^{-2} \, dt. \tag{E.3}$$

An almost-everywhere set may depend on the fixed R, m, n, k, j; C_M must be common. This is an *unproved sufficient condition*, not a newly assumed premise of the organizing raw local limit assertion. It asserts absence of singular conditional roof mass as well as an upper density bound. Neither Theorem B.2 nor Theorem D.2 establishes it.

Proposition E.2 (The exact normalization a pointwise bridge must pay)**.** *If* (E.3) *holds, the original roof measure restricted to any clearance witness, with* $2\varepsilon \le \delta_*$*, is dominated on the central target by a density* d_{clr} *with*

$$\sup_{R,n,k} m^2 \|d_{\mathrm{clr}}\|_{L^\infty(\mathcal{C}_{m,R,n,k}(M))} \le \frac{C_M \varepsilon}{\pi R_- c_*(1-r)}. \tag{E.4}$$

The same upper bound holds for any positive source already dominated by that original collision measure, restricted to these witnesses.

Proof. The witness-union indicator is at most $\sum_j N_{2\varepsilon r^{d'_j}}(T_R^j x)$ off the fixed null set. Apply (E.2) to every term, then the measure domination (E.3). The resulting density bound is $(4\pi R c_*)^{-1} C_M m^{-2} \sum_j 2\varepsilon r^{d'_j}$. The sum is at most $4\varepsilon/(1-r)$, proving the stated constant. Positive domination preserves the bound for the indicated source. At the original outer schedule this would tend to zero after the collision limsup and then $B \to \infty$; no window has been changed. □

On a regular physical branch, the new parametrization has the exact chain rule

$$DX_{R,j} = DT_R^{-j}(X_R)\, DX_R, \qquad D(L_{m,R} \circ X_{R,j}) = DL_{m,R}(X_{R,j})\, DX_{R,j}.$$

The one-flight lower bound for $|b'_z|$ does not lower-bound this long-time roof derivative. Inverse-iterate growth, cancellations and critical angular roofs must be controlled separately. A bound after integration in u can allow angular folds which a uniform bound on each $\mathcal{K}_{j,u}$ would exclude. No supremum is passed through that disintegration here.

A proof of the stronger condition (E.3) is not necessary if a suitably summed kernel bound or comparison-capacity argument can be proved instead. The proposition only makes explicit which normalization and which measure must be controlled. It is not counted as an unconditional closure or a second payment of a source already paid in revision 112.

E.2. **Compatibility with the complete original source.** All identities through (15.3) and the full pool $P_{r_B}^{\mathrm{ng}}$ are unchanged. We have not subtracted the clearance source from $\mathsf{U}_B^{\mathrm{low}}$ using only a weak norm. Consequently the exact remaining target is still (15.8), with the original order of limits. The new results treat the mechanism which is invisible when $J_\varepsilon = \varnothing$: genuine unvisited-disk near misses. They do not pay low-severity shallow incidence without clearance, prove abundance of the full nongrazing comparison pool, or settle positive critical-entrance and section-cut pairing traces.

For precision concerning a minor referee point, the inequality $\alpha f(N) < u_m$ already proved in Corollary 15.2 implies $N < 2\sqrt{u_m/\alpha}$ on the retained source. The independent quadratic count theorem is retained as a finite-count theorem, not presented as another missing payment for that already restricted source. The term "low" in all inherited source symbols means low *incidence* load only. In the monotone exhaustion of Proposition 10.4, every marked weight is explicitly assigned zero on the omitted collision-area null set; its extended integral is taken only over the stated regular charts. This convention never sets a positive one-sided pairing trace to zero.

Appendix F. Scope of the retained comparison statements

For clarity, we record the precise scope of the earlier allocation statements. None of their algebraic maximality assertions is used as a substitute for the physical estimate in Theorem 7.1.

Proposition F.1 (Fixed-window scalar maximality and old-family dominance)**.** *Fix a Borel comparison subspace,* $0 < r < \Delta$*, and* $K > 0$*. Consider nonnegative measurable weights* w_+, w_-

with $w_+ + w_- \le 1$, supported on the respective one-sided window conditions in (6.2)*. Among scalar accepted heights $Z(t)$ satisfying*

$$0 \le Z(t) \le U(t), \qquad Z(t) \le K \int (w_+ + w_-) p_\vee \, d\theta,$$

the greatest is $V_{r,K}(t) = \min\{U(t), KP_r(t)\}$. The allocation is global at the roof and does not couple individual source sheets to comparison sheets.

For a compatible old fixed family with window width r_{old}, physical overlap N and block cost κ, the explicit conditions

$$0 < r \le r_{\mathrm{old}}, \qquad \ell_{\mathrm{comparison}} < \Delta, \qquad K = N\kappa$$

imply $V^{\mathrm{old}}(t) \le V_{r,N\kappa}(t)$ on the same original source and comparison space.

Proof. The support and one-copy conditions give $w_+ + w_- \le Q_r$. Hence every Z is bounded by the stated minimum. On $P_r > 0$ choose $w_\pm = V_{r,K} q_\pm / (KP_r)$; use zero weights otherwise. This is a measurable attaining certificate. The separate source multiplier is $\beta_{r,K}$ defined in Section 7. For the old family, retain only active source blocks at t. Their comparison supports lie in the r-pool by the displayed width condition. Summing the blockwise budget and using physical overlap at most N gives $V^{\mathrm{old}} \le N\kappa P_r$; disjoint original source blocks also give $V^{\mathrm{old}} \le U$. The minimum proves the assertion, with countable families obtained by monotone convergence on their common roof versions. □

Proposition F.2 (Strictness relative to roof-independent whole-interval blocks)**.** *In the interval-source model, the roofwise certificate class is strictly larger than the old class of roof-independent angular source blocks that requires a comparison interval to contain the translate of the* entire *source interval by $[0, r]$ or $[-r, 0]$. This statement compares certificate classes and does not assert a Lorentz realization.*

Proof. Let $s = 1/1000$ and $r = 1/10$. On two disjoint angular sets of measure one give both endpoint traces value one. Let their physical intervals be $(0, s)$ and $(3s/8, r + 7s/8)$, and select only the first as source. Take $\Delta > r + s/2$ and $K = 1$. For $3s/8 < t < 7s/8$ the second interval contains $[t, t + r]$, so $P_r(t) \ge U(t) = 1$. Neither physical interval has length $r + s$, which is necessary for the old translated-whole-source inclusion. Angular subdivision does not alter these intervals. Thus the old class cannot allocate a nonzero whole-source block while the roofwise class accepts the source on the indicated roof interval. □

F.1. **Limit order and the remaining estimate.** The constructions in this article are evaluated first at finite m, R, n, k with B, r_B, h_B, K_B already fixed. Positive physical sums are completed before taking the essential roof supremum, the radius and exact-label suprema, and the collision limsup. Only then is B sent to infinity. The assertion that every positive-length segment contains a sufficiently small window at a fixed count supplies no uniform modulus for this order.

The remaining estimate is precisely

$$\lim_{B\to\infty} \limsup_{m\to\infty} \sup_{R,n,k} m^2 \operatorname*{ess\,sup}_{t\in\mathcal{C}_{m,R,n,k}(M)} \left(\mathsf{U}^{\mathrm{micro}}_{B,m}(t) - K_B P^{\mathrm{ng}}_{r_B,m}(t)\right)_+ = 0.$$

No inference to this estimate is made from bounded wordwise optical weights, local integrability, a finite diagnostic, or a positive finite-count incidence. Theorem 7.1 proves the entire visited-grazing payment in this order; all remaining positive classes in the displayed expression are still part of the original pointwise problem.

Appendix G. Optical cofactors, multiple marks and simultaneous endpoints

We supply explicit matrix details for the optical identity and isolate the one-sided endpoint statement used in the trace theorem. These calculations take place on the original finite itinerary. They do not assert a bound for a sum over itineraries.

Lemma G.1 (The signed corner cofactor)**.** *For $m \geq 1$, with the empty determinant convention when $m = 1$,*

$$G_{0m} = (-1)^{m-1} \frac{R^2 c_0 c_m}{(\prod_{i=0}^{m-1} \tau_i) \det J}. \tag{G.1}$$

In particular the absolute value, not the signed mixed entry, gives $\omega = |G_{0m}|/R$.

Proof. Suppose first that $m > 1$ and put $n = m - 1$. The minor obtained by deleting the last row and first column of the tridiagonal matrix J is triangular, with diagonal entries $\tau_1^{-1}, \dots, \tau_{n-1}^{-1}$. Cramer's rule therefore gives

$$(J^{-1})_{1n} = (-1)^{n+1} \frac{\prod_{i=1}^{n-1} \tau_i^{-1}}{\det J}.$$

For $n = 1$ its numerator is the empty product and the formula says $(J^{-1})_{11} = 1/J_{11}$. The only endpoint–interior couplings are $H_{01} = R^2 c_0 c_1/\tau_0$ and $H_{nm} = R^2 c_n c_m/\tau_{m-1}$, whereas $H_{0m} = 0$. Since $H_{II}^{-1} = R^{-2} C_I^{-1} J^{-1} C_I^{-1}$, the Schur complement gives

$$G_{0m} = -\frac{R^2 c_0 c_m}{\tau_0 \tau_{m-1}} (J^{-1})_{1n},$$

which proves (G.1). If $m = 1$, the mixed entry follows directly from the single square in (4.2) and has positive sign.

For clarity, the equality between the determinant denominator and $\mathfrak{b}_m$ can also be checked without a sign convention for optical coordinates. Both quantities have initial values $B_0 = 0$, $B_1 = \tau_0$ and satisfy

$$B_{l+1} = \left(1 + \tau_l \kappa_l + \frac{\tau_l}{\tau_{l-1}}\right) B_l - \frac{\tau_l}{\tau_{l-1}} B_{l-1}.$$

For the determinant this is the tridiagonal recurrence after multiplication by the flight lengths. For the optical product it follows by eliminating the lower entry of its second column. This verifies the denominator identity used in Lemma 4.3. □

Put $T_{a,b} = \sum_{i=a}^{b-1} \tau_i$ for $0 \leq a < b \leq m$. For $S = \{j_1 < \cdots < j_s\} \subset \{1, \dots, m-1\}$ set $j_0 = 0$ and $j_{s+1} = m$.

Proposition G.2 (Positive expansion and simultaneous optical marks)**.** *The exact denominator has the positive expansion*

$$\mathfrak{b}_m = \sum_{S \subset \{1,\dots,m-1\}} \left(\prod_{j \in S} \kappa_j \right) \prod_{\nu=0}^{|S|} T_{j_\nu, j_{\nu+1}}. \tag{G.2}$$

For every fixed set S of s distinct interior marks,

$$\frac{\omega}{\prod_{j \in S} c_j} \leq \frac{R_+^{s+1}}{2^s \tau_-^{s+1}} \varrho^{m-s-1}. \tag{G.3}$$

For $S = \varnothing$ this is the first optical bound in (4.8)*; for $|S| = 1$ it is the second.*

Proof. Let $E = \left(\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix}\right)$. Expand every factor $\mathsf{C}(\kappa) = I + \kappa E$ in the product (4.6). Between two selected marks the shears combine according to $\mathsf{S}(a)\mathsf{S}(b) = \mathsf{S}(a+b)$, and $E\mathsf{S}(T)E = TE$. The top-right entry of the term with marked set S is exactly the corresponding summand in (G.2). The empty term is $T_{0,m}$. Every coefficient is positive.

To obtain the stronger estimate retaining the unmarked reflections, expand only the factors at the prescribed marks. The coefficient of $\prod_{j\in S}\kappa_j$, with all other factors retained, is the product of the top-right entries of the $s+1$ intervening optical blocks. Indeed $E = e_2 e_1^{\mathsf{T}}$, so repeated multiplication factors that coefficient into these entries. The block with $j_{\nu+1} - j_\nu$ flights has top-right entry at least $\tau_- \varrho^{-(j_{\nu+1}-j_\nu-1)}$, by the same positive-column argument as in (4.9). Keeping only this nonnegative term gives

$$\mathfrak{b}_m \geq \left(\prod_{j\in S} \frac{2}{Rc_j}\right) \tau_-^{s+1} \varrho^{-(m-s-1)}.$$

Substitute this in $\omega = Rc_0c_m/\mathfrak{b}_m$, and use $c_0c_m \leq 1$ and $R \leq R_+$, to prove (G.3). □

This estimate controls simultaneous vanishing incidences on a fixed finite trajectory. Its constant depends on the number of marks, not on their locations. No word-count factor has been suppressed, and no pressure conclusion is drawn by summing its right side.

Lemma G.3 (One-sided limits at simultaneous physical mechanisms)**.** *Fix a positive-length pairing segment and approach one of its ends from the regular interior. The collision points, velocities, flight lengths, visited incidences and third-disk clearances have limits along the entire approach, even if several of the following occur simultaneously: visited grazing, unvisited tangency, a section boundary and a critical incoming endpoint. The limiting reduced Hessian and coarea factor are those in* (5.3) *and* (5.4)*. No continuation of the exact event through its boundary is asserted.*

Proof. Lemma 5.1 gives limits for q and z along the entire segment, not only a subsequence. Hence the initial collision point and $p_0 = -z\cos\theta/R$ converge. The section cap gives the outgoing normal component $c_0 > 0$ at the limit, so the initial velocity converges as well.

Suppose inductively that the departure point x and unit velocity v of a designated flight converge. For its designated next disk with centre O, the entry time is

$$\tau = -(x-O)\cdot v - \sqrt{((x-O)\cdot v)^2 - |x-O|^2 + R^2}.$$

This is the entry root on the actual regular itinerary. Its radicand is nonnegative along the approach and has a nonnegative limit. The root is continuous also when the radicand tends to zero; $\tau \geq \tau_- > 0$ prevents a zero-length flight. The arrival point therefore converges. Reflection by $v \mapsto v - 2(v\cdot n)n$ is continuous even at tangency, so the next velocity converges. Induction applies a finite number of times. An earlier disk may become tangent only at the limiting flight; this does not alter the designated entry-root limit taken from the original event.

At fixed itinerary and radius, the bounded flight lengths restrict all possibly nearby third-disk centres to a finite set. Distance from a compact line segment to each such disk is continuous in the endpoints. Their finite minimum, and hence the original clearance, has a limit. A section decision may change at the limiting state, but its coordinate limit is still obtained from the same side. None of these continuity arguments requires the boundary mechanisms to occur separately. Formula (5.3), with Z^{-1} uniformly bounded, then gives the Hessian limit even with several vanishing incidences. The fixed-direction coarea formula gives the asserted limit for χ. □

Corollary G.4 (An explicit uniform trace modulus)**.** *A permissible constant in* (5.6) *is*

$$C_{\mathrm{H}} = \frac{500 C_{\mathrm{reg}} \sqrt{2\Lambda_\circ}}{4\pi c_* \lambda_\circ^2}. \tag{G.4}$$

It is common to all original sheets, collision counts, radii and exact labels. Its inherited dependence is precisely the regular variation constant C_{reg}; the remaining constants are the fixed geometry and section constants displayed in this article.

Proof. Corollary 4.4 and $\chi \geq \lambda_\circ$ give $AI_m \leq 500\varrho^{(m-2)_+}/(4\pi c_* \lambda_\circ)$. Integrate the regular derivative bound between u and v, and apply Lemma 5.1 to that subsegment:

$$\int_u^v z^{-1}\, ds \leq \lambda_\circ^{-1} \sqrt{2\Lambda_\circ (v-u)}.$$

Multiply the constants and pass to the already proved endpoints. □

Any nonempty collection of visited endpoint grazings belongs to Γ, and one vanishing marked incidence is already sufficient for its zero source statement. Proposition G.2 additionally controls their simultaneous product. This does not alter the distinct treatment of an unvisited third-disk tangency.

Appendix H. Dependency, reference and normalization audit

The direct clearance proofs use the original positive disk separation, the uniform finite-horizon hypothesis, circular curvature, elastic reflection and elementary change of variables. The arbitrary-time stable-test estimate additionally uses Input D.1. Neither uses uniform pressure, a twisted spectral gap, or the unproved localized-shortage estimate to prove its own premise.

H.1. **The precise imported curve estimate.** In the inspected 10 October 2013 author version of Demers–Zhang [6], Lemma 3.2(b), printed page 14, bounds the sum of suprema of the forward arclength Jacobians on homogeneous inverse-image pieces, uniformly for W and for maps in the stipulated class. The geometric-family statement is Theorem 2.7 in that version; the varying-arclength qualification is Remark 2.9(b). The inherited module 241's numbering must therefore be matched to a specified version rather than silently treated as verified. Its stronger finite-testing and pressure argument is not needed for Theorem D.2.

For this paper the coordinates, Euclidean arclength Jacobian, short-curve curvature bound, homogeneous partition, fixed subdivision length, and map-family constants must all agree with Input D.1. Changing boundary perimeter with R requires the coordinate transport and its uniform equivalence; changing the subdivision threshold changes C_g by a bounded factor. These interfaces remain explicit retained hypotheses here, rather than a claim of an independent specialist reconstruction.

In the inspected 19 September 2022 author version of Baladi–Demers [7], Lemma 2.2, printed pages 7–8, is a distortion statement across stable curves in the same homogeneous backward cell. Its closure convention excludes true singularity curves. It is not a four-record, roof-conditioned m^{-2} estimate. The present proofs make no such inference.

H.2. **Checks which enter the proofs.** The radius factor in (B.4) is R, not R^2 or $1/R$. The two impact signs and the full direction circle give the factor $4\pi\delta$ before collision normalization. Exactly one section factor c_*^{-1} occurs in (E.2). The foot has physical time coordinate t_z, so the sign in (C.3) is a minus for both the boundary motion and the direction motion. Increasing-slope curves, not forward stable curves, are used for the forward event.

The uniform lower bounds used above are strictly positive on the whole closed radius interval:

$$g_* - \delta_* = 9/200, \qquad \kappa_- - (1 - R_+)^{-1} = 600/2491, \qquad a_0 = 9/94.$$

The occurrence of N_* in the curve estimate, but not in the exact flux identity, is deliberate: the former is a one-flight union bound on a test curve, whereas the latter is a global one-copy change of section. Empty future blocks are included by $l = 0$. The denominator $1 - r^{1-a}$ is finite only for $a < 1$; no estimate at the endpoint $a = 1$ is claimed.

The source identities in the retained proof bodies are unchanged. All positive sums are completed before essential suprema and the collision limsup. Taking $B \to \infty$ in a stable-test estimate would not by itself permit taking that limit in the target-roof norm. Fixed-m simulations and symbolic identities check algebra and implementation only; they do not certify either the inherited uniform dynamical inputs or the remaining raw pointwise theorem.

H.3. **What is retained, and what has not been reconstructed.** The complete active revision-112 manuscript and its ten active core proof bodies are retained in this manuscript, and an unmodified copy of the downloaded active packet is supplied in `provenance/v112-active`. The original repository additionally contains the recursive revision-111 archive, with source tree `790e627efe3a89a4549123d7e9acacf24089c287`. That recursive archive was not exported by the completed CI artifact available for this revision. Its identity is preserved; it has not been represented as a locally reconstructed, independently audited set of 267 historical proofs. The included source manifest and preservation audit distinguish byte-verified local preservation from remote tree pinning. In particular the historical spectral, arithmetic, induced-return and four-coordinate inverse-problem inputs are not replaced by the new clearance lemmas.

Appendix I. Critical-roof and fold audit: normalization and ordered closure

The new identities have different kinds of uniformity. The clearance-to-endpoint determinant is an identity for each finite physical itinerary and every marked flight. The fold and finite-type bounds are uniform over Borel selectors on a specified chart, with specified derivative constants. The critical coarea error is uniform over selectors but depends on the chart's third-derivative and weight-modulus constants. The critical coefficient sum is over all regular physical words at fixed collision count, relative to the retained uniform pressure input; it is unscaled. None of these qualifiers can be replaced by another.

I.1. **Two distinct uses of reversibility.** At a normal endpoint, the actual reflected ray reverses its incoming velocity. This proves the reversing-periodic-orbit assertion without adding a reflection at an unvisited mark. In collision coordinates, time reversal also gives the derivative identity $\mathcal{M} = JB^{-1}JB$. The first assertion concerns actual points and their section tests; the second concerns the tangent map on the regular domain. At an unvisited tangency, only the one-sided reduced action coefficient is used. A smooth extension of that action does not define the physical map through the forbidden side.

Even at a regular reversing orbit the original section need not be reversal invariant. The occupation identity is $A_{2m} = A_m + A_m^{\mathcal{I}}$ when both original endpoints are in the section, not automatically $2n$. The coefficient $\mathfrak{J}$ describes nearby collision area pushed to the roof. It is not a conditional law on a measure-zero periodic orbit.

I.2. **The exact small-scale error and its range.** The radius r_0 in Theorem 17.3 is a local endpoint-chart radius. The height increment is $s = t - t_x$, and its allowed range is $0 < s < r_0^2/3$. Neither r_0 nor s replaces the original microscopic observation window. In particular the explicit error constant

$$2\pi\sqrt{3}\{2L_w + Kw(0)\}$$

has not been shown summable after multiplying the full physical source by m^2. Decreasing r_0 separately for each orbit does not establish that summability. A general Borel selector retains

the exact fraction $a_E(s)$; only a verified finite tangent-sector structure justifies a limiting angular fraction.

The local fold exponent $1/2$ is produced by integrating a square-root conditional singularity across a transverse clearance direction. A higher finite type has its stated exponent $1/\ell$ and derivative budget. Full critical roofs have a different two-dimensional local coarea law; the fold theorem is not applied there when its clearance transversality fails. A countable disjoint source-chart covering preserves positivity and exact multiplicity, but does not itself bound the sum of the chart constants.

I.3. **The unchanged closure statement.** All original positive coefficients and the complete nongrazing pool are still those of Section 15. The remaining statement is (15.8), with the full physical sum first, then the exact-label and radius suprema and the original roof essential supremum, then the collision limsup, and the outer limit last. A successful continuation must control clearance and critical charts, positive one-sided pairing traces, the low-incidence source, and the actual available comparison capacity in that topology. The new calculations neither subtract these unpaid classes nor introduce a new name for the shortage.

The accompanying programs check the signed generating derivative, stability determinant, normalization, fold constants, selector error and actual horizontal period-two physical rays. The latter are not asserted to lie in the large-collision central roof window. Exact rational checks of their optical Schur complements and finite ray-derivative comparisons test independent implementations; they are not substitutes for the written continuous proofs or the unresolved uniform dynamical estimates.

Appendix J. Cross-flight normalization and source-scope audit

The physical order used in Section 19 is

$$\text{first mark foot} \xrightarrow{a} q_1 \xrightarrow{\ell_1} \cdots \xrightarrow{\ell_{h-1}} q_h \xrightarrow{b} \text{second mark foot}.$$

Only $q_1, \dots, q_h$ are reflections. The incident collision before the first foot and the incident collision after the second foot are also present in the original full trajectory. Neither is deleted from the record; neither auxiliary marked circle changes the clock.

The finite-horizon quantity τ_+ denotes the fixed uniform bound on the length of every real free flight in the original radius family, as specified in the retained model. This revision does not certify a new numerical value for it. All new constants display that dependency. In particular, with

$$L_{\mathrm{pair}} = \tau_+ + 97/100, \qquad L_{\mathrm{end}} = \tau_+ + 191/200,$$

N_{pair} and $N_\circ$ are the numbers of nonzero lattice vectors of length at most L_{pair} and L_{end}, respectively. For either L, they can be evaluated exactly as

$$\#\{(a,b) \in \mathbb{Z}^2 \setminus \{(0,0)\} : a^2 + ab + b^2 \le L^2\}.$$

A conservative bound, without any enumeration assumption, is $\lfloor (2L+1)^2 \rfloor - 1$: the disjoint open radius-$1/2$ disks around these centers and the origin lie in the radius-$(L+1/2)$ disk; area comparison proves the claim. This is an end-candidate count, never a bound on the number of full physical itineraries.

The two-mark same-flight identity counts *ordered* distinct centers. For chronological pairs the expansion $\mathscr{N}(\mathscr{N}-1) = \sum_j N_j(N_j-1) + 2\sum_{j<k} N_j N_k$ accounts explicitly for both orders. A repeated center at two times still supplies two chronological marks. The exact physical prefactor $1/(4\pi R)$ precedes every clearance substitution, while the original section restriction contributes $1/c_*$ only when passing to ν_R^*. Neither the factor two nor the section normalization is absorbed into a supposed independent probability.

In the inherited identity $E_\delta^- = \mathcal{I}E_\delta^+ = T_R E_\delta^+$, both expressions on the right denote *set images* on the regular invertible collision space. Thus $\mathbf{1}_{E_\delta^-}(x) = \mathbf{1}_{E_\delta^+}(T_R^{-1}x)$ almost everywhere; the displayed image is not $T_R^{-1}E_\delta^+$. The null-set convention is a bulk-measure convention, not a rule that positive one-sided endpoint traces vanish. Similarly, the critical coarea density in Section 17 is the contribution of its stated coordinate ball and selector. It is not automatically the complete global density near that roof value.

The new proof chain has three distinct levels. The foot standoff, Jacobi identity, tangential-end uniqueness, exact coarea and stable metric comparison are proved directly in Sections 19–20. The uniform all-separation density and complete factorial bounds additionally require the explicitly retained unconditioned physical pressure inequality $\sup_{R,l} P_l(R) \le Q_*$. The raw pointwise theorem still requires (15.8). No conclusion at the second level is substituted for the third. The native historical proof bodies 260–276 are preserved, and modules 277–278 alone are the new numbered proof modules in this author revision.

Appendix K. Angular-source dependency, metric and limit audit

The new geometric source formulas use only lattice separation, actual circle reflection, the inherited physical finite-horizon bound and the exact one-mark parametrization. The direct source norm in (21.8) is $\sum_C \|\rho_C\|_\infty$, not a sum of masses and not a global collision-space L^∞ norm. It contains both signed first marks. The second mark also has two signs, already paid in C_t; no additional sign or root factor is to be appended to H_*. The one-mark normalization is inserted only after the root sum, as $1/(4\pi R)$. The original source normalization appears once, as c_*^{-1}, inside a positive exact-event domination.

External curve growth and physical coordinates. The inspected author version of [6] dates from 10 October 2013. Its Lemma 3.2(b) is the homogeneous stable curve-Jacobian sum; its proof specifies (H2)–(H4). Theorem 2.7 states uniform properties for its geometric family, and Remark 2.9(b) discusses a small change of boundary arclength. We use none of these statements as an exact-label roof local limit theorem or as a sum over all physical cylinders. Input 22.1 states exactly the forward counterpart needed here; source-curve membership and a common curvature allowance are essential parts of its invocation. Theorem 21.1 verifies a concrete bound K_0 for that membership check. It does not independently re-prove expansion and distortion for the admissible class.

Time reversal is $\mathcal{I}(s,\varphi) = (s,-\varphi)$, an isometry of the original collision-coordinate arclength, and $\mathcal{I}T_R\mathcal{I} = T_R^{-1}$ on the regular set. Thus forward pieces of an unstable curve correspond to backward pieces of its reflected stable curve, with equal arclength Jacobians. This conversion does not reverse the original measured return record in a theorem statement; it only verifies the curve estimate.

For a local common boundary coordinate at reference radius R_0, write $A_R(s,\varphi) = (R_0 s/R, \varphi)$. If $m_A \le |A_R v|/|v|$ and $|A_R v|/|v| \le M_A$, then conjugating a curve map gives

$$J^{\mathrm{common}}T_R^l(x) = J^{\mathrm{physical}}T_R^l(x)\frac{|A_R v_l|/|v_l|}{|A_R v_0|/|v_0|}.$$

There is only an endpoint ratio, at most M_A/m_A, not a factor $(M_A/m_A)^l$. A compact finite local cover of $[R_-, R_+]$ gives common constants whenever the cited geometric curve hypotheses are verified in those local charts. This is a coordinate comparison, not an affine deformation of the triangular physical lattice. The input is kept explicit rather than asserting that compactness alone verifies (H2)–(H4) or the larger pressure and pairing chain.

Boundary and quantifier conventions. All regular first-hit intervals are genuinely physical. Future singular cuts are exhausted through the one-copy curve partition. This removes zero angular or curve measure in bulk integrals only. It says nothing about the positive one-sided trace of a paired source at a singular endpoint. Those traces and critical ball selectors remain as in the inherited optical proofs. Likewise the set-image convention is unchanged: $\mathcal{I}E_\delta^+$ and $T_R E_\delta^+$ mean images on the regular set, whereas $T_R^{-1}(E)$ denotes its preimage; inherited equalities are used with their specified null-set convention.

The curve-growth supremum ranges over all stated radii, all nonnegative iterate counts and the entire admissible curve class. The new kernel holds for every fixed positive first clearance and all positive integer separations. The root sum is countable and positive before any essential supremum. These facts do not commute a roof supremum with an integral over first clearance. The original central roof window and arithmetic labels remain fixed. The collision limsup is taken before the outer B limit. The explicit remaining proof obligation is (15.8), including first-order clearance, critical and boundary sums, incidence-only source and original nongrazing capacity. None is assigned zero mass or zero height merely because the angular-source theorem has been proved.

Appendix L. Current dependency and citation notes

The preserved proof bodies retain their source-stage hypotheses and scope explanations. A later theorem changes the current status only of the conclusion it actually proves. Thus the historical statement that a full geometric pressure input remains to be paid is superseded by Theorem 70.4 and Corollary 70.5. This includes the original small-capacity word-supremum interface (65.19). It does not establish abundance of the different, originally conditioned same-roof comparison pool. The conditional consequences of Input 11.1 can now use its three native geometric conclusions with the constants stated in Corollary 70.5.

Theorem 67.4 supersedes the absence of a full-source unscaled roof-height bound in the earlier weak-roof argument. Its interior variation, all-offset regular unvisited boundaries, complete critical sectors, cap endpoints and inherited zero visited-grazing trace are explicitly accounted for. The proof does not assert zero trace for a different pairing construction or a bound for its singular multiplier. Theorem 68.6 pays every actual finite section side in the specified roof variation topology. Theorem 69.2 restores the exact occupation, lattice labels, initial conditioning and terminal test. Its joint marked Fourier estimate is (69.8); it is uniform in the mark frequencies but has only an $|s|^{-1}$ roof bound.

Theorems 64.1, 65.3 and 66.3 are inherited from the complete immediate predecessor. They respectively give the all-offset regular marked-boundary budget, all-class physical capacity moment, and leading positive-visited incoming critical-fan payment. Their proof bodies and provenance are unchanged. The new endpoint argument does not relabel these results as new contributions, or claim a uniform local Taylor remainder solely from a leading coefficient.

The new Theorem 71.1 retains selected source length in the complete physical branch-supremum sum. Its own enlarged curve class, cutoff and constants are fixed in (71.1)–(71.2). Theorem 72.4 combines this transport with all actual outgoing boundary charges, the exact cap Jacobian cancellation and the full localized singular multiplier. Non-grazing critical fan values are retained as incoming traces. Theorem 73.1 pays all sufficiently deep clearance witnesses at the original m^2 roof-height scale. This supersedes the earlier absence of a normalized payment for that precise subsource; it does not pay the shallow source. Theorem 73.2 proves that the original shortage is equivalent to (73.12), with exactly the same full comparison pool and order of limits.

The original ordered condition (0.1) therefore remains open precisely at the displayed endpoint-layer shortage and any other retained independent input. All positive physical contributions

must be summed before the essential height and label/radius suprema; the outer widths are fixed before the collision limsup, and the outer parameter tends to infinity last. A bounded-variation density can have arbitrarily narrow positive spikes. Even a common unscaled variation bound and vanishing total mass do not imply the collision-normalized roof height needed there. The current mathematical verification includes explicit examples of this distinction. The new theorem provides a rate before taking its collision limsup, instead of deriving height from mass alone. The unchanged spectral/arithmetic reference, near-unit peaks, conditional microscopic-return denominator and inverse nondegeneracy retain their own requirements.

Theorem 74.1 adds the variation of the complete exact-label vector, with internal section-jump cost two. The actual return-tower proof in Lemma 75.1 needs no exponential tail or ergodicity of the full collision space. The positive likelihood theorem retains all collision counts in its conclusion and handles arbitrary bounded trajectory selectors without differentiating them. Its polynomial normalized absolute error does not supply a relative error for an arbitrarily rare selected event. Corollary 75.8 uses the actual selected mass and never transfers a denominator lower bound from a larger event by restriction.

Theorem 76.2 and Corollary 76.3 give the stated weak $C^1 \to (C^1)^*$ marked operator bounds. The endpoint-action derivative contains the factor $e^{\mathsf{T}}G^{-1}e$ in its denominator; the full inverse-gradient moment is conditioned exactly once. These statements do not prove strong anisotropic spectral estimates, an integrable high-frequency tail, or a long-time forward-map Jacobian bound.

For the inspected 45-page arXiv v1 PDF of Demers–Zhang [6], Lemma 3.2(b) is on PDF page 12. This corrects the page-14 locator in preserved module 283 for that particular inspected version. The short homogeneous curve class, one-step expansion and distortion hypotheses (H2)–(H4), and the dependence of its constant on these data and the subdivision length, are the inputs verified in native 281–284. The canonical sources in Section 68 use the enlarged class and its own constants from 284. They are not silently assigned the growth constant of a smaller class. This use is separate from the strong and weak anisotropic spaces and full hypotheses (H1)–(H5) of the external spectral perturbation theorem.

For the inspected 62-page arXiv v3 PDF of Baladi–Demers [7], Proposition 3.14 on PDF pages 25–27 uses finite Cantor rectangles and a mixing crossing time. Its potential and thermodynamic homogeneity conditions, including $qt_0/2 \geq 2$, are not automatically the native $q = 2$ partition or radius-uniform data. The inherited Theorem 70.4 has a different complete native proof; no radius-uniform consequence is attributed to that external proposition without verification of its hypotheses.

Dolgopyat–Nándori [8], Definition 2.2 on PDF pages 3–4 of the inspected 35-page arXiv v1 version, formulates a tested mixed local limit with its observable, minimality, subgroup, covariance and coboundary assumptions. Its bounded continuous and compactly supported test functions do not themselves give the exact-label raw roof essential-height theorem here. The original arithmetic reference is retained. General convex duality [9] is background; the special disk-word duality has its own complete native proof.

The fresh literature audit also distinguishes the 53-page author version of Demers–Zhang, where Lemma 3.2(b) is on page 14 and the full renewal is Lemma 5.5 on pages 35–36, from the 45-page arXiv v1, where that renewal is Lemma 6.2 on pages 32–33. Theorem 71.1 gives a native proof including the last-long-ancestor term $k = n$; it does not rely on a printed terminal-index convention. The present likelihood kernel, return tower and elementary conditional-variation arguments are proved directly. No external mixed local limit theorem is used to infer their remaining raw pointwise consequences.

The source, literature and proof ledgers distinguish retrieved external statements, complete author proofs, conditional interfaces, internal reconstruction and auxiliary finite checks. Actual

build and CI receipts are separately pinned to the mathematical inputs and the executed commit. None is represented as a journal decision or an independent human specialist certification.

Appendix M. Dependency scope of the physical jump-spectrum continuation

The preceding appendix is retained byte for byte from the immediate complete predecessor. Its source-stage descriptions of new results refer to that historical continuation. The current priority, source identity and proof status are those recorded in the introduction and the present appendix.

The present complete derivative and jump-spectrum interface. Theorems 80.3 and 81.1 identify the exact singular derivative and its marked Fourier energy for the full current smooth observation class. The finite physical graph is inherited from native 164, the scalar tame-germ consequences from native 40, 92 and 93, and the signed-current principle from native 153. The new proofs require the complete signed source and norm budget of native 323, 324, 330 and 332, whose continuous dynamical inputs remain author proofs with their stated hypotheses, rather than independently certified facts. The finite side decomposition uses fixed-radius closures and the definable-family cell theorems [10, Theorems 2.9–2.10 and 6.6–6.7]. It is applied to geometry, not to arbitrary C^1 weights.

The equality of variation masses in (80.5) concerns the derivative after signed cancellations. It does not identify a positive majorant with that signed measure. In particular a critical incoming coefficient can meet a negative constant-roof side coefficient. All same-label roof coincidences are combined before squaring in (81.4). The fixed-count frequency limit is never exchanged with the original collision limsup or a radius supremum.

The normalized central jump bound is necessary for the raw target. Its converse is not asserted: a continuous narrow peak has zero jump energy. The original complete local derivative charge of native 334 still includes both the integrable derivative and all its actual atoms. The old zero-clearance payment of native 322 transfers to the new energy through (81.15); the positive-width and other remaining charges are not thereby paid. No claim of full raw or inverse closure is added by the present two modules.

References

[1] Q. Qi,*Collision records, arithmetic, and raw local inversion in a triangular Lorentz gas*, A2-DYN clearance-roof author revision, October 15, 2026. Complete source and 47-page manuscript committed at `8a5a7c2be5630bf5919e9f74ee79de01af0dbb85`; active article Git tree `c2e9e83f39985a34690300433f4b40de4f1f88ac`.

[2] Q. Qi, *Collision records, arithmetic, and raw local inversion in a triangular Lorentz gas*, A2-DYN revision 111, October 14, 2026. Historical source archive pinned at Git tree `790e627efe3a89a4549123d7e9acacf24089c287`; its scoped source objects are retained in the verified historical DAG, without expanding recursive copies. Author commit `8c617cafa5e5493e26817401e866b2415522f7c0`.

[3] Q. Qi, *Collision records, arithmetic, and raw local inversion in a triangular Lorentz gas*, A2-DYN revision 110, October 14, 2026. Historical source, with recovered anchors retained in the provenance packet. Author commit `0abce3fcfbc735ff5cc6d085279abea5f32a7a22`.

[4] Q. Qi, *Collision records, arithmetic, and raw local inversion in a triangular Lorentz gas*, A2-DYN revision 109, October 14, 2026. Historical source, with recovered anchors retained in the provenance packet. Author commit `ab0ad79b9c101b82c80e33511529588d101a45c0`.

[5] Q. Qi, *Collision records, arithmetic, and raw local inversion in a triangular Lorentz gas*, A2-DYN revision 108, October 14, 2026. Historical manuscript and supplement; retained scoped source objects are recoverable from the historical DAG. Author commit `2144665f89362ca80116f1999d99c80875833dcf`.

[6] M. F. Demers and H.-K. Zhang, *A functional analytic approach to perturbations of the Lorentz gas*, Communications in Mathematical Physics **324** (2013), 767–830. The prior audit inspected the author version dated 10 October 2013; the fresh version check is recorded in the literature audit.

[7] V. Baladi and M. F. Demers, *Thermodynamic formalism for dispersing billiards*, Journal of Modern Dynamics **18** (2022), 415–493. The prior audit inspected the author version dated 19 September 2022; the fresh primary-source check is recorded in the literature audit.

[8] D. Dolgopyat and P. Nándori, *On mixing and the local central limit theorem for hyperbolic flows*, Ergodic Theory and Dynamical Systems **40** (2020), 142–174, Definition 2.2 and Section 6.4, Proposition 6.3. The exact inspected version and scope are recorded in the literature audit.

[9] S. Boyd and L. Vandenberghe, *Convex Optimization: Duality*, official lecture notes, `https://web.stanford.edu/class/ee364a/lectures/duality.pdf`. Used for general background; the special disk-word duality is proved in full in the present article.

[10] M. Coste, *An Introduction to O-minimal Geometry*, Dottorato di Ricerca in Matematica, Università di Pisa, Istituti Editoriali e Poligrafici Internazionali, Pisa, 2000. Theorems 2.9–2.10 and 6.6–6.7. Author-hosted text: `https://perso.univ-rennes1.fr/michel.coste/polyens/OMIN.pdf`.