

CONDUCTOR STRATA AND NONREDUCED MULTIPLICATION FAILURE SCHEMES

QIAN QI

ABSTRACT. We study multiplication failure as a coherent-cokernel problem, including its nonreduced structure. For a finite locally free algebra, stable multiplication admits a canonical generated subalgebra; stratifying its action on the quotient gives conductor flags compatible with arbitrary base change on each stratum. In codimension two the controlling quotients have rank three or four, with a quadratic-algebra tower in the latter case. For the truncated local algebra in e variables of order h , the entire generating $(e+1)$ -plane family has stable failure ideal equal to the $\binom{e+h-1}{e+1}$ th power of a determinant divisor. This determines its primary structure and every ordinary power. A triangular conductor argument transports these schemes to global multiplication on $\mathbb{P}^e$ in the dimension-independent range $n \geq 2h-1$, sharp uniformly over generating ranks. We retain the sharp rank-dependent curve comparison, the hyperplane theorem controlled by rank-two quotients, the contact primary classifications and normalization, and the reduced three-plane realization of Haiman's diagonal ideal. The primary assertions always specify their parameter spaces; coherent cokernel comparisons, rather than primary decompositions of arbitrary specializations, are what commute universally with base change.

1. INTRODUCTION

The multiplication map of a linear series can fail to be surjective with a scheme structure which changes even when its support does not. We study this structure for series containing the sections vanishing on a finite divisor. The method has two parts: a global conductor theorem identifies the actual cokernel with a finite-algebra cokernel, and a quotient-algebra description determines the latter scheme without passing to its radical.

Two results take this comparison beyond reduced contacts and rank-two quotients. First, the stable generated algebra carries a representation on the multiplication defect. Its exact-rank strata identify the conductor quotient as a flag $C \subset Q$ with C acting faithfully on Q/C (Theorem 5.1). In codimension two there are two mechanisms: a rank-three quotient with its scalar line, or a rank-four quotient which is quadratic over a quadratic algebra (Theorem 5.3). The transition between them can itself be nonreduced.

Second, let Z be the order- h fat point of $\mathbb{P}^e$, with $e \geq 2$ and $h \geq 3$. For every $n \geq 2h-1$, take all series containing $H^0(\mathcal{I}_Z(n))$ and restricting to a generating $(e+1)$ -plane on Z . In symmetric degrees $m \geq h-1$ their failure ideal is

$$\mathcal{I}_\Delta^{\binom{e+h-1}{e+1}}, \tag{1.1}$$

where Δ is the integral determinant divisor of the map from the augmentation kernel of the restricted series to $\mathfrak{m}/\mathfrak{m}^2$. This is the full primary ideal, and all its powers remain primary (Corollary 7.3). The proof separates a filtered substitution determinant from a triangular global conductor argument. It gives three-plane results on every $\mathbb{C}[x, y]/(x, y)^h$ and works in arbitrary

Date: September 22, 2026.

2020 Mathematics Subject Classification. 14M12, 14H50, 13C40, 14B05, 14N15.

Key words and phrases. Multiplication of sections, finite quotient algebras, Hilbert schemes, primary decomposition, nilpotent structures.

embedding dimension; no diagonal-ideal theorem is used. The global range depends on the fat-point order, rather than its binomially growing length.

Let $D \subset \mathbb{P}^1$ have length d , put $E_{n,D} = H^0(\mathcal{O}_D(n))$ and $W_D = H^0(\mathcal{O}(n)(-D))$, and let $A \subset E_{n,D}$ be a generating k -plane. Write $U_A = \text{res}_D^{-1}(A)$. For $1 \leq k < d$ our conductor range is $n \geq 2d - k$. In every symmetric degree $m \geq 2$ it gives

$$\text{coker}(\text{Sym}^m \mathcal{U}_A \rightarrow V_{mn}) \simeq \text{coker}(\text{Sym}^m \mathcal{A} \rightarrow E_{mn,D}). \quad (1.2)$$

The isomorphism holds on the entire relative generating Grassmannian and after arbitrary base change. The range is sharp uniformly in D, A, m : at $n = 2d - k - 1$ an explicit monomial series already violates it in degree two. The exact fibrewise obstruction is the first cohomology of an evaluation kernel. On a curve of genus g , intrinsic cohomological hypotheses give the same transport; the uniform sufficient bound is $\deg L \geq 2d + 2g - 1$.

The defect-one structural theorem concerns generating hyperplanes in an arbitrary finite locally free algebra, not only pencils or powers of one element. Its failure scheme in all degrees $m \geq 2$ is the space of rank-two quotient algebras with a generating line. The identification is an isomorphism of functors on arbitrary, including nonreduced, test schemes. It follows from the conductor of a unital hyperplane subalgebra and a cyclic presentation of the multiplication cokernel (Theorem 4.2).

For $D = \sum_i d_i p_i$ this gives a complete primary classification of all generating $(d - 1)$ -planes. The components supported on $2p_i$ have transverse algebra

$$G_{d_i} = \mathbb{C}[u, v]/(F_{d_i}, F_{d_i+1}), \quad F_0 = 0, \quad F_1 = 1, \quad F_{j+1} = uF_j + vF_{j-1};$$

those supported on $p_i + p_j$ have transverse algebra $\mathbb{C}[x, y]/(x^{d_i}, y^{d_j})$. Their lengths are $\binom{d_i}{2}$ and $d_i d_j$, but their exact nilpotency indices are $2d_i - 3$ and $d_i + d_j - 1$. The reduced components are disjoint smooth curves on the generating open. The moving scheme is smooth and is a generating open of a projective-line bundle over $\text{Sym}^2 C \times \text{Sym}^{d-2} C$. These results apply to global multiplication on $\mathbb{P}^1$ for $n \geq d + 1$. They classify the whole contact-hyperplane family for every multiplicity partition, not a collection of transverse examples.

There is also a complete stable description for generating three-planes on a reduced divisor. On unit frames its ideal is

$$\bigcap_{i < j} (x_i - x_j, y_i - y_j).$$

The equality, and the analogous equality for every power of the ideal, use Haiman's diagonal-ideal theorem explicitly. The step specific to multiplication is an exact total-degree truncation and the conductor comparison; the radical-ideal theorem is a classical input, not a new assertion here.

The pencil family and the distinction between fibres and total spaces. The earlier contact-pencil classification remains part of the article, now in the improved range. We record its complete statement to fix the bases of its primary assertions.

Theorem 1.1 (Contact-pencil failure schemes). *Let $d \geq 4$ and $n \geq 2d - 2$. On the generating open $G_D^\circ \subset \text{Gr}(2, E_{n,D})$, let $\mathcal{J}_m$ be the zeroth Fitting ideal of the global multiplication cokernel. For $2 \leq m < d - 1$, this ideal is zero: the failure scheme is the whole parameter space. For $m \geq d - 1$ it is a Cartier divisor, and, on a fixed divisor $D = \sum_i d_i p_i$, its complete primary decomposition is*

$$\mathcal{J}_m = \bigcap_{i: d_i \geq 2} \mathcal{I}_{T_i}^{\binom{d_i}{2}} \cap \bigcap_{i < j} \mathcal{I}_{E_{ij}}^{d_i d_j}. \quad (1.3)$$

Here T_i and E_{ij} are failure to separate $2p_i$ and $p_i + p_j$. These are distinct smooth irreducible divisors; there are no embedded associated points. The exact local equations, all coranks and

infinitesimal layers are given in Theorems 9.2 and 9.3. Over a moving multiplicity stratum, primary descent is governed by Lemma 8.1.

With D unrestricted, the pencil failure divisor is integral, and its normalization is the smooth finite incidence marking a scalar length-two subdivisor. The normalization and its singular-support test are given in Theorem 10.2 and Proposition 10.3.

On a unit pencil frame, with $D = (g)$ and ratio v , the determinant is the classical index form $\Delta(g, v) = \det[1, v, \dots, v^{d-1}]$. Its relation to the characteristic polynomial χ_v is

$$\text{disc}(\chi_v) = \text{disc}(g)\Delta(g, v)^2.$$

The equation of failure is $\Delta = 0$, not the discriminant on the left: the latter also vanishes when g has repeated roots, even if v generates the algebra. The quotient and index-form constructions are related to the schemes of generators in [29]. The Grassmannian cohomology presentation of G_r is likewise classical. What must be proved here is the exact identification with the multiplication schemes, including primary fibres and base change.

The dependence of the arguments is

$$\begin{aligned} \text{evaluation kernel} &\implies \text{conductor cokernel,} \\ \text{unital hyperplane conductor} &\implies \text{rank-two quotients,} \\ \text{rank-two quotients} &\implies \text{primary fibres and smooth total scheme,} \\ \text{index form} &\implies \text{pencil primary fibres and normalization,} \\ \text{polarized Cayley-Hamilton + Haiman} &\implies \text{three-plane schemes.} \end{aligned}$$

None of these arguments depends on the statistical interpretation, the quadratic excess classification, or the unrestricted hyperplane embedded structure. Those developments, including all their proofs and hypotheses, are retained in the full complementary manuscript. The present family is not identified with the unrestricted ambient Grassmannian. Section 12 gives the precise classical inputs and the remaining limitation in the nearest-source comparison.

2. SHARP CONDUCTOR REDUCTION AND A CURVE CRITERION

All schemes in this article are over $\mathbb{C}$. Put $V_r = H^0(\mathbb{P}^1, \mathcal{O}(r))$, with $V_r = 0$ for $r < 0$. For a subspace U of sections, U^j denotes its multiplication image, not its abstract symmetric power.

Lemma 2.1 (The evaluation-kernel bound). *Let $U \subset V_n$ be base-point-free, of codimension c , and write $M_U = \ker(U \otimes \mathcal{O} \rightarrow \mathcal{O}(n))$. There are integers $a_i \geq 1$, with $\sum_i a_i = n$, such that $M_U = \bigoplus_{i=1}^{n-c} \mathcal{O}(-a_i)$. If $a = \max_i a_i$, then for $t \geq 0$,*

$$\text{coker}(U \otimes V_t \rightarrow V_{n+t}) \simeq H^1(M_U(t)), \quad UV_t = V_{n+t} \iff t \geq a - 1.$$

In particular $a \leq c + 1$, so $UV_t = V_{n+t}$ for $t \geq c$.

Proof. The splitting theorem for vector bundles on $\mathbb{P}^1$ applies to the locally free evaluation kernel. Its degree is $-n$ and its rank is $n - c$. Moreover $H^0(M_U) = 0$, because $U \rightarrow H^0(\mathcal{O}(n))$ is injective. Thus all its summands have negative degree. Twisting the evaluation sequence by $\mathcal{O}(t)$ and using $H^1(\mathcal{O}(t)) = 0$ gives the asserted cokernel. The formula for $H^1(\mathcal{O}(t - a_i))$ gives the equivalence. Finally the other $n - c - 1$ positive integers have sum at least $n - c - 1$, whence $a \leq n - (n - c - 1) = c + 1$. $\square$

Theorem 2.2 (Sharp uniform conductor range). *Let $D \subset \mathbb{P}^1$ have length $d \geq 2$, let $1 \leq k < d$, and suppose*

$$n \geq 2d - k. \tag{2.1}$$

Let $A \subset H^0(\mathcal{O}_D(n))$ be a generating k -plane, and put $W = H^0(\mathcal{O}(n)(-D))$ and $U = \text{res}_D^{-1}(A)$. For every $m \geq 2$,

$$WU^{m-1} = H^0(\mathcal{O}(mn)(-D)). \quad (2.2)$$

On the entire relative generating Grassmannian over $\text{Hilb}^d(\mathbb{P}^1)$, restriction gives

$$\text{coker}(\text{Sym}^m \mathcal{U} \rightarrow V_{mn} \otimes \mathcal{O}) \simeq \text{coker}(\text{Sym}^m \mathcal{A} \rightarrow \pi_* \mathcal{O}_D(mn)). \quad (2.3)$$

These are isomorphisms of coherent sheaves after arbitrary base change; all their Fitting ideals agree. For a particular geometric fibre, the weaker condition $n - d \geq \max_i a_i(U) - 1$ suffices for the same conclusion. Write $X_{d,k}^\circ$ for this relative generating Grassmannian. The bound (2.1) is sharp as a bound uniform in D, A and simultaneously in all $m \geq 2$. For $k = d$ the cokernel assertion is immediate from $U = V_n$ in the range $n \geq d - 1$; the saturation assertion in that case holds for $n \geq d$.

Proof. Write $D = (g)$, so $W = gV_{n-d}$. The series U is base-point-free: W generates away from D and A generates on D . Its codimension in V_n is $c = d - k$. Lemma 2.1 and $n - d \geq c$ give $WU = gV_{2n-d}$. Further multiplication by U uses input degrees $2n - d, 3n - d, \dots$, all at least c , and proves (2.2). The same argument uses $a - 1$ in place of c on a particular fibre.

All restriction targets and kernels are vector bundles: their first cohomology groups vanish in this range. The map $\mathcal{W} \otimes \text{Sym}^{m-1} \mathcal{U} \rightarrow \pi_* I_D(mn)$ is onto on every geometric fibre, hence is a surjection of vector bundles. It remains onto after every base change. Locally split $0 \rightarrow \mathcal{W} \rightarrow \mathcal{U} \rightarrow \mathcal{A} \rightarrow 0$. The kernel of the map on symmetric powers is the image of $\mathcal{W} \otimes \text{Sym}^{m-1} \mathcal{U}$. Quotienting the restriction square by these kernels proves (2.3). Fitting base change now applies [25].

For sharpness take $n = 2d - k - 1$, $D = d\{0\}$, and, in the affine coordinate t , take the degree- n homogenizations of

$$U = \langle 1, t, \dots, t^{k-1}, t^d, t^{d+1}, \dots, t^n \rangle.$$

This is base-point-free and is the inverse image of a generating k -plane on D . Exponents of its quadratic products lie in

$$[0, 2k - 2] \cup [d, n + k - 1] \cup [2d, 2n].$$

None is $2d - 1$. Thus $t^{2d-1} \notin U^2$, although it belongs to $H^0(I_D(2n))$. Its nonzero cokernel class restricts to zero on D , so even the cokernel comparison fails at this boundary in degree two. This does not assert a sharp threshold separately for each $m > 2$. $\square$

The numerical bound can be replaced by a cohomological hypothesis on a curve. Here and below an ideal such as I_D is the ideal on the curve, whereas a Fitting ideal is on the parameter space.

Proposition 2.3 (An exact cohomological test). *Let C be a smooth projective curve, let L be a line bundle, and let $U \subset H^0(L)$ be base-point-free. Put $W = H^0(L(-D))$. If $H^1(L(-D)) = 0$, then*

$$\text{coker}(W \otimes U \rightarrow H^0(L^2(-D))) \simeq H^1(M_U \otimes L(-D)).$$

More generally, for $B_j = U^j \subset H^0(L^j)$ and its evaluation kernel M_{B_j} , the obstruction to $WB_j = H^0(L^{j+1}(-D))$ is exactly $H^1(M_{B_j} \otimes L(-D))$.

Proof. Tensor the evaluation sequence for B_j by $L(-D)$ and take cohomology. The next term is $B_j \otimes H^1(L(-D)) = 0$. $\square$

Theorem 2.4 (Conductor transport on curves). *Let C have genus g , and let D be effective of length $d \geq 2$. Suppose $L(-D)$ is globally generated, $\text{Sym}^2 H^0(L(-D)) \rightarrow H^0(L^2(-2D))$ is surjective, and*

$$H^1(L^j(-2D)) = 0 \quad (j \geq 1). \quad (2.4)$$

For every generating $A \subset H^0(L|_D)$ and every $m \geq 2$, $U = \text{res}_D^{-1}(A)$ satisfies

$$WU^{m-1} = H^0(L^m(-D)), \quad \text{coker}(\text{Sym}^m U \rightarrow H^0(L^m)) \simeq \text{coker}(\text{Sym}^m A \rightarrow H^0(L^m|_D)).$$

In particular these assertions hold if

$$\deg L \geq 2d + 2g - 1. \quad (2.5)$$

For the relative assertion, let $f : C \rightarrow S$ be smooth and projective of relative dimension one, with S locally Noetherian over $\mathbb{C}$, let D be a relative effective Cartier divisor finite flat of degree d , and let L be invertible. Assume the preceding hypotheses on every geometric fibre. For each $j \geq 1$ and $b = 0, 1, 2$, put $E_{j,b} = f_* L^j(-bD)$. These are locally free, $R^1 f_* L^j(-bD) = 0$, and their formation commutes with arbitrary base change. The same base-change property holds for the finite direct images of $L^j|_D$ and $L^j(-D)|_D$. On the relative generating Grassmannian form $\mathcal{U}$ as the inverse image of $\mathcal{A}$ in $E_{1,0}$. The displayed saturation and cokernel assertions then hold as maps of these bundles and sheaves, after every base change.

Proof. The hypotheses imply $H^1(L^j(-D)) = 0$ by the exact sequence with quotient on D . They make U base-point-free. It contains a pencil without base points, obtained by avoiding the finitely many zeros of one section. The pencil sequence, tensored by $L^j(-2D)$, shows that multiplication onto $H^0(L^{j+1}(-2D))$ is surjective for $j \geq 2$: its obstruction is $H^1(L^{j-1}(-2D))$. Starting with the assumed surjectivity for W^2 , this proves $W^2 U^{m-2} = H^0(L^m(-2D))$.

The map $W \rightarrow H^0(L(-D)|_D)$ is onto by (2.4). Choose $a \in A$ a unit on D and lift it to $u \in U$. Multiplication by u^{m-1} identifies the first normal quotients in these degrees. Consequently Wu^{m-1} fills that quotient in $H^0(L^m(-D))$. Together with the double-vanishing space just proved, this gives the assertion about WU^{m-1} . The locally split restriction square gives the cokernel isomorphism as before.

Under (2.5), the line bundle $L(-D)$ has degree at least $2g + 1$. The classical normal-generation theorem in this degree range gives the required quadratic surjectivity and global generation; see [14, p. 73, Introduction]. Serre duality gives (2.4), since $\deg L(-2D) \geq 2g - 1$. For the relative assertion, the assumed fibrewise H^1 -vanishing for $b = 2$ implies the same vanishing for $b = 1, 0$ by restriction to D . Cohomology and base change for these flat line bundles on a proper flat relative curve gives the named direct images, with no higher cohomology; finite flat pushforward gives the two restriction bundles. In particular $0 \rightarrow E_{1,1} \rightarrow E_{1,0} \rightarrow f_*(L|_D) \rightarrow 0$ is a locally split exact sequence. The quadratic map $\text{Sym}^2 E_{1,1} \rightarrow E_{2,2}$ and each map $E_{1,1} \otimes \text{Sym}^{m-1} \mathcal{U} \rightarrow E_{m,1}$ are surjective on every geometric fibre by the fixed-curve proof. They are thus surjections of vector bundles and remain so under arbitrary pullback. The same locally split restriction square proves the relative cokernel assertion; no base-change assumption on an unflattened image U^j is needed. $\square$

For example on any elliptic curve, a line bundle of degree at least $2d + 1$ transports all the finite-contact schemes below. The curve statement is not a sharpness assertion in positive genus. Its intrinsic hypotheses, and the exact obstruction in Proposition 2.3, distinguish a failure of regularity from a failure of the finite-contact multiplication.

3. GENERATING OPENS AND FINITE PRESENTATIONS

Here “generating” refers to generation as an ideal or invertible module. It does not assert generation as a unital algebra.

Lemma 3.1 (The relative generating open). *Let B/S be a finite locally free commutative algebra over a $\mathbb{C}$ -scheme, and let $A \subset B$ be a subbundle. The following conditions are equivalent: $B \otimes A \rightarrow B$ is surjective; on every geometric fibre A has nonzero image in every residue field of B ; and the open $A^\times$ of sections of A which are units of B is smooth and surjective over S .*

These conditions define an open in the Grassmannian. The assertions have the same form for a subbundle of an invertible B -module, using generating sections in place of units.

Proof. Surjectivity is equivalent to its geometric-fibre version by Nakayama's lemma. On a finite algebra over an algebraically closed field, an ideal is the unit ideal exactly when its images in all residue fields are nonzero. These evaluations cut out finitely many proper hyperplanes in A . Their complement is nonempty over an infinite field and consists precisely of units. On the total space of the vector bundle A , invertibility is the open condition $\det(b \mapsto ab) \neq 0$. This open is smooth over S , and the preceding fibre calculation proves surjectivity. Conversely a unit section after a faithfully flat cover generates the unit ideal, which descends. The Grassmannian open is the complement of the support of the cokernel of $B \otimes A \rightarrow B$. Local trivializations of an invertible module prove the last assertion. $\square$

Lemma 3.2 (Total-degree truncation). *Let B be a rank- d finite locally free algebra over a $\mathbb{C}$ -algebra R , and let $b_1, \dots, b_q \in B$. The R -subalgebra they generate is spanned as an R -module by the monomials of total degree at most $d - 1$. This assertion commutes with base change, with polynomial reduction coefficients on a free chart.*

Proof. Set $w = \sum_j t_j b_j$ in $B[t_1, \dots, t_q]$ and apply Cayley–Hamilton to multiplication by w . The coefficient of t^α , for $|\alpha| = d$, in w^d is $\binom{d}{\alpha} b^\alpha$. All other terms have B -monomial degree less than d : coefficients of the characteristic polynomial are scalars in $R[t]$. The multinomial coefficient is invertible in R . This expresses each degree- d monomial using lower degrees. Multiplying these identities and inducting reduces every higher monomial. The adjugate identity for Cayley–Hamilton makes the coefficients polynomial in the multiplication table. $\square$

4. GENERATING HYPERPLANES AND LENGTH-TWO QUOTIENTS

We now solve a multigenerator problem without assuming that the finite algebra is monogenic, reduced, or constant. If B is a finite locally free commutative algebra of rank d over a scheme S , let $\text{Hilb}^2(B/S)$ denote the functor of quotient algebras $B_T \twoheadrightarrow Q$ that are locally free of rank two over T . It is the relative Hilbert scheme of two points of $\text{Spec}_S B$. A generating line in Q means a line subbundle whose Q -span is Q .

Lemma 4.1 (The conductor of a unital hyperplane). *Unital subalgebras $S_0 \subset B_T$ which are hyperplane subbundles are naturally equivalent to rank-two quotient algebras of B_T . The maps are*

$$S_0 \longmapsto B_T/J, \quad J = \{b \in B_T : bB_T \subset S_0\}, \quad (B_T \twoheadrightarrow Q) \longmapsto \text{preimage}(\mathcal{O}_T \cdot 1).$$

The conductor J is an ideal subbundle of rank $d - 2$, and its formation commutes with arbitrary base change.

Proof. Work locally with $T = \text{Spec } R$, trivialize B/S_0 , and write it as the target of $\phi : B \rightarrow R$. Choose e with $\phi(e) = 1$. Since $1 \in S_0$, the bilinear form $\beta(b, c) = \phi(bc)$ has $\beta|_{S_0 \times S_0} = 0$ and $\beta(1, e) = 1$. An element of its radical first lies in S_0 , by pairing with 1; it then lies in the kernel of the split surjection

$$S_0 \longrightarrow R, \quad s \longmapsto \phi(se),$$

which sends 1 to 1. Thus the radical has rank $d - 2$, and B/J has basis the classes of $1, e$. Associativity shows that the radical is an ideal: $\beta(jb, c) = \beta(j, bc) = 0$. It is precisely the displayed conductor. This split-kernel description proves base change and gives $S_0/J = R \cdot 1$.

Conversely, in a rank-two locally free algebra Q , the unit is a unimodular vector, so $Q/R \cdot 1$ is a line bundle. The inverse image of the scalars is a unital hyperplane subalgebra. No nonzero ideal of Q can be contained in $R \cdot 1$: in a basis $1, e$, if $c \cdot 1$ belongs to such an ideal, then ce must be scalar, so $c = 0$. Its conductor is therefore exactly the original kernel. The two constructions are inverse and are independent of all trivializations. $\square$

Theorem 4.2 (The universal hyperplane failure scheme). *Let B/S be finite locally free of rank $d \geq 3$ over a $\mathbb{C}$ -scheme. On $X = \text{Gr}_S(d-1, B)^\circ$, the generating open, let $\mathcal{A}$ be the tautological hyperplane and put*

$$\mathcal{C}_m = \text{coker}(\text{Sym}^m \mathcal{A} \rightarrow B), \quad Y_m = V(\text{Fitt}_0 \mathcal{C}_m).$$

For every $m \geq 2$ the ideal of Y_m is the same. The common scheme Y is naturally isomorphic to

$$\{(B \rightarrow Q, \ell) : \text{rank } Q = 2, \ell \subset Q \text{ a generating line}\}. \quad (4.1)$$

The map to X sends (Q, ℓ) to the inverse image of ℓ . If $i : Y \hookrightarrow X$ and $B_Y \twoheadrightarrow Q \supset \ell$ is the universal quotient with its generating line, the canonical formula is

$$\mathcal{C}_m \simeq i_*(Q/\ell^{(m)}), \quad \ell^{(m)} = \text{im}(\ell^{\otimes m} \rightarrow Q). \quad (4.2)$$

Both $\ell^{(m)}$ and its quotient are line bundles on Y . In particular all positive-index Fitting ideals are the unit ideal. The isomorphism of schemes and all these assertions commute with arbitrary base change. The same statement holds for an invertible B -module in place of B , with target its m th tensor power.

Proof. Locally for the smooth topology, choose a unit $a \in \mathcal{A}$. Such choices cover X : on each geometric fibre avoid the finitely many evaluation hyperplanes. Replace $\mathcal{A}$ by $A' = a^{-1}\mathcal{A}$, and the target by multiplication by a^{-m} . Now $1 \in A'$. Split $B = A' \oplus Re$ locally. Since $(A')^m$ contains A' , elementary row and column operations present the cokernel by a single row. For $m = 2$ its entries are exactly the coefficients of the products ss' in B/A' , for $s, s' \in A'$. Call their ideal I . Thus Y_2 represents the condition that A' is a subalgebra, with its full scheme structure, not just its support. Modulo I all higher products lie in A' . Conversely every quadratic product is an m -fold product by inserting 1's. Hence the residual ideals equal I for every $m \geq 2$, and the cokernel is locally R/I .

Lemma 4.1 supplies the unique rank-two quotient on Y . Its kernel J also lies in $\mathcal{A} = aA'$, since J is an ideal and $aJ = J$. The image of $\mathcal{A}$ is a generating line in Q . The construction does not depend on a . Indeed another unit $b \in \mathcal{A}$ gives $b/a \in A'$, and its inverse belongs to A' by Cayley–Hamilton in B (the constant coefficient is a unit). Thus $(b/a)^{-1}A' = A'$ and the conductor is unchanged. This also proves descent from the smooth cover. Conversely the inverse image of a generating line in a rank-two quotient becomes the inverse image of the scalars after division by a unit, so it satisfies precisely the residual equations. These constructions are inverse on every test scheme, proving the scheme isomorphism. On Y , the normalized subalgebra A' has conductor J , and $J \subset a^m A' = (\mathcal{A})^m$ since a is a unit and J an ideal. Passing to Q identifies $(\mathcal{A})^m/J$ with $\ell^{(m)}$. This gives the canonical map in (4.2); the local cyclic presentations identify it as an isomorphism. Multiplication by a local unit spanning ℓ makes $\ell^{(m)}$ a direct-summand line, so the quotient is indeed a line bundle. For an invertible B -module P , put $P_Q = P \otimes_B Q$; the same formula is $i_*(P_Q^{\otimes m}/\ell^{(m)})$, where now $\ell \subset P_Q$. Trivialization of P gives the local proof and this intrinsic expression supplies its gluing. $\square$

4.1. Every fixed multiplicity partition. For $r \geq 2$ define polynomials

$$F_0 = 0, \quad F_1 = 1, \quad F_2 = u, \quad F_{j+1} = uF_j + vF_{j-1}, \quad G_r = \mathbb{C}[u, v]/(F_r, F_{r+1}). \quad (4.3)$$

For positive r, s put $H_{r,s} = \mathbb{C}[x, y]/(x^r, y^s)$.

Theorem 4.3 (Complete primary schemes for contact hyperplanes). *Let $D = \sum_i d_i p_i$ have length $d \geq 3$. Consider the entire generating open $\text{Gr}(d-1, H^0(\mathcal{O}_D))^\circ$. For every $m \geq 2$, its multiplication failure scheme is the disjoint union*

$$Y_D \simeq \coprod_{i:d_i \geq 2} (\mathbb{A}^1 \times \text{Spec } G_{d_i}) \amalg \coprod_{i < j} (\mathbb{G}_m \times \text{Spec } H_{d_i, d_j}). \quad (4.4)$$

The identifications use local coordinates on D and trivializations of the restricted line bundle. Each displayed summand is a primary component on this fixed-divisor parameter space. There are no embedded associated points. All reduced components are smooth and pairwise disjoint. Their transverse lengths and exact nilradical indices are, respectively,

length-two support	length	nilpotency index
$2p_i$	$\binom{d_i}{2}$	$2d_i - 3$
$p_i + p_j$	$d_i d_j$	$d_i + d_j - 1$.

The convention for a reduced component is index one. Every failure fibre has cokernel dimension one. On binary inverse-image subseries these are exactly the multiplication failure schemes whenever $n \geq d + 1$; on curves they are the same schemes under Theorem 2.4.

Proof. A degree-two quotient of the product algebra $\prod_i \mathbb{C}[z_i]/(z_i^{d_i})$ decomposes according to the ranks of its idempotent factors. These ranks are locally constant even on a nonreduced base. Its possibilities are rank two at a single factor, or rank one at each of two distinct factors. This is a decomposition into open and closed Hilbert subschemes, not merely a decomposition of closed points.

For rank two in $\mathbb{C}[z]/(z^r)$, the elements $1, z$ form a basis in every geometric quotient, hence on every family by Nakayama. Write $z^2 = uz + v$. Then

$$z^j = F_j(u, v)z + vF_{j-1}(u, v).$$

The relation $z^r = 0$ is equivalent to $F_r = vF_{r-1} = 0$, equivalently $F_r = F_{r+1} = 0$. Thus this Hilbert scheme is $\text{Spec } G_r$. A rank-one quotient at a factor of length r is $\text{Spec } \mathbb{C}[x]/(x^r)$, so the two-factor case is $\text{Spec } H_{r,s}$. A generating line in the first rank-two algebra is uniquely spanned by $1 + \tau z$; in the split two-factor algebra it is uniquely spanned by $(1, t)$, with t a unit. Theorem 4.2 now proves (4.4) over arbitrary test schemes.

The algebra G_r is Artin local: if $v \neq 0$, two consecutive zero values in the recurrence force $F_1 = 0$, and if $v = 0$ they force $u = 0$. The two equations form a regular sequence in $\mathbb{C}[u, v]$. With $\deg u = 1$, $\deg v = 2$, its Hilbert series is

$$\frac{(1 - t^{r-1})(1 - t^r)}{(1 - t)(1 - t^2)}. \quad (4.5)$$

It has length $\binom{r}{2}$ and weighted socle degree $2r - 4$. To see the exact ordinary nilpotency, identify u and $-v$ with the elementary symmetric functions in two variables. Then $F_j = h_{j-1}$ and

$$G_r \simeq H^{2*}(\text{Gr}(2, r), \mathbb{C}), \quad (4.6)$$

where the cohomological degree is divided by two. This is the classical presentation by h_{r-1}, h_r , not a new presentation of Grassmannian cohomology; the rectangular Schur basis is recalled in [31, Theorem 2.7 and the following discussion]. Successive multiplication by $u = s_{(1)}$ adds one box. The only partition of size $2r - 4$ in the 2 by $r - 2$ rectangle is $(r - 2, r - 2)$; the ballot-path count of its standard tableaux is $\frac{1}{r-1} \binom{2r-4}{r-2}$. Consequently

$$u^{2r-4} = \frac{1}{r-1} \binom{2r-4}{r-2} (-v)^{r-2} \neq 0.$$

Every monomial of ordinary degree $2r - 3$ has weighted degree at least $2r - 3$ and vanishes. This proves the index $2r - 3$, including $r = 2$. In $H_{r,s}$ the top monomial is $x^{r-1}y^{s-1}$, giving length rs and index $r + s - 1$.

Each summand is a complete intersection with reduced support a smooth integral curve, so it is primary and has no embedded points. Equivalently, over a local Artin coefficient algebra, a polynomial with nonzero reduction is a nonzerodivisor, as one checks on the successive powers

of the maximal ideal. The same holds after inverting t . This proves the primary assertions also globally on the indicated affine curves. The cokernel assertion follows from Theorem 4.2. The two transports follow from Theorems 2.2 and 2.4. $\square$

Theorem 4.4 (The smooth moving scheme). *Let $d \geq 3$. On the relative generating Grassmannian of $(d-1)$ -planes over $\text{Hilb}^d(C)$, for a smooth connected projective curve C , the finite-contact failure scheme is a generating open of a projective-line bundle over*

$$\text{Sym}^2 C \times \text{Sym}^{d-2} C, \quad (Z, Q) \mapsto D = Z + Q.$$

It is smooth, irreducible, and of dimension $d+1$. Its projection to $\text{Sym}^d C$ is flat of relative dimension one. The finite map of divisor spaces has degree $\binom{d}{2}$; the primary lengths in Theorem 4.3 sum to this degree on every fixed fibre. Under either conductor theorem, these are statements about the original global multiplication scheme.

Proof. A nested pair $Z \leq D$ of effective divisors on a smooth curve is uniquely $D = Z + Q$. This holds in families: a finite flat subscheme of a smooth relative curve is a relative Cartier divisor, and inclusions of their invertible ideals give the residual divisor. The incidence in Theorem 4.2 is therefore exactly the following canonical open projective bundle. On $T = \text{Sym}^2 C \times \text{Sym}^{d-2} C$ let $\mathcal{Z} \subset C \times T$ be the universal degree-two divisor and set $F = (\pi_{\mathcal{Z}})_* \mathcal{O}_{\mathcal{Z}}$. Use the projectivization $\mathbb{P}_T^{\text{lines}}(F)$ parametrizing line subbundles of F , and retain the open where the tautological line ℓ satisfies $F\ell = F$. This is the claimed bundle, not an unspecified local form. For a restricted line bundle replace F by $(\pi_{\mathcal{Z}})_*(L|_{\mathcal{Z}})$ and use generation as an F -module. The open is nonempty on every fibre of its smooth irreducible base, and hence is smooth and irreducible of dimension $d+1$.

The map of divisor spaces is finite (proper with finite geometric fibres). Both spaces are smooth of dimension d ; the finite source is Cohen–Macaulay and the target regular, so the map is flat. Over a reduced divisor it has $\binom{d}{2}$ points. This proves the degree and flatness assertions; alternatively, the fibre lengths computed above give $\sum_i \binom{d_i}{2} + \sum_{i < j} d_i d_j = \binom{d}{2}$. The open projective bundle is smooth of relative dimension one, so its composite projection is flat of that relative dimension. $\square$

4.2. A direct residual presentation. For comparison with the multiplication matrix, normalize a unit and write, on the unital hyperplane chart in $\mathbb{C}[z]/(z^r)$,

$$A' = \langle 1, z^i - c_i z : 2 \leq i \leq r-1 \rangle.$$

Put $c_1 = 1$ and $c_j = 0$ for $j \geq r$. After eliminating the pivot columns, the remaining entries of the quadratic matrix are

$$c_{i+j} - c_i c_{j+1} - c_j c_{i+1} + c_2 c_i c_j \quad (2 \leq i \leq j \leq r-1). \quad (4.7)$$

For $r \geq 4$, the relations with $i = 2$ give $c_{j+2} = u c_{j+1} + v c_j$, with $u = c_2$, $v = c_3 - c_2^2$. They eliminate $c_4, \dots, c_{r-1}$ and leave F_r, F_{r+1} . Every other relation follows from the recurrence identity

$$F_{i+j} = F_i F_{j+1} + F_j F_{i+1} - u F_i F_j,$$

proved by induction in j . For $r = 3$ the single residual entry is c_2^3 , agreeing with $G_3 \simeq \mathbb{C}[u]/(u^3)$. For $r = 4$ one obtains

$$G_4 = \mathbb{C}[u, v]/(u^3 + 2uv, u^2v + v^2), \quad \text{length } G_4 = 6, \quad \mathfrak{m}^5 = 0 \neq \mathfrak{m}^4.$$

Thus the multigenerator transverse scheme is not a powered smooth divisor, even though its length equals the corresponding pencil weight. The smooth moving scheme has these nonreduced fibres; nonreducedness of a fibre is not an embedded component of the total scheme. Nor does this determine the embedded primary ideal along the evaluation curve in the unrestricted, non-generating hyperplane problem retained in the complementary material.

4.3. A noncurvilinear algebra and an embedded component. The general finite-algebra theorem is not restricted to the primary patterns arising from divisors on a smooth curve.

Proposition 4.5 (An exact embedded chart). *Let $B = \mathbb{C}[x, y]/(x^2, y^2)$. On the unit-normalized hyperplane chart*

$$A' = \langle 1, y - ax, xy - bx \rangle \subset B \otimes \mathbb{C}[a, b],$$

the multiplication failure ideal, for every $m \geq 2$, is

$$I = (ab, b^2) = (b) \cap (a, b)^2.$$

Its associated primes are (b) and (a, b) . Its nilradical is a one-dimensional square-zero module supported at the embedded point. These equations give the pullback of the full failure scheme on the generating Grassmannian, not only its reduced support.

Proof. Use the quotient functional $\phi(1) = 0$, $\phi(x) = 1$, $\phi(y) = a$, $\phi(xy) = b$. Products involving 1 are pivots. The other quadratic products have residual values $-2ab, -b^2, 0$. This gives I ; Theorem 4.2 gives every higher degree. The displayed primary decomposition follows by intersecting the monomial ideals. The nilradical is generated by b , whose annihilator modulo I is (a, b) . The normalization by a unit is the same smooth-cover calculation used in that theorem. In particular adding the freely chosen unit does not remove the embedded associated prime. $\square$

Thus the quotient description distinguishes smooth moving contact schemes from finite-algebra families with embedded structure. The absence of embedded points in Theorem 4.3 is a consequence of the curvilinear contact algebra and its explicit Hilbert schemes, not a general assertion about finite algebras.

5. STABLE IMAGES AND HIGHER-DEFECT CONDUCTOR STRATA

The rank-two quotient in Theorem 4.2 is specific to a hyperplane. In larger codimension the appropriate object is a quotient algebra together with a subalgebra acting faithfully on the quotient module. The rank of this action is an additional stratification invariant. We keep its scheme structure throughout.

For a map u of vector bundles write $I_j(u)$ for its ideal of j -minors. Its *exact-rank s locus* means $V(I_{s+1}(u)) \setminus V(I_s(u))$, with this locally closed scheme structure. On it the image and cokernel are vector bundles and commute with arbitrary base change: after inverting a pivot, elementary operations reduce the matrix to $\text{diag}(I_s, 0)$. This is stronger than requiring constant rank on geometric points.

Theorem 5.1 (Stable image and conductor stratification). *Let B be a commutative finite locally free algebra of rank $d \geq 2$ over a locally Noetherian $\mathbb{C}$ -scheme S , and let $\mathcal{A}$ be the universal bundle on $X = \text{Gr}_S(k, B)^\circ$, with $1 \leq k \leq d$. Put $\mu_m : \text{Sym}^m \mathcal{A} \rightarrow B_X$. All Fitting ideals of $\text{coker } \mu_m$ are independent of $m \geq d - 1$. On the exact-rank s locus X_s of μ_{d-1} there is a canonical unital subalgebra bundle $E \subset B_{X_s}$ of rank s . There is an invertible E -module $L \subset B_{X_s}$, also a subbundle over X_s , such that*

$$E = \mathcal{O}[a^{-1}\mathcal{A}], \quad L = E\mathcal{A} = aE, \quad \text{im } \mu_m = L^m \quad (m \geq d - 1) \quad (5.1)$$

on any local unit cover $a \in \mathcal{A} \cap B^\times$.

Suppose $r = d - s > 0$ and put $M = B/E$. The action

$$\theta : E \longrightarrow \mathcal{E}nd_{\mathcal{O}}(M)$$

defines exact-rank strata $X_{s,t}$, with $1 \leq t \leq r^2$. On such a stratum the conductor and its quotient flag are

$$J = \ker \theta = \{b \in B : bB \subset E\}, \quad C = E/J \subset Q = B/J, \quad \text{rank } C = t, \quad \text{rank } Q = r + t. \quad (5.2)$$

Here J is an ideal subbundle of B , and $C \rightarrow \mathcal{E}nd(Q/C)$ is a subbundle injection. Each of these constructions commutes with arbitrary base change within the specified stratum.

More generally, the functor of unital codimension- r subalgebra bundles with action of exact rank t is equivalent to the functor of flags $B \twoheadrightarrow Q \supset C$ of these ranks, where C is unital, Q/C is locally free of rank r , and the action of C on Q/C is a subbundle injection. The inverse sends the flag to the preimage of C .

Proof. Choose a unit a smoothly locally, as in Lemma 3.1, and set $W = a^{-1}\mathcal{A}$. Then $1 \in W$. Lemma 3.2 says that $W^{d-1} = \mathcal{O}[W]$ and that $W^m = W^{d-1}$ for $m \geq d-1$, over the coefficient ring itself. Multiplication by a^m identifies the original image with $a^m\mathcal{O}[W]$. The resulting cokernels are locally isomorphic in these degrees, proving equality of every Fitting ideal before any rank restriction.

On X_s , the pivot description makes $E = \mathcal{O}[W]$ a subbundle with locally free quotient. If b is a second unit in $\mathcal{A}$, then $b/a \in E$ is a unit of B . Cayley–Hamilton for multiplication by b/a , whose constant coefficient is a unit, puts its inverse in E . Thus $\mathcal{O}[b^{-1}\mathcal{A}] \subset E$; reversing a, b gives equality. This proves descent of E . The image $E\mathcal{A}$ is locally aE and therefore descends to the stated invertible E -module and subbundle L . It also proves (5.1) universally.

Because E is a subalgebra, it acts on $M = B/E$. Its action kernel consists of the $e \in E$ with $eB \subset E$. An element b of the displayed conductor already lies in E , by multiplication by 1; hence the two kernels agree. If $jB \subset E$, then for $b, c \in B$ one has $(jb)c = j(bc) \in E$, so J is an ideal of B . On an exact-rank t stratum, pivot reduction for θ makes J locally free of rank $s - t$ and C a subbundle of $\mathcal{E}nd(M)$. The exact sequence $0 \rightarrow C \rightarrow B/J \rightarrow M \rightarrow 0$ proves that Q is locally free of rank $r + t$. All the kernels and images just used are split as $\mathcal{O}$ -modules locally; this proves their base-change assertions. The unit acts as the identity on M , so $t \geq 1$; the other bound comes from $\text{rank } \mathcal{E}nd(M) = r^2$.

Conversely, given a flag as in the statement, its inverse image E is a subalgebra bundle and $B/E \simeq Q/C$. The kernel of its action is precisely the kernel of $B \rightarrow Q$: a larger kernel would give an element of C annihilating Q/C , contrary to the subbundle injection. Thus the conductor construction recovers the flag, including over nonreduced test schemes. These mutually inverse constructions respect all pullbacks within the exact-rank functors. $\square$

The word “stratum” cannot be removed from the base-change assertion for J . The multiplication cokernel is a universally defined coherent sheaf on X ; its conductor need not be a bundle on X . Example 5.4 below exhibits the distinction.

Proposition 5.2 (The extreme-corank failure scheme). *On the Grassmannian of rank- $(d - r)$ subbundles $W \subset B$ containing 1, with $1 \leq r < d$, the unital subalgebra scheme is*

$$V(\text{Fitt}_{r-1} \text{ coker}(\text{Sym}^m W \rightarrow B)), \quad m \geq 2.$$

The defining ideal is independent of m , including its nonreduced structure. Its conductor exact-rank t subschemes are the quotient-flag functors of Theorem 5.1.

Proof. Since $1 \in W$, the image W^m contains W . Split $B = W \oplus M$ locally and remove the identity pivot columns. The residual presentation has r rows, so its $(r - 1)$ st Fitting ideal is the ideal of all entries. In degree two these entries are precisely the coefficients of $W \cdot W$ in B/W . They vanish if and only if W is a subalgebra, on every test scheme. Modulo their ideal, every higher product remains in W . Conversely each quadratic product is an m -fold product by insertion of 1’s. The entry ideals are equal. The last assertion is the functorial equivalence just proved. $\square$

For $r = 1$ this is the usual zeroth-Fitting failure scheme. For $r > 1$ it is the *extreme-corank* locus, not the whole locus where multiplication is nonsurjective. This distinction prevents a false extension of the hyperplane theorem by changing only the quotient rank.

5.1. The two quotient mechanisms in codimension two.

Theorem 5.3 (Codimension-two quotient flags). *Let $E \subset B$ be a unital subalgebra bundle with $M = B/E$ of rank two over a locally Noetherian $\mathbb{C}$ -scheme. For its action θ one has $I_3(\theta) = 0$ scheme-theoretically. The exact-rank-one closed locus and the exact-rank-two open locus have the following functorial descriptions.*

On the first, E is the inverse image of the scalar subalgebra in a locally free rank-three quotient algebra of B . On the second, E is the inverse image of C in a quotient $B \rightarrow Q$, where C is a rank-two algebra over the base and Q is a locally free rank-two algebra over C . In particular Q has rank four over the base. In both cases the kernel of $B \rightarrow Q$ is the conductor, and these descriptions commute with arbitrary base change on their respective strata.

Proof. Locally trivialize M . Since 2 is invertible, write every matrix as a scalar plus a trace-zero matrix. The trace-zero parts of the commuting matrices $\theta(E)$ still commute. The bracket map

$$\bigwedge^2 \mathfrak{sl}_2 \longrightarrow \mathfrak{sl}_2$$

is an isomorphism over $\mathbb{C}$: for the usual basis H, U, V its values are $2U, -2V, H$. Thus the wedge of any two trace-zero columns is zero over the whole coefficient ring, not just over its residue fields. Adjoining the scalar identity column proves $I_3(\theta) = 0$. The identity also gives $I_1(\theta) = \mathcal{O}$.

On the exact-rank-one locus the image is the scalar algebra. Theorem 5.1 gives $C = \mathcal{O}$ and $\text{rank } Q = 3$. Conversely the scalar line in any rank-three algebra has faithful action on its rank-two quotient module: the action is ordinary scalar multiplication. Its inverse image has exactly this conductor.

On the exact-rank-two locus, $C \subset \mathcal{E}nd(M)$ is a rank-two subalgebra bundle. Locally write $C = \langle 1, T \rangle$. At every geometric point T is nonscalar. A nonscalar two by two matrix has a cyclic vector: its minimal polynomial has degree two. After a local lift of such a vector, the map $C \rightarrow M, c \mapsto cv$, has invertible determinant. Thus M is locally free of rank one as a C -module. The exact sequence of C -modules $0 \rightarrow C \rightarrow Q \rightarrow M \rightarrow 0$ consequently splits locally, so Q is locally free of rank two over C .

Conversely, if Q is a locally free rank-two C -algebra, its unit is unimodular as a C -section. Hence Q/C is locally free of rank one over C . Its regular C -action is faithful and is a subbundle injection into its endomorphism bundle over the original base (check on geometric fibres and use a pivot). The quotient-flag equivalence therefore gives exactly the second stratum. All arguments are functorial and use split modules on the indicated strata. $\square$

Remark 5.4. Consider $R = \mathbb{C}[s]$, $B = R[z]/(z^5)$, and the direct summand

$$E = \langle 1, sz^2 + z^3, z^4 \rangle_R.$$

It is a subalgebra since $(sz^2 + z^3)^2 = s^2z^4$. Modulo E take the basis $\bar{z}, \bar{z}^2$; then $\bar{z}^3 = -s\bar{z}^2$. The action of $sz^2 + z^3$ sends $\bar{z}$ to $-s^2\bar{z}^2$ and $\bar{z}^2$ to zero, while z^4 acts as zero. Thus the exact-rank-one subscheme is $s^2 = 0$, not merely $s = 0$. Over R the conductor is Rz^4 ; at the reduced fibre $s = 0$ it is $\langle z^3, z^4 \rangle$. Quotient rank drops from four to three. This simultaneously records the nonreduced rank stratum and the failure of conductor formation to commute with unrestricted specialization. The fibre subalgebras belong to the classical codimension-two polynomial-algebra classification of [34, Theorem 39]; the calculation here retains the relative action and its determinantal thickening.

6. THREE GENERATORS ON A REDUCED CONTACT DIVISOR

This theorem realizes Haiman's diagonal ideal as the stable Fitting ideal of an actual multiplication map. The all-powers identity is Haiman's input, not a theorem deduced here from multiplication. The resulting global-section failure schemes and their coherent cokernels are obtained using the conductor comparison.

Theorem 6.1 (The complete stable three-plane ideal). *Let $D = p_1 + \cdots + p_d$ be reduced, $d \geq 4$, and let A vary on the entire generating open $\mathrm{Gr}(3, H^0(\mathcal{O}_D))^\circ$. On the smooth unit-frame cover write $A = \langle a, ax, ay \rangle$; thus the matrix with rows $(1, x_i, y_i)$ has rank three. For $m \geq d - 1$, the zeroth Fitting ideal J_m is independent of m , and*

$$J_m = \bigcap_{i < j} P_{ij}, \quad P_{ij} = (x_i - x_j, y_i - y_j). \quad (6.1)$$

For every $q \geq 1$ one has the stronger identity

$$J_m^q = \bigcap_{i < j} P_{ij}^q. \quad (6.2)$$

These are complete primary decompositions, including all component intersections. Thus J_m is radical, all its components are smooth of codimension two on this cover and on the fixed- D Grassmannian, and neither it nor any of its powers has embedded associated points. The corank is d minus the number of distinct pairs (x_i, y_i) . For global binary subseries the same statements hold for $n \geq 2d - 3$.

Proof. After dividing the target by a^m , the multiplication matrix has as columns all monomials in x, y of total degree at most m , evaluated at the d labelled points. Lemma 3.2 identifies, over the entire coefficient ring, the ideal of its d -minors for $m \geq d - 1$ with the ideal generated by determinants of arbitrary monomial columns. Those determinants span the alternating polynomials in the d pairs of variables: antisymmetrize a monomial and use linearity. Haiman's theorem [30, Corollary 3.8.3] states that the ideal generated by these alternants has every power equal to the intersection of the corresponding powers of the pair-diagonal ideals. It gives (6.1) and (6.2) before any localization; hence they hold on the rank-three open. In that corollary Haiman's n is our number d of labelled points, and Haiman's exponent d is our power q . These two indices are independent; no restriction on q is imposed. This use of the diagonal-ideal theorem is essential.

Each P_{ij} is a smooth prime ideal of codimension two; its powers are primary (use two normal coordinates in a regular local ring). The intersections therefore have exactly the indicated associated primes. Smooth faithful flat descent gives the fixed- D assertions. Polynomial functions separate any finite set of distinct affine points, so their image dimension is the number of distinct pairs. The truncation lemma gives the stated corank in this finite degree. Finally apply Theorem 2.2 with $k = 3$. $\square$

The stable moving scheme over the reduced-divisor stratum is integral, though it need not be smooth or unbranched. Its normalization marks one equal pair and is smooth and finite. Indeed the ordered configuration cover is etale and the permutation group is transitive on pairs. Upstairs the reduced scheme is the union in (6.1); its normalization is the disjoint union of the smooth pair diagonals. This construction descends. A point is smooth precisely when only one pair coincides; two or more failed pairs give distinct local branches after etale base change. No assertion about nonreduced divisors is inferred from this etale calculation; those have their own quotient-algebra structure in Section 4 when $k = d - 1$.

7. NONREDUCED MULTIGENERATOR SCHEMES AND FAT-POINT TRANSPORT

We next determine a multiplication failure scheme with arbitrarily large embedding dimension and nilpotence order. It is not a diagonal arrangement and its proof does not use Haiman's theorem. Fix integers $e \geq 2$ and $h \geq 3$, and put

$$B_{e,h} = \mathbb{C}[z_1, \dots, z_e]/(z_1, \dots, z_e)^h, \quad d = \binom{e+h-1}{e}, \quad N = \binom{e+h-1}{e+1}. \quad (7.1)$$

Write $\mathfrak{m}$ for its augmentation ideal. On $X = \mathrm{Gr}(e+1, B_{e,h})^\circ$ the augmentation of the universal bundle $\mathcal{A}$ is surjective. Its kernel K has rank e . Let $\Delta \subset X$ be the determinant divisor of the canonical map

$$K \longrightarrow (\mathfrak{m}/\mathfrak{m}^2) \otimes \mathcal{O}_X. \quad (7.2)$$

Thus this divisor is defined without a choice of unit or coordinates.

Theorem 7.1 (Exact primary law for truncated local algebras). *For every $m \geq h-1$, on the entire generating Grassmannian X ,*

$$\mathrm{Fitt}_0 \mathrm{coker}(\mathrm{Sym}^m \mathcal{A} \rightarrow B_{e,h}) = \mathcal{I}_\Delta^N. \quad (7.3)$$

The divisor Δ is nonempty, integral, and Cartier. The displayed ideal is primary, its exact nilradical index is N , and its generic transverse length is N . For every $q \geq 1$ its q th ordinary power has the complete primary description $\mathcal{I}_\Delta^{Nq}$; there are no embedded associated points. On a smooth unit-frame cover, Δ is a generic e by e determinant, and its singular locus is given by matrix rank at most $e-2$.

More precisely, choose a unit $a \in \mathcal{A}$ and write $\mathcal{A} = \langle a, ax_1, \dots, ax_e \rangle$, with $x_i \in \mathfrak{m}$. After multiplying the target by a^{-m} , the cokernel in (7.3) is the cokernel of the square substitution map

$$T_x : B_{e,h} \longrightarrow B_{e,h}, \quad z_i \longmapsto x_i.$$

If M is the matrix of linear coefficients of the x_i , then $\det T_x = (\det M)^N$. These presentation and ideal identities hold after arbitrary base change. The primary assertions refer to the universal parameter space X over $\mathbb{C}$, not to every specialization. For $m < h-1$ the zeroth Fitting ideal is zero, so the stated stable threshold is sharp for this family.

Proof. A unit exists locally because the augmentation of $\mathcal{A}$ is onto; subtracting constants from the remaining normalized basis vectors puts all x_i in $\mathfrak{m}$. The normalized multiplication matrix consists of the columns x^α with $|\alpha| \leq m$. Columns of total degree at least h vanish. When $m \geq h-1$ the remaining columns are exactly the d columns of T_x ; all other columns are zero. This proves the cokernel assertion, over any coefficient algebra.

The map T_x preserves the $\mathfrak{m}$ -adic filtration and its block on $\mathfrak{m}^j/\mathfrak{m}^{j+1}$ is $\mathrm{Sym}^j M$. Hence

$$\det T_x = \prod_{j=1}^{h-1} \det(\mathrm{Sym}^j M) = (\det M)^{\sum_{j=1}^{h-1} \binom{e+j-1}{e}} = (\det M)^N.$$

For completeness, the middle identity follows by taking M diagonal: among all degree- j monomials each diagonal entry occurs with total exponent $\binom{j}{e} \binom{e+j-1}{e-1} = \binom{e+j-1}{e}$. The identity holds for arbitrary matrices because both sides are polynomial functions and diagonalizable matrices with distinct eigenvalues form a dense open over $\mathbb{C}$. It is consequently a polynomial identity and remains valid on nonreduced bases. No condition on the coefficients of degree at least two was used.

The determinant of (7.2) differs from $\det M$ by a unit: the vectors ax_i form a basis of K , and their linear terms are the linear terms of x_i multiplied by the nonzero augmentation of a . Thus the ideal equality descends canonically. To check its geometric structure, send A to $K = A \cap \mathfrak{m}$. This identifies X with the total space of the quotient bundle $\mathfrak{m}/K$ over $\mathrm{Gr}(e, \mathfrak{m})$: a lift of the augmented line has the unique form $1 + v$ modulo K . On its frame cover the entries of the

linear-coefficient matrix M and the higher coefficients are independent variables, subject only to the open condition that the full $d - 1$ by e matrix have rank e . The generic determinant is irreducible. Its zero scheme remains nonempty on this open, as witnessed by

$$x_1 = z_1, \dots, x_{e-1} = z_{e-1}, \quad x_e = z_1^2.$$

It is therefore integral on the frame cover and descends to an integral Cartier divisor on the smooth scheme X .

Locally its equation is a prime element in a regular local ring. Every positive power is primary; alternatively, a hypersurface in a regular scheme has no embedded associated points and its support here has a single generic point. The radical of (δ^N) is (δ) , and δ^{N-1} is nonzero modulo δ^N at its generic point. This proves the exact nilradical index and length, and the same argument proves the assertion for all powers. The partial derivatives of a determinant are its cofactors, giving the stated singular locus. Finally $\text{rank Sym}^m \mathcal{A} = \binom{e+m}{e} < d$ if $m < h - 1$; there are no maximal minors. For $A = \langle 1, z_1, \dots, z_e \rangle$ the stable map is surjective, so this threshold cannot be lowered. $\square$

For $e = 2$ this is a genuine three-plane theorem for every $\mathbb{C}[x, y]/(x, y)^h$: the contact length is $\binom{h+1}{2}$ and the failure multiplicity is $\binom{h+1}{3}$. Already $h = 3$ gives a length-six, embedding-dimension-two algebra whose failure divisor has multiplicity four. These are schemes on a full generating Grassmannian, not selected transverse slices. They are distinct from the curvilinear hyperplane schemes and the reduced pair-diagonal ideal.

7.1. A global conductor theorem in arbitrary projective dimension. The preceding algebra is not the algebra of a divisor on a curve. To apply its scheme calculation to global sections, we require a separate conductor theorem. Its range depends on the order h of the fat point, not on its length d .

Theorem 7.2 (Fat-point conductor transport). *Let $e \geq 1$, $h \geq 2$, let $p \in \mathbb{P}^e$, and let Z be the fat point with ideal $\mathcal{I}_p^h$. Suppose $n \geq 2h - 1$. For any generating k -plane $A \subset H^0(Z, \mathcal{O}_Z(n))$, let*

$$W = H^0(\mathbb{P}^e, \mathcal{I}_p^h(n)), \quad U = \text{res}_Z^{-1}(A).$$

For every $m \geq 2$ one has

$$WU^{m-1} = H^0(\mathbb{P}^e, \mathcal{I}_p^h(mn)). \quad (7.4)$$

Restriction induces an isomorphism of coherent cokernels on the entire relative generating Grassmannian, also with p varying,

$$\text{coker}(\text{Sym}^m \mathcal{U} \rightarrow H^0(\mathcal{O}(mn)) \otimes \mathcal{O}) \simeq \text{coker}(\text{Sym}^m \mathcal{A} \rightarrow \pi_* \mathcal{O}_Z(mn)). \quad (7.5)$$

It and all its Fitting ideal equalities commute with arbitrary base change. The bound is sharp uniformly over all generating dimensions k and all $m \geq 2$; no rank-specific sharpness is asserted.

Proof. First fix $p = [1 : 0 : \dots : 0]$ and work on its affine chart with coordinates $z_1, \dots, z_e$. A section of $\mathcal{O}(n)$ is a polynomial of degree at most n , and W has basis the monomials of total degree between h and n . Choose $u \in U$ with constant coefficient u_0 a unit. This is possible over a local parameter base because the augmentation of A and restriction onto A are surjective.

For every integer t with $h \leq t \leq mn$, one can choose $1 \leq j \leq m$ with $jh \leq t \leq jn$. Indeed the integral intervals $[jh, jn]$ have no gaps: $jn + 1 \geq (j + 1)h$ follows from $n \geq 2h - 1$. Every monomial z^α of total degree t can be factored into j monomials of degrees between h and n . Their product, multiplied by u^{m-j} , belongs to WU^{m-1} and has the form

$$u_0^{m-j} z^\alpha + \text{monomials of total degree strictly greater than } t.$$

Its degree is at most mn . Choosing this column for every monomial of degree between h and mn gives a square triangular matrix with units on the diagonal. It is invertible over the coefficient ring.

These columns are thus a basis of the whole space on the right of (7.4). The reverse containment is immediate.

Restriction in degrees n and mn is surjective, since those degrees are at least $h - 1$. Its kernels and targets are vector bundles, with monomial bases on the current chart. In the locally split sequence $0 \rightarrow W \rightarrow U \rightarrow A \rightarrow 0$, the kernel on symmetric powers is the image of $W \otimes \text{Sym}^{m-1} U$. Equation (7.4) identifies its multiplication image with the kernel of target restriction. Taking the quotient of this restriction square proves (7.5). The triangular proof works over every coefficient ring once u_0 is a unit. For moving p , choose projective coordinates smoothly locally; restriction and the cokernel isomorphism are intrinsic, so they descend. All the split restriction sequences and surjective maps remain so after arbitrary pullback.

At the boundary $n = 2h - 2$, take $A = \langle 1 \rangle$. Then $U = \langle 1 \rangle + W$, and its quadratic monomials have degrees in $\{0\} \cup [h, 2h - 2] \cup [2h, 4h - 4]$. They omit z_1^{2h-1} , although it lies in the conductor kernel in degree $2n$. Its nonzero global cokernel class restricts to zero on Z . Thus the cokernel comparison already fails in degree two. $\square$

Corollary 7.3 (Global noncurvilinear primary schemes). *For $e \geq 2$, $h \geq 3$, and $n \geq 2h - 1$, consider all inverse-image series U of generating $(e+1)$ -planes on $Z = (\mathcal{I}_p^h)$. For every $m \geq h - 1$, their global multiplication failure scheme has the exact primary ideal $\mathcal{I}_\Delta^{\binom{e+h-1}{e+1}}$. Every power is primary and there are no embedded associated points on this fixed- p generating parameter space. In degrees $2 \leq m < h - 1$ the failure ideal is zero. The same primary assertions hold on the relative parameter space with $p \in \mathbb{P}^e$ varying; the coherent cokernel comparison itself holds after every base change.*

Proof. Apply Theorems 7.1 and 7.2. For moving p , coordinate charts give the relative determinant Schubert divisor with free higher-jet parameters. It is reduced and Cartier, and its fibres over the integral base $\mathbb{P}^e$ are nonempty integral divisors of constant dimension. These locally trivial descriptions make the total divisor integral as well. The total parameter space is smooth, so the same local primary-power argument applies. No primary-decomposition assertion is inferred for arbitrary nonreduced specializations of this parameter space. $\square$

8. PRIMARY DESCENT OVER MULTIPLICITY STRATA

The following descent statement concerns ideals, not just the permutation action on their generic points.

Lemma 8.1 (Etale descent of weighted prime divisors). *Let X be a regular integral Noetherian $\mathbb{C}$ -scheme and $p : \tilde{X} \rightarrow X$ a finite etale Galois cover, with group G . Let $\{D_\alpha\}$ be a finite G -stable collection of distinct integral effective Cartier divisors on $\tilde{X}$. Work componentwise if the cover is disconnected. Suppose the positive integers b_α are constant on G -orbits. For an orbit O , there is a unique integral effective Cartier divisor D_O on X whose ideal satisfies*

$$p^* \mathcal{I}_{D_O} = \prod_{\alpha \in O} \mathcal{I}_{D_\alpha} = \bigcap_{\alpha \in O} \mathcal{I}_{D_\alpha}.$$

If a descended ideal $\mathcal{J}$ pulls back to $\prod_\alpha \mathcal{I}_{D_\alpha}^{b_\alpha}$, then its complete primary decomposition is

$$\mathcal{J} = \prod_O \mathcal{I}_{D_O}^{b_O} = \bigcap_O \mathcal{I}_{D_O}^{b_O}. \quad (8.1)$$

There are no embedded associated points, and the formula holds at intersections as well as generically. The descended prime divisor need not be smooth or geometrically unbranched.

Proof. In each regular local ring upstairs, equations for distinct prime Cartier divisors are distinct prime factors. Unique factorization gives product equal to intersection, with or without positive

powers. The reduced orbit union has an invariant invertible ideal subsheaf of $\mathcal{O}_{\tilde{X}}$ with its canonical descent data. Faithfully flat descent gives an invertible ideal subsheaf of $\mathcal{O}_X$. It defines an effective Cartier divisor. Its support is the image of any one member of the orbit, hence is irreducible, since the map is finite. Reducedness descends faithfully flat, so this divisor is integral.

The same factorization upstairs, and flatness for finite intersections, identify the pullbacks of all three ideals in (8.1). Faithfulness makes them equal downstairs. Locally downstairs each $\mathcal{I}_{D_O}$ is generated by a prime element of a regular local ring; its b_O th power is primary. Their intersection is irredundant after discarding components absent from the chart, so it is the complete primary decomposition and has no embedded primes. No normal-crossing hypothesis was used. An orbit can have several members through a point upstairs, giving several etale branches of the integral descended divisor. This explains the final distinction. $\square$

Apply the lemma to the relative generating pencil Grassmannian over a multiplicity stratum. Ordering the distinct support points gives a finite etale cover with permutations among equal multiplicities. On it the divisors T_i, E_{ij} are the relative rank-one Schubert divisors. Their ideals are prime Cartier, and their weights are $\binom{d_i}{2}, d_i d_j$. Formula (8.1) supplies the whole descended primary decomposition. In particular it remains valid at a braid intersection where several values agree. The fixed labelled divisors are smooth; the descended orbit divisors are asserted integral, not automatically smooth or unbranched.

This argument applies to the earlier pencil calculation. For the contact-hyperplane family, Theorem 4.2 gives a scheme isomorphism with the quotient incidence and does not require a conjectural primary descent rule.

9. CONTACT PENCILS: EQUATIONS, PRIMARY COMPONENTS, AND INFINITESIMAL LAYERS

Fix $d \geq 4$ and $n \geq 2d - 2$. In this section the divisor D is fixed, although it need not be reduced. Set $G_D^\circ = \text{Gr}(2, E_{n,D})^\circ$. It is a smooth irreducible variety of dimension $2d - 4$. The conductor construction embeds it as the open generating locus in the Grassmannian of subseries containing W_D . Let $\mathcal{J}_m$ be the zeroth Fitting ideal of $\text{coker}(\text{Sym}^m \mathcal{U} \rightarrow V_{mn})$ on this variety.

Trivialize $\mathcal{O}_D(n)$, and put $R = H^0(\mathcal{O}_D)$. There is a smooth surjective frame map

$$\mathcal{F}_D = \{(a, v) : a \in R^\times, v \in R \setminus \mathbb{C}\} \longrightarrow G_D^\circ, \quad (a, v) \longmapsto \langle a, av \rangle. \quad (9.1)$$

It is the open part of the frame bundle where the first basis vector is a unit. Its relative dimension is four. All calculations on $\mathcal{F}_D$ below are therefore exact faithfully flat local calculations for G_D° , not transverse slices inferred from pointwise ranks.

Lemma 9.1 (The power-basis presentation). *On $\mathcal{F}_D$ the multiplication cokernel is presented by*

$$K_m(v) = [1, v, v^2, \dots, v^m] : \mathcal{O}^{m+1} \longrightarrow R \otimes \mathcal{O}. \quad (9.2)$$

For $m \geq d - 1$, every later column is an $\mathcal{O}$ -linear combination of the first d columns, and

$$\mathcal{J}_m \mathcal{O}_{\mathcal{F}_D} = (\Delta_D(v)), \quad \Delta_D(v) = \det[1, v, \dots, v^{d-1}]. \quad (9.3)$$

For $2 \leq m < d - 1$, $\mathcal{J}_m = 0$.

Proof. By Theorem 2.2, it suffices to multiply sections in the pencil $\langle a, av \rangle$ on D . Its image is generated by $a^m v^j$ for $0 \leq j \leq m$. Multiplication by a^{-m} is an invertible target operation and gives (9.2). Multiplication by v is an endomorphism of the rank- d free module $R \otimes \mathcal{O}$. Its monic characteristic polynomial annihilates it over the coefficient ring, by the adjugate proof of Cayley–Hamilton. Thus v^d and every higher power lie in the span of the first d powers, with regular coefficients. Column operations reduce the presentation to $[K_{d-1}(v), 0]$. This proves (9.3), as an identity of ideals and not merely of radicals. If $m + 1 < d$, all d -minors are zero. Faithful flatness descends this assertion. $\square$

Write $D = \sum_{i=1}^s d_i p_i$, with distinct p_i . Choose local parameters z_i and the Chinese-remainder identification

$$R = \prod_{i=1}^s \mathbb{C}[z_i]/(z_i^{d_i}), \quad v_i = \lambda_i + \sum_{r=1}^{d_i-1} c_{ir} z_i^r.$$

All λ_i, c_{ir} are independent affine coordinates before removing the scalar locus. For $d_i \geq 2$ define T_i by failure of A to separate $2p_i$; for $i < j$ define E_{ij} by failure to separate $p_i + p_j$. On the frame space they have equations $c_{i1} = 0$ and $\lambda_j - \lambda_i = 0$, respectively.

Theorem 9.2 (Full fixed-contact classification). *For every $m \geq d - 1$, in the ordered jet basis $1, z_i, \dots, z_i^{d_i-1}$,*

$$\Delta_D(v) = \prod_{i:d_i \geq 2} c_{i1}^{\binom{d_i}{2}} \prod_{i < j} (\lambda_j - \lambda_i)^{d_i d_j}. \quad (9.4)$$

The complete primary decomposition on G_D° is

$$\mathcal{J}_m = \bigcap_{i:d_i \geq 2} \mathcal{I}_{T_i}^{\binom{d_i}{2}} \cap \bigcap_{i < j} \mathcal{I}_{E_{ij}}^{d_i d_j}. \quad (9.5)$$

All displayed divisors are nonempty, smooth, irreducible, and distinct. They are precisely the associated subvarieties; no embedded associated points occur. The failure divisor has dimension $2d - 5$ and divisor class $\binom{d}{2}$ times the restricted Pluecker class.

Every multiplication corank, for every $m \geq 2$, is determined as follows. Put $e_i = \text{ord}_{z_i}(v_i - \lambda_i)$, with $e_i = d_i$ for a constant v_i , and put $\ell_i = \lceil d_i/e_i \rceil$. For each distinct value λ among the λ_i , set $L_\lambda = \max_{i:\lambda_i=\lambda} \ell_i$, and let $s(v) = \sum_\lambda L_\lambda$. Then

$$\text{corank}(\text{Sym}^m U_A \longrightarrow V_{mn}) = d - \min\{m + 1, s(v)\}. \quad (9.6)$$

On the frame space every higher Fitting ideal is, exactly, the corresponding minor ideal of (9.2).

Proof. For a polynomial P of degree at most $d - 1$, Taylor expansion gives

$$P(v_i(z_i)) = \sum_{r=0}^{d_i-1} \frac{P^{(r)}(\lambda_i)}{r!} (v_i(z_i) - \lambda_i)^r \pmod{z_i^{d_i}}.$$

The linear change from the divided derivatives of P at λ_i to the coefficients of $P(v_i(z_i))$ is triangular, with diagonal $1, c_{i1}, \dots, c_{i1}^{d_i-1}$. Its determinant is $c_{i1}^{\binom{d_i}{2}}$, including when this coefficient is zero. The determinant of the divided-derivative evaluation map

$$P \longmapsto \left(\frac{P^{(r)}(\lambda_i)}{r!} \right)_{i, 0 \leq r < d_i}$$

in the basis $1, T, \dots, T^{d-1}$ is the confluent Vandermonde $\prod_{i < j} (\lambda_j - \lambda_i)^{d_i d_j}$. Here is an explicit normalization of that identity. Start with d distinct variables, cluster d_i of them near λ_i , and in each cluster replace successive evaluation rows by divided-difference rows. The ordinary Vandermonde determinant is divided by the product of the differences *within* each cluster. In the cluster limit the rows are the divided derivatives above, while the remaining cross-cluster factors tend to the displayed product. This proves the identity on a dense open set and hence as a polynomial identity, including all collisions. Combining the two determinants proves (9.4) without inverting any of its factors.

For any length-two $Z \leq D$, restriction $E_{n,D} \rightarrow E_{n,Z}$ is surjective. The locus in $\text{Gr}(2, E_{n,D})$ where its restriction to A has rank at most one is the corresponding irreducible Schubert divisor. Its rank-zero locus is excluded by generation on D . Its rank-one locus is smooth: choose a basis of A with one nonzero image, and vary the second basis vector in a complementary target direction;

the determinant then has a nonzero linear derivative. For irreducibility one may also mark the image line in $E_{n,Z}$; the resulting incidence is a Grassmannian bundle over $\mathbb{P}(E_{n,Z})$, and projects birationally onto the rank-one locus. The generating open is nonempty for each divisor when $d \geq 4$. On frames this follows directly by choosing $a = 1$ and a nonscalar v with the single prescribed linear equality. The same coordinates show that the divisors are distinct.

On each affine frame chart the coordinate ring is a localization of a polynomial ring. The distinct linear factors in (9.4) generate distinct prime ideals there. Unique factorization gives $(\prod f_\alpha^{b_\alpha}) = \bigcap (f_\alpha^{b_\alpha})$; each ideal $(f_\alpha^{b_\alpha})$ is primary. These identities descend by faithful flatness to (9.5). Equivalently, at each point of the smooth base the local ring is regular, each displayed divisor is smooth, and its powers are primary. This also proves absence of embedded primes. The dimension follows since this is a nonzero Cartier divisor. Each component is a Pluecker hyperplane section, and

$$\sum_i \binom{d_i}{2} + \sum_{i < j} d_i d_j = \binom{d}{2},$$

which proves the class formula.

Finally, in $\mathbb{C}[z_i]/(z_i^{d_i})$ the element $v_i - \lambda_i$ has nilpotency index ℓ_i . Its minimal polynomial is $(T - \lambda_i)^{\ell_i}$. In the product algebra the minimal polynomial is therefore

$$P_v(T) = \prod_\lambda (T - \lambda)^{L_\lambda}. \quad (9.7)$$

No polynomial of smaller degree can annihilate v , and division by this monic polynomial shows that $1, v, \dots, v^{s(v)-1}$ form a basis of $\mathbb{C}[v]$. Consequently the rank of $K_m(v)$ is $\min\{m+1, s(v)\}$. The cokernel isomorphism of Theorem 2.2 gives (9.6) and the assertion about Fitting ideals. $\square$

Theorem 9.3 (Exact infinitesimal layers and singularities). *Let $m \geq d-1$. At a point of a smooth frame chart discard the nonvanishing factors of (9.4). Write the remaining distinct linear factors as f_α , with multiplicities b_α , and put*

$$F = \prod_\alpha f_\alpha^{b_\alpha}, \quad P = \prod_\alpha f_\alpha.$$

In the regular local ring B of that chart the failure ring is $B/(F)$ and its nilradical is $\mathcal{N} = (P)/(F)$. Its exact nilpotency index is $\max_\alpha b_\alpha$, with index one for a reduced local ring. For every $q \geq 1$, there is an isomorphism of B -modules, with indicated generator P^q ,

$$\mathcal{N}^q / \mathcal{N}^{q+1} \simeq B / \left(P, \prod_\alpha f_\alpha^{(b_\alpha - q)_+} \right). \quad (9.8)$$

The completed local equation is the same weighted linear arrangement, with unused frame and higher-jet coordinates as smooth parameters. Its tangent cone is defined by the homogeneous product of the active linear factors with these same multiplicities. Its singular scheme on this chart has ideal $(F, \partial F)$, and its singular support is the union of the components with $b_\alpha \geq 2$ and the pairwise intersections of distinct components. These descriptions descend smoothly to G_D° .

Proof. The local ring in question is a localization of a polynomial ring in the unit coordinates of a and the independent jet coordinates of v . The factors f_α are prime, distinct, and vanish at the selected point. Unique factorization shows that $P^q \in (F)$ if and only if $q \geq b_\alpha$ for every α . This proves the nilpotency assertion, including at multiple intersections.

The homomorphism $B \rightarrow \mathcal{N}^q / \mathcal{N}^{q+1}$ sending 1 to the class of P^q is onto. Its kernel is

$$((P^{q+1}, F) : P^q) = (P) + (F : P^q) = \left(P, \prod_\alpha f_\alpha^{(b_\alpha - q)_+} \right).$$

For the middle equality, if $rP^q = sP^{q+1} + tF$, subtract sP from r ; cancellation gives membership in $(F : P^q)$. The last colon is computed by prime-factor exponents. This proves (9.8). Globally the chosen generator carries the corresponding line-bundle trivialization; the displayed isomorphism is a local one and makes no claim of a preferred global generator.

At the chosen point every active factor is linear in centered coordinates. Every discarded factor is a unit, so the completed ideal is exactly the displayed product, not a truncation of it. The initial ideal is generated by the same homogeneous product, which proves the tangent-cone statement. The hypersurface cotangent presentation gives $(F, \partial F)$. At a point on exactly one component of multiplicity one the differential is a unit times $df_\alpha \neq 0$. At every point on a multiple component or on two distinct components it is zero. This proves the singular support assertion. Smooth faithfully flat pullback preserves these local statements; in particular the four frame parameters do not alter the nilpotency index or the singular-support criterion. $\square$

For illustration, length four already gives subseries of codimension two, in every V_n with $n \geq 6$. For every $m \geq 3$ their fixed-contact primary equations are as follows. Put $\delta_{ij} = \lambda_j - \lambda_i$; only first-jet coefficients of multiple points are displayed.

Multiplicities of D	Exact equation on frames	Largest local nilpotency index
$(1, 1, 1, 1)$	$\prod_{i < j} \delta_{ij}$	1
$(2, 1, 1)$	$c_{11}\delta_{12}^2\delta_{13}^2\delta_{23}$	2
$(2, 2)$	$c_{11}c_{21}\delta_{12}^4$	4
$(3, 1)$	$c_{11}^3\delta_{12}^3$	3
(4)	c_{11}^6	6

The reduced arrangement need not have normal crossings: three coincident values give three braid hyperplanes in a two-dimensional normal space. The equation, rather than a normal-crossing label, records that singularity. Formula (9.6) also distinguishes points with the same support of failure but different higher-jet vanishing orders.

10. MOVING CONTACTS AND THE NORMALIZATION OF THE FAILURE DIVISOR

Let $X_d^\circ = X_{d,2}^\circ$ be the relative generating Grassmannian in Theorem 2.2. It is smooth and irreducible of dimension $3d - 4$. The fixed-divisor schemes of the preceding section are its scheme-theoretic fibres, not its irreducible components. For $m \geq d - 1$ denote its failure scheme by Y_m .

10.1. Components over every multiplicity stratum. For a partition $\mathbf{d} = (d_1, \dots, d_s)$ of d , let $B_{\mathbf{d}} \subset \text{Sym}^d \mathbb{P}^1$ be the locally closed stratum of divisors with exactly this multiplicity partition, and restrict X_d° to it. Let M be the set of distinct multiplicities, and let $r_e = \#\{i : d_i = e\}$ for $e \in M$.

Corollary 10.1 (Complete component descent on multiplicity strata). *For $m \geq d - 1$, the failure scheme over $B_{\mathbf{d}}$ has*

$$\#\{e \in M : e \geq 2\} + \binom{\#M}{2} + \#\{e \in M : r_e \geq 2\} \quad (10.1)$$

irreducible components, all of dimension $s + 2d - 5$, and no embedded associated points. A ramification component is indexed by each $e \geq 2$ in M and has primary multiplicity $\binom{e}{2}$. An identification component is indexed by each unordered pair of multiplicities e, f that occurs at distinct support points and has primary multiplicity ef . The ideal of the scheme is the intersection of these prime-divisor ideals raised to the stated powers. On the labelled etale cover its full local equations and nilradical layers are exactly those of Theorem 9.3.

Proof. The ordered configuration space of s distinct points of $\mathbb{P}^1$ is an integral finite etale cover of B_d , with deck group $\prod_{e \in M} \mathfrak{S}_{r_e}$. The relative first-jet and two-point restriction maps are surjective over this cover. Consequently each labelled T_i and E_{ij} is an irreducible relative rank-one Schubert divisor. The local factorization (9.4) holds with the support coordinates free, so there are no further components. The deck group permutes T_i exactly according to the value d_i , and permutes E_{ij} exactly according to the unordered pair $\{d_i, d_j\}$. The images of the orbit unions are precisely the irreducible components downstairs: each is the image of an integral labelled component, and two such images agree exactly for components in the same orbit. This gives (10.1).

Lemma 8.1 applies to this regular ambient space and its invariant weighted divisors, including all branch intersections. It proves that each component downstairs is a prime Cartier divisor; irreducibility here does not assert smoothness or geometric unbranchedness. Pulling its reduced ideal back to the labelled cover gives the reduced product of its labelled branch equations. Their multiplicity is constant on an orbit, so the primary ideal identity in (9.5) descends faithfully flatly. This also proves absence of embedded points and gives the stated local layers. The base stratum has dimension s , giving the dimension formula. A descended component can have intersecting local branches; smoothness of each labelled divisor is not a claim that every descended component is smooth. $\square$

Theorem 10.2 (An integral failure divisor with smooth normalization). *For $d \geq 4$, $n \geq 2d - 2$, and $m \geq d - 1$, the schemes Y_m are all the same integral Cartier divisor Y of dimension $3d - 5$. In particular Y has no embedded associated points. Its normalization is*

$$\tilde{Y} = \{(D, A, Z) : (D, A) \in X_d^\circ, Z \leq D, \deg Z = 2, \text{rank}(A \rightarrow H^0(\mathcal{O}_Z(n))) = 1\}. \quad (10.2)$$

The forgetful map $\tilde{Y} \rightarrow Y$ is finite and birational, and $\tilde{Y}$ is smooth and irreducible.

More explicitly, on an affine support chart write $D = (g(t))$ with g monic of degree d , and on a frame chart put $v = b/a \in \mathbb{C}[t]/(g)$, represented by a polynomial of degree less than d . The failure equation is

$$\Delta(g, v) = \det[1, v, \dots, v^{d-1}]_{\mathbb{C}[t]/(g)} = 0. \quad (10.3)$$

The normalization on this chart is parametrized by

$$g = hq, \quad v = \lambda + hw, \quad (10.4)$$

where h, q are monic of degrees $2, d - 2$, respectively, $\lambda \in \mathbb{C}$, and $\deg w \leq d - 3$. The scalar-pencil locus is omitted; the unit a is a free smooth parameter.

Proof. Theorem 2.2 applies on the whole relative space. The proof of Lemma 9.1 applies over the polynomial coefficient ring of a varying monic g , not only on its fibres. Thus every Y_m has the identical ideal (10.3) on a smooth covering by frame charts. This determinant is not identically zero: $v = t$ makes its matrix the identity for every g .

The space of pairs $Z \leq D$ with lengths $2, d$ is $\text{Sym}^2 \mathbb{P}^1 \times \text{Sym}^{d-2} \mathbb{P}^1$; the map sends (Z, Q) to $D = Z + Q$. It is finite. Indeed it is proper, and a fixed effective divisor has only finitely many degree-two subdivisors, obtained by choosing multiplicities at its finitely many support points. Restriction $E_{n,D} \rightarrow E_{n,Z}$ is a surjection of vector bundles. The rank-at-most-one condition on a pencil is a relative Schubert divisor; its rank-zero part is excluded by the generating hypothesis. This makes (10.2) a closed subscheme of the finite base change to X_d° , so its projection is finite.

Smoothness and irreducibility can be checked explicitly. A divisor with finite support is contained in an affine coordinate chart. On that chart Z is defined by a unique monic quadratic h , and $Z \leq D$ means $g = hq$. Since a is a unit, rank one on Z is equivalent to $v \bmod h$ being a scalar λ . Division by the monic h gives uniquely $v = \lambda + hw$, with the stated degree bound. Thus the framed incidence is an open part of affine space with coordinates the coefficients of h, q, w , the scalar λ , and the unit a . This proves smoothness. Globally irreducibility also follows

from the relative rank-one Schubert description over the integral pair-divisor base; each fibre is irreducible, and its generating open is nonempty. These charts describe the same incidence, so no extra component appears at infinity.

An element v fails to generate the length- d algebra $R = H^0(\mathcal{O}_D)$ precisely when it either takes the same value at two distinct support points or has zero first derivative at a multiple support point. This follows at once from the minimal polynomial (9.7): its degree is d if and only if the support values are distinct and each local $\ell_i = d_i$. These two failure mechanisms are precisely the existence of a length-two $Z \leq D$ on which v is constant. Therefore the image of the finite incidence is the whole support of (10.3). That support is irreducible.

To determine its multiplicity, work over divisors with distinct support, label their roots after an étale base change, and use evaluation coordinates for v . The fixed-fibre formula becomes a product of the distinct differences $\lambda_j - \lambda_i$. At a point with exactly one equal pair the determinant has a nonzero linear derivative. Hence its multiplicity along its unique irreducible support is one. Since the ambient space is smooth, a Cartier hypersurface has no embedded associated points; a generically reduced such hypersurface is reduced. One may see the last assertions locally from unique factorization in the regular local ring, or from the Cohen–Macaulay and $R_0 + S_1$ criterion [24, Tag 031R]. Thus Y is integral.

Over the dense open subset with reduced D and precisely one equal pair there is exactly one possible Z . The finite projection is therefore birational in characteristic zero. Its source is smooth and hence normal, so it is the normalization. The affine parametrization is exactly (10.4), as already proved. The dimension of the divisor is $3d - 5$. $\square$

Proposition 10.3 (Exact singular-support test across collisions). *At a closed point $(D, A) \in Y$, call a length-two $Z \leq D$ scalar if A has rank one on Z . The point is smooth on Y if and only if there is precisely one scalar Z and the following linear system has no nonzero solution. In the associated factorization (10.4), the unknown is $r(t)$ of degree less than two, and the system is*

$$h \mid qr, \quad wr \bmod h \text{ is constant.} \quad (10.5)$$

In particular, this criterion is effective at nonreduced D and is not merely the distinct-support pair-count criterion. The full singular scheme on a frame chart has ideal

$$(\Delta, \partial\Delta/\partial g_0, \dots, \partial\Delta/\partial g_{d-1}, \partial\Delta/\partial v_0, \dots, \partial\Delta/\partial v_{d-1}). \quad (10.6)$$

Proof. If the normalization has more than one point above (D, A) , the local ring of Y cannot be regular. Suppose there is just one. We first justify the differential criterion for a finite normalization $\nu : N \rightarrow Y$ with N smooth. A regular local ring is normal, so a smooth point of Y has an isomorphic normalization near it. Conversely let $A_0 \subset B_0$ be the local finite extension at a point with a unique preimage and common residue field $\mathbb{C}$. Injectivity of $d\nu$ means

$$\mathfrak{m}_{B_0}/(\mathfrak{m}_{B_0}^2 + \mathfrak{m}_{A_0}B_0) = 0.$$

Nakayama's lemma first gives $\mathfrak{m}_{B_0} = \mathfrak{m}_{A_0}B_0$. It then gives, for the finite A_0 -module B_0 , that 1 generates B_0 , since the quotient by $\mathfrak{m}_{A_0}$ is $\mathbb{C}$. The injective map $A_0 \rightarrow B_0$ is consequently surjective. Thus $A_0 = B_0$ is regular. Local finiteness here follows by localizing the finite morphism; uniqueness of the preimage makes the resulting semilocal ring local.

In the affine normalization coordinates, a tangent vector with $\dot{g} = \dot{v} = 0$ satisfies

$$0 = q\dot{h} + h\dot{q}, \quad 0 = \dot{\lambda} + w\dot{h} + h\dot{w}.$$

Set $r = \dot{h}$, so $\deg r < 2$ because h is monic. The first equation is soluble exactly when $h \mid qr$, and then determines $\dot{q} = -qr/h$. The second is soluble exactly when the remainder of wr modulo h is constant; it determines $\dot{\lambda}$ as the negative of that constant and determines $\dot{w}$ by division. The resulting degrees satisfy the required bounds. If $r = 0$ all four variations are zero. Thus the differential has nonzero kernel exactly when (10.5) has a nonzero solution. The unit frame

parameter has zero variation in a kernel and creates no additional one. Smooth frame descent proves the criterion on Y . Finally (10.6) is the hypersurface Jacobian presentation in its smooth affine ambient space. $\square$

Remark 10.4 (A smooth total point with a triple primary fibre). Take $d = 4$, $g = t^3(t - 1)$, and $A = \langle 1, t^2 \rangle$ on $D = (g)$. The only scalar length-two subdivisor is $h = t^2$. In (10.4) one has $q = t(t - 1)$ and $w = 1$. For $r = r_0 + r_1 t$, divisibility $h \mid qr$ forces $r_0 = 0$, while constancy of $wr \bmod h$ forces $r_1 = 0$. Hence the total divisor is smooth here. On the fixed-divisor fibre, the two support values are distinct and (9.4) is a unit times c_{11}^3 . Its exact local nilpotency index is three. Thus even a smooth point of the integral total divisor can have a nonreduced primary fibre.

Proposition 10.5 (Discriminants and specialization). *Let $\chi_v(T)$ be the characteristic polynomial of multiplication by v in $\mathbb{C}[t]/(g)$. With the usual monic polynomial normalization of discriminants,*

$$\text{disc}(\chi_v) = \text{disc}(g) \Delta(g, v)^2. \quad (10.7)$$

This is a polynomial identity on the entire coefficient space. The equation $\Delta = 0$, not $\text{disc}(\chi_v) = 0$, defines the failure divisor, including when g has multiple roots. Its fibre over each fixed D is exactly the primary scheme in (9.5).

Proof. When g has distinct roots $t_1, \dots, t_d$, the evaluation matrix changes the basis $1, t, \dots, t^{d-1}$ to point values. Therefore

$$\Delta(g, v) = \frac{\prod_{i < j} (v(t_j) - v(t_i))}{\prod_{i < j} (t_j - t_i)}.$$

Squaring gives (10.7), because $\chi_v(T) = \prod_i (T - v(t_i))$ on this open set. Both sides are polynomial in the coefficients, so the identity holds everywhere. For $v = t$ the determinant is one even when g is nonreduced; this also shows why the factor $\text{disc}(g)$ must not be included in the failure equation. The scheme-theoretic fibre statement follows from Fitting base change in Theorem 2.2, or by specializing the polynomial determinant and applying (9.4). $\square$

Proof of Theorem 1.1. The fixed-contact assertions are Theorems 9.2 and 9.3. The moving-contact assertions are Theorem 10.2 and Proposition 10.3. All four concern the original multiplication map, by the coherent cokernel isomorphism of Theorem 2.2. $\square$

11. PENCIL COLLISIONS AND COMPARISON OF LOCAL TYPES

The normalization criterion above separates phenomena which a set-theoretic discriminant calculation conflates. In the affine normalization write $g = hq$ and $v = \lambda + hw$, with h monic of degree two. The kernel test is

$$h \mid qr, \quad wr \bmod h \text{ constant}, \quad \deg r < 2.$$

The following three cases give concrete local distinctions.

If g has distinct roots, the normalization is unramified; a unique equal value pair gives a smooth point, and two disjoint equal pairs give an ordinary double crossing. Three equal values give the three braid factors $\delta_{12}\delta_{13}(\delta_{13} - \delta_{12})$, not a normal-crossing divisor in a two-dimensional normal space. These are exact equations because root labelling is étale here.

For $g = t^3(t - 1)$ and $v = t^2$, the scalar subdivisor $h = t^2$ is unique. One has $q = t(t - 1)$ and $w = 1$. Divisibility forces the constant term of r to vanish and the second condition forces its linear term to vanish. The total pencil divisor is smooth, although its fixed- D fibre has primary equation a unit times c_{11}^3 .

For $g = t^4$ and $v = t^2$, again $h = t^2$ is the unique scalar subdivisor. Now $q = t^2$ and $w = 1$. The constant $r = 1$ satisfies the kernel test. Thus the total pencil divisor is singular and its

normalization is ramified at the unique preimage. This last example is asserted to have a unique normalization point, not an unproved analytic cusp normal form. In contrast, the moving contact-hyperplane scheme is smooth at all these divisor collisions by Theorem 4.4. Its fixed fourfold-contact transverse algebra is G_4 , with length six and nilpotency index five, rather than the pencil's sixth power of a prime divisor.

12. RELATION TO EARLIER CONSTRUCTIONS

The determinant $\det[1, v, \dots, v^{d-1}]$ is a classical local index form. Arpin, Bozlee, Herr and Smith construct schemes of monogenerators and polygenerators for finite locally free algebras [29]. Their Proposition 3.6 and Definition 3.12 identify the index-form open and its closed complement; Proposition 3.14 gives the polygenerator open by minors. Thus neither the generator functor nor its index determinant is claimed as new here. Our use concerns the coherent cokernel of a global section-multiplication map and, for generating hyperplanes, its exact closed incidence with a rank-two quotient. Lemma 4.1 and Theorem 4.2 prove that closed-scheme identification on nonreduced bases, rather than deriving it from agreement of generator opens.

The algebra G_r is the classical two-row Grassmannian cohomology ring; [31, Theorem 2.7] provides the rectangular Schur basis in a more general quotient. Its role here is an exact transverse algebra of multiplication failure, determined by a rank-two quotient of $\mathbb{C}[z]/(z^r)$. The explicit residual equations (4.7) verify this identification without a tangent-cone approximation. The diagonal-ideal and all-powers identities in the three-plane result are precisely [30, Corollary 3.8.3]; the argument here supplies the finite total-degree presentation and conductor transport, not a new proof of that theorem.

Normal generation in sufficiently positive degree on a curve is classical [14]. The curve conductor theorem uses it only for $L(-D)$, and then proves the precise quotient comparison by filling the double-vanishing space and its first normal layer. On $\mathbb{P}^1$ the evaluation-kernel argument gives the sharper rank-dependent bound and its boundary example.

For pencils, confluent Vandermonde factorization and the discriminant identity are classical algebra. The moving normalization is the explicit scalar-pair incidence of that index-form divisor. Its finite map and differential test are proved in Section 10; the normalization is not offered as a construction independent of index-form geometry. The contact-hyperplane incidence is different: it is the failure scheme itself, not just its normalization, and its total space is smooth even at collisions.

Subalgebra varieties, quotient flags, and the additional comparisons. Iovanov and Sistko [32, Theorem 0.1] classify maximal subalgebras of finite-dimensional algebras over fields under their Schur hypothesis. In the commutative algebraically closed case, the alternatives include identifying two residue factors and imposing a first-order condition at one factor. These closed-point mechanisms are classical and are not claimed here as new. Sistko [33, Definition 3.1, Theorems 1.1 and 4.1] constructs subalgebra varieties and computes the maximal-subalgebra vanishing ideal for basic algebras. The distinction needed here is between such a variety and the full relative quadratic-equation functor: for $\mathbb{C}[z]/(z^r)$ the latter already has the transverse algebra G_r , of length $\binom{r}{2}$. Agreement on the reduced variety does not identify this thickening.

The Hilbert functor used in Section 4 is the standard functor of locally free quotient algebras; its representability is not a contribution of this article. Likewise, nested effective divisors on a curve are classically expressed by a divisor and its residual divisor. What is established here is the identification of these functors with specified multiplication Fitting schemes, the canonical cokernel line (4.2), and the global-section transports. These are separate assertions from the classical classification of geometric hyperplanes.

The faithful-action viewpoint is also classical; compare [33, Proposition 3.4] for subalgebras of matrix algebras. Our conductor flags record the *specified inclusion* into B , the action on B/E , and its exact-rank scheme structure. A conductor rank bound alone is not the flag equivalence, nor does either imply unrestricted base change across an action-rank jump. The one-variable codimension-two classification is already given in [34, Theorem 39]. We use it to locate the fibres of Example 5.4, not as a new classification of polynomial subalgebras. The quotient-tower statement concerns arbitrary finite B and its relative exact-rank functors.

In [29, Example 4.16], the two-generator jet calculation has curvilinear target $k[\varepsilon]/\varepsilon^{m+1}$. Our target in (7.1) has embedding dimension e . The symmetric-power determinant identity in the proof is classical linear algebra. Its role is to determine the full primary power of the tangent determinant, uniformly in e, h , and then identify that power with global section multiplication on $\mathbb{P}^e$ using Theorem 7.2. Merely knowing the polygenerator open would not determine this exponent or the global degree range. These comparisons identify the inputs and additional claims; they are not a certificate of exhaustive priority against all literature.

Ballico's failure-locus work [4, 5] is a necessary nearest-source comparison. The accessible statements of [5, Theorem 0.2, Proposition 2.2, Remark 2.3] concern failure on finite linear sections of a fixed completely embedded variety. The complete theorem pages of [4] have not been obtained in this revision, so their exact relation to the present scheme statements remains unverified. We make no inference of non-anticipation from that access limitation. The results here are submitted with complete proofs and explicit classical inputs; priority of the resulting package is a separate question for source comparison and editorial assessment.

REFERENCES

- [1] R. Balan, P. Casazza, and D. Edidin, On signal reconstruction without phase, *Appl. Comput. Harmon. Anal.* **20** (2006), 345–356.
- [2] E. Ballico and Ph. Ellia, The maximal rank conjecture for nonspecial curves in $\mathbb{P}^3$, *Invent. Math.* **79** (1985), 541–555.
- [3] E. Ballico and Ph. Ellia, The maximal rank conjecture for non-special curves in $\mathbb{P}^n$, *Math. Z.* **196** (1987), 355–367.
- [4] E. Ballico, On the Failure Locus of Higher Order Properties of Embeddings in Projective Spaces, *Math. Nachr.* **163** (1993), 5–13, doi:10.1002/mana.19931630102.
- [5] E. Ballico, On the failure cycles for the quadratic normality of a projective variety, *Pacific J. Math.* **172** (1996), 307–313.
- [6] S. Basu, R. Pollack, and M.-F. Roy, *Algorithms in Real Algebraic Geometry*, second ed., Springer, 2006.
- [7] A. Conca, M. Mostafazadehfard, A. K. Singh, and M. Varbaro, Hankel determinantal rings have rational singularities, *Adv. Math.* **335** (2018), 111–129; arXiv:1709.05685, Sections 1–2.
- [8] J. A. Eagon and D. G. Northcott, Ideals defined by matrices and a certain complex associated with them, *Proc. Roy. Soc. London Ser. A* **269** (1962), 188–204.
- [9] D. Eisenbud, C. Huneke, and B. Ulrich, Heights of ideals of minors, *Amer. J. Math.* **126** (2004), 417–438; arXiv:math/0209184.
- [10] A. Frühbis-Krüger and M. Zach, Determinantal singularities, arXiv:2106.04855v1, 2021; Lemma 1.3, Remark 1.18, Section 1.3.2, and Proposition 2.12.
- [11] W. Fulton, *Intersection Theory*, second ed., Springer, 1998, Theorem 14.4.
- [12] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, Resultants, and Multidimensional Determinants*, Birkhäuser, 1994.
- [13] C. J. Geyer, On the asymptotics of constrained M -estimation, *Ann. Statist.* **22** (1994), 1993–2010.
- [14] M. Green and R. Lazarsfeld, On the projective normality of complete linear series on an algebraic curve, *Invent. Math.* **83** (1986), 73–90.
- [15] A. B. Hempel, P. J. Goulart, and J. Lygeros, Inverse parametric quadratic programming and an application to hybrid control, *IFAC Proceedings Volumes* **45**(17) (2012), 68–73.
- [16] G. Kempf and D. Laksov, The determinantal formula of Schubert calculus, *Acta Math.* **132** (1974), 153–162.
- [17] J. Kohonen, A meet-in-the-middle algorithm for finding extremal restricted additive 2-bases, *J. Integer Seq.* **17** (2014), Article 14.6.8.

- [18] J. Kohonen, An improved lower bound for finite additive 2-bases, *J. Number Theory* **174** (2017), 518–524.
- [19] A. R. Kustin, Canonical complexes associated to a matrix, arXiv:1509.04367, 2015, discussion following (1.0.2) and Section 2.9.
- [20] E. Larson, The maximal rank conjecture, arXiv:1711.04906, Theorem 1.2.
- [21] B. Mourrain and V. Y. Pan, Multivariate polynomials, duality, and structured matrices, *J. Complexity* **16** (2000), 110–180.
- [22] N. A. Nguyen, S. Olaru, P. Rodriguez-Ayerbe, M. Hovd, and I. Necoara, Constructive solution of inverse parametric linear/quadratic programming problems, *J. Optim. Theory Appl.* **172** (2017), 623–648.
- [23] C. Scheiderer, Extreme points of Gram spectrahedra of binary forms, *Discrete Comput. Geom.* **67** (2022), 1174–1190.
- [24] The Stacks Project Authors, *The Stacks Project*, Tag 031R, Lemma 10.157.3, the $R_0 + S_1$ reducedness criterion; consulted September 22, 2026.
- [25] The Stacks Project Authors, *The Stacks Project*, Section 15.8, Tags 07Z8 and 07ZA, presentation independence and base change of Fitting ideals; consulted September 22, 2026.
- [26] A. W. van der Vaart, *Asymptotic Statistics*, Cambridge University Press, 1998.
- [27] G. Yu, Upper bounds for finite additive 2-bases, *Proc. Amer. Math. Soc.* **137** (2009), 11–18.
- [28] The Stacks Project Authors, *The Stacks Project*, Section 10.63, Tag 00L9; Lemmas 10.63.3 and 10.63.5 and Proposition 10.63.6, associated primes; consulted September 22, 2026.
- [29] S. Arpin, S. Bozlee, L. Herr, and H. Smith, *The scheme of monogenic generators I: Representability*, arXiv:2108.07185v2 (2022), especially Proposition 3.6, Definition 3.12, and Proposition 3.14.
- [30] M. Haiman, *Hilbert schemes, polygraphs, and the Macdonald positivity conjecture*, *J. Amer. Math. Soc.* **14** (2001), 941–1006; arXiv:math/0010246v2, Corollary 3.8.3.
- [31] D. Grinberg, *A basis for a quotient of symmetric polynomials*, arXiv:1910.00207v2 (2021), Theorem 2.7 and the subsequent discussion.
- [32] M. C. Iovanov and A. Sistko, *Maximal subalgebras of finite-dimensional algebras*, arXiv:1705.00762v2 (2017), Theorem 0.1.
- [33] A. Sistko, *Automorphism groups of finite-dimensional algebras acting on subalgebra varieties*, arXiv:1809.09760v2 (2019), Definition 3.1, Proposition 3.4, and Theorems 1.1 and 4.1.
- [34] R. Grönkvist, E. Leffler, A. Torstensson, and V. Ufnarovski, *Describing subalgebras of $K[x]$ using derivatives*, arXiv:2107.11916v1 (2021), Theorem 39, pp. 94–95.