

1	Contest	1
2	Mathematics	1
3	Numerical	2
4	Number theory	2
5	Combinatorial	2
6	Graph	2
7	Geometry	5

Contest (1)

template.cpp	14 lines
<pre>#include <bits/stdc++.h> using namespace std; #define rep(i, a, b) for(int i = a; i < (b); ++i) #define all(x) begin(x), end(x) #define sz(x) (int)(x).size() typedef long long ll; typedef pair<int, int> pii; typedef vector<int> vi; int main() { cin.tie(0)->sync_with_stdio(0); cin.exceptions(cin.failbit); }</pre>	
.bashrc	3 lines
<pre>alias c='g++ -Wall -Wconversion -Wfatal-errors -g -std=c++17 \ -fsanitize=undefined,address' xmodmap -e 'clear lock' -e 'keycode 66=less greater' #caps =◇</pre>	
.vimrc	6 lines
<pre>set cin aw ai is ts=4 sw=4 tm=50 nu noeb bg=dark ru cul sy on im jk <esc> im kj <esc> no ; : " Select region and then type :Hash to hash your selection. " Useful for verifying that there aren't mistypes. ca Hash w !cpp -dD -P -fpreprocessed \ tr -d '[:space:]' \ \ md5sum \ cut -c-6</pre>	
hash.sh	3 lines
<pre># Hashes a file, ignoring all whitespace and comments. Use for # verifying that code was correctly typed. cpp -dD -P -fpreprocessed tr -d '[:space:]' md5sum cut -c-6</pre>	

Mathematics (2)

2.1 Trigonometry

$\sin(v + w) = \sin v \cos w + \cos v \sin w$
 $\cos(v + w) = \cos v \cos w - \sin v \sin w$

$$\tan(v + w) = \frac{\tan v + \tan w}{1 - \tan v \tan w}$$
$$\sin v + \sin w = 2 \sin \frac{v + w}{2} \cos \frac{v - w}{2}$$
$$\cos v + \cos w = 2 \cos \frac{v + w}{2} \cos \frac{v - w}{2}$$

$$(V + W) \tan(v - w)/2 = (V - W) \tan(v + w)/2$$

where V, W are lengths of sides opposite angles v, w .

$$a \cos x + b \sin x = r \cos(x - \phi)$$
$$a \sin x + b \cos x = r \sin(x + \phi)$$

where $r = \sqrt{a^2 + b^2}, \phi = \text{atan2}(b, a)$.

2.2 Geometry

2.2.1 Triangles

Side lengths: a, b, c
Semiperimeter: $p = \frac{a + b + c}{2}$
Area: $A = \sqrt{p(p - a)(p - b)(p - c)}$
Circumradius: $R = \frac{abc}{4A}$
Inradius: $r = \frac{A}{p}$
Length of median (divides triangle into two equal-area triangles):
 $m_a = \frac{1}{2}\sqrt{2b^2 + 2c^2 - a^2}$
Length of bisector (divides angles in two):

$$s_a = \sqrt{bc \left[1 - \left(\frac{a}{b + c} \right)^2 \right]}$$

Law of sines: $\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c} = \frac{1}{2R}$
Law of cosines: $a^2 = b^2 + c^2 - 2bc \cos \alpha$
Law of tangents: $\frac{a + b}{a - b} = \frac{\tan \frac{\alpha + \beta}{2}}{\tan \frac{\alpha - \beta}{2}}$


2.2.2 Quadrilaterals

With side lengths a, b, c, d , diagonals e, f , diagonals angle θ , area A and magic flux $F = b^2 + d^2 - a^2 - c^2$:

$$4A = 2ef \cdot \sin \theta = F \tan \theta = \sqrt{4e^2 f^2 - F^2}$$

For cyclic quadrilaterals the sum of opposite angles is 180° ,
 $ef = ac + bd$, and $A = \sqrt{(p - a)(p - b)(p - c)(p - d)}$.

2.2.3 Spherical coordinates



$$x = r \sin \theta \cos \phi \quad r = \sqrt{x^2 + y^2 + z^2}$$
$$y = r \sin \theta \sin \phi \quad \theta = \text{acos}(z / \sqrt{x^2 + y^2 + z^2})$$
$$z = r \cos \theta \quad \phi = \text{atan2}(y, x)$$

2.3 Derivatives/Integrals

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1 - x^2}} \quad \frac{d}{dx} \arccos x = -\frac{1}{\sqrt{1 - x^2}}$$
$$\frac{d}{dx} \tan x = 1 + \tan^2 x \quad \frac{d}{dx} \arctan x = \frac{1}{1 + x^2}$$
$$\int \tan ax = -\frac{\ln |\cos ax|}{a} \quad \int x \sin ax = \frac{\sin ax - ax \cos ax}{a^2}$$
$$\int e^{-x^2} = \frac{\sqrt{\pi}}{2} \text{erf}(x) \quad \int x e^{ax} dx = \frac{e^{ax}}{a^2} (ax - 1)$$

Integration by parts:

$$\int_a^b f(x)g(x)dx = [F(x)g(x)]_a^b - \int_a^b F(x)g'(x)dx$$

2.4 Sums

$$c^a + c^{a+1} + \dots + c^b = \frac{c^{b+1} - c^a}{c - 1}, c \neq 1$$

$$1 + 2 + 3 + \dots + n = \frac{n(n + 1)}{2}$$
$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(2n + 1)(n + 1)}{6}$$
$$1^3 + 2^3 + 3^3 + \dots + n^3 = \frac{n^2(n + 1)^2}{4}$$
$$1^4 + 2^4 + 3^4 + \dots + n^4 = \frac{n(n + 1)(2n + 1)(3n^2 + 3n - 1)}{30}$$

2.5 Series

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots, (-\infty < x < \infty)$$
$$\ln(1 + x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots, (-1 < x \leq 1)$$
$$\sqrt{1 + x} = 1 + \frac{x}{2} - \frac{x^2}{8} + \frac{2x^3}{32} - \frac{5x^4}{128} + \dots, (-1 \leq x \leq 1)$$