

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots, \quad (-\infty < x < \infty)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots, \quad (-\infty < x < \infty)$$

Numerical (3)

3.1 Polynomials and recurrences

Tridiagonal.h
Description: $x = \text{tridiagonal}(d, p, q, b)$ solves the equation system

$$\begin{pmatrix} b_0 \\ b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_{n-1} \end{pmatrix} = \begin{pmatrix} d_0 & p_0 & 0 & 0 & \cdots & 0 \\ q_0 & d_1 & p_1 & 0 & \cdots & 0 \\ 0 & q_1 & d_2 & p_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & q_{n-3} & d_{n-2} & p_{n-2} \\ 0 & 0 & \cdots & 0 & q_{n-2} & d_{n-1} \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_{n-1} \end{pmatrix}.$$

This is useful for solving problems on the type
$$a_i = b_i a_{i-1} + c_i a_{i+1} + d_i, \quad 1 \leq i \leq n,$$
where a_0, a_{n+1}, b_i, c_i and d_i are known. a can then be obtained from
$$\{a_i\} = \text{tridiagonal}(\{1, -1, -1, \dots, -1, 1\}, \{0, c_1, c_2, \dots, c_n\}, \{b_1, b_2, \dots, b_n, 0\}, \{a_0, d_1, d_2, \dots, d_n, a_{n+1}\}).$$

Fails if the solution is not unique.
If $|d_i| > |p_i| + |q_{i-1}|$ for all i , or $|d_i| > |p_{i-1}| + |q_i|$, or the matrix is positive definite, the algorithm is numerically stable and neither `tr` nor the check for `diag[i] == 0` is needed.
Time: $\mathcal{O}(N)$

```

typedef double T;
vector<T> tridiagonal(vector<T> diag, const vector<T>& super,
    const vector<T>& sub, vector<T> b) {
    int n = sz(b); vi tr(n);
    rep(i,0,n-1) {
        if (abs(diag[i]) < 1e-9 * abs(super[i])) { // diag[i] == 0
            b[i+1] -= b[i] * diag[i+1] / super[i];
            if (i+2 < n) b[i+2] -= b[i] * sub[i+1] / super[i];
            diag[i+1] = sub[i]; tr[++i] = 1;
        } else {
            diag[i+1] -= super[i]*sub[i]/diag[i];
            b[i+1] -= b[i]*sub[i]/diag[i];
        }
    }
    for (int i = n; i--;) {
        if (tr[i]) {
            swap(b[i], b[i-1]);
            diag[i-1] = diag[i];
            b[i] /= super[i-1];
        } else {
            b[i] /= diag[i];
            if (i) b[i-1] -= b[i]*super[i-1];
        }
    }
    return b;
}

```

Number theory (4)

4.1 Pythagorean Triples

The Pythagorean triples are uniquely generated by

$$a = k \cdot (m^2 - n^2), \quad b = k \cdot (2mn), \quad c = k \cdot (m^2 + n^2),$$

template .bashrc .vimrc hash Tridiagonal IntPerm

with $m > n > 0, k > 0, m \perp n$, and either m or n even.

Combinatorial (5)

5.1 Permutations

IntPerm.h
Description: Permutation -> integer conversion. (Not order preserving.)
Integer -> permutation can use a lookup table.
Time: $\mathcal{O}(n)$

044568, 6 lines

```

int permToInt(vi& v) {
    int use = 0, i = 0, r = 0;
    for(int x:v) r = r * ++i + __builtin_popcount(use & ~(1<<x)),
        use |= 1 << x; // (note: minus, not ~!)
    return r;
}

```

5.1.1 Derangements

Permutations of a set such that none of the elements appear in their original position.

$$D(n) = (n-1)(D(n-1) + D(n-2)) = nD(n-1) + (-1)^n = \left\lfloor \frac{n!}{e} \right\rfloor$$

5.2 Partitions and subsets

5.2.1 Partition function

Number of ways of writing n as a sum of positive integers, disregarding the order of the summands.

$$p(0) = 1, \quad p(n) = \sum_{k \in \mathbb{Z} \setminus \{0\}} (-1)^{k+1} p(n - k(3k-1)/2)$$

$$p(n) \sim 0.145/n \cdot \exp(2.56\sqrt{n})$$

n	0	1	2	3	4	5	6	7	8	9	20	50	100
$p(n)$	1	1	2	3	5	7	11	15	22	30	627	$\sim 2\text{e}5$	$\sim 2\text{e}8$

5.2.2 Lucas' Theorem

Let n, m be non-negative integers and p a prime. Write $n = n_k p^k + \dots + n_1 p + n_0$ and $m = m_k p^k + \dots + m_1 p + m_0$. Then $\binom{n}{m} \equiv \prod_{i=0}^k \binom{n_i}{m_i} \pmod{p}$.

5.3 General purpose numbers

5.3.1 Stirling numbers of the first kind

Number of permutations on n items with k cycles.

$$c(n, k) = c(n-1, k-1) + (n-1)c(n-1, k), \quad c(0, 0) = 1$$

$$\sum_{k=0}^n c(n, k) x^k = x(x+1) \dots (x+n-1)$$

$$c(8, k) = 8, 0, 5040, 13068, 13132, 6769, 1960, 322, 28, 1$$

$$c(n, 2) = 0, 0, 1, 3, 11, 50, 274, 1764, 13068, 109584, \dots$$

5.3.2 Eulerian numbers

Number of permutations $\pi \in S_n$ in which exactly k elements are greater than the previous element. k j :s s.t. $\pi(j) > \pi(j+1)$, $k+1$ j :s s.t. $\pi(j) \geq j$, k j :s s.t. $\pi(j) > j$.

$$E(n, k) = (n-k)E(n-1, k-1) + (k+1)E(n-1, k)$$

$$E(n, 0) = E(n, n-1) = 1$$

$$E(n, k) = \sum_{j=0}^k (-1)^j \binom{n+1}{j} (k+1-j)^n$$

5.3.3 Stirling numbers of the second kind

Partitions of n distinct elements into exactly k groups.

$$S(n, k) = S(n-1, k-1) + kS(n-1, k)$$

$$S(n, 1) = S(n, n) = 1$$

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

5.3.4 Bell numbers

Total number of partitions of n distinct elements. $B(n) = 1, 1, 2, 5, 15, 52, 203, 877, 4140, 21147, \dots$ For p prime,

$$B(p^m + n) \equiv mB(n) + B(n+1) \pmod{p}$$

5.3.5 Labeled unrooted trees

on n vertices: n^{n-2}
on k existing trees of size n_i : $n_1 n_2 \dots n_k n^{k-2}$
with degrees d_i : $(n-2)! / ((d_1-1)! \dots (d_n-1)!)$

5.3.6 Catalan numbers

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \binom{2n}{n} - \binom{2n}{n+1} = \frac{(2n)!}{(n+1)!n!}$$

$$C_0 = 1, \quad C_{n+1} = \frac{2(2n+1)}{n+2} C_n, \quad C_{n+1} = \sum C_i C_{n-i}$$

$$C_n = 1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, \dots$$

- sub-diagonal monotone paths in an $n \times n$ grid.
- strings with n pairs of parenthesis, correctly nested.
- binary trees with with $n+1$ leaves (0 or 2 children).
- ordered trees with $n+1$ vertices.
- ways a convex polygon with $n+2$ sides can be cut into triangles by connecting vertices with straight lines.
- permutations of $[n]$ with no 3-term increasing subseq.

Graph (6)

6.1 Fundamentals