

# Assignment-5

Trishan Mondal, Soumya Dasgupta, Aarattrick Basu

## Problem 5.14

We will begin with the assumption, the underlying field  $k$  is infinite and algebraically closed (according to contexts). The property of lines passing through points is a projective property. So we can take a suitable projective transformation so that  $P_1 = [0 : 0 : 1]$ . Thus, any line passing through this looks like  $ax + by = 0$  where  $a, b \in k$ . The set of lines passing through  $P_1$  is

$$A = \{x + my : m \in k\} \cup \{y = 0\}$$

Since, the field is infinite, there is infinitely many elements in  $A$ . Given two points in  $\mathbb{P}^2$  there is a unique line passing through  $P_1$  and that point. Thus the set of lines

$$L = \{\ell \text{ pass through } P_1 \text{ and } P_i : 2 \leq i \leq n\} \subset A$$

is finite. So there are only finitely many line in the above set. But in  $A$  there are infinitely elements. So, there are infinitely many elements in  $A \setminus L$ .

Since  $P_1$  is a simple point of  $F$ , there is a tangent  $T$  at  $P$  so that the tangent  $T$  don't contained in  $V(F)$  (or  $F$ ). From the problem 5.12 we can say,

$$\sum I(P; F \cap T) = n$$

where  $n = \deg F$ . Thus, If we take  $P_2, \dots, P_m$  be the other intersection points (here  $m \leq n$ ) of  $T$  and  $F$ , by the previous calculation we can say there exists infinitely many lines through  $P$  don't intersect  $F$  at  $P_i$  ( $i > 1$ ). These lines are transversal to  $F$ . ■

## Problem 5.18

Let us consider the general equation of conic in  $\mathbb{P}^2$ , that is

$$Ax^2 + By^2 + Cz^2 + Exy + Fyz + Gzx = 0$$

Since the point  $[0 : 0 : 1]$  and  $[0 : 1 : 0]$ ,  $[1 : 0 : 0]$  passes through the above conic we can say,  $A = B = C = 0$ . Thus the equation of conic reduces to  $Exy + Fyz + Gzx = 0$ . Also the points  $[1 : 1 : 1]$  and  $[1 : 2 : 3]$  passes through the curve. So we have the following linear equations,

$$\begin{aligned} E + F + G &= 0 \\ 2E + 6F + 3G &= 0 \\ \implies \begin{pmatrix} 1 & 1 & 1 \\ 2 & 6 & 3 \end{pmatrix} \begin{pmatrix} E \\ F \\ G \end{pmatrix} &= 0 \end{aligned}$$

Note that the rows of the above matrix are linearly independent. So the null space of it must have dimension 1. Note that  $(3, -4, 1)^T$  is a solution to the above matrix equation. Since the dimension of null space is 1 we can say any other solution must be a scalar multiple of  $(3, -4, 1)^T$ . So the equation of conic passing through the five points is  $\lambda(3xy - 4yz + zx) = 0$ . This will represent a unique conic in  $\mathbb{P}^2$ . By construction the conic is unique! ■