

Rings and Modules

TRISHAN MONDAL

Assignment-3

Question 1.

- (i) Consider a finitely generated R -module M . If $M_1 \subseteq M_2 \subseteq M$ be submodules such that $M/M_1 \cong M/M_2$ as R -modules, prove that $M_1 = M_2$. If M is not finitely generated, then is the above necessarily true?
- (ii) If $M \cong N$ as R -modules, is it necessarily true that the annihilator ideals are the same; that is, $\text{ann}_R(M) = \text{ann}_R(N)$?

Solution.

- (i). From the Isomorphism theorems we know that,

$$(M/M_1)/(M_2/M_1) \cong (M/M_2)$$

Which is isomorphic to M/M_1 . So, $(M/M_1)/(M_2/M_1) \cong (M/M_1)$. Now we will use [Problem 2.\(i\)](#) to prove, $M_2/M_1 = (0)$. We know the natural surjection, $\pi : M/M_1 \rightarrow (M/M_1)/(M_2/M_1)$ induces an surjective map from M/M_1 to M/M_1 . Since M is finitely generated M/M_1 is finitely generated. So, π induces an Isomorphism. And hence, $\ker(\pi) = M_2/M_1 = (0)$. Since $M_1 \subseteq M_2$ we can say $M_1 = M_2$.

If M is not finitely generated, then the statement is not necessarily true. For example, Consider $M = \bigoplus_{i \in \mathbb{N}} \mathbb{Z}/2\mathbb{Z}$ as a \mathbb{Z} module. Now assume $M_1 = \mathbb{Z}/2\mathbb{Z}$ and $M_2 = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ are submodules of M and satisfying the given condition. We can see that, $M/M_1 \cong \bigoplus_{i \in \mathbb{N}} \mathbb{Z}/2\mathbb{Z} \cong M/M_2$ but $M_2 \neq M_1$.

- (ii). Let, φ be the module Isomorphism between M and N . Let, $r \in \text{ann}_R(M)$ then $rM = 0$ which means, $\varphi(rM) = r\varphi(M) = rN = 0$. So, $r \in \text{ann}_R(N)$. So, $\text{ann}_R(M) \subseteq \text{ann}_R(N)$. Similarly we can get a module homomorphism between N and M . We can show $\text{ann}_R(N) \subseteq \text{ann}_R(M)$. And hence $\text{ann}_R(M) = \text{ann}_R(N)$. ■

Question 2.

- (i) Let M be a finitely generated R -module, and let $\varphi : M \rightarrow M$ be a surjective R -module homomorphism. Prove that φ must be an isomorphism.
- (ii) Recall that R is said to be Noetherian if all ideals are finitely generated. Equivalently, every ascending sequence of ideals $I_1 \subseteq I_2 \subseteq \dots$ stabilizes; that is, there exists $n_0 \geq 1$ such that $I_n = I_{n_0}$ for all $n > n_0$. If R is Noetherian, and $\theta : R \rightarrow R$ is a surjective ring homomorphism, prove that θ must be an isomorphism.

Solution.

(i). M is a finitely generated R -module. So, M can be treated as finitely generated $R[t]$ module by defining $t \cdot m = \varphi(m)$. Since, φ is module homomorphism it's not hard to see this multiplication gives us, M is module over $R[t]$.

We can see that $t \cdot M = \varphi(M) = M$ since, φ is an surjective homomorphism. Similarly $t^n \cdot M = M$. From here we can say that $(t) \cdot M = M$. Now we will propose a lemma.

Lemma. If M is finitely generated module R -module. If A is an ideal of R such that $M = AM$ then we will get $a \in A$ such that $(1 + a) \cdot M = 0$.

Proof. Let, $\{x_1, \dots, x_n\}$ be generators of M . Since, $AM = M$ we can say,

$$\begin{aligned}
 x_1 &= a_{11}x_1 + \dots + a_{1n}x_n \\
 x_2 &= a_{21}x_1 + \dots + a_{2n}x_n \\
 &\vdots \\
 x_n &= a_{n1}x_1 + \dots + a_{nn}x_n \\
 \Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} &= \begin{pmatrix} a_{11} & \dots & a_{1n} \\ a_{21} & \dots & a_{2n} \\ \vdots & \vdots & \vdots \\ a_{n1} & \dots & a_{nn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \\
 \Rightarrow \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} &= \underbrace{\begin{pmatrix} 1 - a_{11} & \dots & -a_{1n} \\ -a_{21} & \dots & -a_{2n} \\ \vdots & \vdots & \vdots \\ -a_{n1} & \dots & 1 - a_{nn} \end{pmatrix}}_{\text{call this matrix } A} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}
 \end{aligned}$$

By multiplying $\text{adj}(A)$ from left to both side of the above equation we will get, $(\det A)x_i = 0$ for all $i = 1, \dots, n$. Now, notice that the determinant of A is in form $1 + a$ for some $a \in A$ as all $a_{ij} \in A$. From here we can see that $(\det A)M = 0 = (1 + a)M$. \square

Now we will proceed to the main problem. As we have already noted $(t) \cdot M = M$ there must exists a polynomial $g \in (t)$ such that, $(1 + g) \cdot M = 0$. If $g = \sum_{i=1}^n a_i t^i$ we can say, $1_M = -\sum a_i \varphi^i$ which means $\psi \circ \varphi = 1_M$ as a module homomorphism where $\psi = -\sum_{i=1}^n a_i \varphi^{i-1}$. So, φ needs to be injective hence φ is an isomorphism. \blacksquare

(ii). $\theta : R \rightarrow R$ is a Surjective map. we can see that, $\ker(\theta) \subseteq \ker(\theta^2)$. So, we can get a chain of ideals $\ker(\theta) \subseteq \ker(\theta^2) \subseteq \dots \subseteq \ker(\theta^n) \subseteq \dots$. Since, R is noetherian, we can Say the above chain terminates. Let say, $\ker(\theta^n) = \ker(\theta^{n_0}) \forall n > n_0$.

Claim: $\ker(\theta^{n_0}) = \{0\}$

Proof: If $f : R \rightarrow R$ is a surjective homomorphism with $\ker(f) = \ker(f^2)$, then if $x \in \ker f$ We have, $f(x) = 0$ we can say, $f(y) = x$ for some $y \in R$ as f is Surjective. so, $f^2(y) = 0 \Rightarrow y \in \ker(f^2)$ So, $y \in \ker(f) = \ker(f^2)$ which means, $f(y) = x = 0$. So, $\ker(f) = \{0\}$. If we take $f = \theta^{n_0}$ then we get $\ker \theta^{n_0} = \{0\}$. \square

We can use the above claim to get $\ker \theta$ is $\{0\}$. which readily gives us θ is an isomorphism. ■

Question 3.

- (i) If M is a finitely generated R -module where R is Noetherian, prove that all submodules of M are necessarily finitely generated.
- (ii) If $N_1 \subseteq N_2$ are R -modules such that N_1 and N_2/N_1 are finitely generated as R -modules, then N_2 must be finitely generated as well.

Solution.

(i). We will proceed by induction on the number of generators of M . If M is $\langle v \rangle$, then $\varphi: R \rightarrow M$ defined by $r \mapsto rv$ is an epimorphism, and so $M \cong R/\ker(\phi)$, and is Noetherian as the quotient of a Noetherian ring. So every submodule of M is finitely generated. Suppose the statement holds for all R -modules with k generators, for $k \leq n$. Let $M = \langle v_1, \dots, v_{n+1} \rangle$. Let $M' = \langle v_1, \dots, v_n \rangle$. Consider N be a submodule of M . Now, assume $T = N \cap M'$. So T is a submodule of M' , hence finitely generated. Now by Isomorphism theorems of module we get,

$$N/T = N/(N \cap M') \cong (N + M')/M'.$$

Now, $(N + M')/M'$ is a M/M' submodule. Since M/M' is cyclic we can say $(N + M')/M'$ is finitely generated (from $n = 1$ case). So, both T and N/T is finitely generated and by [Problem 3.\(ii\)](#) we can say N is finitely generated.

(ii) Given $N_1 \subseteq N_2$. N_1 and N_2/N_1 are finitely generated R -module We can Say. there is a surjective homomorphism $\varphi_1: R^n \rightarrow N_1$ and. $\varphi_2: R^m \rightarrow N_2/N_1$ for some, $n, m \in \mathbb{N}$. Consider the following exact sequence.

$$0 \longrightarrow N_1 \xrightarrow{i} N_2 \xrightarrow{\pi} N_2/N_1 \longrightarrow 0$$

We claim that the above exact sequence can be extended to the following commutative diagram with both rows being exact.

$$\begin{array}{ccccccc} 0 & \longrightarrow & N_1 & \xrightarrow{i} & N_2 & \xrightarrow{\pi} & N_2/N_1 \longrightarrow 0 \\ & & \uparrow \varphi_1 & & \uparrow \varphi & & \uparrow \varphi_2 \\ 0 & \longrightarrow & R^n & \xrightarrow{i'} & R^{m+n} & \xrightarrow{\pi'} & R^m \longrightarrow 0 \end{array}$$

Let, $\{e_1, e_2, \dots, e_n\}$ be generating set of R^n and $\{e_1, \dots, e_{n+m}\}$ be generating set of R^{m+n} . Consider, $i'(e_j) = e_j$ for $j = 1, 2, \dots, n$ and take. $\pi'(e_j) = e_j$ [$j > n$]. This construction allows the bottom row to be exact. $i(\varphi_1(e_1)), \dots, i(\varphi_1(e_n))$ and, $\{e_{n+1}, \dots, e_{n+m}\}$ maps to $\pi^{-1}(\varphi_2(e_{n+1})), \dots, \pi^{-1}(\varphi_2(e_{n+m}))$.

Now we will show that φ is surjective. For any $n \in N_2$ take $\varphi_2^{-1}(\pi(n))$ which is in R^m (as φ_2 is surjective). Now by our construction π' is surjective homomorphism. So, there is a pre image of n in R^{m+n} that is $\pi'^{-1}(\varphi_2^{-1}(\pi(n)))$.

So, we can write $N_2 \cong R^{m+n}/\ker \varphi$. Since, R^{m+n} is finitely generated quotient of it is also finitely generated. So, N_2 is finitely generated.

Question 4.

Prove that every R -module is free if, and only if, R is a field.

Solution. Let's assume every R module is free. Consider $x \neq 0$ is an element of R . Clearly, $R/(x)$ is also a R -module and by definition it must be free. So it must have a R -basis. If B is a basis then, $x \cdot B = \{0\}$. Hence we must get a linear combination of basis elements with each coefficient being $x(\neq 0)$. So, there can not have any such basis. Only possibility is $B = \{0\}$. But then $R \cong Rx$ which means we must get a non zero element y such that $yx = 1$ (Since R is commutative ring and contains 1). So each element of R has inverse and hence R is a Field.

If R is a field then any R -module is a vector space over R and we know every vector space has a basis and every element of the vector space can be written as linear combination of finite basis elements. So every R -module is free. ■

Question 5.(Generalization of a theorem of Cohen due to Jothilingam)

If M is a finitely generated R -module. If PM is a finitely generated submodule of M for each prime ideal P of R , show that all submodules of M are finitely generated as well.

Solution. Let, $\Sigma = \{\text{Set of all submodules of } M \text{ which are not finitely generated}\}$. We can give this set partial order \subseteq , $I \subseteq J$ iff I is contained in J . Let, $M_1 \subseteq M_2 \subseteq M_3 \cdots$ be a chain of submodules of M . Consider the union of these modules $\mathfrak{M} = \cup M_i$. If the union was finitely generated then all the generators of \mathfrak{M} must belong to some M_n . Then any elements of M_n can be generated by these finite generators which is a contradiction. We know every partially ordered sets which has upper bound has a unique maximal element N in Σ . Let, N be the maximal submodule of M which is not finitely generated.

Claim : $\text{ann}_R(M/N)$ is a prime ideal of R .

Proof. Call the ideal $\text{ann}_R(M/N) = I$. Define, $N(s) = \{x \in M : xs \in N\}$. Clearly this is a R -Module. If I is not prime then there is $ab \in I$ such that $a, b \notin I$. We can see that $N + aM$ and $N + bM$ strictly contains N as both a, b don't annihilate M/N . By maximality of N we can say that both $N + aM, N + bM$ are finitely generated. Notice that $N \subseteq N(s)$. We can see $N(a), N(b)$ strictly contains N . Because $N \subset bM$ as. $abM \subseteq N$, $bM \subseteq N(a)$ and hence $N \subset N + bM \subseteq N(a)$. So we can say that $N(a)$ is also finitely generated.

Since $N + aM$ is finitely generated, let $\{n_i + am_i\}$ this set generates $N + aM$. Let, $a \in N \subset N + aM$, we can write, $a = \sum a_i(n_i + am_i) = \sum a_i n_i + a \sum a_i m_i$ Which means $a \sum a_i \in N$ and hence $a \sum a_i \in N(a)$. So, $a \in Rn_1 + \cdots + Rn_k + aN(a)$ Which means $N \subseteq Rn_1 + \cdots + Rn_k + aN(a) \subseteq N$ and hence N is finitely generated which is a contradiction. □

Take an element $x \in M \setminus N$ and put $J = (N : x)$. There are two possibilities, $I = J$ or $I \subset J$.

Case 1: $P = J$, we consider the submodule $N + xR$. By our maximality assumption on N , the submodule $N + xR$ is finitely generated. Let $\{n_i + xt_i : 1 \leq i \leq k\}$ be a set of generators for $N + xR$. Here $n_i \in N$ and $t_i \in R$ for all $1 \leq i \leq k$. Thus, each $a \in N$ can be expressed as

$$a = \sum_{i=1}^k (n_i + xt_i) r_i = \sum_{i=1}^k n_i r_i + x \left(\sum_{i=1}^k t_i r_i \right)$$

with $r_i \in R$ for each $1 \leq i \leq k$. This expression shows that $x \left(\sum_{i=1}^k t_i r_i \right) \in N$ and thus $\left(\sum_{i=1}^k t_i r_i \right) \in J = I$. Hence

$$N \subseteq n_1 R + n_2 R + \dots + n_k R + xP.$$

On the other hand, we know that $xP \subset N$; therefore the last inclusion is actually an equality:

$$N = n_1 R + n_2 R + \dots + n_k R + xI$$

Taking into consideration the fact that $xI \subseteq MP \subset N$, we obtain

$$\begin{aligned} N &= n_1 R + n_2 R + \dots + n_k R + xI \\ &\subseteq n_1 R + n_2 R + \dots + n_k R + IM \\ &\subseteq n_1 R + n_2 R + \dots + n_k R + N \subseteq N. \end{aligned}$$

It follows that

$$N = n_1 R + n_2 R + \dots + n_k R + IM$$

Since IM is finitely generated, the above identity shows that N is finitely generated; which is a contradiction.

Case 2: Now, let us consider the case $I \subset J$. In this case, there exists an element $s \in R \setminus I$ such that $xs \in N$. Thus N is properly contained in both $N(s)$ and $N + sM$. Thus N is finitely generated (we can trace the same proof we did for **Claim**). Which leads us to a contradiction. Hence $\Sigma = \phi$. So all submodule of M is finitely generated. ■

Question 6. Let,

$$0 \rightarrow M' \xrightarrow{\alpha} M \xrightarrow{\beta} M'' \rightarrow 0$$

be a short exact sequence of R -modules which splits. Recall this means α is $1 - 1$, β is onto, $\text{Ker}(\beta) = \text{Image}(\alpha)$, and there exists a 'splitting' (an R module homomorphism) $s : M'' \rightarrow M$ such that $\beta \circ s = \text{Id}_{M''}$. Prove that the set of all splittings $s : M'' \rightarrow M$ is in bijection with the set $\text{Hom}_R(M'', M')$ of all R -module homomorphisms from M'' to M' .

Solution. Consider $\text{Hom}(M'', M)^*$ be the set of all homomorphism from M'' to M that is 'splitting' for the given exact sequence. Let, s is given as initial split for the exact sequence. We will give a set theoretic bijection $F : \text{Hom}(M'', M') \rightarrow \text{Hom}(M'', M)^*$ defined as $F : \varphi \mapsto s + \alpha \circ \varphi$.

Let, $s' \in \text{Hom}(M'', M)^*$ clearly $\beta \circ (s - s') = 0$. From here we can say that, $\text{Im}(s - s') \subseteq \text{Ker } \beta = \text{Im } \alpha$. We can say, $s - s' = \alpha \circ \varphi$ for some $\varphi \in \text{Hom}(M'', M')$. So, $F(\varphi) = s'$. Hence F is surjective.

Also, let, $s + \alpha \circ \varphi = s + \alpha \circ \psi$. We can say, $\alpha \circ \varphi(x) = \alpha \circ \psi(x)$ for all $x \in M''$. Since α is injective we can say that, $\varphi = \psi$. And hence F is injective. So, F is bijective. ■

Question 7. (Local-Global principle)

For a multiplicative subset S of R and an R -module M , define a relation on the set $M \times S$ by

$$(m_1, s_1) \sim (m_2, s_2)$$

if, and only if, there exists $s \in S$ such that $s(s_1m_2 - s_2m_1) = 0$. This is an equivalence relation (assume this - you can verify it but not show the calculation here). Denoting the equivalence class of (m, s) by $\frac{m}{s}$, the equivalence classes form an $S^{-1}R$ -module under the addition

$$\frac{m_1}{s_1} + \frac{m_2}{s_2} = \frac{s_1m_2 + s_2m_1}{s_1s_2}$$

and the scalar multiplication

$$\frac{r}{s} \cdot \frac{m}{t} = \frac{rm}{st}$$

Assume this also - verify it for yourself or look up the routine proof.

Prove that if $S^{-1}M$ is the zero $S^{-1}R$ -module with $S = R - \mathfrak{m}$ for each maximal ideal \mathfrak{m} , prove that $M = (0)$.

Solution. Let, there is a $0 \neq x \in M$. consider \mathfrak{m} be the maximal ideal containing $\text{ann}_R(x)$. Now Consider $S = R - \mathfrak{m}$. If 0 is the only element in $S^{-1}M$ then, $\frac{x}{1} \sim 0$. which means there is $s \in S$ such that $sx = 0$. But then, $s \in \text{ann}_R(x)$ and $s \in R - \mathfrak{m}$. which is a contradiction. ■