## file4

## January 9, 2017

```
In [1]: [x**0.5 \text{ for } x \text{ in } range(100) \text{ if } x**2<25]
Out[1]: [0.0, 1.0, 1.4142135623730951, 1.7320508075688772, 2.0]
In [5]: L=[]
         for x in range(100):
             if x**2<25:
                 L.append (x**0.5)
         L
Out[5]: [0.0, 1.0, 1.4142135623730951, 1.7320508075688772, 2.0]
In [10]: M=list(filter(lambda x: x**2 < 25, range(100)))
          L=list (map (lambda x: x**0.5, M))
          L
Out[10]: [0.0, 1.0, 1.4142135623730951, 1.7320508075688772, 2.0]
In [33]: mx=5
          X=[x/(mx-1) for x in range(mx)]
          Χ
Out[33]: [0.0, 0.25, 0.5, 0.75, 1.0]
In [40]: a=[4,5,0,0,1]
          aX=sum([X[i]*a[i] for i in range(mx)])
          aX/a0
Out[40]: 0.225
In [ ]:
In [7]: mx=5
        my=6
        X=[[x/(mx-1) \text{ for } x \text{ in } range(mx)] \text{ for } y \text{ in } range(my)]
         Χ
```

```
Out[7]: [[0.0, 0.25, 0.5, 0.75, 1.0],
         [0.0, 0.25, 0.5, 0.75, 1.0],
         [0.0, 0.25, 0.5, 0.75, 1.0],
         [0.0, 0.25, 0.5, 0.75, 1.0],
         [0.0, 0.25, 0.5, 0.75, 1.0],
         [0.0, 0.25, 0.5, 0.75, 1.0]]
In [13]: mx=5
         my = 6
         Y=[y/(my-1) \text{ for } x \text{ in } range(mx)] \text{ for } y \text{ in } range(my)]
Out[13]: [[0.0, 0.0, 0.0, 0.0, 0.0],
          [0.2, 0.2, 0.2, 0.2, 0.2],
           [0.4, 0.4, 0.4, 0.4, 0.4]
           [0.6, 0.6, 0.6, 0.6, 0.6],
           [0.8, 0.8, 0.8, 0.8, 0.8],
           [1.0, 1.0, 1.0, 1.0, 1.0]]
In [16]: a=[[1,2,3,4,5],
            [5, 6, 7, 8, 9],
            [9,10,11,12,13],
            [13, 14, 15, 16, 17],
            [10, 25, 40, 25, 10],
            [10, 25, 40, 25, 10]]
         sum([sum(c) for c in a])
Out[16]: 400
In [15]: X
Out[15]: [[0.0, 0.25, 0.5, 0.75, 1.0],
          [0.0, 0.25, 0.5, 0.75, 1.0],
           [0.0, 0.25, 0.5, 0.75, 1.0],
          [0.0, 0.25, 0.5, 0.75, 1.0],
           [0.0, 0.25, 0.5, 0.75, 1.0],
           [0.0, 0.25, 0.5, 0.75, 1.0]]
In [17]: aX1=sum([sum([a[i][j]*X[i][j] for i in range(my)]) for j in range(mx)])
         aX1
Out[17]: 210.0
In [18]: a0=sum([sum([a[i][j] for i in range(my)]) for j in range(mx)])
         a0
Out[18]: 400
In [19]: aX2=sum([sum([a[i][j]*X[i][j]*X[i][j])  for i in range(my)]) for j in range
         aX2
```

```
Out[19]: 148.75
In [20]: ax=aX1/a0
          ax
Out[20]: 0.525
In [21]: aY1=sum([sum([a[i][j]*Y[i][j] for i in range(my)]) for j in range(mx)])
Out[21]: 272.0
In [22]: ay=aY1/a0
          ay
Out[22]: 0.68
In [15]: P=[[x,y,z] for x in range (1,100) for y in range (x,100) for z in range (y,100)
In [9]: def centroids(A):
             my=len(A)
             mx=len(A[0])
             A0=sum([sum([A[i][j] for i in range(my)]) for j in range(mx)])
             X = [[x/(mx-1) \text{ for } x \text{ in } range(mx)] \text{ for } y \text{ in } range(my)]
             Y=[[y/(my-1) \text{ for } x \text{ in } range(mx)] \text{ for } y \text{ in } range(my)]
             AX=sum([sum([A[i][j]*X[i][j] for i in range(my)]) for j in range(mx)])
             AY=sum([sum([A[i][j]*Y[i][j] for i in range(my)]) for j in range(mx)])
             Ax=AX/A0
             Ay = AY/A0
             return (Ax, Ay)
In [16]: centroids(P)
Out [16]: (0.61111111111111112, 0.6151167426677631)
In [29]: P = [[x,y,z] \text{ for } x \text{ in } range(1,26) \setminus
             for y in range (x, 26)
             for z in range (y, 26) if x**2 + y**2 == z**2
          Р
Out[29]: [[3, 4, 5],
           [5, 12, 13],
           [6, 8, 10],
           [7, 24, 25],
           [8, 15, 17],
           [9, 12, 15],
           [12, 16, 20],
           [15, 20, 25]]
In [19]: P
```

```
Out[19]: [[3, 4, 5], [5, 12, 13], [6, 8, 10], [8, 15, 17], [9, 12, 15]]

In [22]: Q=[[x,y,z] for x in range(1,240) for y in range(x,240) for z in range(y,240) centroids(Q)
```

Out [22]: (0.6168428878242194, 0.6161335621366104)

In [ ]: