

VOLTERRA AND FLIESS SERIES EXPANSION

Françoise Lamnabhi-Lagarigue

Laboratoire des Signaux et Systèmes, CNRS, Supelec, France

Keywords: nonlinear system, Volterra series, Fliess generating power series, Volterra kernel, response to typical inputs, formal computing

Contents

- 1. Introduction
- 2. Functional representation of nonlinear systems
 - 2.1. Volterra Functional Series
 - 2.2. On the Convergence of Volterra Series
- 3. Recursive computation of the kernels.
 - 3.1. Exponential Input Method
 - 3.2. Differential Geometry Approach
 - 3.3. Algebraic Approach
 - 3.4. Links between Volterra and Fliess Series
 - 3.5. Efficient Computation of Volterra Kernels
- 4. Computation of the response to typical inputs
 - 4.1. Transfer Function Approach: Association of Variables
 - 4.2. Algebraic Approach
- Glossary
- Bibliography
- Biographical Sketch

Summary

After recalling the definition of the Volterra series expansion and some of its convergence issues, we will study various methods in order to derive the Volterra kernels and the response to typical inputs. In particular, an algorithm, by means of which a large class of nonlinear systems can be analysed, is described.

This algorithm is based on an algebraic approach to Volterra functional expansions using noncommutative generating power series; this approach allows a natural generalization, to the nonlinear domain, of the symbolic operational calculus of Heaviside, widely used in linear system theory. Moreover, it has the advantage, compared with the method using the multidimensional Laplace transforms, of allowing an easy implementation on a computer.

1. Introduction

The input-output behavior of a large class of nonlinear systems can be described by the Volterra functional expansion

$$y(t) = \int_0^t h_1(t, \tau_1) u(\tau_1) d\tau_1 + \int_0^t \int_0^{\tau_1} h_2(t, \tau_1, \tau_2) u(\tau_1) u(\tau_2) d\tau_2 d\tau_1 + \dots,$$

where $y(t)$ is the circuit output and $u(t)$ is the circuit input (assumed to be here scalar for simplicity sake). h_n is the n th-order Volterra kernel. This expansion is a generalisation of the well-known convolution integral

$$y(t) = \int_0^t h_1(t, \tau_1) u(\tau_1) d\tau_1$$

used in linear system theory.

These expansions are used in every branch of nonlinear system theory: identification and modeling, realization, stability, optimal control, stochastic differential equations, and filtering.

Although the Volterra series has been successfully used in many applications, it has not received a great deal of attention from engineers and designers. The reason for this seems to be the tedious computations involved in the determination of Volterra kernels. The algebraic approach to nonlinear functional expansions based on noncommutative generating power series, offers a powerful and systematic tool for analyzing a large class of nonlinear systems.

After recalling the definition of the Volterra series expansion and some of its convergence issues, we will study various methods in order to derive the Volterra kernels and the response to typical inputs. The analysis is then applied to the study of weakly nonlinear circuits in order to derive distortion rates or intermodulation products.

2. Functional Representation of Nonlinear Systems

2.1. Volterra Functional Series

For the simplicity of presentation, we shall consider time-invariant systems. If a system is linear and time-invariant, then the output $y(t)$ can be expressed as the convolution of the input $u(t)$ with the system unit impulse response $h(t)$:

$$y(t) = \int_{-\infty}^{\infty} h(\tau) u(t - \tau) d\tau \quad (1)$$

The system unit impulse response $h(t)$ completely characterizes the linear time-invariant system since, once known, the response to any input can be determined from (1). A system is said to be causal if the output at any given time does not depend on future values of the input. That is, for any time t_1 ,

$$y(t_1) = \int_{-\infty}^0 h(\tau) u(t_1 - \tau) d\tau = 0.$$

This will be so if and only if

$$h(\tau) = 0, \quad \text{for} \quad \tau < 0.$$

The extension of (1) to nonlinear time-invariant systems with memory is the Volterra series

$$y(t) = h_0 + \sum_{n=1}^{\infty} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} h_n(\tau_1, \tau_2, \dots, \tau_n) u(t-\tau_1) u(t-\tau_2) \cdots u(t-\tau_n) d\tau_1 d\tau_2 \cdots d\tau_n \quad (2)$$

This functional form was first studied by Volterra. Much of his work in this area is summarized in his book. The functions $h_n(\tau_1, \tau_2, \dots, \tau_n)$ are called the Volterra kernels of the system. A nonlinear system which can be represented by a Volterra series is completely characterized by its Volterra kernels.

Also, with an argument similar to that of linear systems, it can be shown that the nonlinear system is causal if and only if

$$h_n(\tau_1, \tau_2, \dots, \tau_n) = 0, \quad \text{for } \tau_j < 0, \quad j = 1, \dots, n.$$

It is well known that, without loss of generality, the kernels can be assumed to be symmetric. In fact any kernel $h_n(\tau_1, \tau_2, \dots, \tau_n)$ can be replaced by a symmetric one by setting

$$h_n^{\text{sym}}(\tau_1, \tau_2, \dots, \tau_n) = \frac{1}{n!} \sum_{(\tau_{i_1}, \tau_{i_2}, \dots, \tau_{i_n}) \in S} h_n(\tau_{i_1}, \tau_{i_2}, \dots, \tau_{i_n}),$$

where S is the set of all permutations of τ_1, \dots, τ_n .




Figure 1: An example of a nonlinear system

The multiple Laplace transform $L[.]$ of the n th-order Volterra kernel $n > 0$ (one-sided in each variable)

$$H_n(s_1, \dots, s_n) = \int_0^{\infty} \cdots \int_0^{\infty} h_n(\tau_1, \dots, \tau_n) e^{-s_1 \tau_1} \cdots e^{-s_n \tau_n} d\tau_1 d\tau_2 \cdots d\tau_n$$

is called the n th-order transfer function. Since $h_n(\tau_1, \dots, \tau_n)$ is symmetric, so is $H_n(s_1, \dots, s_n)$.

2.2. On the Convergence of Volterra Series

The Volterra series is a power series with memory. This can be seen by changing the input by a gain factor c so that the new input is $cu(t)$. By using (2), the new output is

$$y(t) = h_0 + \sum_{n=1}^{\infty} c^n \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} h_n(\tau_1, \tau_2, \dots, \tau_n) u(t - \tau_1) u(t - \tau_2) \cdots u(t - \tau_n) d\tau_1 d\tau_2 \cdots d\tau_n,$$

which is a power series in the amplitude factor c . It is a series with memory since the integrals are convolutions. As a consequence of its power series character, there are some limitations associated with the application of the Volterra series to nonlinear problems. One major limitation is the convergence of this series.

In order to illustrate this let us consider the system of Figure 1 where the system L is a linear system with the unit impulse response $h(t)$

$$z(t) = \int_{-\infty}^{\infty} h(\tau) u(t - \tau) d\tau \quad (3)$$

and the system N is a nonlinear no-memory system with the input-output relation

$$y(t) = N[z(t)] = \frac{z(t)}{1 + z^2(t)}.$$

The Taylor series expansion of this expression is

$$y(t) = \sum_{n=0}^{\infty} (-1)^n [z(t)]^{2n+1} \quad (4)$$

which converges only for $z^2(t) < 1$. The Volterra series representation of the overall system T is now easily derived by substituting (3) for (4) to obtain

$$y(t) = \sum_{n=0}^{\infty} (-1)^n \left[\int_{-\infty}^{\infty} h(\tau) u(t - \tau) d\tau \right]^{2n+1},$$

in which the Volterra kernels are

$$h_{2n+1}(\tau_1, \dots, \tau_{2n+1}) = (-1)^n h(\tau_1) h(\tau_2) \dots h(\tau_{2n+1})$$

and

$$h_{2n}(\tau_1, \dots, \tau_{2n}) = 0, \quad n \geq 0.$$

Since the Taylor series converges only for $z^2(t) < 1$, the above Volterra series will diverge at those times for which $|z(t)| \geq 1$. The Volterra series thus is valid only for the class of inputs $u(t)$ for which the amplitude of $z(t)$ is less than one.

Now let N be replaced by the following nonlinear no-memory system

$$y(t) = \text{Esign}[z(t)].$$

Clearly, the system T cannot be represented by a Volterra series. It is therefore evident that generally many types of nonlinear systems, such as those that include saturating elements, cannot be characterized by a Volterra series that converges for all inputs.

Under certain conditions, a functional $y(t) = T[x(t)]$ can be approximated to any desired degree of accuracy by a finite series of the form of equation (2). Such a functional is called continuous. In particular it is easy to show that the functional relation between the solution (output) and the forcing function (input) of a nonlinear differential equation with constant coefficients which satisfies the Lipschitz conditions is continuous.

If $T[x(t)]$ can exactly be represented by a converging infinite series of the form of equation (2), it is called *analytic* or *weak*. Conditions for convergence are discussed by Volterra and Brillant. Brillant also notes that two special types of systems, for which the functional relation between input and output is analytic, are a linear system and a nonlinear no-memory systems with a power series relation between input and output. He then shows that various combinations such as cascading, adding, or multiplying such systems results in an analytic system.

In practice, most of the analogue circuits used in communication systems, such as modulators, mixers, amplifiers, harmonic oscillators, etc..., are of a weak nature and therefore analysed and designed in the frequency domain. For such weakly nonlinear circuits (having, say, distortion components of 20dB or more below the fundamental one), the Volterra series technique can be readily used in the frequency domain to obtain results both quantitatively and qualitatively.

Given an input-output map described by a nonlinear control system $\dot{x} = f(x, u)$ and a nonlinear output $y = h(x)$, there exists simple means for obtaining a series representation of the output $y(t)$ in terms of the input $u(t)$. When the control enters linearly, $\dot{x} = f(x) + ug(x)$, the method yields the existence of a Volterra series representation.

3. Recursive Computation of the Kernels

Several methods have been developed in the literature for determining the kernels or the associated transfer functions.

Among them, the method of exponential inputs is particularly used. After recalling this method we describe a differential geometry approach and an algebraic approach based on generating power series when the system is described by a set of differential equations. We shall see that the algebraic approach has the advantage of being easily implementable on a computer by using algebraic computing software.

3.1. Exponential Input Method

Let us consider the Volterra series expansion of a nonlinear system of the form

$$y(t) = \sum_{n=1}^{\infty} \int_0^{t_1} \cdots \int_0^{t_n} h_n(\tau_1, \tau_2, \dots, \tau_n) u(t-\tau_1) u(t-\tau_2) \cdots u(t-\tau_n) d\tau_1 d\tau_2 \cdots d\tau_n. \quad (5)$$

Let the input $u(t)$ be a sum of exponentials

$$u(t) = e^{s_1 t} + e^{s_2 t} + \cdots + e^{s_k t},$$

where s_1, s_2, \dots, s_k are rationally independent. This means that there are no rational numbers $\alpha_1, \alpha_2, \dots, \alpha_k$ such that the sum $\alpha_1 s_1 + \alpha_2 s_2 + \cdots + \alpha_k s_k$ is rational. Then (5) becomes

$$y(t) = \sum_{n=1}^{\infty} \left[\sum_{k_1=1}^k \cdots \sum_{k_n=1}^k H_n(s_{k_1}, \dots, s_{k_n}) e^{(s_{k_1} + \cdots + s_{k_n})t} \right]. \quad (6)$$

If each s_i occurs in $(s_{k_1}, \dots, s_{k_n})$, m_i times, then there are

$$\frac{n!}{m_1! m_2! \cdots m_k!}$$

identical terms in the expression between brackets. Thus (5) can be written in the form

$$y(t) = \sum_{n=1}^{\infty} \sum_m \frac{n!}{m_1! m_2! \cdots m_k!} H_n(s_{k_1}, \dots, s_{k_n}) e^{(s_{k_1} + \cdots + s_{k_n})t}, \quad (7)$$

where m under the summation sign indicates that the sum includes all the distinct vectors (m_1, \dots, m_k) such that $\sum_{i=1}^k m_i = n$. Note that if $m_1 = m_2 = \cdots = m_k = 1$ then the amplitude associated with the exponential component $e^{(s_{k_1} + \cdots + s_{k_n})t}$ is simply $k! H_k(s_1, \dots, s_k)$.




Figure 2: A simple nonlinear circuit

This suggests a recursive procedure for determining all the nonlinear transfer functions from the behavior of a system.

Let us apply the method to the simple nonlinear circuit of Figure 2 consisting of a capacitor, a linear resistor and a nonlinear resistor in parallel with the current source $i(t)$.

The nonlinear differential equation relating the current excitation $i(t)$ and the voltage $v(t)$ across the capacitor is given by

$$\dot{v} + k_1 v + k_2 v^2 = i \quad (8)$$

Let $i(t) = e^{st}$. Equating the coefficients of e^{st} on both sides of (8) after the substitution of (7) for $v(t)$ we get

$$H_1(s) = \frac{1}{s + k_1}.$$

In order to determine $H_2(s_1, s_2)$ let us take $i(t) = e^{s_1 t} + e^{s_2 t}$ and identify the coefficient of the term $2!e^{(s_1+s_2)t}$ after the substitution of (7) for $v(t)$ in both sides of (8). We obtain $H_2(s_1, s_2)$ in term of $H_1(s)$ as follows

$$H_2(s_1, s_2) = -k_2 H_1(s_1) H_1(s_2) H_1(s_1 + s_2).$$

Similarly the third-order transfer function is obtained by injecting a sum of three exponentials inputs

$$i(t) = e^{s_1 t} + e^{s_2 t} + e^{s_3 t}.$$

It follows

$$H_3(s_1, s_2, s_3) = -\frac{2}{3} [H_2(s_1, s_2) H_1(s_3) + H_2(s_2, s_3) H_1(s_1) + H_2(s_1, s_3) H_1(s_2)] H_1(s_1 + s_2 + s_3)$$

Repeating this process indefinitely gives higher order nonlinear transfer functions in terms of lower-order nonlinear transfer functions.

-
-
-

TO ACCESS ALL THE **23 PAGES** OF THIS CHAPTER,
[Click here](#)

Bibliography

Barrett J. (1963). The use of functionals in the analysis of nonlinear physical systems, *Journal of Electronics and Control*, 15, pp. 567--615. [This report develops a method of analysis applicable equally to linear and nonlinear systems. The main method discussed is the expansion of the input-output relation in a functional power series.]

Bedrosian E. and Rice S. (1971). The output properties of Volterra systems (nonlinear systems with memory) driven by harmonic and Gaussian input, *Proceedings IEEE*, 59, pp. 1688--1707. [For a wide class of systems troublesome distortions often occur. These distortions can be computed with the help of Volterra series. In this paper, results, both old and new, which will aid the reader in applying Volterra-series-type analyses to systems driven by sine waves or Gaussian noise, are presented.]

Boyd S., Chua L.O. and Desoer C.A. (1984). Analytical foundations of Volterra series, *IMA J. of Mathematical Control & Information*, Oxford Univ. Press, 1, pp. 243—282 [In this paper it is studied the *analysis* involved with Volterra series. It is addressed system-theoretic issues ranging from bounds on the gain and incremental gain of Volterra series operators to the existence of Volterra series operator inverses, and mathematical topics such as the relation between Volterra series operators and Taylor series. A general steady-state theorem for Volterra series operators is proven, and then it is established a general formula for the spectrum of the output of a Volterra series operator in terms of the spectrum of a periodic input.]

Boyd S. and Chua L.O. (1985). Fading memory and the problem of approximating nonlinear operators with Volterra series, *IEEE CAS* 32(11), pp. 1150--1161. [This paper shows that any time-invariant continuous nonlinear operator can be approximated by a Volterra series. Moreover it is shown that the approximating operator can be realized as a finite-dimensional linear dynamical system with a nonlinear readout map.]

Bussgang J., Ehrman L. and Graham J.W. (1974). Analysis of nonlinear systems with multiples inputs, *Proceedings IEEE*, 62, pp. 1088--1119. [In this paper analytical modelling of communication receivers to account for their nonlinear response to multiple input signals is discussed. The method is based on the application of the Wiener-Volterra analysis of nonlinear functionals.]

Brockett R.W. (1976). Volterra series and geometric control theory, *Automatica*, 12, pp. 167-176. Brockett R.W. and Gilbert E.G. (1976). An addendum to Volterra series and geometric control theory, *Automatica*, 12, p. 635. [In these papers it is shown that controlled differential equations which are analytic in the state and linear in the control have solutions which can be expanded in a Volterra series provided there is no finite escape time. Necessary and sufficient conditions for a Volterra series to be realizable by a linear-analytic system are also given.]

Chua L.O. and Ng C.Y. (1979). Frequency domain analysis of nonlinear systems: general theory & formulation of transfer functions, *Electronics Circuits and Systems*, 3(4), pp. 165--185 & 3(6) pp. 257—269. [A unified study of the applications of Volterra functional series to nonlinear system analysis is presented with special emphasis on frequency-domain results. In particular, an in-depth analysis of the harmonic and intermodulation frequency components due to each Volterra kernel of a given order is

presented and explicit formulas are given.]

Crouch P.E. (1981) Dynamical realizations of finite Volterra series, *SIAM J. Control and Optimization*, 19(2), pp. 177--202. [In this paper, realizations of finite Volterra series are viewed as nonlinear analytic input-output systems, with state space described by an analytic manifold. It is shown that a minimal realization of a finite Volterra series can be described as a cascade of linear subsystems with polynomial link maps.]

Crouch P.E. and Lamnabhi-Lagarigue F. (1989). Algebraic and multiple integral identities, *Acta Applicanda Mathematica*, 15, pp. 235-274. [Identities involving symmetric products in an abstract algebra are developed. This abstract setting allow to derive identities involving multiple integrals of functions of many variables which are useful in system theory.]

Flake R.H. (1963). Volterra series representation of nonlinear systems, *IEEE Trans. Appl. Ind.*, 81, pp. 330-335. [This paper presents an approach for determining the kernels or the associated transfer functions.]

Fliess M. (1981). Fonctionnelles causales non linéaires et indéterminées non commutatives, *Bull. Soc. Math. France*, 109, pp. 3-40. [This paper introduces the algebraic approach to nonlinear functional expansions based on noncommutative generating power series.]

Fliess M., Lamnabhi M. and Lamnabhi-Lagarigue F. (1983). Algebraic approach to nonlinear functional expansions, *IEEE CS*, 30, pp. 550--570. [In this paper, a new theory of functional is presented which makes use of formal power series in several noncommutative variables and iterated integrals. The symbolic calculus introduced replaces the so-called association of variables.]

Fliess M. and Lamnabhi-Lagarigue F. (1982). Application of a new functional expansion to the cubic anharmonic oscillator, *J. Math. Physics*, 23, pp. 495--502. [A representation of causal functionals has been introduced which makes use of noncommutative generating power series and iterated integrals. This technique allows the solutions of nonlinear differential equations with forcing terms and both the deterministic and the stochastic cases to be obtained in a simple and natural way.]

George D. (1959). Continuous nonlinear systems, MIT RLE Technical Report No. 355. [This PhD thesis introduces the technique of association of variables.]

Gilbert E.G. (1977). Functional expansions for the response of nonlinear differential systems, *IEEE AC*, 22, pp. 909--921. [This paper analyses functional approximations of some nonlinear differential input-output equations].

Hassine L., Toffano Z., Lamnabhi-Lagarigue F., Joindot I. and Destrez A. (1994). Volterra functional series for noise in semiconductor lasers, *IEEE J. Quantum Electron.*, 30(11), pp. 2534--2546. [In this paper, a full analysis of noise in semi-conductor lasers has been presented. Time-covariance and cross-covariance functions of all variables governing single-mode semi-conductor laser behavior are given.]

Lamnabhi M. (1986). Functional analysis of nonlinear circuits: a generating power series approach, *IEE Proceedings*, 133, pp. 375-384. [An algorithm based on an algebraic approach to Volterra functional expansions using noncommutative generating power series is derived. This approach allow a natural generalization, to the nonlinear domain, of the symbolic operational calculus of Heaviside.]

Lamnabhi-Lagarigue F. and Lamnabhi M. (1979). Détermination algébrique des noyaux de Volterra associés à certains systèmes non linéaires, *Ricerche di Automatica*, 10, pp. 17-26. [Using non-commutative formal power series, an algebraic closed form of Volterra kernels of nonlinear differential systems with multiple inputs has been derived.]

Lamnabhi-Lagarigue F. and Stefani G. (1990). Singular optimal control problems: on the necessary conditions for optimality, *SIAM J. Contr. Optim.*, 28, pp. 823-840. [Higher-order necessary conditions for the optimality of a totally singular arc in the case of nonlinear systems not necessarily linear in the control variables have been obtained. New third order conditions are derived.]

Leo Van Hemmen J., Kistler W. M. and Thomas E. G. F. (2000). Calculation of Volterra kernels for solutions of nonlinear differential equations, *SIAM J. Appl. Math.*, 61(1), pp. 1--21. [In this paper it is considered vector-valued autonomous differential equations of the form $\dot{x} = f(x) + v$ with analytic f and investigate the nonanticipative solution operator v maps to $x = Av$ in terms of its Volterra series. It is shown that Volterra kernels of order > 1 occurring in the series expansion of the solution operator A are

continuous functions, and it is established recurrence relations between the kernels allowing their explicit calculation.]

Lesiak C. and Krener A.J. (1978). The existence and uniqueness of Volterra series for nonlinear systems, IEEE AC, 23, pp. 1090--1095. [In particular, the uniqueness of Volterra series representations is discussed in this paper. This work generalizes Brockett's technique and Gilbert's work]

Martin A. (1991). Calcul d'approximations de la solution d'un syst\eme non lin'eaire utilisant le logiciel SCRATCHPAD, in Algebraic Computing in Control, Lect. Note Contr. Inform. Sc., Springer Verlag (G.~Jacob and F.~Lamnabhi-Lagarigue Eds.), 165, 1991. [Computations of the solutions of some polynomial differential equations are derived using an algebraic computing language.]

Rugh W.J. (1981). Nonlinear System Theory, Baltimore, MD : The John Hopkins Univ. Press, and Schetzen M. (1980). The Volterra and Wiener Theories of Nonlinear Systems, Wiley. [In these books, the input-output behaviour of a large class of nonlinear circuits has been analyzed by using the Volterra functional expansion.]

Tang Y.S., Mees A.I. and Chua L.O. (1983). Hopf bifurcation via Volterra series, IEEE AC, 28(1), pp. 42--53.[The Hopf bifurcation theorem gives a method of predicting oscillations which appear in a nonlinear system when a parameter is varied. The paper shows the links with this theorem and the Volterra series expansion associated with the solutions.]

Van Trees H.I. (1964). Functional techniques for the analysis of nonlinear behaviour of phased-locked loops, Proceedings IEE, 52, pp. 894--911. [This paper shows the use of functional techniques for the analysis of some electronic circuits.]

Volterra V. (1930). Theory of Functionals (translated from the Spanish), London: Blackie (reprinted by Dover, New York, 1959). [In this pioneering book, it is introduced the Volterra series expansions.]

Wiener N. (1958) Nonlinear Problems in Random Theory, New York : Wiley. [In this book, mathematical foundations of functional expansions in the stochastic case are presented; see also the paper "Response of a linear device to noise" by the same author and published in the MIT Cambridge report on Radiation Lab. Techn. 129, 1942.]

Biographical Sketch

Francoise Lamnabhi-Lagarigue is Directeur de Recherche au Centre National de la Recherche Scientifique since 1993. She obtained the "maîtrise des sciences mathématiques pures" degree at the Université Paul Sabatier (Toulouse) in 1976 and she held a CNRS position in 1980. She obtained her Docteur d'Etat es Sciences Physiques from Université Paris Sud in 1985. Some recent activities include the following :

- Editor of International Journal of Control;
- Responsible (1998-2001) of an agreement with the Groupe PSA on the Global Chassis Control project;
- Coordinator (1998-2002) of the Nonlinear Control Network (NCN)
<http://www.supelec.fr/lss/NCN>;
- Director of the Marie Curie Control Training Site (CTS) <http://www.supelec.fr/lss/CTS>;
- Expert evaluator at the European Commission;
- Nominated member of the Board of Governors of the IEEE Control Systems Society for the year 2002 and elected Member for 2003-2005.

Her main research interests lie in the fields of nonlinear systems including system analysis and control design. Several important contributions should be emphasized. The analysis of nonlinear systems using functional expansions based on the Fliess generating power series and the development of formal computing for the Volterra series expansion associated with the output of nonlinear systems. Then singular optimal control and singular tracking for nonlinear systems have been studied. In the same time, an exact combinatorial formula for the derivatives of the output in terms of the derivatives of the inputs has been derived. More recently, her research interests include performance and robustness issues in

nonlinear control, identification of nonlinear systems, and the application of these techniques to power systems, to vehicle global chassis control, to magnetic suspension, and to hydraulic actuators.

Francoise Lamnabhi-Lagarigue is author and co-author of more than 100 publications which include one book, 50 journal papers and chapters of book, a Special issue of the International Journal on Control (Recent advances in the control of nonlinear systems, Vol 71(5), 1998), the co-edition of 6 books (the more recent ones being: - Stability and Stabilization of Nonlinear Systems, LNCIS 246, Springer-Verlag, 1999; - Nonlinear Control in the Year 2000, LNCIS 258 et LNCIS 259, Springer-Verlag, 2000; - Advances in Control of Nonlinear Systems, LNCIS 264, Springer-Verlag, 2001; - Commande des Systemes Non Lineaires, 2 volumes, Traite IC2, Hermes, 2002), about 45 published conference papers, 2 theses and various reports and European proposals.