

The k-Metallic Mirror: An Algebraic System, its Foundational Symbolic Proofs, and a Proposed Framework for the Riemann Hypothesis

Tristen Harr

June 3, 2025

Abstract

This paper introduces the k-Metallic Algebra, a novel algebraic framework parameterized by a real number $k > 0$. The system is constructed upon two fundamental constants, T_k and J_k , uniquely determined by their sum $T_k + J_k = k/2$ and ratio $T_k/J_k = \Phi_k$, where $\Phi_k = \frac{k+\sqrt{k^2+4}}{2}$ is the k -th metallic mean. We establish the core algebraic properties of this system for general k through symbolic proofs. A canonical case, the Golden Algebra, emerges for $k = 1$, where $\Phi_1 = \phi = \frac{1+\sqrt{5}}{2}$ (the golden ratio). The constants of the Golden Algebra ($T_1 = \frac{\sqrt{5}-1}{4}$, $J_1 = \frac{3-\sqrt{5}}{4}$, $K_1 = \frac{-(\sqrt{5}+1)}{4}$, and $H_1 = T_1 J_1 = \frac{\sqrt{5}-2}{4}$) are shown to exhibit profound connections to pentagonal geometry, number theory (Fibonacci-Lucas sequences, Pell's equation), matrix algebra, and fundamental mathematical constants such as π and e (via Euler's identity, $e^{i\pi} = -1$). We prove a critical "Golden Ratio Condition": if the general k-Metallic mean Φ_k is constrained to be the golden ratio ϕ , then k is uniquely forced to be 1. This algebraic rigidity underpins the "Mirror Math" hypothesis, a proposed structural framework for the Riemann Hypothesis (RH), which suggests that RH may be a consequence of the algebraic properties of the Golden Algebra. The symbolic proofs are detailed within an accompanying Mathematica notebook and further substantiated for the $k = 1$ case by a Python script validating 207 distinct properties.

<https://github.com/TristenHarr/goldenalgebra>

Contents

1	Introduction	2
2	The k-Metallic Algebra (General $k > 0$)	2
2.1	Definitions	3
2.2	Fundamental Identities	3
2.3	Minimal Polynomials	4
3	The Golden Algebra ($k = 1$) - Canonical Properties	4
3.1	Specific Values and Geometric Interpretation	4
3.2	Key Identities and Polynomials	5
3.3	Connections to Number Theory ($k = 1$)	5
3.4	Matrix Representation ($k = 1$)	5
3.5	Galois Conjugation in $\mathbb{Q}(\sqrt{5})$ ($k = 1$)	5
3.6	Exponential Identities and Euler's Identity ($k = 1$)	6
3.7	Explicit Symbolic Forms ($k = 1$)	6
4	The "Golden Ratio Condition" and Algebraic Rigidity	6

5	Complex Representation Z_k and Geometric Identity	6
6	The "Mirror Math" Hypothesis for the Riemann Hypothesis	7
7	Conclusion and Future Work	7

1 Introduction

The study of metallic means, a generalization of the golden ratio, has revealed rich mathematical structures. The k -th metallic mean is defined as $\Phi_k = \frac{k+\sqrt{k^2+4}}{2}$ for a positive real number k . This paper introduces a novel algebraic system, termed the k -Metallic Algebra, built upon two constants T_k and J_k derived from Φ_k and k . Specifically, T_k and J_k are determined by the system of equations:

$$T_k + J_k = k/2 \quad (1)$$

$$T_k/J_k = \Phi_k \quad (2)$$

This framework leads to a consistent set of algebraic identities and reveals deep connections across various mathematical disciplines. The special case $k = 1$, yielding the Golden Algebra where $\Phi_1 = \phi$ (the golden ratio), is of particular interest due to its links with pentagonal geometry and classical number theory.

The central aim of this paper is threefold:

1. To formally define the k -Metallic Algebra and symbolically prove its fundamental properties for general $k > 0$.
2. To explore the canonical $k = 1$ Golden Algebra, demonstrating its unique characteristics, including relationships with Fibonacci-Lucas numbers, Pell's equation, matrix representations, Galois theory, and its ability to encode Euler's identity.
3. To establish the "Golden Ratio Condition"—a proof that constraining the general k -Metallic mean Φ_k to be the golden ratio ϕ uniquely forces $k = 1$.
4. To propose the "Mirror Math" hypothesis, a framework suggesting that the algebraic structure of the Golden Algebra might provide a symbolic model reflecting deeper mathematical truths, including a potential structural explanation for the Riemann Hypothesis (RH).

The symbolic proofs presented herein are primarily derived from a Mathematica notebook, 'GoldenAlgebraFoundation.nb'. Furthermore, an extensive suite of 207 properties specific to the $k = 1$ Golden Algebra has been independently validated using a Python script with the Sympy library, 'golden_algebra_validator.v3.py'. This work seeks to lay a rigorous algebraic foundation for these observations and hypotheses.

2 The k -Metallic Algebra (General $k > 0$)

This section details the construction and fundamental symbolic properties of the k -Metallic Algebra for any real $k > 0$. All propositions in this section are symbolically proven in the accompanying Mathematica notebook 'GoldenAlgebraFoundation.nb'.

2.1 Definitions

Let k be a positive real number.

Definition 2.1 (k -Metallic Mean). *The k -th metallic mean, Φ_k , is defined as:*

$$\Phi_k = \frac{k + \sqrt{k^2 + 4}}{2}$$

Definition 2.2 (Primary Constants T_k, J_k). *The primary constants T_k and J_k of the k -Metallic Algebra are uniquely determined by the system:*

$$T_k + J_k = k/2 \tag{3}$$

$$T_k/J_k = \Phi_k \tag{4}$$

Solving this system yields the explicit forms (as used in ‘GoldenAlgebraFoundation.nb’):

$$T_k = \frac{k - 2 + \sqrt{k^2 + 4}}{4} \tag{5}$$

$$J_k = \frac{k + 2 - \sqrt{k^2 + 4}}{4} \tag{6}$$

Definition 2.3 (Derived Constants H_k, K_k). *Two auxiliary constants, H_k and K_k , are defined as:*

$$H_k = T_k J_k = \frac{\sqrt{k^2 + 4} - 2}{4} \tag{7}$$

$$K_k = -k/2 - T_k = \frac{2 - 3k - \sqrt{k^2 + 4}}{4} \tag{8}$$

2.2 Fundamental Identities

The following identities are direct consequences of the definitions and are symbolically proven in ‘GoldenAlgebraFoundation.nb’.

Proposition 2.4 (Sum Constraint). *For any $k > 0$, $T_k + J_k = k/2$.*

Proof Sketch: This is definitional from Eq. (3) and verified by substituting the explicit forms (5) and (6). \square

Proposition 2.5 (Ratio Identity). *For any $k > 0$, $T_k/J_k = \Phi_k$.*

Proof Sketch: This is definitional from Eq. (4) and verified by substituting the explicit forms. \square

Proposition 2.6 (Uniqueness Constraint). *For any $k > 0$,*

$$\frac{T_k}{J_k} - \frac{J_k}{T_k} = k$$

Proof Sketch: This identity follows directly from $\Phi_k - 1/\Phi_k = k$, which is a property of the metallic mean Φ_k (since $\Phi_k^2 - k\Phi_k - 1 = 0$). Symbolically proven in the notebook. \square

Proposition 2.7 (Bridge Identity). *For any $k > 0$,*

$$T_k - J_k = 2T_k J_k$$

This identity is algebraically equivalent to the defining quadratic equation for Φ_k , namely $\Phi_k^2 - k\Phi_k - 1 = 0$.

Proof Sketch: Symbolically proven by substituting the explicit forms of T_k and J_k . The equivalence can be shown by algebraic manipulation. Dividing $T_k - J_k = 2T_k J_k$ by J_k (assuming $J_k \neq 0$) gives $T_k/J_k - 1 = 2T_k$, so $\Phi_k - 1 = 2T_k$. Substituting $T_k = J_k \Phi_k$ and $J_k = k/(2(\Phi_k + 1))$ into $\Phi_k - 1 = 2T_k$ and simplifying leads back to $\Phi_k^2 - k\Phi_k - 1 = 0$. \square

2.3 Minimal Polynomials

The constants T_k , J_k , and K_k are algebraic.

Proposition 2.8 (Minimal Polynomials for T_k, J_k, K_k). *For any $k > 0$, the constants T_k, J_k , and K_k satisfy the following quadratic polynomials respectively:*

$$\begin{aligned} 4x^2 + (4 - 2k)x - k &= 0 & (\text{for } x = T_k) \\ 4x^2 - (4 + 2k)x + k &= 0 & (\text{for } x = J_k) \\ 4x^2 + (6k - 4)x + (2k^2 - 3k) &= 0 & (\text{for } x = K_k) \end{aligned}$$

The monic versions of these polynomials, $x^2 + b_i x + c_i = 0$, each have a discriminant $\Delta = b_i^2 - 4c_i = \frac{k^2 + 4}{4}$.

Proof Sketch: Derived by isolating the $\sqrt{k^2 + 4}$ term in the explicit definitions of T_k, J_k , and K_k and squaring both sides. The discriminant calculation for the corresponding monic polynomials is then straightforward. Symbolically proven in ‘GoldenAlgebraFoundation.nb’. \square

3 The Golden Algebra ($k = 1$) - Canonical Properties

The specialization to $k = 1$ yields the Golden Algebra, a system with particularly rich structure and connections. All properties in this section are symbolically proven in ‘GoldenAlgebraFoundation.nb’ and further extensively validated by ‘golden_algebra_validator_v3.py’.

3.1 Specific Values and Geometric Interpretation

For $k = 1$, the constants become:

- $\Phi_1 = \phi = \frac{1+\sqrt{5}}{2}$ (The Golden Ratio)
- $T_1 = \frac{\sqrt{5}-1}{4}$
- $J_1 = \frac{3-\sqrt{5}}{4}$
- $H_1 = T_1 J_1 = \frac{\sqrt{5}-2}{4}$
- $K_1 = \frac{-(\sqrt{5}+1)}{4}$

The geometric significance is established by:

Proposition 3.1. *The constants T_1 and K_1 are directly related to pentagonal geometry:*

$$T_1 = \cos(2\pi/5) \quad \text{and} \quad K_1 = \cos(4\pi/5)$$

3.2 Key Identities and Polynomials

Proposition 3.2 (Fundamental $k = 1$ Identities). *The Golden Algebra constants satisfy:*

1. $T_1 + J_1 = 1/2$
2. $T_1 - J_1 = 2H_1$
3. $T_1/J_1 - J_1/T_1 = 1$

Proposition 3.3 (Pentagon Polynomials for $k = 1$). *The constants T_1, K_1, J_1 satisfy specific quadratic ("Pentagon") polynomials:*

- T_1 and K_1 are roots of $4x^2 + 2x - 1 = 0$.
- J_1 is a root of $4x^2 - 6x + 1 = 0$.

3.3 Connections to Number Theory ($k = 1$)

Proven in Section 6 of 'GoldenAlgebraFoundation.nb'.

Proposition 3.4 (Fibonacci and Lucas Numbers). *The Golden Algebra constants are intrinsically linked to Fibonacci (F_n) and Lucas (L_n) numbers:*

1. $T_1/J_1 = \phi$.
2. The Binet-type formulas are recovered:

$$F_n = \frac{\phi^n - (-1/\phi)^n}{\sqrt{5}} = \frac{(T_1/J_1)^n - (-J_1/T_1)^n}{\sqrt{5}}$$

$$L_n = \phi^n + (-1/\phi)^n = (T_1/J_1)^n + (-J_1/T_1)^n$$

Proposition 3.5 (Pell's Equation $x^2 - 5y^2 = 1$). *Connections to the Pell equation for $D = 5$:*

1. $\sqrt{5} = 4T_1 + 1$.
2. The fundamental unit $u = 9 + 4\sqrt{5}$ is related to T_1 by $u = 13 + 16T_1$.

3.4 Matrix Representation ($k = 1$)

Proven in Section 8 of 'GoldenAlgebraFoundation.nb'.

Definition 3.6 (Golden Matrix G). *The Golden Matrix G is defined as $G = \begin{pmatrix} T_1 & -J_1 \\ J_1 & T_1 \end{pmatrix}$.*

Proposition 3.7. *The Golden Matrix G has the following properties:*

1. $\text{Tr}(G) = 2T_1 = \frac{\sqrt{5}-1}{2} = \phi'$ (golden ratio conjugate).
2. $\text{Det}(G) = T_1^2 + J_1^2 = \frac{5-2\sqrt{5}}{4}$.

3.5 Galois Conjugation in $\mathbb{Q}(\sqrt{5})$ ($k = 1$)

Proven in Section 7 of 'GoldenAlgebraFoundation.nb'. Let $\sigma : \sqrt{5} \mapsto -\sqrt{5}$ be the non-trivial Galois automorphism.

Proposition 3.8. *The action of σ on the Golden Algebra constants is:*

- $\sigma(T_1) = K_1$ and $\sigma(K_1) = T_1$.
- $\sigma(J_1) = (3 + \sqrt{5})/4$.
- $\sigma(H_1) = \sigma(T_1)\sigma(J_1) = K_1\sigma(J_1) = \frac{-2-\sqrt{5}}{4}$.
- $\sigma(\phi) = 1 - \phi = -1/\phi$.

3.6 Exponential Identities and Euler's Identity ($k = 1$)

Proven in Section 10 of 'GoldenAlgebraFoundation.nb'.

Proposition 3.9 ($k = 1$ Exponential Identity). *The general exponential identity $e^{in\pi T_k} = e^{in\pi k/2} \cdot \overline{e^{in\pi J_k}}$ specializes for $k = 1$ to:*

$$e^{in\pi T_1} = e^{in\pi/2} \cdot \overline{e^{in\pi J_1}}$$

An alternative form, using $T_1 + J_1 = 1/2$, is $e^{in\pi(T_1+J_1)} = e^{in\pi/2}$.

Corollary 3.10 (Connection to Euler's Identity). *Setting $n = 1/T_1$ in the identity $e^{in\pi T_1} = e^{in\pi/2} \cdot \overline{e^{in\pi J_1}}$ leads to $e^{i\pi} = -1$.*

Proof Sketch: The LHS becomes $e^{i\pi}$. The RHS becomes $e^{i\pi/(2T_1)} e^{-i\pi J_1/T_1}$. The equality of the exponents (modulo $2m\pi$ for integer m), $\pi = \frac{\pi}{2T_1} - \frac{\pi J_1}{T_1} + 2m\pi$, simplifies to $1 = \frac{1}{2T_1} - \frac{J_1}{T_1} + 2m$. This further simplifies to $2T_1 = 1 - 2J_1 + 4mT_1$, or $2(T_1 + J_1) - 1 = 4mT_1$. Since $T_1 + J_1 = 1/2$, the LHS is 0, so $4mT_1 = 0$. As $T_1 \neq 0$, this implies $m = 0$. Thus, the exponents are equal, confirming $e^{i\pi} = e^{i\pi}$, and hence Euler's identity is consistently encoded. \square

3.7 Explicit Symbolic Forms ($k = 1$)

Proven in Section 9 of 'GoldenAlgebraFoundation.nb'.

Proposition 3.11. *The definitions of T_k, J_k, K_k, H_k yield the following exact forms when $k = 1$: $T_1 = (\sqrt{5} - 1)/4$, $J_1 = (3 - \sqrt{5})/4$, $K_1 = -(\sqrt{5} + 1)/4$, $H_1 = (\sqrt{5} - 2)/4$.*

4 The "Golden Ratio Condition" and Algebraic Rigidity

A pivotal result for the "Mirror Math" hypothesis, symbolically proven in Section 3 of 'GoldenAlgebraFoundation.nb'.

Theorem 4.1 (Uniqueness of $k = 1$ under Golden Ratio Constraint). *Let $\Phi_k = \frac{k+\sqrt{k^2+4}}{2}$ be the k -Metallic Mean for $k > 0$. If Φ_k is constrained to be the Golden Ratio $\phi = \frac{1+\sqrt{5}}{2}$ (i.e., Φ_k satisfies the polynomial $x^2 - x - 1 = 0$), then k is uniquely determined to be $k = 1$.*

Proof Sketch: The k -Metallic Mean Φ_k satisfies $\Phi_k^2 - k\Phi_k - 1 = 0$. If Φ_k also satisfies $\Phi_k^2 - \Phi_k - 1 = 0$, then by subtraction, $(k - 1)\Phi_k = 0$. Since $\Phi_k > 0$ for $k > 0$, it follows that $k - 1 = 0$, so $k = 1$. This is demonstrated in the notebook using 'Solve'. \square

Remark 4.2. *Similarly, as shown by combining results from Section 1.3 and 2.3 of the notebook, if T_k is constrained to satisfy the Pentagon Polynomial $4x^2 + 2x - 1 = 0$ (which is characteristic of T_1), k is also uniquely forced to be 1.*

5 Complex Representation Z_k and Geometric Identity

This section refers to results from Section 5 of 'GoldenAlgebraFoundation.nb'.

Definition 5.1 (Complex Representation Z_k). *For each $k > 0$, we define $Z_k = T_k + iJ_k$. The argument is $\Theta_k = \text{Arg}(Z_k) = \arctan(J_k/T_k) = \arctan(1/\Phi_k)$.*

Proposition 5.2 (Fundamental Geometric Identity). *The parameter k of the k -Metallic Algebra is related to the argument Θ_k by:*

$$k = 2 \cot(2\Theta_k)$$

Proof Sketch: Symbolically proven in ‘GoldenAlgebraFoundation.nb’. This identity links the algebraic parameter k directly to the geometry of Z_k^2 in the complex plane. For $k = 1$, $1 = 2 \cot(2\Theta_1) \implies \tan(2\Theta_1) = 2$. For $k = 2$, $2 = 2 \cot(2\Theta_2) \implies \Theta_2 = \pi/8$. \square

6 The "Mirror Math" Hypothesis for the Riemann Hypothesis

The unique properties of the Golden Algebra ($k = 1$), particularly its algebraic rigidity demonstrated by Theorem 4.1, form the basis for a structural hypothesis concerning the Riemann Hypothesis (RH). This framework is outlined in Section 4 of ‘GoldenAlgebraFoundation.nb’.

Postulate 6.1 (Spectral Correspondence). *For any non-trivial zero $s_0 = \beta_0 + i\gamma_0$ of the Riemann zeta function $\zeta(s)$, its real part $\beta_0 = \Re(s_0)$ is proposed to correspond to $k_0/2$ within the k -Metallic Algebra framework. Thus, $k_0 = 2\Re(s_0)$.*

Postulate 6.2 (The Golden Ratio Condition). *For the k_0 -metallic system (derived from s_0 via Postulate 6.1) to serve as a "faithful mirror" of fundamental mathematical structures, its characteristic metallic mean, $\Phi_{k_0} = T_{k_0}/J_{k_0}$, *must* be the Golden Ratio ϕ . This implies Φ_{k_0} must satisfy the minimal polynomial of ϕ : $\Phi_{k_0}^2 - \Phi_{k_0} - 1 = 0$.*

Theorem 6.3 (Implication for Riemann Hypothesis). *If Postulate 6.1 and Postulate 6.2 hold, then all non-trivial zeros of the Riemann zeta function must lie on the critical line $\Re(s_0) = 1/2$.*

Proof. By Postulate 6.2, $\Phi_{k_0} = \phi$. From Theorem 4.1, if $\Phi_{k_0} = \phi$, then $k_0 = 1$. By Postulate 6.1, $\Re(s_0) = k_0/2$. Substituting $k_0 = 1$, we obtain $\Re(s_0) = 1/2$. \square

Remark 6.4. *This framework proposes an algebraic path to the Riemann Hypothesis, contingent on the mathematical substantiation of Postulates 6.1 and 6.2, particularly the fundamental reason for imposing the Golden Ratio Condition (Postulate 6.2).*

7 Conclusion and Future Work

The Mathematica notebook ‘GoldenAlgebraFoundation.nb’ has systematically defined the k -Metallic Algebra and symbolically proven its core properties for general $k > 0$. It further detailed the specialization to the $k = 1$ Golden Algebra, demonstrating its unique arithmetic, geometric, and algebraic characteristics, including connections to fundamental number theory and constants like e and π . Crucially, the notebook established that imposing conditions characteristic of the Golden Algebra (such as the Golden Ratio Condition on Φ_k) uniquely forces $k = 1$. These findings provide the necessary algebraic underpinnings for the "Mirror Math" hypothesis, which postulates a structural connection between the Golden Algebra and the Riemann Hypothesis. The Python script ‘golden_algebra_validator_v3.py’ offers extensive independent validation of 207 properties for the $k = 1$ case, underscoring its rich and consistent structure.

The primary direction for future research is the rigorous mathematical justification of Postulate 6.2 (The Golden Ratio Condition) from first principles, aiming to demonstrate why this specific algebraic constraint should apply when modeling phenomena related to the Riemann zeta function. Further work also includes exploring predictive models for zeta zero parameters based on this framework and investigating the properties and applications of k -Metallic Algebras for $k \neq 1$.

Acknowledgements

Thank you to my parents, my family, my friends, and all the people who believe in me. Also thanks shoulders of giants for letting me stand on you. Well.. assuming I am not crazy. Time will tell.